Detection and quantification of key dental pathogens through wastewater

monitoring

Olivia N. Birch[a], Sang C. Par[a], Justin C. Greaves[a]

[a]Department of Environmental and Occupational Health, School of Public Health

Indiana University-Bloomington

Corresponding Author:

Justin C.J. Greaves

jcgreave@iu.edu

Indiana University-School of Public Health

2719 E 10th Street, Innovation Center

Bloomington Indiana, 47408

Keywords

Wastewater surveillance, Oral pathogens, Water treatment, Persistence, Gum disease,

Dental caries

Abstract

Wastewater-based epidemiology (WBE) has been widely used to track viral pathogens

like SARS-CoV-2 and polio, but its potential for monitoring common dental bacterial

pathogens that infect the oral cavity has yet to be explored. *Streptococcus mutans* and

Porphyromonas gingivalis are key oral bacterial pathogens that cause highly prevalent

dental diseases worldwide, such as dental caries and gingivitis. Our main objective for

this study was to investigate the presence and prevalence of these oral bacteria in

wastewater to determine the feasibility of using WBE for oral pathogens. We measured

S. mutans and *P. gingivalis* nucleic acids in weekly samples for 24 months at a local

wastewater treatment plant. A total of 100 untreated wastewater samples were collected

once a week between June 2023 and May 2025. Samples were concentrated, extracted

for DNA, then tested for each bacterium. Our results showed that 89% and 58% were

positive for *S. mutans* and *P. gingivalis*, respectively, which shows that wastewater

surveillance is appropriate for oral bacteria. Average concentrations were 4.57 $\log_{10}$

genome copies/L and 3.03 $\log_{10}$ genome copies/L for *S. mutans* and *P. gingivalis*,

respectively. Detections of oral bacteria were observed in the primary and final effluent

but concentrations were significantly lower in the final effluent than untreated

wastewater. The high levels of oral bacteria in wastewater indicated a potential

transmission mechanism for these bacteria through water, specifically for *S. mutans*.

Additionally, this study underscores the unique potential for WBE to be used in the

surveillance of oral bacterial pathogens.

1.0 Introduction

Dental caries, or tooth decay, is one of the most prevalent chronic diseases worldwide, affecting nearly 2.3 billion people with permanent tooth decay and over 530 million children with primary tooth loss, according to the World Health Organization (Organization, 2017). *S. mutans* is a major culprit in dental caries, playing a significant role in plaque formation and acid production that demineralizes tooth enamel (Caufield, 2000; Forssten et al., 2010; Lemos et al., 2019). *P. gingivalis*, on the other hand, is a key pathogen in periodontal disease, a severe gum infection that can lead to tooth loss and systemic health complications which can include cardiovascular disease and diabetes (Marutescu et al., 2023; Rafiei et al., 2017). The high global burden of these oral health issues underscores the importance of developing innovative disease surveillance methods to understand the spread of these diseases.

Wastewater Based Epidemiology (WBE) has become increasingly popular in recent years as a tool to monitor the spread of endemic and emerging pathogens passively in a community. WBE has already been successfully employed to monitor various pathogens, such as SARS-CoV-2 and antimicrobial resistance genes, yet its application to oral bacteria remains largely unexplored (Greaves et al., 2020b; Rodriguez et al., 2024). Given the widespread presence of these oral bacteria in human populations, understanding their prevalence in wastewater can provide insights into community-level oral health trends and transmission dynamics. Traditional surveillance methods, such as clinical examinations and self-reported surveys, have limitations in reach, accuracy, and cost-effectiveness (Pitiphat et al., 2002; Sekundo et al., 2019). WBE, on the other hand,

allows for large-scale, community-wide monitoring without the need for individual

participation, offering a more comprehensive and real-time picture of oral health

burdens (Cohen et al., 2024; Daughton, 2020). If a strong correlation between

wastewater bacterial levels and oral disease prevalence is established, this approach

could serve as an early warning system for emerging trends in oral health conditions,

prompting timely public health interventions (Kasprzyk-Hordern et al., 2022; Mao et al.,

2020). However, there is no current public health surveillance system for the oral

diseases caused by these specific dominant pathogens at the national or local level.

Hence, our current understanding on the prevalence of these diseases within the past 5

years has been severely limited. Therefore, the use of WBE is especially timely in filling

the gaps in information about these oral pathogens.

Beyond monitoring, the detection of oral bacteria in wastewater raises concerns about

their environmental persistence and potential transmission pathways (Chopra et al.,

2020; Galarde-López et al., 2024). Although wastewater treatment processes are

designed to remove or reduce microbial contaminants, some pathogens can survive or

be reintroduced into the environment through treated effluent and biosolids by attaching

to particles (Chahal et al., 2016; Marutescu et al., 2023). If *S. mutans* and *P. gingivalis*

demonstrate resilience through treatment processes, there is a risk of environmental

exposure, which could have implications for public health (Huang et al., 2024; Li et al.,

2022; Rams et al., 2023; Sweeney et al., 2004; Teoh et al., 2018). Further studies are

needed to assess whether these bacteria can colonize environmental surfaces,

influence microbial ecosystems, or contribute to antimicrobial resistance.

In this study, we determined the concentration of *S. mutans* and *P. gingivalis* in

wastewater samples over one year. We also determined the concentration of *S. mutans*

and *P. gingivalis* in wastewater primary influent, primary effluent, and secondary

effluent. Lastly, we observed the prevalence and persistence of *S. mutans* over time in

influent wastewater by running a month-long decay experiment. By measuring the

prevalence of *S. mutans* and *P. gingivalis* in untreated wastewater and evaluating their

reduction through treatment processes, this study aims to bridge a significant

knowledge gap in WBE. If successful, integrating oral bacteria surveillance into WBE

frameworks could revolutionize how we monitor, predict, and mitigate oral health

diseases at a population level. Public health agencies could use wastewater data to

design targeted interventions, such as fluoridation programs, oral health education

campaigns, or improved access to dental care in communities with high oral bacterial

loads (Melbye and Armfield, 2013).

2.0 Materials and Methods

<u>Study site & temporal sampling</u>:

100 sewage samples were collected from Blucher Poole Wastewater Treatment Plant, a

local wastewater treatment plant (WWTP) in Bloomington, Indiana. Blucher Poole has a

flow of more than 4.5 million gallons per day and caters to the northern part of Monroe

County. This treatment plant was selected to represent the majority of the Bloomington,

Indiana, population and to observe bacterial shedding rates in wastewater. These

samples were collected weekly starting from June 2023 to May 2025. A second local

WWTP was selected, Dillman Road Wastewater Treatment Plant, so that we were able

to compare and contrast between the bacterial quantification of the two plants. Dillman

Road is located on the south side of Bloomington, Indiana and has a peak hydraulic

capacity at 30 million gallons daily (Bloomington, 2023). For this WWTP, 13 weekly

sewage samples from December 2023 to February 2024 were collected and transported

to the lab. 24-hour composite influent (prior to primary treatment) wastewater samples

were collected in 500-milliliter sterile polypropylene bottles and transported in a cooler

and frozen to -20°C within 2 hours of collection. Once received, these samples are

immediately prepared for filtration.

<u>Effluent Sampling</u>

Composite samples of primary and secondary/final effluent wastewater were collected

from the local WWTP and placed at -20°C for storage until sample processing. These

samples were collected during a 2-week intensive study and in the month of February

when concentrations were highest. For the primary influent, a 500 mL 24-hour

composite sample was collected, this is the sewage sample that comes directly into the

treatment plant and has not received any treatment. We received a 100 mL sample of

the 24-hour composite primary effluent sample, which consists of the sewage sample

after primary sedimentation has occurred. For the 24-hour composite secondary/final

effluent, 500 mL was collected in 500 mL sterile polypropylene bottles. This sample is

the sewage sample post aeration, activated sludge, and secondary sedimentation. All

samples were transported in the same manner as mentioned previously.

Decay Experiments:

A decay experiment was conducted on Streptococcus mutans in wastewater over the

course of a month to observe its persistence. The experiment was conducted in

triplicates of 400 milliliters of composite influent wastewater in 500-milliliter flasks.

Flasks were then covered with parafilm and left in the dark for a month. 25 mL samples

were taken at each of the following time points: 2 hours, 1 day, 4 days, 7 days, 11 days,

and 27 days. Before taking each sample, the flasks were swirled to homogenize the

wastewater and bacteria inside. Samples were then processed using the methods

below and quantified for *S. mutans*.

Concentration and DNA Extraction:

For final effluent samples, 250 mL of the final effluent was taken to be filtered and for all

other samples, 50 mL was prepared to be filtered. All samples were acidified to a pH of

3.5. A known concentration of bovine coronavirus solution was also added to each

sample as a tracer. For filtration, an electronegative mixed cellulose ester (MCE) 47mm

diameter and 0.45 µm pore size filter was used to filter samples based on previous

studies (Bivins et al., 2020b; Greaves et al., 2022; Rodriguez et al., 2024). Once

filtration through the filter was complete, the filters were then transferred into 2.0 mL

PowerBead tubes and either further extracted immediately or stored at -20 °C for

extraction later. The DNA from the samples was then extracted in accordance to the

provided manufacturer instructions using the Allprep PowerViral DNA/RNA extraction kit

(Cat. No. 28000-50 Qiagen, Germantown, MD,USA). This kit resulted in the elution of

DNA in 100 µL of RNase-free water provided within the kit. Once eluted these samples

were then stored at -20°C in preparation for quantification.

<u>Bacterial and Viral Quantification:</u>

Bacterial quantification of the samples was performed using digital polymerase chain

reaction (dPCR) and the QIAcuity™ Four Platform System (Cat. No 911042, Qiagen,

Germantown, MD, USA), the QIAcuity Nanoplate 26k (Cat. No 250001, Germantown,

MD, USA), and the QIAcuity™ Probe PCR Kit (Cat. No. 250132 Qiagen, Germantown,

MD, USA) as described in previous studies (Rodriguez et al., 2024). The environmental

mix for each molecular testing consisted of 1X Qiacuity Probe PCR mix, 08 µM of each

forward and reverse primer, 0.4 µM of each probe, and 10 µL of the target DNA for a

total volume of 40 µL. The molecular primers and probes used for the detection of *S.*

mutans and *P. gingivalis* are described in **Table S1**. While the primer and probe sets

were chosen from peer-reviewed, previously validated assays, their specificity was

formally assessed through in silico screening using NCBI BLAST. Primer and probe

sets were also able to detect a diluted culture of each bacteria. All samples were tested

using a duplex, where *S. mutans* was detected in the green channel with the FAM probe

and *P. gingivalis* was tested in the yellow channel with the HEX probe. The primers and

probes were manufactured by IDT (Coralville, IA, USA). The Zen™ internal quencher

(IDTDNA.com) and the 3' Black Hole ™ quencher were used to double-quench the

probes. PCR was run using the following conditions: an initial denaturation step at

95 °C for 2 minutes then 45 cycles of 95 °C for 5 seconds and 60 °C for 30 seconds.

The lowest detection limit was estimated for each PCR assay using DNA controls and a

dilution series which resulted in a calculated 380 copies/L for both *S. mutans* and *P.*

gingivalis. CrAssphage was used as a molecular fecal indicator and quantified with

techniques, probes, and primers published previously (Bivins et al., 2020a; Greaves et

al., 2020a; Wu et al., 2020) The threshold for each PCR reaction was marked by the

top of the negative band of the negative control for the specific probe being tested.

<u>Data analysis and statistics:</u>

The following formula was used to calculate the concentration of oral bacteria in gene

copes/L:

183
$$\frac{gc}{L} = \frac{gc}{uL} X \frac{Vol\ PCR\ reaction\ (40uL)}{Vol.\ NA\ analyzed\ (5uL)} X \frac{Vol.\ NA\ eluted\ (50uL)}{vol\ sample\ (0.05L)}$$

184   The analyses for our data was conducted using ANOVA and was performed on

185   GaphPad Prism v 10 (Boston, MA, USA) where independent factors were taken into

186   consideration. Graphs and figures were made using GraphPad Prism v 10 (Boston,

187   MA, USA). **For calculating the rate of decay, the decay curve was fit to the**

188   **following formula:**

189
$$C_T = C_0 e^{-kt}$$

190   In this equation, C_T represents the concentration of the pathogen at time *T*, C_0

191   represents the initial concentration of the pathogen at time 0, and *k* is representative of

192   the decay rate.

3.0 Results

<u>Bacterial and Viral Detection and quantification:</u>

Figure 1 shows the average monthly wastewater flow rates into the main local

wastewater treatment plant tested in this study. Values throughout the year show the

lowest value being in the summer and the highest values in the winter months. In

addition to monitoring flow rates, the fecal indicators crAssphage was measured in all

the samples throughout the duration of the experiments and concentrations remained

stable over the two-year period (**Figure S1**). CrAssphage had an average detected

concentration of 8.50 $\log_{10}$ GC/L. Statistical tests were done using a one-way ANOVA

followed by tukey multiple comparisons to measure changes in monthly concentrations

crAssphage, and results showed no statistical differences between crAssphage

concentrations over the different months of the experiment.

Figure 1. Average wastewater flow rate per month for the local wastewater treatment
plant monitored.

       Figure 2 shows the monthly average concentrations over the span of two years

for *S. mutans* and *P. gingivalis* in the main local wastewater treatment plant. Bacterial

concentrations were normalized by wastewater flow rates to accommodate for

population changes. The highest concentration of *S. mutans* was observed in February

of year 1 with a calculated monthly average concentration of 5.01 $\log_{10}$ GC/L and the

lowest detected concentration being in August of year 2 with 3.39 $\log_{10}$ GC/L. For *P.*

gingivalis, the month with the highest average concentration was November 2023 with a

calculated average monthly concentration of 3.81 $\log_{10}$ GC/L and the month with the

lowest values was September 2024 where none of the samples had detections and the

values were assumed to be the detection limit. Over the course of the entire selected

timeline, the average detected concentration of *S. mutans* and *P. gingivalis* was 4.57

$\log_{10}$ GC/L and 3.03 $\log_{10}$ GC/L respectively. **Table 1** breaks down the concentration

values for *S. mutans* and *P. gingivalis* into four different time periods within the year,

corresponding with Spring (March-May), Summer (June-August), Fall (September-

November), and Winter (December-February). Values are also the average of the two

years. For *S. mutans* the highest average detected concentration was in the winter with

a value of 4.78 $\log_{10}$ GC/L and a positive detection rate of 100%. Statistical tests

showed winter concentrations were significantly higher than summer and spring

concentrations for *S. mutans*. *P. gingivalis* also had the highest average detected

concentration in the winter with a value of 3.12 $\log_{10}$ GC/L and a positive detection rate

of 84% which was also the highest detection rate across all seasons. No significant

differences were found between the concentrations across all seasons for *P. gingivalis*.

Figure 2. Concentrations of oral bacteria in wastewater. Top two figures show monthly
average concentrations of *S.mutans* (A) and *P. gingivalis* (B) in Bloomington
wastewater over the two-year duration normalized by wastewater flow. The bars are
standard deviation for four samples collected each month (n=4-5). Middle figures show
average concentrations for *S.mutans* (C) and *P. gingivalis* (D) in each season over the
two years. Bottom figures show violin plots for both oral bacteria in year 1 and year 2 (E
for *S.mutans* and F for *P. gingivalis*).
238

244 **Table 1:** Detection rates and average concentrations of *S. mutans* and *P. gingivalis* in
245 wastewater samples.
246

Target	Seasons	Sample period	No. of Positive Samples (%)	Concentration (Log$_{10}$ copies/L)
S. mutans	Spring 2024 & 2025	March-May	19 (76)	4.37
	Summer 2023 & 2024	June-August	20 (80)	4.34
	Fall 2023 & 2024	Sept-Oct	25 (100)	4.35
	Winter 2024 & 2025	Jan-Feb & Dec previous year	25 (100)	4.78
	Total		89 (89)	4.57
P. gingivalis	Spring 2024 & 2025	March-May	12 (48)	2.93
	Summer 2023 & 2024	June-August	12 (48)	2.75
	Fall 2023 & 2024	Sept-Oct	13 (52)	3.04
	Winter 2024 & 2025	Jan-Feb & Dec previous year	21 (84)	3.12
	Total		58 (58)	3.03

247
248

A comparison was also done between year 1 (June 2023 to May 2024) and year

2 (June 2024 to May 2025) through using Pearson's correlation r. Statistical analysis

showed no correlation between the different years for both bacteria (**Figure S3**).

Correlation analysis for crAssphage for the different years showed a strong positive

correlation (r=0.77). A comparison was made between two different WWTP, Blucher

Poole (BP) and Dillman Road (DR), for the two bacteria in the winter of year 2, which

can be observed in **Figure S2**. For both WWTPs, *S. mutans* was observed to have a

larger spread in concentrations than *P. gingivalis* which can be observed in the top two

graphs within **Figure S2**. More samples of *P. gingvalis* were disregarded due to not

being detected, in comparison to *S. mutans*. For BP, the average concentration of *S.*

mutans was 4.93 $\log_{10}$ GC/L and for *P. gingivalis* it was 3.14 $\log_{10}$ GC/L. The average

concentration of *P. gingivalis* for DR was higher at 3.53 $\log_{10}$ GC/L whereas *S. mutans*

was relatively similar to BP with an average concentration of 4.86 $\log_{10}$ GC/L. *P.*

gingivalis was statistically higher in the DR WWTP compared to the BP WWTP whereas

there were no statistical differences between the concentrations of *S. mutans* in the DR

and BP WWTPs.

Effluent sampling:

In **Figure 3** the concentrations of *S. mutans* and *P. gingivalis* in primary influent, primary

effluent, and secondary effluent are shown. For both species of oral bacteria, they were

most prevalent in the primary influent samples with *S. mutans* having an average

concentration of 3.87 $\log_{10}$ GC/L and for *P. gingivalis*, a concentration of 3.21 $\log_{10}$

GC/L. The difference in concentration between the primary influent and the primary

effluent samples was larger for *P. gingivalis* than it was for *S. mutans* with calculated

differences of 1.08 $\log_{10}$ GC/L and 0.106 $\log_{10}$ GC/L, respectively. The final effluent

samples still resulted in some detected concentrations, with *S. mutans* having an

average of 3.08 $\log_{10}$ GC/L and *P. gingivalis* having an average of 1.99 $\log_{10}$ GC/L. The

difference between primary influent and secondary effluent for both *S. mutans* and *P.*

gingivalis was significant ($p < 0.0001$). For *S. mutans*, the average concentration for

primary effluent and secondary effluent was determined to be significant as well

($p < 0.0001$). The difference between the concentration of *P. gingivalis* in primary influent

and primary effluent samples was significant (p=0.009). There was determined to be no

significance in the concentration of *S. mutans* in the primary influent and the primary

effluent samples (p=0.816) as well as there being no significant difference in the

concentration of *P. gingivalis* between the primary effluent and secondary effluent

samples (p=0.064).

Figure 3. Occurrence of *S. mutans* and *P. gingivalis* in wastewater treatment plant
primary influent, primary effluent, and secondary effluent in 4 replicate samples.
286

Decay Rate:

The decay rate of *S. mutans* at room temperature in the absence of sunlight was

observed over the span of one month, with a visual representation of the data being

shown in **Figure 4**. There was positive detection for *S. mutans* all the way up to 27

days with an average value of 4.05 $\log_{10}$ GC/L. Across all three replicates, the average

k_{obs} for *S. mutans* was -0.4786 d^{-1}. For *P. gingivalis*, the decay rate was unable to be

observed due to the low initial concentration of the bacteria in wastewater.

Figure 4. Decay of *S. mutans* in wastewater primary influent at room temperature in the
absence of sunlight.
296

4.0 Discussion

To the best of the authors' knowledge, this study is the first study to detect and

quantify *S. mutans* and *P. gingivalis* in wastewater samples. These two pathogens

cause two of the most common dental diseases that infect a majority of populations

worldwide. Even though these pathogens are highly prevalent in US populations,

routine surveillance on the prevalence of these different pathogens or the oral disease

they cause is not done, especially since the outbreak of the COVID-19 pandemic (Dani

et al., 2016; Yan et al., 2021). Through our study, we were able to show that both

pathogens were detected in more than 60% of the 65 samples tested in wastewater

over the 4 different seasons in a year. The high detection of these oral pathogens in

wastewater samples from a mid-size city (75,000 residents) shows that wastewater-

based epidemiology (WBE) is highly feasible for these dental pathogens and oral

health. WBE may also serve as a low-cost method to monitor the spread and

prevalence of these dental diseases when traditional monitoring methods are not done.

S. mutans had consistently higher concentrations than *P. gingivalis* in wastewater

throughout the sampling period. In some samples, concentrations of *S. mutans* reached

to more than 10 times higher than *P. gingivalis*. With numerous studies estimating

higher percentages of tooth decay (90%) than gum disease (50%) in adults (Research,

2022; United States. Public Health Service. Office of the Surgeon General and National

Institute of Dental and Craniofacial Research (U.S.), 2021), our differential observations

of *S. mutans* (which associates with tooth decay) and *P. gingivalis* (which associates

with gum disease) is consistent with past clinical prevalences of these diseases in US

populations (Griffen et al., 1998; Rafiei et al., 2018; Thorild et al., 2002; Zubaidah et al.,

2022). Higher prevalences of a disease should result in higher concentrations in

wastewater (Rabe et al., 2023; Weidhaas et al., 2021). However, concentrations of *S.*

mutans could also be higher in wastewater due to the shedding dynamics from teeth as

opposed to the shedding dynamics of *P. gingivalis* from gums (Bugueno et al., 2020;

Liao et al., 2014; Liu et al., 2017). While studies have not specifically tested shedding of

each pathogen, previous studies have observed statistically higher concentrations of *S.*

mutans than *P. gingivalis* in dental samples from healthy patients, alluding to higher

shedding of *S. mutans* than *P. gingivalis* (Almeida et al., 2020; Dani et al., 2016; Davis

et al., 2012). Differential shedding could be due to the stark difference in oral hygiene

when it comes to the gums and teeth where cleaning of the teeth is done more

frequently than cleaning of the gums (Albertsson and Dijken, 2010; Deinzer et al., 2019;

Goryawala et al., 2016; Zimmermann et al., 2015). The variations in shedding might

also be simply attributed to the ease with which they attach or detach from different

surfaces, such as enamel versus gums (Gibbons, 1984; Mei et al., 2009; Vanhoute,

1983).

In addition to being shed in saliva, oral pathogens may also be swallowed, survive

passage through the gastrointestinal tract, and be excreted in feces. Several studies

have detected the DNA of oral microbes, including S. mutans, in human feces,

particularly through 16S rRNA gene sequencing (Abdelbary et al., 2022; Lin et al.,

2024). This fecal shedding pathway may significantly influence the concentration of

these bacteria in wastewater. For example, *S. mutans* may be more resilient than *P.*

gingivalis during gastrointestinal transit, potentially resulting in higher levels of S.

mutans in fecal matter. While feces are generally considered the primary route of

microbial contribution to wastewater, no dedicated studies have confirmed the specific

shedding patterns of *S. mutans* and *P. gingivalis* (Newton et al., 2015). As a result,

there remains uncertainty about how their wastewater concentrations correlate with

actual infection prevalence in the population. Future studies should examine the

shedding dynamics of each oral pathogen so that improved connections between dental

health and wastewater concentrations can be developed.

WBE has been shown to be a very useful tool for understanding disease trends,

specifically over the different seasons of a year (Brinkman et al., 2017; Kuhn et al.,

2023). When trends for different seasons are determined, health agencies can then pick

specific seasons throughout the year to focus mitigation efforts on (dos Santos et al.,

2024; Mao et al., 2020). Our results showed seasonal trends for *S. mutans* and *P.*

gingivalis, with both oral bacteria having their highest average concentrations in the

winters over the two-year period. However, results for *P. gingivalis* were not significant,

potentially due to the low detection rate of this pathogen. Nevertheless, detections in

winter for *P. gingivalis* were also much higher than the detections in other months,

suggesting potential seasonality similar to *S. mutans*.

S. mutans showed significantly higher concentrations in the winter than in the

spring and summer. The highest temperature months of July and August exhibited

notably lower concentrations compared to the other months of the year, especially in

year 2. These results highlight that *S. mutans* and dental caries could be more prevalent

in months with colder weather. Prior studies have observed diminished oral health in

colder months due to drier mouths and immune system changes in the winter (Fares,

2013; Fisman, 2007). Drier mouths due to increased indoor heater use can reduce the

volume of saliva (known to prevent bacterial growth), which in turn can cause the

extended growth of pathogens like *S. mutans* (King et al., 2000; Ligtenberg et al., 2020;

Weng et al., 2022). Colder months can also lead to diet changes, lowering nutrient

levels, or less hydration which could potentially contribute to tooth decay and poorer

oral health in colder months compared to hotter months (Fares, 2013; Moynihan and

Petersen, 2004; Sheiham, 2001). Although we observed higher average concentrations

of oral pathogens during the winter months across both years, several outlier months in

each season deviated from this trend, suggesting that additional factors may be

influencing oral health dynamics. These findings are also region-specific; other areas

may exhibit different seasonal patterns for these two oral pathogens. Future studies

should integrate oral health indicators with behavioral and environmental factors to

uncover the underlying drivers of seasonal variation.

Though average concentrations in the different seasons across years show

potential seasonal trends, trends of bacteria in each year were not well correlated

(**Figure S3**). Specifically for S. mutans, our data observed significantly lower

concentrations in year 2 than in year 1 suggesting reduction in oral pathogen burden in

the community. P. gingivalis has statistically similar concentrations across both years

but concentrations over time were also not well correlated. A potential reason for

reduction in oral health burden is that COVID-19-related dental clinic closures and

delayed care likely worsened oral health, but as services resumed, individual dental

health may have improved, potentially lowering oral health-related markers in

wastewater over time (Brian and Weintraub, 2020; Dill et al., 2023). Prior clinical

studies have also showed lowering oral health burden over the past decade which

follow the trend in our study (Dai et al., 2025; Lagerweij and van Loveren, 2015; Qin et

al., 2022). Additionally, year to year patterns can significantly change depending on

healthcare access, economic conditions, and public health policies that affect

community oral health in a year (Cooray et al., 2025; Dunleavy et al., 2024; Lee et al.,

2021).

Our study results showed that *S. mutans* concentrations in wastewater from both

WWTP's were not statistically different within the wintertime even though both WWTPs

serve different population numbers and types. These results suggest that *S. mutans* can

be prevalent in different WWTP and populations across geographical areas. The higher

observed concentration of *S. mutans* compared to *P. gingivalis* in the DP WWTP is

consistent with what was observed for BR WWTP. Our results do show statistically

higher concentrations of *P. gingivalis* in the DR WWTP compared to the BP WWTP

(p=0.0273). BP WWTP primarily serves the university campus community, which is

mostly comprised of young students between the ages of 18 and 24. DR WWTP serves

the rest of the local population which contains children and vulnerable groups that may

not be present in the other WWTP. As oral health significantly impacts those that are

most vulnerable, such as children (below 18) and elderly (above 65), it is expected that

there would be higher levels of reduced oral health in the DR WWTP which contains

these vulnerable groups (Bloomington, 2023; Weng et al., 2022; Xiao et al., 2020).

However, to specifically determine whether these factors affect oral health, further

analysis is needed on samples across the sewershed and in other geographic locations.

Fate in the wastewater treatment process

Throughout the wastewater treatment process, both *S. mutans* and *P. gingivalis*

were detectable but at significantly lower concentrations. These data suggest that the

treatment process is reducing the oral bacterial population, but not completely. The

primary treatment step was able to significantly reduce *P. gingivalis* but the secondary

treatment step (both activated sludge and secondary sedimentation process) was

needed to reduce *S. mutans*. These differences are potentially due to how likely each

microbe is to stick to fecal particles in wastewater. Multiple prior studies have shown a

reduction of various pathogens in wastewater through the primary and secondary

treatment processes (Delanka-Pedige et al., 2020; Frigon et al., 2013; Fu et al., 2010;

Skouteris et al., 2020). The same studies also show that certain pathogens are able to

survive through the process and enter the environment at lower levels, similar to what

we are observing for *P. gingivalis* and *S. mutans* (Delanka-Pedige et al., 2020; Frigon et

al., 2013; Fu et al., 2010; Skouteris et al., 2020) (Cuevas-Ferrando et al., 2022). Beyond

assessing the fate of oral bacteria in the wastewater treatment plant, our study

assessed the decay of oral bacteria in wastewater. With *P. gingivalis* having very low

concentrations, assessing its decay in water would be difficult. Furthermore, as *P.*

gingivalis is an anaerobic bacteria, its persistence in wastewater would be lower than *S.*

mutans due to the presence of oxygen in wastewater (Diaz and Rogers, 2004; Lemos et

al., 2019; Rafiei et al., 2018; Skouteris et al., 2020). Our calculated decay rates (0.48 d^{-}

^{1}) for *S. mutans* in wastewater are comparable with other types of common fecal

bacteria found in wastewater and the environment, such as E. coli (-0.14 to -1.13 day^{-1})

(Chan et al., 2015; Nakhle et al., 2021; Tiwari et al., 2023) and enterococci (-0.19 to -1.2

day^{-1})(Greaves et al., 2020b; Sagarduy et al., 2019; Tiwari et al., 2019; Tiwari et al.,

2023). This environmental and wastewater treatment plant survival, therefore, highlight

the potential novel transmission pathway of these oral pathogens through the

environment in contaminated waters.

<u>Limitations</u>

A key limitation of our study is the inability to compare our wastewater data with clinical

diseases that these pathogens cause (periodontal diseases and dental caries). The

main reason why we cannot access this data is because there is no clinical surveillance

of these pathogens in our region which highlights how important wastewater

surveillance can be to the monitoring of oral health. Future studies should try to

incorporate clinical studies or survey data with wastewater surveillance to better

connect wastewater data with the oral health of the community. Another major limitation

includes the use of only molecular methods to detect the various oral pathogens in this

study. While molecular methods offer fast and precise measurements for a diverse

range of targets, they do not assess viability of specific organism. Subsequent studies

should incorporate culturable or other viability assessment methods to measure the

viable concentration of oral pathogens, especially when investigating persistence.

5.0 Conclusion

In conclusion, this study demonstrated the feasibility of using WBE for oral pathogen

surveillance in a community. Additionally, our study showed a potential environmental

transmission pathway for *S. mutans* through water. Overall, this research has the

potential to provide critical insights into oral health trends, improve disease surveillance,

and enhance public health strategies. By leveraging wastewater analysis, we can move

toward a more proactive and non-invasive approach to oral disease monitoring,

ultimately contributing to better health outcomes and reducing the global burden of

dental and periodontal diseases.

462
463

6.0 References

Abdelbary, M.M.H., Hatting, M., Bott, A., Dahlhausen, A., Keller, D., Trautwein, C. and Conrads, G. 2022. The oral-gut axis: Salivary and fecal microbiome dysbiosis in patients with inflammatory bowel disease. Front Cell Infect Microbiol 12, 1010853.

Albertsson, K.W. and Dijken, J.W.V. 2010. Awareness of toothbrushing and dentifrice habits in regularly dental care receiving adults. Swed Dent J 34(2), 71-78.

Almeida, V.D.M., Azevedo, J., Leal, H.F., de Queiroz, A.T.L., da Silva, H.P. and Reis, J.N. 2020. Bacterial diversity and prevalence of antibiotic resistance genes in the oral microbiome. Plos One 15(9).

Bivins, A., Crank, K., Greaves, J., North, D., Wu, Z. and Bibby, K. 2020a. Cross-assembly phage and pepper mild mottle virus as viral water quality monitoring tools—potential, research gaps, and way forward. Current Opinion in Environmental Science & Health 16, 54-61.

Bivins, A., Greaves, J., Fischer, R., Yinda, K.C., Ahmed, W., Kitajima, M., Munster, V.J. and Bibby, K. 2020b. Persistence of SARS-CoV-2 in Water and Wastewater. Environ Sci Technol Lett 7(12), 937-942.

Bloomington, C.o. 2023 Dillman Road Wastewater Treatment Plant.

Brian, Z. and Weintraub, J.A. 2020. Oral Health and COVID-19: Increasing the Need for Prevention and Access. Prev Chronic Dis 17, E82.

Brinkman, N.E., Fout, G.S. and Keely, S.P. 2017. Retrospective Surveillance of Wastewater To Examine Seasonal Dynamics of Enterovirus Infections. Msphere 2(3).

Bugueno, I.M., Zobairi El-Ghazouani, F., Batool, F., El Itawi, H., Angles-Cano, E., Benkirane-Jessel, N., Toti, F. and Huck, O. 2020. Porphyromonas gingivalis triggers the shedding of inflammatory endothelial microvesicles that act as autocrine effectors of endothelial dysfunction. Sci Rep 10(1), 1778.

Caufield, P.W. 2000. Dental caries, an infectious and transmissible disease. J Dent Res 79(5), 1218-1218.

Chahal, C., van den Akker, B., Young, F., Franco, C., Blackbeard, J. and Monis, P. 2016. Pathogen and Particle Associations in Wastewater: Significance and Implications for Treatment and Disinfection Processes. Adv Appl Microbiol 97, 63-119.

Chan, Y.M., Thoe, W. and Lee, J.H.W. 2015. Field and laboratory studies of *Escherichia coli* decay rate in subtropical coastal water. J Hydro-Environ Res 9(1), 1-14.

Chopra, A., Bhat, S.G. and Sivaraman, K. 2020. Porphyromonas gingivalis adopts intricate and unique molecular mechanisms to survive and persist within the host: a critical update. J Oral Microbiol 12(1).

Cohen, A., Vikesland, P., Pruden, A., Lee, L.M., Krometis, L.A., Darling, A., Yancey, M., Helmick, M., Singh, R., Gonzalez, R., Meit, M., Degen, M. and Taniuchi, M. 2024. Making waves: The benefits and challenges of responsibly implementing wastewater-based surveillance for rural communities. Water Res 250.

Cooray, U., Singh, A., Aida, J., Tsakos, G. and Peres, M.A. 2025. Impact of Poverty Reduction on Oral Health Outcomes among US Adults. J Dent Res, 220345251323183.

Cuevas-Ferrando, E., Pérez-Cataluña, A., Falcó, I., Randazzo, W. and Sánchez, G. 2022. Monitoring Human Viral Pathogens Reveals Potential Hazard for Treated Wastewater Discharge or Reuse. Front Microbiol 13.

Dai, X., Dai, M., Liang, Y., Li, X. and Zhao, W. 2025. Global burden and trends of oral disorders among adolescent and young adult (10-24 years old) from 1990 to 2021. BMC Oral Health 25(1), 486.

Dani, S., Prabhu, A., Chaitra, K.R., Desai, N.C., Patil, S.R. and Rajeev, R. 2016. Assessment
of *Streptococcus mutans* in healthy versus gingivitis and chronic periodontitis: A clinico-
microbiological study. Contemp Clin Dent 7(4), 529-534.
Daughton, C.G. 2020. Wastewater surveillance for population -wide Covid-19: The present and
future. Science of the Total Environment 736.
Davis, J.E., Freel, N., Findley, A., Tomlin, K., Howard, K.M., Seran, C.C., Cruz, P. and Kingsley,
K. 2012. A molecular survey of *S. mutans* and *P. gingivalis* oral microbial burden in
human saliva using Relative Endpoint Polymerase Chain Reaction (RE-PCR) within the
population of a Nevada dental school revealed disparities among minorities. Bmc Oral
Health 12.
Deinzer, R., Cordes, O., Weber, J., Hassebrauck, L., Weik, U., Krämer, N., Pieper, K. and
Margraf-Stiksrud, J. 2019. Toothbrushing behavior in children - an observational study
of toothbrushing performance in 12 year olds. Bmc Oral Health 19.
Delanka-Pedige, H.M.K., Munasinghe-Arachchige, S.P., Zhang, Y.Y. and Nirmalakhandan, N.
2020. Bacteria and virus reduction in secondary treatment: Potential for minimizing post
disinfectant demand. Water Res 177.
Diaz, P.I. and Rogers, A.H. 2004. The effect of oxygen on the growth and physiology of. Oral
Microbiol Immun 19(2), 88-94.
Dill, J., Fontana, M., Bader, J., Chaffin, J., Strock, S. and Eckert, G. 2023. Impact of the
COVID-19 pandemic on oral health care use in the United States through December
2021. J Am Dent Assoc 154(10), 937-947 e933.
dos Santos, M.C., Silva, A.C.C., Teixeira, C.D., Prazeres, F.P.M., dos Santos, R.F., Rolo, C.D.,
Santos, E.D., da Fonseca, M.S., Valente, C.O., Hodel, K.V.S., Fonseca, L.M.D., Fiuza,
B.S.D., Bueno, R.D., de Andrade, J.B. and Machado, B.A.S. 2024. Wastewater
surveillance for viral pathogens: A tool for public health. Heliyon 10(13).
Dunleavy, G., Verma, N., Raghupathy, R., Jain, S., Hofmeister, J., Cook, R., Vujicic, M.,
Kebschull, M., Chapple, I., West, N. and Pitts, N. 2024. Inequalities in oral health:
estimating the longitudinal economic burden of dental caries by deprivation status in six
countries. BMC Public Health 24(1), 3239.
Fares, A. 2013. Factors Influencing the Seasonal Patterns of Infectious Diseases. Int J
Preventive Med 4(2), 128-132.
Fisman, D.N. 2007. Seasonality of infectious diseases. Annu Rev Publ Health 28, 127-143.
Forssten, S.D., Björklund, M. and Ouwehand, A.C. 2010. , Caries and Simulation Models.
Nutrients 2(3), 290-298.
Frigon, D., Biswal, B.K., Mazza, A., Masson, L. and Gehr, R. 2013. Biological and
Physicochemical Wastewater Treatment Processes Reduce the Prevalence of Virulent.
Appl Environ Microb 79(3), 835-844.
Fu, C.Y., Xie, X., Huang, J.J., Zhang, T., Wu, Q.Y., Chen, J.N. and Hu, H.Y. 2010. Monitoring
and evaluation of removal of pathogens at municipal wastewater treatment plants. Water
Sci Technol 61(6), 1589-1599.
Galarde-López, M., Velazquez-Meza, M.E., Godoy-Lozano, E.E., Carrillo-Quiroz, B.A., Cornejo-
Juárez, P., Sassoé-González, A., Ponce-de-León, A., Saturno-Hernández, P. and
Alpuche-Aranda, C.M. 2024. Presence and Persistence of Bacteria before and after
Hospital Wastewater Treatment. Microorganisms 12(6).
Gibbons, R.J. 1984. Adherent Interactions Which May Affect Microbial Ecology in the Mouth. J
Dent Res 63(3), 378-385.
Goryawala, S.N., Chavda, P., Udhani, S., Pathak, N.V., Pathak, S. and Ojha, R. 2016. A
survey on oral hygiene methods practiced by patients attending Dentistry Department at
a Tertiary Care Hospital from Central Gujarat. J Int Soc Prev Commu 6(2), 115-119.

Greaves, J., North, D. and Bibby, K. 2022. Particle association and size fraction of molecular
viral fecal pollution indicators in wastewater. Environmental Science: Water Research &
Technology 8(9), 1814-1821.
Greaves, J., Stone, D., Wu, Z. and Bibby, K. 2020a. Persistence of emerging viral fecal
indicators in large-scale freshwater mesocosms. Water Research X 9.
Greaves, J., Stone, D., Wu, Z. and Bibby, K. 2020b. Persistence of emerging viral fecal
indicators in large-scale freshwater mesocosms. Water Res X 9, 100067.
Griffen, A.L., Becker, M.R., Lyons, S.R., Moeschberger, M.L. and Leys, E.J. 1998. Prevalence
of *Porphyromonas gingivalis* and periodontal health status. J Clin Microbiol 36(11),
3239-3242.
Huang, S., Huang, J., Du, J.Y., Li, Y.J., Wu, M.J., Chen, S., Zhan, L. and Huang, X.J. 2024.
The LiaSR Two-Component System Regulates Resistance to Chlorhexidine in.
Microorganisms 12(3).
Kasprzyk-Hordern, B., Adams, B., Adewale, I.D., Agunbiade, F.O., Akinyemi, M., Archer, E.,
Badru, F.A., Barnett, J., Bishop, I.J., Di Lorenzo, M., Estrela, P., Faraway, J., Fasona,
M.J., Fayomi, S.A., Feil, E.J., Hyatt, L.J., Irewale, A.T., Kjeldsen, T., Lasisi, A.K.S.,
Loiselle, S., Louw, T.M., Metcalfe, B., Nmormah, S.A., Oluseyi, T.O., Smith, T.R.,
Snyman, M.C., Sogbanmu, T.O., Stanton-Fraser, D., Surujlal-Naicker, S., Wilson, P.R.,
Wolfaardt, G. and Yinka-Banjo, C.O. 2022. Wastewater-based epidemiology in hazard
forecasting and early-warning systems for global health risks. Environment International
161.
King, J.A., Rosal, M.C., Ma, Y.S., Reed, G., Kelly, T.A., Stanek, E.J. and Ockene, I.S. 2000.
Sequence and seasonal effects of salivary cortisol. Behav Med 26(2), 67-73.
Kuhn, K.G., Shukla, R., Mannell, M., Graves, G.M., Miller, A.C., Vogel, J., Malloy, K.,
Deshpande, G., Florea, G., Shelton, K., Jeffries, E., De Leon, K.B. and Stevenson, B.
2023. Using Wastewater Surveillance to Monitor Gastrointestinal Pathogen Infections in
the State of Oklahoma. Microorganisms 11(9).
Lagerweij, M.D. and van Loveren, C. 2015. Declining Caries Trends: Are We Satisfied? Curr
Oral Health Rep 2(4), 212-217.
Lee, W.C., Li, C.Y., Serag, H., Tabrizi, M. and Kuo, Y.F. 2021. Exploring the Impact of ACA on
Rural-Urban Disparity in Oral Health Services Among US Noninstitutionalized Adults. J
Rural Health 37(1), 103-113.
Lemos, J.A., Palmer, S.R., Zeng, L., Wen, Z.T., Kajfasz, J.K., Freires, I.A., Abranches, J. and
Brady, L.J. 2019. The Biology of. Microbiol Spectr 7(1).
Li, X.W., Wang, Y.F., Jiang, X.L., Zeng, Y.H., Zhao, X.R., Washio, J., Takahashi, N. and Zhang,
L.L. 2022. Investigation of drug resistance of caries-related streptococci to antimicrobial
peptide GH12. Front Cell Infect Mi 12.
Liao, S.M., Klein, M.I., Heim, K.P., Fan, Y.W., Bitoun, J.P., Ahn, S.J., Burne, R.A., Koo, H.,
Brady, L.J. and Wen, Z.Z.T. 2014. Extracellular DNA Is Upregulated during Growth in
Biofilms, Actively Released via Membrane Vesicles, and Influenced by Components of
the Protein Secretion Machinery. J Bacteriol 196(13), 2355-2366.
Ligtenberg, A.J.M., Meuffels, M. and Veerman, E.C.I. 2020. Effects of environmental
temperature on saliva flow rate and secretion of protein, amylase and mucin 5B. Arch
Oral Biol 109, 104593.
Lin, G., Kageyama, S., Maeda, A., Sakamoto, E., Ma, J., Asakawa, M., Furuta, M., Yamashita,
Y. and Takeshita, T. 2024. Oral-to-rectum microbial transmission in orthopedic patients
without a history of intestinal disorders. Front Cell Infect Microbiol 14, 1358684.
Liu, J., Sun, L.P., Liu, W., Guo, L.H., Liu, Z.H., Wei, X. and Ling, J.Q. 2017. A Nuclease from
Streptococcus mutans Facilitates Biofilm Dispersal and Escape from Killing by
Neutrophil Extracellular Traps. Front Cell Infect Mi 7.

Mao, K., Zhang, K.K., Du, W., Ali, W., Feng, X.B. and Zhang, H. 2020. The potential of
wastewater-based epidemiology as surveillance and early warning of infectious disease
outbreaks. Curr Opin Env Sci Hl 17, 1-7.
Marutescu, L.G., Popa, M., Gheorghe-Barbu, I., Barbu, I.C., Rodriguez-Molina, D., Berglund, F.,
Blaak, H., Flach, C.F., Kemper, M.A., Spiessberger, B., Wengenroth, L., Larsson,
D.G.J., Nowak, D., Radon, K., Husman, A.M.D., Wieser, A., Schmitt, H., Gradisteanu,
G.P., Vrancianu, C.O. and Chifiriuc, M.C. 2023. Wastewater treatment plants, an
"escape gate" for ESCAPE pathogens. Front Microbiol 14.
Mei, L., Busscher, H.J., van der Mei, H.C., Chen, Y.X., de Vries, J. and Ren, Y.J. 2009. Oral
bacterial adhesion forces to biomaterial surfaces constituting the bracket-adhesive-
enamel junction in orthodontic treatment. Eur J Oral Sci 117(4), 419-426.
Melbye, M.L.R. and Armfield, J.M. 2013. The dentist's role in promoting community water
fluoridation A call to action for dentists and educators. J Am Dent Assoc 144(1), 65-73.
Moynihan, P. and Petersen, P.E. 2004. Diet, nutrition and the prevention of dental diseases.
Public Health Nutrition 7(1a), 201-226.
Nakhle, P., Boithias, L., Pando-Bahuon, A., Thammahacksa, C., Gallion, N., Sounyafong, P.,
Silvera, N., Latsachack, K., Soulileuth, B., Rochelle-Newall, E.J., Marcangeli, Y., Pierret,
A. and Ribolzi, O. 2021. Decay Rate of *Escherichia coli* in a Mountainous Tropical
Headwater Wetland. Water-Sui 13(15).
Newton, R.J., McLellan, S.L., Dila, D.K., Vineis, J.H., Morrison, H.G., Eren, A.M. and Sogin,
M.L. 2015. Sewage reflects the microbiomes of human populations. mBio 6(2), e02574.
Organization, W.H. 2017 Sugars and dental caries.
Pitiphat, W., Garcia, R.I., Douglass, C.W. and Joshipura, K.J. 2002. Validation of self-reported
oral health measures. J Public Health Dent 62(2), 122-128.
Qin, X., Zi, H. and Zeng, X. 2022. Changes in the global burden of untreated dental caries from
1990 to 2019: A systematic analysis for the Global Burden of Disease study. Heliyon
8(9), e10714.
Rabe, A., Ravuri, S., Burnor, E., Steele, J.A., Kantor, R.S., Choi, S., Forman, S., Batjiaka, R.,
Jain, S., León, T.M., Vugia, D.J. and Yu, A.T. 2023. Correlation between wastewater
and COVID-19 case incidence rates in major California sewersheds across three variant
periods. J Water Health 21(9), 1303-1317.
Rafiei, M., Kiani, F., Sayehmiri, F., Sayehmiri, K., Sheikhi, A. and Zamanian Azodi, M. 2017.
Study of Porphyromonas gingivalis in periodontal diseases: A systematic review and
meta-analysis. Med J Islam Repub Iran 31, 62.
Rafiei, M., Kiani, F., Sayehmiri, K., Sayehmiri, F., Tavirani, M., Dousti, M. and Sheikhi, A. 2018.
Prevalence of Anaerobic Bacteria (P.gingivalis) as Major Microbial Agent in the
Incidence Periodontal Diseases by Meta-analysis. J Dent (Shiraz) 19(3), 232-242.
Rams, T.E., Sautter, J.D. and van Winkelhoff, A.J. 2023. Emergence of Antibiotic-Resistant
Porphyromonas gingivalis in United States Periodontitis Patients. Antibiotics-Basel
12(11).
Research, N.I.o.D.a.C. 2022 Oral Health in America - April 2022 Bulletin.
Rodriguez, R.A., Garza, F.M., Birch, O.N. and Greaves, J.C.J. 2024. Co-occurrence of adeno-
associated virus 2 and human enteric adenovirus (group F) in wastewater after
worldwide outbreaks of acute hepatitis of unknown etiology (AHUE). Sci Total Environ
955, 176806.
Sagarduy, M., Courtois, S., Del Campo, A., Garmendia, J.M. and Petrau, A. 2019. Differential
decay and prediction of persistence of *Enterococcus spp*. and *Escherichia coli* culturable
cells and molecular markers in freshwater and seawater environments. Int J Hyg Envir
Heal 222(4), 695-704.

Sekundo, C., Stock, C., Jürges, H. and Listl, S. 2019. Patients' self-reported measures of oral
health-A validation study on basis of oral health questions used in a large multi-country
survey for populations aged 50+. Gerodontology 36(2), 171-179.
Sheiham, A. 2001. Dietary effects on dental diseases. Public Health Nutr 4(2B), 569-591.
Skouteris, G., Rodriguez-Garcia, G., Reinecke, S.F. and Hampel, U. 2020. The use of pure
oxygen for aeration in aerobic wastewater treatment: A review of its potential and
limitations. Bioresource Technol 312.
Sweeney, L.C., Dave, J., Chambers, P.A. and Heritage, J. 2004. Antibiotic resistance in
general dental practice - a cause for concern? J Antimicrob Chemoth 53(4), 567-576.
Teoh, L., Stewart, K., Marino, R. and McCullough, M. 2018. Antibiotic resistance and relevance
to general dental practice in Australia. Aust Dent J 63(4), 414-421.
Thorild, I., Lindau-Jonson, B. and Twetman, S. 2002. Prevalence of salivary Streptococcus
mutans in mothers and in their preschool children. Int J Paediatr Dent 12(1), 2-7.
Tiwari, A., Kauppinen, A. and Pitkänen, T. 2019. Decay of *Enterococcus faecalis, Vibrio
cholerae* and MS2 Coliphage in a Laboratory Mesocosm Under Brackish Beach
Conditions. Front Public Health 7.
Tiwari, A., Kauppinen, A., Räsänen, P., Salonen, J., Wessels, L., Juntunen, J., Miettinen, I.T.
and Pitkänen, T. 2023. Effects of temperature and light exposure on the decay
characteristics of fecal indicators, norovirus, and *Legionella* in mesocosms simulating
subarctic river water. Science of the Total Environment 859.
United States. Public Health Service. Office of the Surgeon General and National Institute of
Dental and Craniofacial Research (U.S.) 2021 Oral health in America : advances and
challenges : executive summary, pp. 1 online resource (1 PDF file (i, 22 pages)),
National Institute of Dental and Craniofacial Research,, Bethesda (MD).
Vanhoute, J. 1983. Bacterial Adherence in the Mouth. Rev Infect Dis 5, S659-S669.
Weidhaas, J., Aanderud, Z.T., Roper, D.K., VanDerslice, J., Gaddis, E.B., Ostermiller, J.,
Hoffman, K., Jamal, R., Heck, P., Zhang, Y., Torgersen, K., Vander Laan, J. and
LaCross, N. 2021. Correlation of SARS-CoV-2 RNA in wastewater with COVID-19
disease burden in sewersheds. Science of the Total Environment 775.
Weng, C.T., Huang, S.L., Yang, H.W., Kao, C.C., Wei, C.C. and Huang, Y.F. 2022. Oral
microbiota in xerostomia patients-A preliminary study. J Dent Sci 17(1), 324-330.
Wu, Z., Greaves, J., Arp, L., Stone, D. and Bibby, K. 2020. Comparative fate of CrAssphage
with culturable and molecular fecal pollution indicators during activated sludge
wastewater treatment. Environmental International 136.
Xiao, J., Fiscella, K.A. and Gill, S.R. 2020. Oral microbiome: possible harbinger for children's
health. Int J Oral Sci 12(1).
Yan, Y.J., Wang, B.W., Yang, C.M., Wu, C.Y. and Ou-Yang, M. 2021. Autofluorescence
Detection Method for Dental Plaque Bacteria Detection and Classification: Example of
Porphyromonas gingivalis, *Aggregatibacter actinomycetemcomitans*, and *Streptococcus
mutans*. Dent J-Basel 9(7).
Zimmermann, H., Zimmermann, N., Hagenfeld, D., Veile, A., Kim, T.S. and Becher, H. 2015. Is
frequency of tooth brushing a risk factor for periodontitis? A systematic review and meta-
analysis. Community Dent Oral 43(2), 116-127.
Zubaidah, N., Dianawati, N., Ridwan, R.D., Shirakawa, T., Kuntaman, K., Setiawatie, E.M.,
Tanzil, M.I. and Kunarti, S. 2022. The Clinical Pattern and Prevalence of *Streptococcus
mutans* and *Streptococcus sobrinus* among Adult and Children Patients with Dental
Caries. Pesqui Bras Odontope 22.

Figure 4

Figure 1

Figure 2

Figure 3