Assessing ChatGPT's Performance in Delineating Uveitis: An analysis of responses to real-world case presentations

Authors

Muhammad Sohail Halim[1,2], Aly Hamza Khowaja[4]*, Zoha Zahid Fazal[1]*, Tanya Jain[1], Kholood Janjua[1,2], Ammar Aamir Khan[2,3], Anh Ngoc Tram Tran[1,2], Yasir J Sepah[1]

Affiliations

1. Byers Eye Institute, Stanford University, Palo Alto, California
2. Ocular Imaging Research and Reading Center, Sunnyvale, California
3. University of California, Davis, California
4. Aga Khan University Medical College, Karachi, Pakistan

*Both authors contributed equally to this manuscript.

Correspondence

Yasir J Sepah, MBBS

Assistant Professor, Byers Eye Institute

Stanford University School of Medicine

Palo Alto, California, United States 94303

Email: yjs@stanford.edu

Word count

2700 words

Keywords

Artificial intelligence, chatbot, explanation, ocular urgency, patient safety, ophthalmology

Abstract

Background

In the world of Artificial Intelligence (AI), Generative Pretrained Transformer-3 (GPT-3), has gained significant popularity for its demonstrated potential in medical education and diagnostics. However, its understanding of ocular urgencies, particularly uveitis, demands a focused investigation.

Methods

This proof-of-concept study explored the application of ChatGPT, a language model derived from GPT-3, in delineating uveitis based on 24 case presentations. We analyzed ChatGPT's communication quality through 14 qualitative metrics by computing patient data at four different levels to act as prompts. These included patient history, drug history, examination findings, and clinical investigations.

Results

Our results showed that at the initial prompt, ChatGPT's responses were comprehensive for most (8 out of 14) variables and correct but inadequate for some (3 out of 14) variables in the majority (>50.0%) of uveitis cases. *Ethical considerations* was the only variable in terms of which responses consistently showed mixed accuracy and outdated data across all prompts in most (95.8%) uveitis cases. Also, none of the ChatGPT responses were completely inaccurate in terms of any variable at any prompt for any uveitis case.

Conclusion

The results reveal ChatGPT's strengths and limitations in answering queries for patients with uveitis or its differential diagnosis while emphasizing the indispensable role of physicians in ethical decision-making.

Introduction

The ever-expanding sphere of artificial intelligence (AI) has transformed the digital life around us. The development of large language models (LLMs) has been particularly notable, from OpenAI's Generative Pretrained Transformer (GPT) to competing models like Google's Gemini, Anthropic's Claude, and Meta's Llama(1–3). These LLMs can understand the context of each question and generate human-like responses across various domains. AI has also seamlessly integrated itself into the field of healthcare and introduced new dimensions to patient guidance(4) and physician insight(5). Although GPT-3 was not originally developed for use in medicine, its capabilities as an LLM have shown significant promise(6). In fact, ChatGPT, an LLM prototype of GPT-3, has demonstrated over 80% accuracy in suggesting differentials for common clinical diagnoses(7). Current research suggests that that ophthalmic patients are now increasingly likely to utilize AI algorithms, often chatbots, to discuss their immediate symptoms and signs prior to a clinical appointment, allowing for potential triage to optimize healthcare system flow(8,9).

While there have been positive outcomes about its incorporation into clinical practice, the completeness of

the conclusions drawn by ChatGPT while answering questions is still unexplored. This especially holds

true for more intricate medical conditions, including ocular conditions like uveitis. As one of the leading

causes of blindness affecting all age groups globally, uveitis is a challenging diagnosis and often has a

complex presentation(10,11). Additionally, patients often incur delayed management and suboptimal

outcomes owing to the dearth of evidence-based guidelines(12) and scarcity of uveitis specialists

worldwide(13). Hence, attempts to fill the gaps with technological advancements are now being made as

patients often resort to online resources as the first point of consult(14). Since AI chatbots are now leading

the game of answering patient queries, recent studies have explored ChatGPT's capabilities in uveitis

diagnosis, demonstrating AI accuracy rates comparable to that of uveitis specialists(15,16).

With the initial benchmarks for AI performance in uveitis diagnosis set forth, our proof-of-concept research

aims to gather unique insights by examining the completeness and quality of AI-generated communication

on real-world uveitis presentations. We hypothesize that ChatGPT, among other evolving AI models,

provides reasonable explanations to uveitis patients and guides them towards the right direction of

ophthalmic care, in addition to assisting ophthalmologists, especially those without formal uveitis training

or those in remote areas, to effectively manage complex cases. This is to ensure timely diagnoses and

treatment to reduce uveitis-associated blindness rates globally. Our results would also be crucial for

understanding AI's potential in supporting ophthalmic physicians and patients to help mitigate the

increasing burden of vision-threatening morbidities on healthcare resources.

Methods

Data Collection and Processing

We developed a robust methodology, starting with the collection of a diverse set of real-world acute uveitis

cases from medical records of patients seen at a tertiary care hospital with a confirmed diagnosis. Owing to

the rarity of the condition and pilot nature of our study, only 24 cases representing various uveitic entities

and complexities were selected by an expert from the electronic medical records (EMR). Each case was

chosen based on its completeness of documentation and final diagnosis confirmation. All 24 cases of acute

uveitis had thus been diagnosed, investigated, and treated by uveitis specialists. After noting the chief

complaints, past medical, surgical, ocular, family, and social history, baseline slit lamp examination,

clinical investigations (hematological, radiological and ocular where relevant), and treatment advised by

the specialist, the case charts for each patient were prepared based on graded details for AI prompting. After

extracting patient presentations, the data was reviewed and processed to remove any unique identifiers or

protected patient health information (PHI) to maintain patient anonymity. The attending's assessment and
plan were also collected and recorded as the gold standard or benchmark for the AI to be compared against.
Figure 1 illustrates the flowchart of data collection and processing for unique uveitis case creation.

Experimental Design
We tested the GPT-3.5 model, (OpenAI, August 2023 release, California, USA) in this study. We used the
same model and version for each of the uveitis cases to ensure uniformity and standardization in the study.
Each presentation, as documented in EMR, was provided separately to the program after being de-identified
to protect patient data. Our experimental design used a standardized prompt in addition to patient case
findings to guide the AI's output. The standardized prompt skeleton (SPS) read as follows: 'In the context
of the patient's medical history, medication and examination [above/below], please formulate an assessment
to explore potential causes. Additionally, outline a comprehensive plan for diagnostic investigations and
initial management strategies.' To evaluate the model's performance with incomplete data, we tested four
different styles of case presentation for each patient, with each subsequent prompt adding more information
to the vignette provided previously as illustrated in *Figure 2*. These prompts were as follows:

- P1: Patient history
- P2: Patient history + Drug history
- P3: Patient history + Drug history + Physical examination findings
- P4: Patient history + Drug history + Physical examination findings + Clinical investigations
112
We randomized the order of case input using a computer program to mitigate order bias. Each response
by ChatGPT was collected twice and copied onto an online spreadsheet for the rest of the study process.
The details for the experimental design are provided in *Figure 3*.

Data Analysis
We assessed ChatGPT (GPT-3.5) output on qualitative metrics as determined by the study's principal
investigator:

- *Clinical judgment:* Evaluation of the clinical judgment demonstrated in responses. This metric
considered how well both physicians and ChatGPT recognized and interpreted clinical information to
provide appropriate recommendations or plans.
- *Contextual understanding:* The ability of responses to understand the context of the patient's medical
history and present concerns. The responses were assessed to evaluate how well ChatGPT tailored the
answers to specific patient scenarios.

• *Nuanced responses:* The nuances and subtleties that are often present in medical cases. The depth of understanding demonstrated in identifying rare conditions or complex diagnostic considerations was assessed.

• *Patient-centric approach:* Elements such as empathy, bedside manner, and the ability to address patient concerns beyond clinical facts were assessed for this metric.

• *Cohesiveness and organization:* The structure and organization of responses were evaluated. Factors such as whether the information was well-presented, logically structured, and easy for a layperson to follow were assessed.

• *Explanation of medical concepts:* The quality of explanation of medical concepts was assessed. Whether ChatGPT provided clear explanations that were understandable to a patient, or a non-medical audience was considered.

• *Risk-benefit discussions:* Whether the responses adequately discussed the risks and benefits of proposed interventions was evaluated. This metric reflects the thoroughness of the information provided.

• *Ethical considerations:* The inclusion of ethical considerations in the responses, such as patient autonomy, informed consent, and potential conflicts of interest was assessed.

• *Tailored recommendations:* How well ChatGPT provided recommendations that were tailored to the patient's specific medical history, preferences, and needs was evaluated.

• *Handling of uncertainty:* How the response handled uncertainty was considered. ChatGPT's ability to acknowledge insufficient information and provide appropriate next steps was assessed.

• *Use of evidence:* Whether responses referenced evidence-based guidelines, research, or clinical studies was evaluated. This metric indicated the level of reliance on established medical knowledge.

• *Communication style:* The communication style of responses, clarity, conciseness, and the use of language appropriate for the audience was assessed.

• *Awareness of limitations:* Whether ChatGPT was aware of its own limitations was assessed. This included ChatGPT's acknowledgment that it is an AI language model and that the cases might require specialist input.

• *Inclusion of red flags:* Whether the responses included warning signs or red flags that would necessitate urgent medical attention was evaluated.

Each of the above metrics was scored using a four-point scale as below :

• Comprehensive (Score: 4/4)

• Correct but inadequate (Score: 3/4)

• Mixed with correct/incorrect and outdated data (Score: 2/4)

• Completely incorrect (Score: 1/4)

All the qualitative metrics were scored using whole integers with no partial points allowed. Descriptive

statistics were used to analyze ChatGPT's performance, calculating the frequency and percentage of

responses for each score category across the 14 qualitative metrics. This analysis was applied to evaluate

performance across different levels of information provided in the prompts. Response on each prompt was

collected twice to ensure the credibility of ChatGPT 3.5.

Ethical Considerations

The study was conducted in accordance with the Declaration of Helsinki. Approval of retrospective studies

was obtained from the IRB committee of Stanford University (IRB-68008). All data was anonymized and

maintained with confidentiality. Stringent measures were implemented to protect patient privacy. All

potential ethical concerns related to AI involvement in healthcare were considered.

Results

Our results, as illustrated in *Figure 4* and summarized in *Table 1*, showed that ChatGPT's responses were

comprehensive for *communication style* (100.0%), *explanation of medical concepts* (95.8%), *cohesiveness*

and organization (95.8%), *patient-centric approach* (87.5%), *contextual understanding* (79.2%), *nuanced*

responses (66.7%), *clinical judgement* (62.5%), and *tailored recommendation* (58.3%) in most uveitis cases

at the initial prompt. The response accuracy also generally improved for these variables when additional

patient information was prompted. However, ChatGPT's response accuracy was subpar in terms of its *use*

of evidence (25.0%), *handling uncertainty* (16.7%), *risk-benefit discussions* (4.2%), *ethical considerations*

(4.2%), *inclusion of red flags* (4.2%), and *awareness of limitations* (0.0%) for uveitis diagnosis at the initial

prompt. Amongst these variables, response completeness only improved for *use of evidence* (25.0% at P1

to 37.5% at P4) while other variables showed a worsening or stagnant performance with additional prompts.

Notably, ChatGPT's responses in terms of *ethical considerations* were consistently mixed with

correct/incorrect and outdated data in 95.8% of uveitis cases regardless of prompts. However, none of the

responses were completely inaccurate in terms of any variable for any uveitis case at any prompt.

ChatGPT responses showed incredible performance in terms of *patient-centric approach*, with 87.5% of

cases showing accuracy at the initial prompt which scaled up to 100% with the subsequent prompts.

Similarly, in terms of *cohesiveness and organization*, ChatGPT responses were complete for 95.8% of cases

at the initial prompt which increased and stabilized to 100% accuracy with additional prompts.

Communication style was also a strongpoint of ChatGPT responses with 100% comprehensiveness for all

uveitis cases across all prompts. Lastly, the *explanation of medical concepts* was also well done by

ChatGPT for all uveitis case presentations, albeit with a few (4.2%) responses showing correct but
inadequate information at P2 and P4.

With respect to *clinical judgment*, 62.5% of the responses generated were completely accurate while the
rest 37.5% were correct but lacked information at the initial prompt. With additional information provided,
comprehensiveness increased from 50.0% at P2 to 54.2% at P3 but stabilized to 62.5% at the final prompt.
While the information shared by ChatGPT may be inadequate for this variable, no uveitis case prompted
responses with mixed accuracy or complete inaccuracy. Regarding *contextual understanding*, the
comprehensiveness of ChatGPT responses scaled down from 79.2% at P1 to 66.7% at P2 but increased to
70.8% at P3 and readjusted to 79.2% at P4. The remaining responses were correct but inadequate at each
prompt, albeit 4.2% cases showing mixed accuracy at P2. Similarly, *nuanced responses* were initially
comprehensive for 66.7% of uveitis cases but fluctuated, i.e. down to 62.5% at P2, back to 66.7% at P3,
and up to 75.0% at P4, with the remaining responses being correct but inadequate at each prompt.

The *use of evidence* was suboptimal in most ChatGPT responses with 70.8% of uveitis cases having correct
but inadequate information while 4.2% responses had mixed accuracy at the initial prompt. However, the
comprehensiveness increased incrementally from 25.0% at P1 to 37.5% at P4. *Risk-benefit discussions*
consistently showed high inadequacy in the data which increased with each prompt, i.e. 91.7% cases
showing correct but incomplete information at P1 scaling up to `100.0% at P4. ChatGPT responses to uveitis
management also performed poorly in term of *ethical considerations,* with 95.8% of responses showing
mixed accuracy and outdated data across all prompts, and the comprehensiveness reduced from 4.2% to
0.0% with the addition of drug history at P2. ChatGPT was also ill-equipped in terms of generating
responses *handling uncertainty* for uveitis cases, with comprehensiveness equaling 16.7% at all prompts,
albeit at P3 when it reduced further to 12.5%, with the remaining responses being correct but incomplete.
The *awareness of limitations* was also one of the shortcomings of ChatGPT toward uveitis management as
the responses demonstrated high inadequacy in the information presented, and comprehensiveness only
mildly increased from 0.0% to 4.2% with the addition of drug history (P2) and exam findings (P3) but
reduced back to 0.0% at P4. Lastly, ChatGPT was also limited in its response comprehensiveness by the
inclusion of red flags since most cases generated accurate but incomplete information across all prompts,
with some content (4.2%) showing mixed accuracy and outdated data with drug history addition at P2.

Discussion

Our findings demonstrate the promising potential of ChatGPT in summarizing case details, generating
initial differentials, and providing patient-centric explanations for most uveitis presentations, albeit with

some inaccuracies when considering ethicalities, uncertainties, and emergencies. These results are in
congruence with previous findings where ChatGPT provided understandable and relevant suggestions to
improve clinical decision-making for healthcare workers(17). Since ChatGPT also explained medical
concepts in a well-organized and cohesive manner, it can potentially serve as a valuable tool to answer
patients' queries and accurately guide them towards visiting a uveitis specialist when warranted by their
symptoms as concurred by a recent similar study(18). ChatGPT also showed impressive accuracy and
consistency in clinical judgement of uveitis cases, further confirming their utility to assist physicians,
particularly ophthalmologists, in devising evidence- and case-based care as also proven by Chen et. al.(19).

However, responses by ChatGPT served low adequacy in terms of discussing risk-benefits, red flags, and
limitations for uveitis cases despite prompting additional patient information in our study. This indicates
that while ChatGPT may be excellent in giving clinically comprehensive responses, it lacks in suggesting
improved outcomes in medicolegal and ethical aspects, which sustain as areas better handled by physicians.
While AI chatbots could be potentially useful tools to assist clinicians in their decision-making processes,
their performance may not yet be at the same level as experienced clinicians for deployment. Through
comprehensive ethical guidelines, ophthalmologists can thus ensure the responsible use of ChatGPT by
promoting reliable information exchange, protecting patient privacy, and empowering uveitis patients to
make informed decisions about their health(20).

Our results also showed that response completeness fluctuated with additional prompts for some variables.
Possible reasons for this variability, as cited in ophthalmic literature, include inconsistencies in training
data, sensitivity to prompt information, random output generation, or limitations in maintaining context
across queries(21,22). The high proportion of responses mixed with inaccurate or outdated data as found in
our study has also been previously recognized as 'AI hallucinations' and defined as invented information
by the chatbot causing a serious challenge for its use in medical practice(23). This variability is thus proof
that despite its great promise and vast expanse of knowledge, AI models can only be justified as a
supplement rather than a substitute to healthcare professionals(24).

This study has several limitations. Potential confounding factors include: (a) the 'gold standard' set forth
in our study being the attending's plan, which may differ for some aspects of real-world management across
different physicians, (b) variation in how ChatGPT handles changes in prompting structure from other
LLMs, including newer paid versions of ChatGPT, which were never tested in our study, (c) potential
selection bias owing to the uveitis cases shortlisted by a single expert from our team which might be skewed
toward certain complexities or uveitis types, (d) the inability of GPT-3.5 to incorporate external references

or recent literature unless specifically quoted in the prompts, and (e) the non-replicability of the generated
responses using health records if patient data differs in terms of demographic or comorbid variables.

Conclusion

In conclusion, our proof-of-concept study provides insights into the evolving relationship between AI and
ophthalmic care and the need for cautious integration and continuous improvement in specialized medical
domains. AI chatbots such as ChatGPT indeed have potential in educating patients, addressing the shortage
of uveitis specialists in developing countries, and serving as an aid to clinicians regarding uveitis diagnoses
and treatment(25). Fixing inconsistencies within variables identified by our study may thus necessitate
ongoing human oversight, increased dataset size, rigorous training, comprehensive testing, and further
validation of the AI models to better complement ophthalmic care.

Disclosures

Authors' Contributions
MSH and YJS conceptualized the study. YJS supervised the study. AHK and AK collected the data. TJ and
KJ analyzed the data. MSH, AHK, ZZF and ANTT drafted the initial manuscript. AHK and ZZF contributed
equally to making the final edits. All authors reviewed and approved the final draft.
Conflict of Interest
The authors have no conflict of interests to declare.
Funding
This study did not receive any intra- or extramural funding.
Acknowledgments
None.

References

1. Masalkhi M, Ong J, Waisberg E, Lee AG. Google DeepMind's gemini AI versus ChatGPT: a
comparative analysis in ophthalmology. Eye (Lond). 2024 Jun;38(8):1412–7.
2. Mavrych V, Yaqinuddin A, Bolgova O. Claude, ChatGPT, Copilot, and Gemini Performance versus
Students in Different Topics of Neuroscience. Adv Physiol Educ. 2025 Jan 17;
3. Wang Y, Liang L, Li R, Wang Y, Hao C. Comparison of the Performance of ChatGPT, Claude and
Bard in Support of Myopia Prevention and Control. J Multidiscip Healthc. 2024 Aug 13;17:3917–29.

4. Frontiers | Exploring the opportunities and challenges of NLP models in higher education: is Chat
GPT a blessing or a curse? [Internet]. [cited 2025 Jan 19]. Available from:
https://www.frontiersin.org/journals/education/articles/10.3389/feduc.2023.1166682/full

5. Giordano C, Brennan M, Mohamed B, Rashidi P, Modave F, Tighe P. Accessing Artificial
Intelligence for Clinical Decision-Making. Front Digit Health. 2021 Jun 25;3:645232.

6. Levine DM, Tuwani R, Kompa B, Varma A, Finlayson SG, Mehrotra A, et al. The diagnostic and
triage accuracy of the GPT-3 artificial intelligence model: an observational study. Lancet Digit
Health. 2024 Aug;6(8):e555–61.

7. Hirosawa T, Harada Y, Yokose M, Sakamoto T, Kawamura R, Shimizu T. Diagnostic Accuracy of
Differential-Diagnosis Lists Generated by Generative Pretrained Transformer 3 Chatbot for Clinical
Vignettes with Common Chief Complaints: A Pilot Study. Int J Environ Res Public Health. 2023 Feb
15;20(4):3378.

8. David D, Zloto O, Katz G, Huna-Baron R, Vishnevskia-Dai V, Armarnik S, et al. The use of artificial
intelligence based chat bots in ophthalmology triage. Eye. 2024 Nov 26;1–5.

9. Lyons RJ, Arepalli SR, Fromal O, Choi JD, Jain N. Artificial intelligence chatbot performance in
triage of ophthalmic conditions. Can J Ophthalmol. 2024 Aug;59(4):e301–8.

10. Joltikov KA, Lobo-Chan AM. Epidemiology and Risk Factors in Non-infectious Uveitis: A
Systematic Review. Front Med (Lausanne). 2021 Sep 10;8:695904.

11. Hadjadj J, Dechartres A, Chapron T, Assala M, Salah S, Dunogué B, et al. Relevance of diagnostic
investigations in patients with uveitis: Retrospective cohort study on 300 patients. Autoimmunity
Reviews. 2017 May 1;16(5):504–11.

12. Patil A, Gupta S, Venkatesh P, Banerjee M, Kumar V, Chawla R, et al. First contact investigations
and compliance to treatment in patients with uveitis. Int J Ophthalmol. 2023 Nov 18;16(11):1820–6.

13. Tsui E, Crowell EL, Gangaputra S, Moussa K, Shantha JG, Shusko AJ, et al. Current Landscape of
Uveitis Specialists in the United States. J Acad Ophthalmol (2017). 2022 Jul;14(2):e187–92.

14. Rojas-Carabali W, Cifuentes-González C, Gutierrez-Sinisterra L, Heng LY, Tsui E, Gangaputra S, et
al. Managing a patient with uveitis in the era of artificial intelligence: Current approaches, emerging
trends, and future perspectives. Asia-Pacific Journal of Ophthalmology. 2024 Jul 1;13(4):100082.

15. Rojas-Carabali W, Cifuentes-González C, Wei X, Putera I, Sen A, Thng ZX, et al. Evaluating the
Diagnostic Accuracy and Management Recommendations of ChatGPT in Uveitis. Ocul Immunol
Inflamm. 2024 Oct;32(8):1526–31.

16. Rojas-Carabali W, Sen A, Agarwal A, Tan G, Cheung CY, Rousselot A, et al. Chatbots Vs. Human
Experts: Evaluating Diagnostic Performance of Chatbots in Uveitis and the Perspectives on AI
Adoption in Ophthalmology. Ocular Immunology and Inflammation. 2024 Sep 13;32(8):1591–8.

17. Liu S, Wright AP, Patterson BL, Wanderer JP, Turer RW, Nelson SD, et al. Using AI-generated
suggestions from ChatGPT to optimize clinical decision support. J Am Med Inform Assoc. 2023 Apr
22;30(7):1237–45.

18. Mohammadi SS, Khatri A, Jain T, Thng ZX, Yoo W sun, Yavari N, et al. Evaluation of the
Appropriateness and Readability of ChatGPT-4 Responses to Patient Queries on Uveitis.
Ophthalmology Science [Internet]. 2025 Jan 1 [cited 2025 Jan 24];5(1). Available from:
https://www.ophthalmologyscience.org/article/S2666-9145(24)00130-1/fulltext

19. Chen JS, Reddy AJ, Al-Sharif E, Shoji MK, Kalaw FGP, Eslani M, et al. Analysis of ChatGPT
Responses to Ophthalmic Cases: Can ChatGPT Think like an Ophthalmologist? Ophthalmol Sci.
2024 Aug 23;5(1):100600.

20. Wang C, Liu S, Yang H, Guo J, Wu Y, Liu J. Ethical Considerations of Using ChatGPT in Health
Care. Journal of Medical Internet Research. 2023 Aug 11;25:e48009.

21. Cappellani F, Card KR, Shields CL, Pulido JS, Haller JA. Reliability and accuracy of artificial
intelligence ChatGPT in providing information on ophthalmic diseases and management to patients.
Eye (Lond). 2024 May;38(7):1368–73.

22. Antaki F, Touma S, Milad D, El-Khoury J, Duval R. Evaluating the Performance of ChatGPT in
Ophthalmology. Ophthalmol Sci. 2023 May 5;3(4):100324.

23. Hatem R, Simmons B, Thornton JE. A Call to Address AI "Hallucinations" and How Healthcare
Professionals Can Mitigate Their Risks. Cureus. 15(9):e44720.

24. Can ChatGPT Help in the Awareness of Diabetes? | Annals of Biomedical Engineering [Internet].
[cited 2025 Jan 19]. Available from: https://link.springer.com/article/10.1007/s10439-023-03356-1

25. Artificial Intelligence in Global Health: Defining a Collective Path Forward | Global Health | U.S.
Agency for International Development [Internet]. [cited 2025 Jan 19]. Available from:
https://www.usaid.gov/cii/ai-in-global-health

Figure 1: Flowchart of data collection and process for uveitis assessment.

Figure 2: Flowchart depicting four levels of prompts.

Figure 3: Flowchart of experimental design for uveitis assessment.

Figure 4: Graphs showing the comprehensiveness of ChatGPT responses at each level of prompt.

VARIABLES	Comprehensive (%)				Correct But In	
	P1	P2	P3	P4	P1	P2
Clinical Judgment	62.5	50	54.2	62.5	37.5	50
Cohesiveness and Organization	95.8	100	100	100	4.2	0
Contextual Understanding	79.2	66.7	70.8	79.2	20.8	29.2
Nuanced Responses	66.7	62.5	66.7	75	33.3	37.5
Patient-Centric Approach	87.5	100	100	100	12.5	0
Use of Evidence	25	29.2	29.2	37.5	70.8	66.7
Explanation of medical concepts	95.8	100	95.8	100	4.2	0
Risk-Benefit Discussions	4.2	0	0	0	91.7	95.8
Ethical Considerations	4.2	0	0	0	0	4.2
Tailored Recommendation	58.3	50	58.3	66.7	41.7	50
Handling Uncertainty	16.7	16.7	12.5	16.7	83.3	83.3
Awareness of Limitations	0	4.2	4.2	0	100	95.8
Communication Style	100	100	100	100	0	0
Inclusion of Red Flags	4.2	8.3	8.3	4.2	95.8	87.5

VARIABLES	Mixed with correct/incorrect				Completely In	
	P1	P2	P3	P4	P1	P2
Clinical Judgment	0	0	0	0	0	0
Cohesiveness and Organization	0	0	0	0	0	0
Contextual Understanding	0	4.2	0	0	0	0
Nuanced Responses	0	0	0	0	0	0
Patient-Centric Approach	0	0	0	0	0	0
Use of Evidence	4.2	4.2	0	0	0	0
Explanation of medical concepts	0	0	0	0	0	0
Risk-Benefit Discussions	4.2	4.2	0	0	0	0
Ethical Considerations	95.8	95.8	95.8	95.8	0	0
Tailored Recommendation	0	0	0	0	0	0
Handling Uncertainty	0	0	0	0	0	0
Awareness of Limitations	0	0	0	0	0	0
Communication Style	0	0	0	0	0	0
Inclusion of Red Flags	0	4.2	0	0	0	0

nadequate (%)	
P3	**P4**
45.8	37.5
0	0
29.2	20.8
33.3	25
0	0
70.8	62.5
4.2	0
100	100
4.2	4.2
41.7	33.3
87.5	83.3
95.8	100
0	0
91.7	95.8

naccurate (%)	
P3	**P4**
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0
0	0