Full Title: National and Subnational Level Estimates of Maternal Delivery at Home and Their

Predictors in Bangladesh: Evidence from Population-Based Survey 2012-2022

Short Title: Maternal delivery at home in Bangladesh

Authors: Raisha Binte Islam[1], Md. Lutful Kader[1], Somaya Mostarin[2], Mushfika Binta Latif[3],

Amina Anjum[4], Syed Toukir Ahmed Noor[5*]

Affiliations:

[1]Department of Statistics, University of Dhaka, Dhaka-1000, Bangladesh.

[2]Department of Health Education, National Institute of Preventive and Social Medicine

(NIPSOM), Mohakhali, Dhaka-1212, Bangladesh

[3]Department of Economics, York University, Canada.

[4]Department of Health Service Management Policy, National Institute of Preventive and Social

Medicine (NIPSOM), Mohakhali, Dhaka-1212, Bangladesh

[5]Department of Statistics, Shahjalal University of Science and Technology, Sylhet-3114,

Bangladesh.

*Corresponding author
Syed Toukir Ahmed Noor
Department of Statistics,
Shahjalal University of Science and Technology, Sylhet-3114, Bangladesh.
Email: syedtoukir86@gmail.com

Abstract

Background: Despite significant progress in reducing maternal and neonatal mortality, home delivery remains a substantial public health challenge in Bangladesh and many other low- and middle-income countries. While the proportion of home deliveries has markedly decreased in the past decade, pronounced disparities persist across geographic, socioeconomic, and demographic groups. A nuanced understanding of the prevalence and determinants of maternal home delivery is key to designing targeted interventions. This study examines national and subnational variations in maternal home delivery and associated factors in Bangladesh.

Methods: We analyzed data from the Multiple Indicator Cluster Surveys (2012–13 and 2019) and the Bangladesh Demographic and Health Survey (2022), covering 20,770 ever-married women aged 15–49 who gave birth in the preceding two years. District-level prevalence, descriptive statistics, and multivariable logistic regression were used to assess trends and determinants.

Results: Home delivery prevalence declined from 68% in 2012–13 to 35% in 2022. Disparities remain: divisions such as Barisal (48.9%), Chattogram, Sylhet, and Mymensingh showed higher rates, while Dhaka and Khulna had the lowest. At the subnational level, remote areas like Bandarban, Rangamati, and Bhola exhibited higher prevalence. The logistic regression analysis identified several significant predictors, such as women with no formal education, limited ANC visits (≤3), rural residence, lower wealth status, multiparity (≥3 children), and lack of media exposure were more likely to deliver at home.

Conclusions: Despite marked improvement, persistent geographic and socioeconomic inequities highlight the need for targeted interventions. Strengthening healthcare infrastructure in underserved regions, promoting maternal health awareness, scaling up ANC utilization, and reducing financial barriers through subsidies and incentive programs can help further decrease home deliveries. Future research should explore cultural and religious factors to inform context-specific policies for equitable facility-based childbirth.

Keywords: Maternal delivery, socioeconomic disparities, determinants, Bangladesh.

Introduction

Despite a 40% reduction in the global maternal mortality ratio (MMR)- from 328 deaths per 100,000 live births in 2000 to 197 in 2023- over 90% of all maternal deaths continue to occur in low- and lower-middle-income countries [1] . South Asia remains a critical contributor, accounting for nearly one-quarter of global maternal fatalities [2]. As a prominent example, Bangladesh, a lower-middle-income country in South Asia, has made commendable progress in reducing its MMR from 412 deaths per 100,000 live births in 1976 to 123 in 2020 [3]. This progress has been accompanied by significant scale-up in key maternal health interventions, including antenatal care (4+ visits) from 17% in 2004 to 41% in 2022, facility-based delivery from 10% in 2004 to 65% in 2022, skilled birth attendance from 23% in 2007 to 70% in 2022, and postnatal care (by medically trained providers within 48 hours) from 16% in 2004 to 55% in 2022 [4]. Despite these advances, the gains have been unevenly distributed across geographic regions and socioeconomic strata, resulting in persistent disparities in maternal health outcomes. Consequently, Bangladesh remains off track to meet the Sustainable Development Goal target (SDG 3.1), which targets an MMR of fewer than 70 deaths per 100,000 live births by 2030 [5].

To accelerate the progress toward this ambitious goal, there is an urgent need to improve emergency obstetric services and minimize the incidence of maternal delivery at home, which are associated with preventable maternal and neonatal mortality. Globally, maternal delivery at home accounted for an estimated 287,000 maternal and 2.4 million newborn deaths in 2020 alone [6,7]. A study from Nigeria found that non-institutional delivery (delivery outside health facilities, often by unskilled attendants) was associated with a 79.2% chance of maternal death among the cases studied [8]. In Bangladesh, more than one-third of all deliveries still occur at home, with marked regional disparities across administrative divisions [4] Reducing home deliveries is, therefore, a critical public health priority to advance equity and reduce mortality and neonatal mortality [9–11]. Maternal delivery at home is influenced by a complex interplay of sociodemographic, economic, cultural, and health system factors. Key factors include maternal education, quantity of antenatal care visits, parity, household wealth, religion, and geographic accessibility to healthcare facilities [12–14]. In addition, entrenched poverty, traditional and religious norms, poor transportation infrastructure, and the limited autonomy among women in decision-making further exacerbate the preference for delivery at home [15]. Other factors such as low maternal and paternal education, advanced maternal age, multiparity, and low wealth status also contribute to increased home delivery practices [16].

Fear of cesarean sections, perceptions of poor quality of care, and the lack of female doctors

also play substantial roles in influencing delivery preferences [15].

According to the availability, accessibility, acceptability, and quality (AAAQ) framework,

both supply-side (e.g. healthcare infrastructure and provider availability) and demand-side

(e.g. household-level barriers and perceptions) factors must be addressed to reduce maternal

delivery at home [17]. While individual and household-level determinants have been widely

studied, health system constraints- especially at the subnational level- remain largely

underexplored in the context of Bangladesh. Identifying subnational areas (here district in

Bangladesh) with persistently high rates of maternal delivery at home is essential for

designing and implementing evidence-based maternal health interventions.

However, limited research has assessed the sub-national level distribution of maternal delivery

at home in Bangladesh [18–20]. Moreover, little is known about how associated factors of

maternal delivery at home vary over time across urban and rural settings. To our knowledge,

no prior study has investigated subnational variations in maternal delivery at home and

explored the evolving determinants of this practice. Therefore, we aimed to investigate

national and subnational patterns of maternal delivery at home and analyze the associated

factors as a whole as well as in urban and rural areas in Bangladesh. The findings will inform

targeted, context-specific policies and support the implementation of effective maternal health

interventions.

Materials and Methods

Data Sources

This study utilized secondary data derived from three nationally representative, population-

based cross-sectional surveys: the Multiple Indicator Cluster Survey (MICS) conducted in

2012–13 and 2019, and the Bangladesh Demographic and Health Survey (BDHS) conducted

in 2022. The MICS, implemented by the Bangladesh Bureau of Statistics (BBS) with technical

and financial support from the United Nations International Children's Emergency Fund

(UNICEF), provides comprehensive data on household characteristics and maternal and child

health indicators at the district level. The BDHS, administered under the authority of the

National Institute of Population Research and Training (NIPORT), Medical Education and

Family Welfare Division, Ministry of Health and Family Welfare (MOHFW), offers vital

demographic and health statistics, including district-level estimates for the first time in 2022.

 Sampling Design and Sample Size

Both the MICS and BDHS employed a uniform methodology to ensure national

representativeness. Detailed methodological descriptions are available in the respective

technical reports [4,21,22]. Briefly, the MICS applied a multistage stratified cluster sampling

strategy, covering all 64 administrative districts of Bangladesh. Each district was stratified

into urban and rural areas. Enumeration areas (EAs) were selected using the probability

proportional to size (PPS) method within these strata. From each selected EA, 20 households

were systematically sampled [23,24]. The MICS sampled a total of 55,120 in 2012-2013 and

64,400 households in 2019, of which 51,895 and 61,242 were successfully interviewed, with

response rates of 98.5% and 99.4%, respectively. After excluding respondents with no

delivery history or missing birth outcome data, the final analytical sample included 7795 and

9284 women aged 15-49 years with a history of childbirth in 2012-2013 and 2019,

respectively.

The BDHS 2022 utilized a two-stage stratified sampling strategy based on the 2011

Population Census. In the first stage, 675 EAs were selected (237 urban and 438 rural) using

PPS sampling. In the second stage, 45 households were systematically selected from each EA

to ensure adequate representation across residence types, and for the first time, BDHS

provided the subnational level variable, which offers an opportunity to analyze the subnational

level estimate [25]. In 2022, the final response rate of BDHS was 99.6%, with 30,330

households selected and 30,018 households interviewed. The final analytical sample from

BDHS 2022 consisted of 3,691 women of reproductive age (15–49 years) with recent birth

histories. We conducted a complete case analysis; as a result, we omitted all missing

observations in our study. Women aged 15-49 with a live birth in the past 2 years were

considered for all the rounds.

Outcome Variable

The main outcome variable for this study was maternal delivery at home, defined

dichotomously. Deliveries occurring at the respondent's home were coded as "1" (maternal

delivery at home), and those occurring at any health facility were coded as "0".

Explanatory Variables

A total of 13 independent variables were selected on an extensive review of previous literature

[13,26,27], capturing individual-, household-, and community-level determinants. Individual-

level factors included women's age, education (no education, primary, secondary, and higher),

number of antenatal care (ANC) visits (none, 1, 2, 3, and ≥4), parity (1–2, ≥3), media

exposure, religion (Non-Muslim, Muslim), husband's age, and spousal age gap (≤5 years, 6–9

years, and more than 9 years). Household-level factors included education of household's

head, household size and household's socioeconomic status. The community-level factors

encompassed administrative division and place of residence (urban or rural). Socioeconomic

status was assessed using the wealth index, constructed via principal component analysis

(PCA) as provided in MICS and BDHS datasets. Households were classified into five

quintiles: poorest, poorer, middle, richer, and richest [4,21,22].

Statistical analysis

The data were analyzed via Stata version 17.0 (StataCorp, College Station, Texas, USA),

whereas data visualization was performed via R version 4.3.0. To account for the complex

survey design, we incorporated primary sampling units (PSUs) and strata and applied

sampling weights to ensure representativeness. The Stata "svyset" command was used to

adjust for the survey design [28], and all analyses adhered to the STROBE cross-sectional

reporting guidelines [29].

Descriptive statistics were used to summarize the characteristics of the study population,

which were reported as frequencies and proportions in percentages. We conducted bivariate

analysis using the Chi-square test to assess associations between the explanatory variables and

the outcome variable. We employed simple binary logistic regression analysis to estimate

crude odds ratios (ORs) with 95% CIs for each explanatory variable in the unadjusted

analysis. Variables that were statistically significant in the simple logistic regression models

were subsequently included in the multiple logistic regression model to estimate adjusted odds

ratios (AORs) with their respective 95% CIs. Statistical significance was determined at the 5%

level (p-value <0.05). Multicollinearity was checked for all the variables through VIF, and the

value of VIF was <5 for all variables.

To account for the complex survey design, we applied the "svyset" command in Stata,

incorporating sampling weights, strata, and primary sampling units (PSUs). All the analyses

were done using the "svy" prefix to ensure design-based estimates and inference [28]. The

study adhered to the STROBE guidelines for reporting cross-sectional studies [29]. We used

Stata version 17.0 (StataCorp, College Station, Texas, USA) for data analysis, R version 4.3.0

for data visualization and ArcGIS version 10.8 for mapping.

Ethical Consideration

This study utilized deidentified secondary data obtained from the UNICEF

(https://mics.unicef.org/surveys) and DHS (https://www.dhsprogram.com/data/available-

datasets.cfm) websites. Since the data were publicly available and did not contain any

personally identifiable information, ethical approval was not needed. All participants provided

informed consent, and the original surveys were carried out in accordance with ethical

standards and approved by the appropriate institutional review boards.

Results

Sample characteristics

Across all survey years, approximately one-third of women were aged 20-24 years (33.6% in

2012-13, 32.2% in 2019, and 33.8% in 2022). Educational attainment demonstrated a positive

trend over time, the proportion of women completing higher education increased from 12.4%

in 2012 to 18.5% in 2022. Nearly nine in 10 women were Muslim (89.2% in 2012, 90.4% in

2019 and 92.7% in 2022). Media exposure declined slightly over the years, from 76.6% in

2012 to 66.3% in 2022. Utilization of ANC services improved, with the proportion of women

receiving at least four ANC visits increased from 22.7% in 2012 to 40.5% in 2022. Fertility

pattern showed that majority of women had 1–2 children, increasing from 65.6% in 2012 to

71.9% in 2022. The proportion of households headed by individuals with no education

accounted for 56.2% in 2012, 30.3% in 2019 and 15.5% in 2022. The proportion of women in

the poorest wealth quintile declined from 28.3% in 2012 to 20.2% in 2022. Household size

remained relatively stable, with approximately two-thirds of women living in households

comprising five or more members (65.6% in 2012-13, 63% in 2019 and 69.5% in 2022). Most

women resided in rural areas (83.8% in 2012, 80.9% in 2019 and 73% in 2022), and the

Dhaka division consistently had the highest representation across survey years (**Table 1**).

Table 1: Sample characteristics of the respondents, presented in terms of weighted percentage

Variables	2012-13		2019		2022	
	Number	%	Number	%	Number	%
Age of women (in years)						
15-19	906	11.6	1,234	13.3	614	16.6
20-24	2,615	33.6	2,987	32.2	1248	33.8
25-29	2,351	30.2	2,562	27.6	962	26.1
30-34	1,143	14.7	1,640	17.7	592	16.0
35-49	780	10.0	861	9.3	274	7.4
Education of women						
No education	1,621	20.8	846	9.1	183	5.0
Primary	2,294	29.4	2,151	23.2	831	22.5
Secondary	2,917	37.4	4,691	50.5	1995	54.0

Higher	963	12.4	1,596	17.2	682	18.5
Religion						
Non-Muslim	842	10.8	896	9.7	269	7.3
Muslim	6,953	89.2	8,388	90.4	3422	92.7
Media exposure						
No	1,186	23.5	3,308	35.7	1243	33.7
Yes	3,872	76.6	5,967	64.3	2447	66.3
Number of ANC visits						
No visit	2,730	35.2	1,677	18.1	268	7.4
1 visit	1,018	13.1	1,123	12.1	526	14.6
2 visits	1,227	15.8	1,611	17.4	694	19.2
3 visits	1,024	13.2	1,529	16.5	658	18.2
4+ visits	1,758	22.7	3,339	36.0	1462	40.5
Parity						
1-2	5,116	65.6	6,401	69.0	2653	71.9
3+	2,679	34.4	2,883	31.1	1037	28.1
Age of husband (in year)						
15-29	2,514	32.9	2,789	30.3	1237	33.8
30-34	1,960	25.6	2,627	28.5	913	24.9
35-39	1,606	21.0	2,062	22.4	832	22.7
40-44	872	11.4	988	10.7	407	11.1
45-49	409	5.4	505	5.5	191	5.2
50-90	290	3.8	243	2.6	79	2.2
Spousal age gap (in year)						
≤5	2,939	38.0	3,534	38.4	1408	38.5
6-9	2,630	34.0	3,088	33.5	1192	32.6
>9	2,173	28.1	2,593	28.1	1060	29.0
Education of household head						
No education	4,375	56.2	2,812	30.3	566	15.5
Primary	1,009	13.0	2,794	30.1	1059	28.9
Secondary	1,410	18.1	2,602	28.0	1277	34.9
Higher	992	12.7	1,073	11.6	757	20.7
Wealth index						
Richest	1,174	15.1	1,587	17.1	661	17.9
Fourth	1,283	16.5	1,770	19.1	718	19.5
Middle	1,474	18.9	1,810	19.5	787	21.3
Second	1,661	21.3	1,886	20.3	780	21.1
Poorest	2,203	28.3	2,231	24.0	745	20.2
Household size						
≤4	2684	34.4	3,439	37.0	1128	30.6
5+	5,111	65.6	5,845	63.0	2563	69.5
Residence						
Urban	1,261	16.2	1,774	19.1	997	27.0
Rural	6,534	83.8	7,510	80.9	2694	73.0
Division						
Barisal	703	9.0	835	9.0	219	5.9
Chittagong	1,585	20.3	1,926	20.8	793	21.5
Dhaka	2,002	25.7	1,795	19.3	926	25.1
Khulna	1,044	13.4	1,275	13.7	380	10.3
Mymensingh			558	6.0	333	9.0
Rajshahi	715	9.2	956	10.3	381	10.3
Rangpur	972	12.5	1,135	12.2	419	11.4
Sylhet	774	9.9	804	8.7	240	6.5

203 *Variations in maternal delivery at home*

In Bangladesh, the prevalence of maternal delivery at home exhibited a marked decline over the past decade, decreasing from 68.1% [66.0-70.0%] in 2012-2013 to 46.4% [45.0-48.0%] in 2019 and 34.9% [32.0-37.0%] in 2022. Regional variation was evident, with Khulna Division demonstrating the lowest prevalence of maternal delivery at home across all years. In contrast, Barishal had the highest prevalence in 2012-13 and 2022, followed by the Sylhet division (**Figure 1**).

Figure 1: National and divisional estimates of maternal delivery at home in Bangladesh, 2012-2022

Although regional variations were evident, district-level disparities in maternal delivery at home were more pronounced (**Table S1 and Figure 2**). In 2022, the prevalence of maternal delivery at home was >50% in 14 of 64 districts. The highest prevalence was observed in Gopalganj (68.8%), Khagrachhari (68.5%), and Lalmonirhat (66.3%), while the lowest was reported in Bandarban (0%), Chuadanga (0%), and Kushtia (6.5%). In 2012, home-based maternal delivery was most prevalent in hill tract districts such as Bandarban (97.2%) and Rangamati (88.6%), and in haor/coastal districts including Sunamganj (89.3%), Noakhali (88.1%), and Patuakhali (88.0%). By 2019, the burden remained high in hill districts-84.6% in Bandarban, 73.0% in Rangamati, and 76.1% in Khagrachhari. Despite a declining national trend, hill tract and northeastern districts had persistently high prevalence of maternal delivery at home.

Figure 2: Distribution of maternal delivery at home at the subnational level in Bangladesh, 2012--2022

Subnational level variations in maternal delivery at home

Despite national and subnational reduction, maternal delivery at home remained common among women aged 35-49 years, although the prevalence in this subgroup declined disproportionately from 79.1% in 2012-2013 to 49.1% in 2022. Women with no formal education consistently showed the highest prevalence of maternal delivery at home (87.5% in 2012-2013, 75.6% in 2019 and 55.7% in 2022), in contrast to women with higher education, whose prevalence fell from 32.2% to 12.6% over the same period. Muslim women persistently reported higher percentage of maternal delivery at home than non-Muslim women (68.9% vs 58.9% in 2012-2013, 47.5% vs 34.1% in 2019 and 36.4% vs 16.1% in 2022). No media exposure was also associated with a higher prevalence of maternal delivery at home, though the prevalence declined notably from 78.3% in 2012 to 51.6% in 2022. Women receiving four or more ANC visits had significantly lower prevalence of maternal delivery at home than those receiving none (38.6% vs 91.1% in 2012, 23.9% vs 80.5% in 2019 and 19.2% vs 76.3% in 2022). Higher parity (3+ children) was associated with increased maternal delivery at home relative to lower parity (1-2 children) over years (81.0% vs 61.8% in 2012, 60.9% vs 39.9% in 2019, and 50.1% vs 28.9% in 2022). Furthermore, women of households headed by individuals with no formal education, those from the poorest wealth quintile, rural residents exhibited consistently higher prevalence of home-based maternal delivery. No significant differences in the prevalence of maternal delivery at home were observed in 2022 based on partner's current age, spousal age gap or household size (**Table 2**).

247 **Table 2:** Prevalence (weighted percentage) of maternal delivery at home by sociodemographic
248 characteristics in Bangladesh, 2012-2022

Variables	2012-13 % [95%CI]	p-value	2019 % [95%CI]	p-value	2022 % [95%CI]	p-value
Age of women (in years)		<0.001		<0.001		<0.001
15-19	65.2 [60.9,69.3]		45.8 [42.5,49.0]		34 [29.9,38.4]	
20-24	66.7 [64.1,69.1]		42.9 [40.8,45.1]		32.4 [29.1,35.9]	
25-29	67.6 [64.8,70.3]		45.8 [43.6,48.1]		34 [30.3,38.0]	
30-34	67.3 [63.4,70.9]		48 [45.3,50.8]		35.7 [31.2,40.4]	
35-49	79.1 [75.3,82.4]		58.3 [54.5,62.1]		49.1 [41.7,56.6]	
Education of women		<0.001		<0.001		<0.001
No education	87.5 [85.4,89.3]		75.6 [72.0,78.8]		55.7 [45.7,65.2]	
Primary	79.4 [76.9,81.7]		63.9 [61.4,66.3]		52.7 [48.5,57.0]	
Secondary	63.8 [61.3,66.2]		42.5 [40.8,44.2]		33.1 [30.4,35.9]	
Higher	32.2 [28.4,36.3]		19.3 [17.2,21.6]		12.6 [9.9,15.9]	
Religion		<0.001		<0.001		<0.001
Non-Muslim	58.9 [52.7,64.7]		34.1 [29.4,39.2]		16.1 [11.2,22.6]	
Muslim	68.9 [67.2,70.6]		47.5 [46.2,48.9]		36.4 [33.9,38.9]	
Media exposure		<0.001		<0.001		<0.001
No	78.3 [75.2,81.1]		65.1 [62.9,67.3]		51.6 [47.8,55.4]	
Yes	56.3 [53.9,58.7]		37.3 [35.9,38.8]		26.4 [24.0,28.9]	
Number of ANC visits		<0.001		<0.001		<0.001
No visit	91.1 [89.6,92.4]		80.5 [78.2,82.6]		76.3 [70.0,81.6]	
1 visit	72.8 [68.6,76.7]		66.4 [63.2,69.5]		57.1 [52.2,61.9]	
2 visits	68.8 [65.6,71.9]		51.8 [49.0,54.6]		41.3 [37.0,45.7]	
3 visits	60.9 [57.2,64.6]		40.4 [37.6,43.2]		28.3 [23.9,33.2]	
4+ visits	38.6 [35.3,42.1]		23.9 [22.3,25.7]		19.2 [16.8,21.9]	
Parity		<0.001		<0.001		<0.001
1-2	61.8 [59.8,63.8]		39.9 [38.4,41.4]		28.9 [26.7,31.3]	
3+	81 [78.8,83.0]		60.9 [58.7,63.0]		50.1 [46.1,54.1]	
Age of husband (in year)		0.006		<0.001		0.0729
15-29	68.2 [65.5,70.7]		48.7 [46.5,50.9]		33.7 [30.5,37.0]	
30-34	66.5 [63.4,69.6]		42.7 [40.5,44.9]		33.5 [29.4,37.8]	
35-39	66.2 [62.7,69.4]		44.5 [41.9,47.1]		33.9 [30.1,37.9]	
40-44	68 [63.6,72.2]		48.2 [44.6,51.8]		37.8 [32.4,43.4]	
45-49	72.2 [66.3,77.5]		50.9 [46.0,55.8]		39 [30.5,48.3]	
50-90	80.6 [74.2,85.7]		58.5 [51.4,65.2]		50.8 [38.3,63.1]	
Spousal age gap (in year)		0.0019		<0.001		0.12
≤5	70.1 [67.6,72.4]		51.2 [49.3,53.2]		37.3 [34.1,40.7]	
6-9	69.3 [66.6,71.8]		47.5 [45.4,49.6]		33.9 [30.0,37.9]	
>9	64.1 [61.2,67.0]		38.7 [36.6,40.9]		32.4 [28.7,36.4]	
Education of household head		<0.001		<0.001		<0.001
No education	78.6 [76.7,80.3]		60.5 [58.3,62.6]		53.8 [48.7,58.8]	
Primary	72 [68.0,75.8]		52.3 [50.0,54.5]		46.9 [43.1,50.8]	
Secondary	59.5 [55.6,63.2]		36.4 [34.3,38.6]		29 [25.8,32.4]	
Higher	37.3 [33.2,41.6]		20.2 [17.7,22.9]		13.3 [10.8,16.4]	
Wealth index		<0.001		<0.001		<0.001
Richest	34.2 [30.3,38.3]		19.8 [17.6,22.3]		12.1 [9.3,15.6]	
Fourth	58.8 [55.1,62.5]		34.3 [31.8,36.9]		24.6 [21.0,28.7]	
Middle	75.2 [72.4,77.8]		46 [43.4,48.6]		31.6 [27.7,35.8]	
Second	81.6 [79.2,83.8]		58.4 [55.8,60.9]		45.5 [40.7,50.4]	
Poorest	86.5 [84.7,88.2]		73.8 [71.4,76.0]		57.3 [52.4,62.1]	
Household size		<0.001		<0.001		0.435
≤4	63.3 [60.4,66.1]		43.5 [41.5,45.4]		33.5 [30.3,36.9]	
5+	70.8 [68.9,72.7]		48.2 [46.6,49.9]		35.5 [32.8,38.3]	
Residence		<0.001		<0.001		<0.001

Urban	48.6 [43.9,53.4]	32.2 [29.5,35.1]	23.4 [19.7,27.5]
Rural	73.3 [71.6,74.9]	50.4 [48.9,51.9]	39.1 [36.2,42.1]

Determinants of maternal delivery at home

In multivariable analyses, women's religious affiliation, ANC utilization, parity, household head's educational attainment, household wealth status, and administrative division were consistently associated with the likelihood of home delivery across survey years. Muslim women exhibited significantly higher odds of maternal delivery at home compared to non-Muslim women, with the association strengthening over time- most notably in 2022 (AOR 3.20, 95% CI 1.99-5.14), relative to 2019 (AOR 1.99, 95% CI 1.60-2.47) and 2012-2013 (AOR 1.36, 95% CI 1.02-1.82). A clear inverse relationship was observed between ANC visit frequency and home delivery; women with four or more ANC visits had substantially lower odds of home delivery compared to those with no ANC visits, with similar magnitudes across all survey years (2012-13: AOR 0.16, 95% CI 0.12-0.20; 2019: AOR 0.17, 95% CI 0.14-0.20; 2022: AOR 0.16, 95% CI 0.11-0.23). Higher parity (≥3 children) was consistently associated with increased odds of maternal delivery at home (2012-13: AOR 1.73, 95% CI 1.25-2.39; 2019: AOR 1.73, 95% CI 1.46-2.05; 2022: AOR 1.81, 95% CI 1.40-2.35). Women residing in households led by individuals with higher educational attainment were significantly less likely to deliver at home compared to those in households headed by individuals with no formal education (2012-13: AOR 0.65, 95% CI 0.50-0.86; 2019: AOR 0.65, 95% CI 0.51-0.82; 2022: AOR 0.50, 95% CI 0.36-0.71). A strong socioeconomic gradient was evident, wherein women from the poorest households had markedly higher odds of maternal delivery at home compared to those from the wealthiest households (2012-13: AOR 2.54, 95% CI 1.80-3.59; 2019: AOR 3.34, 95% CI 2.59-4.30; 2022: AOR 2.84, 95% CI 1.86-4.33). Regionally, women from Khulna division consistently showed lower odds of home delivery (2012-13: AOR 0.61, 95% CI 0.48-0.78; 2019: AOR 0.59, 95% CI 0.47-0.74; 2022: AOR 0.38, 95% CI 0.26-0.56). Furthermore, women exposed to media were less likely than their counterparts for maternal delivery at home in recent years (2019: AOR 0.78, 95% CI 0.68-0.89; 2022: AOR 0.75, 95% CI 0.61-0.92). In contrast, maternal age, educational attainment, and urban-rural residence were significantly associated with maternal delivery at home in earlier survey rounds (2012–2013 and 2019), but their effects were less prominent in 2022 (**Figure 3 and Table S2**).

277
Figure 3: Determinants of maternal delivery at home in Bangladesh, 2012-2022

The frequency of ANC visits and household wealth were consistently associated with maternal

delivery at home across both urban and rural settings. Specifically, women who attended four

or more ANC visits and those from the poorest households demonstrated a lower likelihood of

maternal delivery at home compared to their counterparts. In rural areas, maternal delivery at

home was significantly associated over time with religious affiliation, media exposure, parity,

and the educational attainment of the household head. However, these associations were not

consistently observed among urban women across survey periods (**Table 3**).

Table 3: Determinants of maternal delivery at home in Bangladesh by place of residence, 2012-2022

| | 2012 | 2019 | 2022 | 2012 | 2019 | 2022 |
| | Urban | Urban | Urban | Rural | Rural | Rural |
Variables	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Age of women						
15-19y	1	1	1	1	1	1
20-24y	1.51[0.73,3.1]	1.16 [0.7-1.93]	1.7[0.88,3.29]	1.03 [0.79,1.34]	0.93 [0.76-1.14]	0.75 [0.55,1.03]
25-29y	0.8[0.32,1.99]	1.08 [0.56-2.07]	0.95 [0.35,2.56]	1 [0.64,1.55]	1.02 [0.78-1.34]	0.86 [0.57,1.29]
30-34y	0.47[(0.12,1.8]	0.41 [0.17-1.01]	1.19 [0.38,3.75]	0.6 [0.33,1.08]	0.92 [0.65-1.32]	0.81 [0.45,1.48]
35-49y	0.64[0.13,3.02]	0.66 [0.22-2.13]	5.63 [1.59,19.92]**	0.52 [0.24,1.09]	0.79 [0.5-1.26]	1.2 [0.58,2.49]
Education of women						
No education	1	1	1	1	1	1
Primary	0.18 [0.07,0.46]***	1.26 [0.69,2.32]	1.58 [0.7,3.54]	1.11 [0.72,1.72]	0.79 [0.61,1.02]	1.2[0.71,2.04]
Secondary	0.29 [(0.12,0.7]**	0.67 [0.38,1.2]	1.37 [0.57,3.28]	0.81 [0.53,1.23]	0.61[0.47,0.79]***	1.03[0.6,1.76]
Higher	0.11 [0.04,0.27]***	0.3 [0.14,0.64]**	1.03 [0.34,3.11]	0.47 [0.29,0.77]**	0.41[0.3,0.56]***	0.65[0.34,1.24]
Religion						
Non-muslim		1	1	1	1	1
Muslim	1.34[0.65,2.77	1.90 [1.1-3.26]*	2.16[0.84,5.56]	1.43[1.02,2.01]*	1.98 [1.56-2.52]***	3.29[1.92,5.64]*
Media exposure						
No	1	1	1	1	1	1
yes	1.72[0.83,3.56]	0.89 [0.62-1.28]	0.85[0.51,1.43]	0.71[0.54,0.93]*	0.78 [0.67-0.88]***	0.73[0.59,0.91]**
Number of ANC visits						
No visit	1	1	1	1	1	1
1 visit	0.42[0.18,0.97]*	0.91 [0.49-1.7]	0.27[0.11,0.66]**	0.4[0.27,0.6]***	0.59 [0.47-0.75]***	0.57[0.37,0.88]*

2 visits	0.69[0.32,1.5]	0.38 [0.22-0.65]***	0.36[0.15,0.9]*	0.37[0.28,0.5]***	0.39 [0.32-0.48]***	0.32[0.21,0.48]***
3 visits	0.38[0.19,0.76]**	0.23 [0.13-0.41]***	0.25[0.09,0.68]**	0.29[0.21,0.4]***	0.29 [(0.23-0.35]***	0.22[0.13,0.36]***
4+ visits	0.12[0.06,0.24]***	0.14 [0.08-0.24]***	0.11[0.05,0.27]***	0.16[0.12,0.22]***	0.18 [0.15-0.22]***	0.17[0.11,0.27]***
Parity						
1-2	1	1	1	1	1	1
3 or more	1.1[0.54,2.23]	1.95 [1.25-3.02]**	1.37[0.73,2.57]	1.86[1.22,2.84]**	1.67 [1.39-2]***	2.04[1.49,2.78]***
Age of husband (in year)						
14-29	1	1	1	1	1	1
30-34	2.59[1.4,4.77]**	0.71 [0.46-1.11]	0.76[0.37,1.55]	1.19[0.88,1.61]	0.93 [0.76-1.14]	1.31[0.94,1.82]
35-39	2.13[0.89,5.08]	0.87 [0.46-1.62]	0.54[0.22,1.32]	1.06[0.66,1.7]	0.95 [0.72-1.26]	0.99[0.64,1.54]
40-44	4.59[1.41,14.98]*	0.87 [0.35-2.16]	0.68[0.25,1.89]	1.08[0.56,2.05]	1.02 [0.7-1.5]	0.83[0.45,1.53]
45-49	1[0.17,5.93]	1.17 [0.39-3.5]	0.63[0.15,2.61]	1.31[0.57,2.99]	1.01 [0.62-1.65]	0.64[0.27,1.53]
50-90	4.43[0.62,31.86]	0.56 [0.14-2.22]	0.22[0.05,1.03]	1.28[0.46,3.53]	1.45 [0.79-2.65]	0.87[0.35,2.12]
Spousal age gap (in year)						
<= 5 year	1	1	1	1	1	1
6-9 year	1.27[0.79,2.02]	0.93 [0.64-1.34]	1.07[0.59,1.91]	1.02[0.8,1.3]	1.07 [0.91-1.26]	0.94[0.68,1.3]
>9 year	0.53[0.25,1.12]	0.73 [0.42-1.25]	1.59[0.78,3.22]	0.94[0.65,1.35]	0.85 [0.67-1.08]	0.98[0.65,1.48]
Education of household head						
No education	1	1	1	1	1	1
Primary	1.05[0.53,2.08]	0.79[0.53,1.19]	0.88[0.47,1.65]	0.83[0.61,1.13]	0.94[0.8,1.09]	0.9[0.67,1.2]
Secondary	0.68[0.38,1.23]	0.64[0.44,0.93]*	0.56[0.32,0.99]*	0.85[0.66,1.11]	0.78[0.67,0.92]**	0.67[0.5,0.91]**
Higher	0.57[0.31,1.07]	0.63[0.36,1.1]	0.35[0.17,0.72]**	0.68[0.5,0.92]*	0.66[0.5,0.86]**	0.54[0.36,0.8]**

Wealth index					
Richest	1	1	1	1	1	1
Fourth	1.82[1.07,3.09]*	1.38 [0.95-2]	1.95[0.9,4.25]	1.4[1.07,1.81]*	1.78 [1.37-2.31]***	1.29[0.82,2.03]
Middle	3.17[1.48,6.82]**	1.39[(0.89-2.17]	2.83[1.48,5.41]**	2.31[1.69,3.17]***	2.23 [1.71-2.9]***	1.22[0.81,1.84]
Second	4.66[1.53,14.22]**	1.98 [1.17-3.34]*	4.29[1.96,9.4]***	2.18[1.51,3.14]***	2.77 [2.09-3.67]***	1.82[1.15,2.88]*
Poorest	4.44[1.84,10.74]**	2.17 [1.13-4.17]*	6.29[2.52,15.74]***	2.35[1.55,3.56]***	4.10 [3.04-5.53]***	2.11[1.31,3.41]**
Household size					
≤4	1	1	1	1	1	1
5+	1.2[0.73,1.96]	0.90 [0.66-1.22]	0.65[0.39,1.07]	1.06[0.83,1.35]	1.00 [0.87-1.15]	0.98[0.78,1.23]
Division					
Dhaka	1	1	1	1	1	1
Barishal	0.92[0.44,1.9]	1.20[0.67,2.15]	0.49[0.23,1.01]	1.83[1.26,2.66]**	1.9[1.45,2.48]***	1.56[0.97,2.51]
Chattogram	0.79[0.48,1.31]	0.96[0.67,1.38]	1.55[0.9,2.68]	1.65[1.14,2.38]**	1.32[1.04,1.68]**	0.94[0.62,1.42]
Khulna	0.37[0.21,0.67]**	0.33[0.17,0.64]**	0.32[0.12,0.9]*	0.74[0.54,1.01]	0.68[0.53,0.87]**	0.39[0.25,0.61]***
Mymensingh		0.42[0.28,0.72]**	1.33(0.71,2.47]		2.1[1.56,2.69]***	1.16[0.78,1.74]
Rajshahi	0.46[0.21,1.03]	0.60[0.36,0.99]*	0.44[0.19,1]*	0.76[0.54,1.07]	0.93[0.74,1.18]	0.75[0.49,1.17]
Rangpur	1.77[0.81,3.9]	1.42[0.77,3.62]	0.66[0.26,1.65]	1.67[1.18,2.37]**	1.24[0.95,1.54]	1.01[0.65,1.57]
Sylhet	1.05[0.45,2.44]	1.49[0.85,2.63]	0.76[0.4,1.47]	1.37[0.91,2.05]	1.55[1.17,2.06]**	1.18[0.77,1.8]

288   **Note:** AOR= Adjusted Odds Ratio, CI= Confidence interval, * p-value<0.05, ** p-value<0.01, *** p-value<0.001

Discussion

This study provides important insights into the changing landscape and persistent disparities of maternal delivery at home in Bangladesh over the past decade. Although the national prevalence of maternal delivery at home has declined markedly from 68.0% in 2012 to 34.9% in 2022, our findings reveal that this progress is not uniform across the country. Maternal delivery at home varied substantially at the district level, with the hill track districts-such as Rangamati, Bandarban, and Khagrachari-continuing to report rates exceeding 60% in 2022, nearly double the national average. These geographic inequities highlight the limited reach of facility-based delivery interventions in remote and marginalized areas, where barriers such as poor infrastructure, entrenched cultural preferences, and restricted access to healthcare persist despite national improvements.

Our analysis also identified several key factors associated with maternal delivery at home, including religion, number of antenatal care (ANC) visits, parity, education of the household head, household wealth, and administrative division. Muslim women were more likely to have a maternal delivery at home compared to non-Muslims. Women with fewer than four ANC visits, those with higher parity, and those from the poorest households were significantly more likely to experience maternal delivery at home. Importantly, our findings show that the determinants of maternal delivery at home are not consistent between urban and rural areas, suggesting that context-specific, modifiable factors must be considered when designing interventions. While household wealth, ANC visits, and parity were the strongest predictors in rural settings, in urban areas, religion and the education level of the household head played a more prominent role. This urban-rural divergence indicates that interventions must be tailored to the unique drivers of maternal delivery at home in each setting-prioritizing economic empowerment and improved ANC access in rural areas, and focusing on education and cultural engagement in urban contexts.

Our findings reveal substantial division- and district-level differences in home delivery rates. Home delivery remained particularly prevalent in the Barisal, Sylhet, and Mymensingh divisions, where approximately half of the deliveries still occurred at home. These regions have historically faced challenges in healthcare infrastructure, literacy rates, and accessibility, which may contribute to the sustained high prevalence of home deliveries [30]. Previous studies have

similarly highlighted that healthcare services in the Mymensingh and Barisal divisions are
inadequate, which may further explain the persistence of home deliveries in these areas [31,32].
Similar findings were also reported in different countries, which reported that home delivery rates
were typically higher in less developed areas with poor transportation and access to healthcare
[33,34]. At the district level, home deliveries were most common in peripheral and remote areas
such as Bandarban, Rangamati, Khagrachari, Bhola, and Habiganj. This pattern aligns with
findings from Nepal, where home delivery rates were notably higher in mountainous regions due
to limited access to healthcare facilities [35].

One important factor influencing home delivery was found to be educational attainment.
Compared with women without formal education, those with greater levels of education had a
much lower likelihood of giving birth at home. This result is in line with other research from
several nations, indicating that education is crucial in influencing maternal healthcare choices
[36,37]. Awareness and knowledge might be the main reasons for this type of scenario, as women
with lower levels of education may possess limited awareness regarding maternal healthcare
facilities and the advantages of in-facility deliveries [37–39]. Education empowers women to
efficiently navigate healthcare institutions and make informed decisions about their health [37].
While the gap in home delivery rates between educated and less educated women has narrowed
over time, further efforts are required to ensure universal access to facility-based deliveries,
particularly among disadvantaged groups.

Home delivery rates are strongly impacted by the household head's educational attainment in
addition to the mother's education. The more educated the household head was, the less likely the
woman was to give birth at home. According to a Nigerian study, households with educated heads
spent more money on healthcare, which increased the rates of institutional delivery [40]. Another
study in Bangladesh revealed that a lower education level of household heads decreases maternal
healthcare utilization in Bangladesh [41]. These findings emphasize the broader role of education
not only in individual decision-making but also in shaping household-level health-seeking
behaviors.

Media exposure was identified as a protective factor against home delivery. Increasing exposure
to media can alter perceptions positively and encourage expectant mothers to choose facility-

based deliveries over home deliveries, as public awareness and health-related behaviors are
strongly influenced by media exposure. Owing to a lack of knowledge, cultural customs, and
restricted access to medical facilities, home deliveries continue in many areas. To promote
institutional deliveries and enhance the health of mothers and newborns, increasing media
exposure can be a useful tactic [42,43]. In Nigeria, similar findings have been reported, where
increased usage of maternal healthcare services was linked to media exposure [44]. Expanding
media-based health awareness programs could be an effective strategy to reduce home delivery
rates further.

Religious beliefs also play a role in home delivery choices. Our research suggests that Muslim
women exhibited a greater tendency for home delivery than their non-Muslim peers did. Cultural
norms, traditional beliefs, and socioeconomic factors may contribute to this trend [45]. A similar
pattern has been observed in India, where women of the Muslim faith exhibited a greater
propensity for home births [36]. However, further research is necessary to investigate the
underlying causes of this difference and to develop culturally relevant treatments that encourage
institutional deliveries within various religious communities.

The utilization of antenatal care (ANC) is a crucial factor influencing home delivery, as women
who visit four or more ANC appointments are much less inclined to deliver at home. This aligns
with findings from Afghanistan, where ANC visits were strongly associated with lower home
deliveries [46]. High-quality ANC services provide essential counseling, health screenings, and
education on the benefits of skilled birth attendance, which collectively influence women's
decisions to seek facility-based delivery over home delivery [47–51]. The consistent relationship
between ANC utilization and a lower likelihood of home delivery highlights the need to
strengthen ANC services and ensure equitable access for all pregnant women.

Parity, or the number of previous childbirths, also influenced home delivery rates. Home
deliveries were more common among women with three or more children. This finding is similar
to those of studies conducted in Bangladesh and Ethiopia, which reported that women who are
first-time mothers or have fewer pregnancies are more likely to choose facility-based births
[30,37]. Likewise, multiparous women in rural Zambia are more inclined to give birth at home
[52]. Women with multiple previous births may feel more confident about delivering at home,

particularly if they have had uncomplicated deliveries in the past. Furthermore, multiparous
women may be more likely to opt for delivery at home because of logistical difficulties, financial
limitations, and a dependence on traditional birth attendants. Addressing these barriers through
community-based interventions and financial support programs could help encourage facility-
based deliveries among high-parity women.

Our findings highlight a strong wealth disparity in home delivery rates. Women from poorer
households were significantly more likely to give birth at home than those from affluent
households were. Similar findings have been reported in several countries, including Nigeria,
Nepal, and India [53–56]. Financial constraints, out-of-pocket fees, and indirect costs such as
transportation and lost wages may deter women from seeking facility-based deliveries. Studies
from Kenya and West Africa similarly report persistent economic disparities in maternal
healthcare utilization, emphasizing the need for financial support mechanisms such as conditional
cash transfers and subsidies to promote institutional deliveries among economically
disadvantaged groups [47,48,57].

Urban☐rural differences in home delivery rates were also evident. Home deliveries were more
common among women in rural areas. According to some studies conducted in Africa and
Indonesia, urban women are significantly more likely than their rural counterparts are to give
birth in a medical facility [58–60]. Women living in rural areas confront unique obstacles to
facility delivery, such as a lack of qualified healthcare personnel, restricted availability of medical
services, and poor healthcare infrastructure [44,49,61,62]. Bridging the urban☐rural gap in
maternal healthcare requires strategic investments in rural healthcare facilities, improved
transportation networks, and community-based healthcare interventions.

Bangladesh has implemented a range of interventions over the past decade-including the
expansion of community clinics, maternal health voucher schemes, and awareness campaigns-that
have contributed to the overall reduction in maternal delivery at home and increased skilled birth
attendance [63]. The Maternal Health Voucher Scheme (MHVS), launched in 2007, has been
particularly impactful in improving access to antenatal care, skilled assistance at delivery, and
postnatal care among disadvantaged women, though administrative and coverage challenges
remain [64,65]. Our study affirms the success of these efforts at the national level but also reveals

persistent gaps: maternal delivery at home remains disproportionately high among marginalized

groups such as women in the hill tracts, high-parity mothers, and the poorest households.

Additionally, our findings highlight that determinants of maternal delivery at home differ between

urban and rural settings, with factors like religion and household head's education being more

influential in urban areas, while wealth, parity, and ANC visits are more critical in rural regions.

These context-specific disparities suggest that, despite overall progress, current interventions have

not fully addressed the unique barriers faced by these populations. To further reduce maternal

delivery at home, particularly in high-prevalence areas, our study recommends mobile and

culturally sensitive outreach in remote districts, greater engagement of religious and community

leaders, incentivization of complete ANC utilization, and adaptation of policies to the distinct

socio-demographic realities of different regions.

Strengths and Limitations

A key strength of this study is its ability to provide valuable insights at both the national and

subnational levels using data from the MICS 2012–13 and 2019, as well as the BDHS 2022. The

use of the most recent BDHS 2022 dataset, which employed rigorous sampling techniques,

enhances the generalizability of the findings to women of reproductive age (15–49 years) across

the country. Additionally, the large sample size and application of appropriate statistical

techniques allowed for robust identification of significant predictors and assessment of inequality

gradients across multiple subgroups. The findings can inform policies aimed at reducing maternal

delivery at home clusters and identifying areas with limited access to facility-based childbirth

services.

However, the study has certain limitations. First, as with most MICS and DHS-based

studies, data are self-reported and thus subject to recall bias and social desirability bias, which

may affect the accuracy of reported reproductive behaviors. Second, due to the cross-sectional

nature of the survey, the study cannot establish causal relationships between maternal delivery at

home and its determinants. Third, some relevant covariates, such as maternal comorbidities,

nutritional status, dietary patterns, physical activity, and psychosocial factors, were not included

in the BDHS and thus could not be examined, even though they may have significant implications

for maternal delivery at home. Fourth, socioeconomic characteristics were measured at the time of
the interview, which may not accurately reflect the woman's status at the time of pregnancy or
childbirth. And finally, although the BDHS 2022 incorporated district-level estimates for the first
time, the limited sample size in several districts has resulted in unstable prevalence rates, with
some districts reporting values of '0,' which is inconsistent with prior nationally representative
surveys such as MICS.

Conclusion

Over the last decade, Bangladesh has made great strides in lowering the country's home delivery
rates. However, one-third of births still occur at home, highlighting ongoing challenges in
ensuring universal access to institutional deliveries. Geographic, socioeconomic, and
demographic disparities persist, underscoring the need for targeted interventions. Strengthening
healthcare infrastructure, particularly in remote and hilly areas, improving maternal health
education, and increasing ANC utilization can further reduce home deliveries. Additionally,
addressing financial barriers through subsidies and incentive programs may encourage greater use
of facility-based care. Culturally sensitive approaches, media-based awareness campaigns, and
community engagement initiatives should be prioritized to achieve equitable maternal healthcare
access. To develop inclusive and successful policies, future studies should look at how religious,
cultural, and economic issues affect decisions about maternal delivery at home. Bangladesh may
achieve universal facility-based birthing by resolving these inequities, which would ultimately
improve the health of expectant mothers and newborns.

Declarations

Declarations

Acknowledgements: We are grateful to the MICS and DHS team for allowing us to conduct the

analysis of this study using the MICS 2012 and 2019 datasets and the BDHS 2022 dataset.

Author Contributions: Conceptualization, R.B.I; Data curation, R.B.I, and M.L.K; Formal

analysis, R.B.I, and M.L.K; Methodology, R.B.I, and S.T.A.N; Visualization, R.B.I, and S.T.A.N;

Investigation, S.M., and A.A; Supervision, S.T.A.N ; Validation, M.B.L; Writing—original draft,

R.B.I, and M.L.K; Writing - review and editing, S.M., A.A., S.T.A.N and M.B.L; All authors

have read and agreed to the published version of the manuscript.

Funding: The authors received no specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

Preprint: Available on medRxiv.

Consent for publication: Not applicable.

Competing interests: There are no potential conflicts (financial, professional, or personal) for

any of the authors to disclose.

Data availability: All data relevant to the study are included in the article or uploaded as

supplemental information. All relevant data are within the manuscript. Raw MICS

(https://mics.unicef.org/) and BDHS (https://dhsprogram. com/) data are publicly available that

can be accessed upon request.

References

1. Trends in maternal mortality 2000 to 2023: estimates by WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division. [cited 13 May 2025]. Available: https://www.who.int/publications/i/item/9789240108462

2. El-Saharty S, Ohno N. South Asia's quest for reduced maternal mortality: What the data show. In: World Bank Blog: Investing in Health. 2015.

3. Hossain AT, Siddique AB, Jabeen S, Khan S, Haider MM, Ameen S, et al. Maternal mortality in Bangladesh: Who, when, why, and where? A national survey-based analysis. J Glob Health. 2023;13. doi:10.7189/JOGH.13.07002

4. The DHS Program - Bangladesh: DHS, 2022 - Final Report (English). [cited 13 May 2025]. Available: https://www.dhsprogram.com/publications/publication-fr386-dhs-final-reports.cfm?cssearch=1551986_1

5. Document Viewer. [cited 13 May 2025]. Available: https://docs.un.org/en/A/RES/70/1

6. UNISEF. Maternal and newborn health. 2023.

7. Kitila SB, Feyissa GT, Olika AK, Wordofa MA. Maternal Healthcare in Low- and Middle-Income Countries: A Scoping Review. Health Serv Insights. 2022;15. doi:10.1177/11786329221100310

8. Oladipo IA, Akinwaare MO. Trends and patterns of maternal deaths from 2015 to 2019, associated factors and pregnancy outcomes in rural Lagos, Nigeria: a cross-sectional study. Pan African Medical Journal. 2023;44.

9. Hernández-Vásquez A, Chacón-Torrico H, Bendezu-Quispe G. Prevalence of home birth among 880,345 women in 67 low- and middle-income countries: A meta-analysis of Demographic and Health Surveys. SSM Popul Health. 2021;16. doi:10.1016/j.ssmph.2021.100955

10. Shrestha SK, Banu B, Khanom K, Ali L, Thapa N, Stray-Pedersen B, et al. Changing trends on the place of delivery: Why do Nepali women give birth at home? Reprod Health. 2012;9: 1–8. doi:10.1186/1742-4755-9-25

11. Sahoo J, Singh SV, Gupta VK, Garg S, Kishore J. Do socio-demographic factors still predict the choice of place of delivery: A cross-sectional study in rural North India. J Epidemiol Glob Health. 2015;5: S27–S34.

12. Shrestha SK, Banu B, Khanom K, Ali L, Thapa N, Stray-Pedersen B, et al. Changing trends on the place of delivery: Why do Nepali women give birth at home? Reprod Health. 2012;9: 1–8. doi:10.1186/1742-4755-9-25

13. Tiruneh SA, Lakew AM, Yigizaw ST, Sisay MM, Tessema ZT. Trends and determinants of home delivery in Ethiopia: Further multivariate decomposition analysis of 2005-2016 Ethiopian Demographic Health Surveys. BMJ Open. 2020;10: 1–10. doi:10.1136/bmjopen-2019-034786

14. Caulfield T, Onyo P, Byrne A, Nduba J, Nyagero J, Morgan A, et al. Factors influencing place of
delivery for pastoralist women in Kenya: A qualitative study. BMC Womens Health. 2016;16: 1–11.
doi:10.1186/s12905-016-0333-3

15. Sarker BK, Rahman M, Rahman T, Hossain J, Reichenbach L, Mitra DK. Reasons for preference of
home delivery with traditional birth attendants (TBAs) in Rural Bangladesh: A qualitative
exploration. PLoS One. 2016;11. doi:10.1371/journal.pone.0146161

16. Islam MM, Shahjahan M. Exploring the reasons and factors influencing the choice of home
delivery of births in rural Bangladesh: a community-based cross-sectional study. J Health Res.
2022;36. doi:10.1108/JHR-07-2020-0284

17. World Health Organization. WORLD HEALTH STATISTICS - MONITORING HEALTH FOR THE SDGs.
World Health Organization. 2016; 1.121.

18. Dey R, Dey SR, Haque M, Rahman AB, Kundu S, Setu SP, et al. Mapping the prevalence and
covariates associated with home delivery in Bangladesh: A multilevel regression analysis. PLoS
One. 2024;19: e0313606.

19. Talukder A, Anik BH, Hossain MI, Haq I, Habib MJ. Socioeconomic and demographic factors for
mothers' delivery at home: a comparative study among BDHS 2007, 2011 and 2014. Asian Journal
of Social Health and Behavior. 2022;5: 10–17.

20. Ahmed KT, Karimuzzaman M, Mahmud S, Rahman L, Hossain MM, Rahman A. Influencing factors
associated with maternal delivery at home in urban areas: a cross-sectional analysis of the
Bangladesh Demographic and Health Survey 2017–2018 data. J Health Popul Nutr. 2023;42: 83.

21. BANGLADESH 2012-13 MICS FINAL REPORT RELEASED | UNICEF MICS. [cited 13 May 2025].
Available: https://mics.unicef.org/news/bangladesh-2012-13-mics-final-report-released

22. Bangladesh 2019 MICS Report - English | PDF | Drinking Water | Sanitation. [cited 13 May 2025].
Available: https://www.scribd.com/document/537704949/Bangladesh-2019-MICS-Report-English

23. Pathey P. Bangladesh multiple indicator cluster survey 2012–2013 Key findings. Bangladesh Bur
Stat UNICEF Bangladesh. 2014;2014: 2014.

24. Noor STA, Shil P, Talukdar A, Aktar S, Uddin MJ. Exploring Factors Influencing Wealth-Related
Disparities in Institutional Delivery: A Decomposition Analysis Using Bangladesh Multiple Indicator
Cluster Survey (MICS) 2019. Public Health Challenges. 2025;4: e70066.

25. National Institute of Population Research and Training (NIPORT), ICF. Bangladesh Demographic
and Health Survey 2022 Final Report. 2024. Available:
https://www.dhsprogram.com/publications/publication-FR386-DHS-Final-Reports.cfm

26. Ahmed KT, Karimuzzaman Md, Mahmud S, Rahman L, Hossain MdM, Rahman A. Influencing
factors associated with maternal delivery at home in urban areas: a cross-sectional analysis of the
Bangladesh Demographic and Health Survey 2017–2018 data. J Health Popul Nutr. 2023;42: 83.
doi:10.1186/s41043-023-00428-9

27.   Enuameh YAK, Okawa S, Asante KP, Kikuchi K, Mahama E, Ansah E, et al. Factors influencing health
facility delivery in predominantly rural communities across the three ecological zones in Ghana: A
cross-sectional study. PLoS One. 2016;11: 1–16. doi:10.1371/journal.pone.0152235

28.   Dey D, Haque MdS, Islam MdM, Aishi UI, Shammy SS, Mayen MdSA, et al. The proper application
of logistic regression model in complex survey data: a systematic review. BMC Med Res Methodol.
2025;25: 15. doi:10.1186/s12874-024-02454-5

29.   von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the
Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for reporting
observational studies. International Journal of Surgery. 2014;12: 1495–1499.
doi:https://doi.org/10.1016/j.ijsu.2014.07.013

30.   Ahmed KT, Karimuzzaman Md, Mahmud S, Rahman L, Hossain MdM, Rahman A. Influencing
factors associated with maternal delivery at home in urban areas: a cross-sectional analysis of the
Bangladesh Demographic and Health Survey 2017–2018 data. J Health Popul Nutr. 2023;42: 83.
doi:10.1186/s41043-023-00428-9

31.   Huda TM, Chowdhury M, El Arifeen S, Dibley MJ. Individual and community level factors
associated with health facility delivery: A cross sectional multilevel analysis in Bangladesh. PLoS
One. 2019;14: e0211113.

32.   Badiuzzaman M, Murshed SM, Rieger M. Improving maternal health care in a post conflict setting:
evidence from Chittagong Hill tracts of Bangladesh. J Dev Stud. 2020;56: 384–400.

33.   Teshale AB, Alem AZ, Yeshaw Y, Kebede SA, Liyew AM, Tesema GA, et al. Exploring spatial
variations and factors associated with skilled birth attendant delivery in Ethiopia: geographically
weighted regression and multilevel analysis. BMC Public Health. 2020;20: 1–19.

34.   Moshi FV, Lymo G, Gibore NS, Kibusi SM. Prevalence and factors associated with home childbirth
with unskilled birth assistance in Dodoma-Tanzania: a cross sectional study. East Afr Health Res J.
2020;4: 92.

35.   Shahabuddin ASM, De Brouwere V, Adhikari R, Delamou A, Bardaj A, Delvaux T. Determinants of
institutional delivery among young married women in Nepal: Evidence from the Nepal
Demographic and Health Survey, 2011. BMJ Open. 2017;7: e012446.

36.   Sahoo J, Singh SV, Gupta VK, Garg S, Kishore J. Do socio-demographic factors still predict the
choice of place of delivery: A cross-sectional study in rural North India. J Epidemiol Glob Health.
2015;5. doi:10.1016/j.jegh.2015.05.002

37.   Tiruneh SA, Lakew AM, Yigizaw ST, Sisay MM, Tessema ZT. Trends and determinants of home
delivery in Ethiopia: Further multivariate decomposition analysis of 2005-2016 Ethiopian
Demographic Health Surveys. BMJ Open. 2020;10: 1–10. doi:10.1136/bmjopen-2019-034786

38.   Straneo M, Fogliati P, Azzimonti G, Mangi S, Kisika F. Where do the rural poor deliver when high
coverage of health facility delivery is achieved? Findings from a community and hospital survey in
Tanzania. PLoS One. 2014;9: 1–17. doi:10.1371/journal.pone.0113995

39. Kruk ME, Hermosilla S, Larson E, Vail D, Chen Q, Mazuguni F, et al. Who is left behind on the road
to universal facility delivery? A cross-sectional multilevel analysis in rural Tanzania. Tropical
Medicine and International Health. 2015;20: 1057–1066. doi:10.1111/tmi.12518

40. Ogundari K, Abdulai A. Determinants of household's education and healthcare spending in
Nigeria: Evidence from survey data. African Development Review. 2014;26: 1–14.

41. Rahman MM, Amin MT, Ferdous Z, Patwary H, Haider MM. Education of household head and
maternal healthcare utilization: the case of Bangladesh. BMC Public Health. 2024;24: 3439.

42. Mills S, Williams JE, Adjuik M, Hodgson A. Use of health professionals for delivery following the
availability of free obstetric care in Northern Ghana. Matern Child Health J. 2008;12: 509–518.
doi:10.1007/s10995-007-0288-y

43. Agha S, Carton TW. Determinants of institutional delivery in rural Jhang, Pakistan. Int J Equity
Health. 2011;10: 1–12. doi:10.1186/1475-9276-10-31

44. Babalola S, Fatusi A. Determinants of use of maternal health services in Nigeria - Looking beyond
individual and household factors. BMC Pregnancy Childbirth. 2009;9: 43. doi:10.1186/1471-2393-
9-43

45. Kamal SMM, Hassan CH, Alam GM. Determinants of institutional delivery among women in
Bangladesh. Asia Pac J Public Health. 2015;27: NP1372–NP1388. doi:10.1177/1010539513486178

46. Rahman M, Saha P, Uddin J. Associations of antenatal care visit with utilization of institutional
delivery care services in Afghanistan: intersections of education, wealth, and household decision-
making autonomy. BMC Pregnancy Childbirth. 2022;22: 255.

47. Colombara D V., Hernández B, Schaefer A, Zyznieuski N, Bryant MF, Desai SS, et al. Institutional
delivery and satisfaction among indigenous and poor women in Guatemala, Mexico, and Panama.
PLoS One. 2016;11: 1–17. doi:10.1371/journal.pone.0154388

48. Tekelab T, Yadecha B, Melka AS. Antenatal care and women's decision making power as
determinants of institutional delivery in rural area of Western Ethiopia. BMC Res Notes. 2015;8:
1–8. doi:10.1186/s13104-015-1708-5

49. Séraphin MN, Ngnie-Teta I, Ayoya MA, Khan MR, Striley CW, Boldon E, et al. Determinants of
Institutional Delivery Among Women of Childbearing Age in Rural Haiti. Matern Child Health J.
2015;19: 1400–1407. doi:10.1007/s10995-014-1646-1

50. Agha S, Williams E. Quality of antenatal care and household wealth as determinants of
institutional delivery in Pakistan: Results of a cross-sectional household survey. Reprod Health.
2016;13: 1–8. doi:10.1186/s12978-016-0201-5

51. Feyissa TR, Genemo GA. Determinants of institutional delivery among childbearing age women in
Western Ethiopia, 2013: Unmatched case control study. PLoS One. 2014;9: 1–7.
doi:10.1371/journal.pone.0097194

52.   Scott NA, Henry EG, Kaiser JL, Mataka K, Rockers PC, Fong RM, et al. Factors affecting home
delivery among women living in remote areas of rural Zambia: a cross-sectional, mixed-methods
analysis. Int J Womens Health. 2018; 589–601.

53.   Newell R, Spillman I, Newell M-L. The use of facilities for labor and delivery: the views of women in
rural Uganda. J Public Health Afr. 2017;8: 592.

54.   Olorunsaiye CZ, Huber LB, Laditka SB, Kulkarni S, Boyd AS. Factors associated with health facility
delivery in West and Central Africa: a multilevel analysis of seven countries. Health Care Women
Int. 2020;41: 3–21.

55.   Singh PK, Rai RK, Singh L. Examining the effect of household wealth and migration status on safe
delivery care in urban India, 1992–2006. 2012.

56.   Devkota B, Maskey J, Pandey AR, Karki D, Godwin P, Gartoulla P, et al. Determinants of home
delivery in Nepal–A disaggregated analysis of marginalised and non-marginalised women from the
2016 Nepal Demographic and Health Survey. PLoS One. 2020;15: e0228440.

57.   Kinuthia J, Kohler P, Okanda J, Otieno G, Odhiambo F, John-Stewart G. A community-based
assessment of correlates of facility delivery among HIV-infected women in western Kenya. BMC
Pregnancy Childbirth. 2015;15: 1–9. doi:10.1186/s12884-015-0467-6

58.   Idris H, Budiastuti A, Razak R, Hasyim H. Delivery Services Utilization Based on Urban Rural Status
in Indonesia. Open Access Maced J Med Sci. 2022;10: 1147–1152.

59.   Newell R, Spillman I, Newell M-L. The use of facilities for labor and delivery: the views of women in
rural Uganda. J Public Health Afr. 2017;8: 592.

60.   Moyer CA, Mustafa A. Drivers and deterrents of facility delivery in sub-Saharan Africa: a
systematic review. Reprod Health. 2013;10: 1–14.

61.   Dahiru T, Oche OM. Determinants of antenatal care, institutional delivery and postnatal care
services utilization in Nigeria. Pan African Medical Journal. 2015;21: 1–17.
doi:10.11604/pamj.2015.21.321.6527

62.   Amano A, Gebeyehu A, Birhanu Z. Institutional delivery service utilization in Munisa Woreda,
South East Ethiopia: a community based cross-sectional study. BMC Pregnancy Childbirth.
2012;12: 1–6. doi:10.1186/1471-2393-12-105

63.   Mahmood SS, Amos M, Hoque S, Mia MN, Chowdhury AH, Hanifi SMA, et al. Does healthcare
voucher provision improve utilisation in the continuum of maternal care for poor pregnant
women? Experience from Bangladesh. Glob Health Action. 2019;12.
doi:10.1080/16549716.2019.1701324,

64.   Chowdhury AH, Hanifi SMA, Iqbal M, Hossain A, Stones W, Amos M, et al. Does maternal health
voucher scheme have association with distance inequality in maternal and newborn care
utilization? Evidence from rural Bangladesh. PLoS One. 2023;18: e0295306.

65.   Talukder MN, Rob U, Musa SAJM, Bajracharya A, Keya KT, Noor FR, et al. Evaluation of the impact
of the voucher program for improving maternal health behavior and status in Bangladesh. 2014.

2012-13	2019	2022

<20%	20%-40%	41%-60%	61%-80%	>80%