Full Title: Spatiotemporal variations in the prevalence and determinants of maternal delivery at

home in Bangladesh, 2012-2022: A population-based cross-sectional study

Short Title: Maternal delivery at home in Bangladesh

Authors: Raisha Binte Islam[1*], Syed Toukir Ahmed Noor[2], Md. Lutful Kader[1], Somaya

Mostarin[3], Mushfika Binta Latif[4], Amina Anjum[5]

Affiliations:

[1]Department of Statistics, University of Dhaka, Dhaka-1000, Bangladesh.

[2]Department of Statistics, Shahjalal University of Science and Technology, Sylhet-3114,

Bangladesh.

[3]Department of Health Education, National Institute of Preventive and Social Medicine

(NIPSOM), Mohakhali, Dhaka-1212, Bangladesh

[4]Department of Economics, York University, Canada.

[5]Department of Health Service Management Policy, National Institute of Preventive and Social

Medicine (NIPSOM), Mohakhali, Dhaka-1212, Bangladesh

*Corresponding author

Raisha Binte Islam

Department of Statistics, University of Dhaka, Dhaka-1000, Bangladesh.

Email: r.b.islam1997@gmail.com

Abstract

Background

Despite remarkable advancements in reducing maternal and neonatal mortality, maternal

delivery at home continue to pose a significant public health challenge in many low- and middle-

income countries including Bangladesh. Although the incidence of maternal delivery at home

has declined substantially over the past decade, pronounced disparities persist across

geographic, socioeconomic, and demographic dimensions in Bangladesh. A nuanced

understanding of the spatiotemporal patterns in the prevalence and determinants of maternal

delivery at home is essential for designing targeted interventions. This study investigates the

spatial variations in the prevalence and determinants of maternal delivery at home in Bangladesh

over time.

Methods

We analyzed data from the Multiple Indicator Cluster Survey conducted in 2012–13 and 2019

as well as the Bangladesh Demographic and Health Survey conducted in 2022 to compare

district-level estimates for maternal delivery at home. Descriptive statistics and multivariable

logistic regression were employed to assess the prevalence and determinants of maternal

delivery at home.

Results

The prevalence of maternal delivery at home declined from 68.1% in 2012-13 to 34.9% in 2022

in Bangladesh. However, significant geographical disparities persist. The prevalence of

maternal delivery at home was higher in the Barisal (48.9%), Chattogram, Sylhet, and

Mymensingh divisions. The lowest rates were observed in Dhaka and Khulna. At the district

level, remote areas such as Bandarban, Rangamati (55.9%), and Bhola presented higher rates of

home deliveries. Multivariable logistic regression analysis identified several significant

predictors of home delivery. Compared with those with secondary or higher education, women

with no formal education were more likely to deliver at home. Limited ANC visits (≤3 visits)

significantly increased the likelihood of home birth. Rural residence, lower household wealth,

multiparity (≥3 children), and lack of media exposure were also significant determinants of

home delivery.

Conclusions

Despite progress in reducing home deliveries, persistent geographic and socioeconomic

disparities highlight the need for targeted interventions. Strengthening healthcare infrastructure

in underserved regions, improving maternal health awareness, increasing ANC utilization, and

addressing financial barriers through subsidies and incentive programs can further reduce the

number of home births. Future research should explore the influence of cultural and religious

factors to develop context-specific policies that promote equitable access to facility-based

childbirth.

Keywords: Maternal delivery, spatiotemporal variations, socioeconomic disparities,

determinants, Bangladesh.

Introduction

Despite a 40% reduction in the global maternal mortality ratio (MMR)- from 328 deaths per 100,000 live births in 2000 to 197 in 2023- over 90% of all maternal deaths continue to occur in low- and lower-middle-income countries [1] . South Asia remains a critical contributor, accounting for nearly one-quarter of global maternal fatalities [2]. As a prominent example, Bangladesh, a lower-middle-income country in South Asia, has made commendable progress in reducing its MMR from 412 deaths per 100,000 live births in 1976 to 123 in 2020 [3]. This progress has been accompanied by significant scale-up in key maternal health interventions, including antenatal care (4+ visits) from 17% in 2004 to 41% in 2022, facility-based delivery from 10% in 2004 to 65% in 2022, skilled birth attendance from 23% in 2007 to 70% in 2022, and postnatal care (by medically trained providers within 48 hours) from 16% in 2004 to 55% in 2022 [4]. Despite these advances, the gains have been unevenly distributed across geographic regions and socioeconomic strata, resulting in persistent disparities in maternal health outcomes. Consequently, Bangladesh remains off track to meet the Sustainable Development Goal target (SDG 3.1), which targets an MMR of fewer than 70 deaths per 100,000 live births by 2030 [5].

To accelerate the progress toward this ambitious goal, there is an urgent need to improve emergency obstetric services and minimize the incidence of maternal delivery at home, which are associated with preventable maternal and neonatal mortality. Globally, maternal delivery at home accounted for an estimated 287,000 maternal and 2.4 million newborn deaths in 2020 alone [6,7]. A study from Nigeria found that non-institutional delivery (delivery outside health facilities, often by unskilled attendants) was associated with a 79.2% chance of maternal death among the cases studied [8]. In Bangladesh, more than one-third of all deliveries still occur at home, with marked regional disparities across administrative divisions [4] Reducing home deliveries is, therefore, a critical public health priority to advance equity and reduce mortality and neonatal mortality [9–11]Maternal delivery at home is influenced by a complex interplay

of sociodemographic, economic, cultural, and health system factors. Key factors include maternal education, quantity of antenatal care visits, parity, household wealth, religion, and geographic accessibility to healthcare facilities [12–14]. In addition, entrenched poverty, traditional and religious norms, poor transportation infrastructure, and the limited autonomy among women in decision-making further exacerbate the preference for delivery at home [15]. Other factors such as low maternal and paternal education, advanced maternal age, multiparity, and low wealth status also contribute to increased home delivery practices [16]. Fear of cesarean sections, perceptions of poor quality of care, and the lack of female doctors also play substantial roles in influencing delivery preferences [15].

According to the availability, accessibility, acceptability, and quality (AAAQ) framework, both supply-side (e.g. healthcare infrastructure and provider availability) and demand-side (e.g. household-level barriers and perceptions) factors must be addressed to reduce maternal delivery at home [17]. While individual and household-level determinants have been widely studied, health system constraints- especially at the subnational level- remain largely underexplored in the context of Bangladesh. Identifying subnational areas (here district in Bangladesh) with persistently high rates of maternal delivery at home is essential for designing and implementing evidence-based maternal health interventions. However, limited research has assessed the temporal trends and geographic distribution of maternal delivery at home in Bangladesh [18–20] Moreover, little is known about how associated factors of maternal delivery at home vary over time across urban and rural settings. To our knowledge, no prior study has comprehensively investigated subnational spatiotemporal variations in maternal delivery at home or explored the evolving determinants of this practice.

Therefore, we aimed to investigate district-level spatiotemporal patterns of maternal delivery at home and analyze the temporal dynamics of associated factors in urban and rural areas in

Bangladesh. The findings will inform targeted, context-specific policies and support the

implementation of effective maternal health interventions.

Materials and Methods

Data Sources

This study utilized secondary data derived from three nationally representative, population-

based cross-sectional surveys: Multiple Indicator Cluster Survey (MICS) conducted in 2012–

13 and 2019, and the Bangladesh Demographic and Health Survey (BDHS) conducted in 2022.

The MICS, implemented by the Bangladesh Bureau of Statistics (BBS) with technical and

financial support from the United Nations International Children's Emergency Fund (UNICEF),

provides comprehensive data on household characteristics and maternal and child health

indicators at the district level. The BDHS, administered under the authority of the National

Institute of Population Research and Training (NIPORT), Medical Education and Family

Welfare Division, Ministry of Health and Family Welfare (MOHFW), offers vital demographic

and health statistics, including district-level estimates for the first time in 2022.

Sampling Design and Sample Size

Both the MICS and BDHS employed a uniform methodology to ensure national

representativeness. Detailed methodological descriptions are available in the respective

technical reports [4,21,22]. Briefly, the MICS applied a multistage stratified cluster sampling

strategy, covering all 64 administrative districts of Bangladesh. Each district was stratified into

urban and rural areas. Enumeration areas (EAs) were selected using the probability proportional

to size (PPS) method within these strata. From each selected EA, 20 households were

systematically sampled [23,24]The MICS sampled a total of 55,120 in 2012-2013 and 64,400

households in 2019, of 51,895 and 61,242 were successfully interviewed, with response rates

of 98.5% and 99.4%, respectively. After excluding respondents with no delivery history or

missing birth outcome data, the final analytical sample included 7795 and 9284 women aged

15-49 years with a history of childbirth in 2012-2013 and 2019 respectively.

The BDHS 2022 utilized a two-stage stratified sampling strategy based on the 2011 Population

Census. In the first stage, 675 EAs were selected (237 urban and 438 rural) using PPS sampling.

In the second stage, 45 households were systematically selected from each EA to ensure

adequate representation across residence types [25]. In 2022, the final response rate of BDHS

was 99.6%, with 30,330 households selected and 30,018 households interviewed. The final

analytical sample from BDHS 2022 consisted of 3,691 women of reproductive age (15–49

years) with recent birth histories.

Outcome Variable

The main outcome variable for this study was maternal delivery at home, defined

dichotomously. Deliveries occurring at the respondent's home were coded as "1" (maternal

delivery at home), and those occurring at any health facility were coded as "0".

Explanatory Variables

A total of 13 independent variables were selected on an extensive review of previous literature

[13,26,27], capturing individual-, household-, and community-level determinants. Individual-

level factors included women's age, education (no education, primary, secondary, and higher),

number of antenatal care (ANC) visits (none, 1, 2, 3, and ≥4), parity (1–2, ≥3), media exposure,

religion (Non-Muslim, Muslim), husband's age, and spousal age gap (≤5 years, 6–9 years, and

more than 9 years). Household-level factors included education of household's head, household

size and household's socioeconomic status. The community-level factors encompassed

administrative division and place of residence (urban or rural). Socioeconomic status was

assessed using the wealth index, constructed via principal component analysis (PCA) as

provided in MICS and BDHS datasets. Households were classified into five quintiles: poorest,

poorer, middle, richer, and richest [4,21,22].

Statistical analysis

The data were analyzed via Stata version 17.0 (StataCorp, College Station, Texas, USA),

whereas data visualization was performed via R version 4.3.0. To account for the complex

survey design, we incorporated primary sampling units (PSUs) and strata and applied sampling

weights to ensure representativeness. The Stata "svyset" command was used to adjust for the

survey design [28], and all analyses adhered to the STROBE cross-sectional reporting

guidelines [29].

Descriptive statistics were used to summarize the characteristics of the study population, which

were reported as frequencies and proportions in percentages. We conducted bivariate analysis

using the Chi-square test to assess associations between the explanatory variables and the

outcome variable. We employed simple binary logistic regression analysis to estimate crude

odds ratios (ORs) with 95% CIs for each explanatory variables in the unadjusted analysis.

Variables that were statistically significant in the simple logistic regression models were

subsequently included in the multiple logistic regression model to estimate adjusted odds ratios

(AORs) with their respective 95% CIs. Statistical significance was determined at the 5% level

(p-value <0.05).

To account for the complex survey design, we applied the "svyset" command in Stata,

incorporating sampling weights, strata and primary sampling units (PSUs). All the analyses

were done using the "svy" prefix to ensure design-based estimates and inference [28]. The study

adhered to the STROBE guidelines for reporting cross-sectional studies [29]. Spatial analysis

was conducted using district-level choropleth maps to visualize geographic disparities in

maternal delivery at home. We used Stata version 17.0 (StataCorp, College Station, Texas,

USA) for data analysis, R version 4.3.0 for data visualization and ArcGIS version 10.8 for

spatial mapping.

Ethical Consideration

This study utilized deidentified secondary data obtained from the UNICEF

(https://mics.unicef.org/surveys) and DHS (https://www.dhsprogram.com/data/available-

datasets.cfm) websites. Since the data were publicly available and did not contain any personally

identifiable information, ethical approval was not needed. All participants provided informed

consent, and the original surveys were carried out in accordance with ethical standards and

approved by the appropriate institutional review boards.

Results

Sample characteristics

Across all survey years, approximately one-third of women were aged 20-24 years (33.6% in

2012-13, 32.2% in 2019, and 33.8% in 2022). Educational attainment demonstrated a positive

trend over time, the proportion of women completing higher education increased from 12.4%

in 2012 to 18.5% in 2022. Nearly nine in 10 women were Muslim (89.2% in 2012, 90.4% in

2019 and 92.7% in 2022). Media exposure declined slightly over the years, from 76.6% in 2012

to 66.3% in 2022. Utilization of ANC services improved, with the proportion of women

receiving at least four ANC increased from 22.7% in 2012 to 40.5% in 2022. Fertility pattern

showed that majority of women had 1–2 children, increasing from 65.6% in 2012 to 71.9% in

2022. The proportion of households headed by individuals with no education accounted for

56.2% in 2012, 30.3% in 2019 and 15.5% in 2022. The proportion of women in the poorest

wealth quintile declined from 28.3% in 2012 to 20.2% in 2022. Household size remained

relatively stable, with approximately two-thirds of women living in households comprising five

or more members (65.6% in 2012-13, 63% in 2019 and 69.5% in 2022). Most women resided

in rural areas (83.8% in 2012, 80.9% in 2019 and 73% in 2022), and the Dhaka division

consistently had the highest representation across survey years (**Table 1**).

Table 1: Sample characteristics of the respondents, presented in terms of weighted percentage

Variables	2012-13		2019		2022	
	Number	%	Number	%	Number	%
Age of women (in years)						
15-19	906	11.6	1,234	13.3	614	16.6
20-24	2,615	33.6	2,987	32.2	1248	33.8
25-29	2,351	30.2	2,562	27.6	962	26.1
30-34	1,143	14.7	1,640	17.7	592	16.0
35-49	780	10.0	861	9.3	274	7.4
Education of women						
No education	1,621	20.8	846	9.1	183	5.0
Primary	2,294	29.4	2,151	23.2	831	22.5
Secondary	2,917	37.4	4,691	50.5	1995	54.0
Higher	963	12.4	1,596	17.2	682	18.5
Religion						
Non-Muslim	842	10.8	896	9.7	269	7.3
Muslim	6,953	89.2	8,388	90.4	3422	92.7
Media exposure						
No	1,186	23.5	3,308	35.7	1243	33.7
Yes	3,872	76.6	5,967	64.3	2447	66.3
Number of ANC visits						
No visit	2,730	35.2	1,677	18.1	268	7.4
1 visit	1,018	13.1	1,123	12.1	526	14.6
2 visits	1,227	15.8	1,611	17.4	694	19.2
3 visits	1,024	13.2	1,529	16.5	658	18.2
4+ visits	1,758	22.7	3,339	36.0	1462	40.5
Parity						
1-2	5,116	65.6	6,401	69.0	2653	71.9
3+	2,679	34.4	2,883	31.1	1037	28.1
Age of husband (in year)						
15-29	2,514	32.9	2,789	30.3	1237	33.8
30-34	1,960	25.6	2,627	28.5	913	24.9
35-39	1,606	21.0	2,062	22.4	832	22.7
40-44	872	11.4	988	10.7	407	11.1
45-49	409	5.4	505	5.5	191	5.2
50-90	290	3.8	243	2.6	79	2.2
Spousal age gap (in year)						
≤5	2,939	38.0	3,534	38.4	1408	38.5
6-9	2,630	34.0	3,088	33.5	1192	32.6
>9	2,173	28.1	2,593	28.1	1060	29.0
Education of household head						
No education	4,375	56.2	2,812	30.3	566	15.5
Primary	1,009	13.0	2,794	30.1	1059	28.9
Secondary	1,410	18.1	2,602	28.0	1277	34.9
Higher	992	12.7	1,073	11.6	757	20.7
Wealth index						
Richest	1,174	15.1	1,587	17.1	661	17.9
Fourth	1,283	16.5	1,770	19.1	718	19.5
Middle	1,474	18.9	1,810	19.5	787	21.3
Second	1,661	21.3	1,886	20.3	780	21.1
Poorest	2,203	28.3	2,231	24.0	745	20.2

Household size						
≤4	2684	34.4	3,439	37.0	1128	30.6
5+	5,111	65.6	5,845	63.0	2563	69.5
Residence						
Urban	1,261	16.2	1,774	19.1	997	27.0
Rural	6,534	83.8	7,510	80.9	2694	73.0
Division						
Barisal	703	9.0	835	9.0	219	5.9
Chittagong	1,585	20.3	1,926	20.8	793	21.5
Dhaka	2,002	25.7	1,795	19.3	926	25.1
Khulna	1,044	13.4	1,275	13.7	380	10.3
Mymensingh			558	6.0	333	9.0
Rajshahi	715	9.2	956	10.3	381	10.3
Rangpur	972	12.5	1,135	12.2	419	11.4
Sylhet	774	9.9	804	8.7	240	6.5

Spatial variations in maternal delivery at home

In Bangladesh, the prevalence of maternal delivery at home exhibited a marked decline over the past decade, decreasing from 68.1% [66.0-70.0%] in 2012-2013 to 46.4% [45.0-48.0%] in 2019 and 34.9% [32.0-37.0%] in 2022. Regional variation was evident, with Khulna Division demonstrating the lowest prevalence of maternal delivery at home across all years. In contrast, Barishal had the highest prevalence in 2012-13 and 2022, followed by the Sylhet division (**Figure 1**). Although regional variations were evident, district-level disparities in maternal delivery at home were more pronounced (**Table S1 and Figure 2**). In 2022, the prevalence of maternal delivery at home was >50% in 14 of 64 districts. The highest prevalence was observed in Gopalganj (68.8%), Khagrachhari (68.5%), and Lalmonirhat (66.3%), while the lowest was reported in Bandarban (0%), Chuadanga (0%), and Kushtia (6.5%). In 2012, home-based maternal delivery was most prevalent in hill tract districts such as Bandarban (97.2%) and Rangamati (88.6%), and in haor/coastal districts including Sunamganj (89.3%), Noakhali (88.1%), and Patuakhali (88.0%). By 2019, the burden remained high in hill districts-84.6% in Bandarban, 73.0% in Rangamati, and 76.1% in Khagrachhari. Despite a declining national trend, hill tract and northeastern districts had persistently high prevalence of maternal delivery at home.

Figure 1: National and divisional estimates of maternal delivery at home in Bangladesh, 2012-2022

Figure 2: Spatial distribution of home delivery in Bangladesh, 2012--2022

Subpopulation-level variations in maternal delivery at home

Despite national and subnational reduction, maternal delivery at home remained common among women aged 35-49 years, although the prevalence in this subgroup declined disproportionately from 79.1% in 2012-2013 to 49.1% in 2022. Women with no formal education consistently showed the highest prevalence of maternal delivery at home (87.5% in 2012-2013, 75.6% in 2019 and 55.7% in 2022), in contrast to women with higher education, whose prevalence fell from 32.2% to 12.6% over the same period. Muslim women persistently reported higher percentage of maternal delivery at home than non-Muslim women (68.9% vs 58.9% in 2012-2013, 47.5% vs 34.1% in 2019 and 36.4% vs 16.1% in 2022). No media exposure was also associated with a higher prevalence of maternal delivery at home, though the prevalence declined notably from 78.3% in 2012 to 51.6% in 2022. Women receiving four or more ANC visits had significantly lower prevalence of maternal delivery at home than those receiving none (38.6% vs 91.1% in 2012, 23.9% vs 80.5% in 2019 and 19.2% vs 76.3% in 2022). Higher parity (3+ children) was associated with increased maternal delivery at home relative to lower parity (1-2 children) over years (81.0% vs 61.8% in 2012, 60.9% vs 39.9% in 2019, and 50.1% vs 28.9% in 2022). Furthermore, women of households headed by individuals with no formal education, those from the poorest wealth quintile, rural residents exhibited

249 consistently higher prevalence of home-based maternal delivery. No significant differences in

250 the prevalence of maternal delivery at home were observed in 2022 based on partner's current

251 age, spousal age gap or household size (**Table 2**).

252 **Table 2:** Prevalence (weighted percentage) of maternal delivery at home by sociodemographic
253 characteristics in Bangladesh, 2012-2022

Variables	2012-13		2019		2022	
	% [95%CI]	p-value	% [95%CI]	p-value	% [95%CI]	p-value
Age of women (in years)		<0.001		<0.001		<0.001
15-19	65.2 [60.9,69.3]		45.8 [42.5,49.0]		34 [29.9,38.4]	
20-24	66.7 [64.1,69.1]		42.9 [40.8,45.1]		32.4 [29.1,35.9]	
25-29	67.6 [64.8,70.3]		45.8 [43.6,48.1]		34 [30.3,38.0]	
30-34	67.3 [63.4,70.9]		48 [45.3,50.8]		35.7 [31.2,40.4]	
35-49	79.1 [75.3,82.4]		58.3 [54.5,62.1]		49.1 [41.7,56.6]	
Education of women		<0.001		<0.001		<0.001
No education	87.5 [85.4,89.3]		75.6 [72.0,78.8]		55.7 [45.7,65.2]	
Primary	79.4 [76.9,81.7]		63.9 [61.4,66.3]		52.7 [48.5,57.0]	
Secondary	63.8 [61.3,66.2]		42.5 [40.8,44.2]		33.1 [30.4,35.9]	
Higher	32.2 [28.4,36.3]		19.3 [17.2,21.6]		12.6 [9.9,15.9]	
Religion		<0.001		<0.001		<0.001
Non-Muslim	58.9 [52.7,64.7]		34.1 [29.4,39.2]		16.1 [11.2,22.6]	
Muslim	68.9 [67.2,70.6]		47.5 [46.2,48.9]		36.4 [33.9,38.9]	
Media exposure		<0.001		<0.001		<0.001
No	78.3 [75.2,81.1]		65.1 [62.9,67.3]		51.6 [47.8,55.4]	
Yes	56.3 [53.9,58.7]		37.3 [35.9,38.8]		26.4 [24.0,28.9]	
Number of ANC visits		<0.001		<0.001		<0.001
No visit	91.1 [89.6,92.4]		80.5 [78.2,82.6]		76.3 [70.0,81.6]	
1 visit	72.8 [68.6,76.7]		66.4 [63.2,69.5]		57.1 [52.2,61.9]	
2 visits	68.8 [65.6,71.9]		51.8 [49.0,54.6]		41.3 [37.0,45.7]	
3 visits	60.9 [57.2,64.6]		40.4 [37.6,43.2]		28.3 [23.9,33.2]	
4+ visits	38.6 [35.3,42.1]		23.9 [22.3,25.7]		19.2 [16.8,21.9]	
Parity		<0.001		<0.001		<0.001
1-2	61.8 [59.8,63.8]		39.9 [38.4,41.4]		28.9 [26.7,31.3]	
3+	81 [78.8,83.0]		60.9 [58.7,63.0]		50.1 [46.1,54.1]	
Age of husband (in year)		0.006		<0.001		0.0729
15-29	68.2 [65.5,70.7]		48.7 [46.5,50.9]		33.7 [30.5,37.0]	
30-34	66.5 [63.4,69.6]		42.7 [40.5,44.9]		33.5 [29.4,37.8]	
35-39	66.2 [62.7,69.4]		44.5 [41.9,47.1]		33.9 [30.1,37.9]	
40-44	68 [63.6,72.2]		48.2 [44.6,51.8]		37.8 [32.4,43.4]	
45-49	72.2 [66.3,77.5]		50.9 [46.0,55.8]		39 [30.5,48.3]	
50-90	80.6 [74.2,85.7]		58.5 [51.4,65.2]		50.8 [38.3,63.1]	
Spousal age gap (in year)		0.0019		<0.001		0.12
≤5	70.1 [67.6,72.4]		51.2 [49.3,53.2]		37.3 [34.1,40.7]	
6-9	69.3 [66.6,71.8]		47.5 [45.4,49.6]		33.9 [30.0,37.9]	
>9	64.1 [61.2,67.0]		38.7 [36.6,40.9]		32.4 [28.7,36.4]	
Education of household head		<0.001		<0.001		<0.001
No education	78.6 [76.7,80.3]		60.5 [58.3,62.6]		53.8 [48.7,58.8]	
Primary	72 [68.0,75.8]		52.3 [50.0,54.5]		46.9 [43.1,50.8]	
Secondary	59.5 [55.6,63.2]		36.4 [34.3,38.6]		29 [25.8,32.4]	
Higher	37.3 [33.2,41.6]		20.2 [17.7,22.9]		13.3 [10.8,16.4]	
Wealth index		<0.001		<0.001		<0.001

Richest	34.2 [30.3,38.3]		19.8 [17.6,22.3]		12.1 [9.3,15.6]	
Fourth	58.8 [55.1,62.5]		34.3 [31.8,36.9]		24.6 [21.0,28.7]	
Middle	75.2 [72.4,77.8]		46 [43.4,48.6]		31.6 [27.7,35.8]	
Second	81.6 [79.2,83.8]		58.4 [55.8,60.9]		45.5 [40.7,50.4]	
Poorest	86.5 [84.7,88.2]		73.8 [71.4,76.0]		57.3 [52.4,62.1]	
Household size		<0.001		<0.001		0.435
≤4	63.3 [60.4,66.1]		43.5 [41.5,45.4]		33.5 [30.3,36.9]	
5+	70.8 [68.9,72.7]		48.2 [46.6,49.9]		35.5 [32.8,38.3]	
Residence		<0.001		<0.001		<0.001
Urban	48.6 [43.9,53.4]		32.2 [29.5,35.1]		23.4 [19.7,27.5]	
Rural	73.3 [71.6,74.9]		50.4 [48.9,51.9]		39.1 [36.2,42.1]	

Determinants of maternal delivery at home

In multivariable analyses, women's religious affiliation, ANC utilization, parity, household head's educational attainment, household wealth status, and administrative division were consistently associated with the likelihood of home delivery across survey years. Muslim women exhibited significantly higher odds of maternal delivery at home compared to non-Muslim women, with the association strengthening over time- most notably in 2022 (AOR 3.20, 95% CI 1.99-5.14), relative to 2019 (AOR 1.99, 95% CI 1.60-2.47) and 2012-2013 (AOR 1.36, 95% CI 1.02-1.82). A clear inverse relationship was observed between ANC visit frequency and home delivery; women with four or more ANC visits had substantially lower odds of home delivery compared to those with no ANC visits, with similar magnitudes across all survey years (2012-13: AOR 0.16, 95% CI 0.12-0.20; 2019: AOR 0.17, 95% CI 0.14-0.20; 2022: AOR 0.16, 95% CI 0.11-0.23). Higher parity (≥3 children) was consistently associated with increased odds of maternal delivery at home (2012-13: AOR 1.73, 95% CI 1.25-2.39; 2019: AOR 1.73, 95% CI 1.46-2.05; 2022: AOR 1.81, 95% CI 1.40-2.35). Women residing in households led by individuals with higher educational attainment were significantly less likely to deliver at home compared to those in households headed by individuals with no formal education (2012-13: AOR 0.65, 95% CI 0.50-0.86; 2019: AOR 0.65, 95% CI 0.51-0.82; 2022: AOR 0.50, 95% CI 0.36-0.71). A strong socioeconomic gradient was evident, wherein women from the poorest households had markedly higher odds of maternal delivery at home compared to those from the

wealthiest households (2012-13: AOR 2.54, 95% CI 1.80-3.59; 2019: AOR 3.34, 95% CI 2.59-

4.30; 2022: AOR 2.84, 95% CI 1.86-4.33). Regionally, women from Khulna division

consistently showed lower odds of home delivery (2012-13: AOR 0.61, 95% CI 0.48-0.78;

2019: AOR 0.59, 95% CI 0.47-0.74; 2022: AOR 0.38, 95% CI 0.26-0.56). Furthermore, women

exposed to media were less likely than their counterparts for maternal delivery at home in recent

years (2019: AOR 0.78, 95% CI 0.68-0.89; 2022: AOR 0.75, 95% CI 0.61-0.92). In contrast,

maternal age, educational attainment, and urban-rural residence were significantly associated

with maternal delivery at home in earlier survey rounds (2012–2013 and 2019), but their effects

were less prominent in 2022 (**Figure 3 and Table S2**).

Figure 3: Determinants of maternal delivery at home in Bangladesh, 2012-2022

The frequency of ANC visits and household wealth were consistently associated with maternal

delivery at home across both urban and rural settings. Specifically, women who attended four

or more ANC visits and those from the poorest households demonstrated a lower likelihood of

maternal delivery at home compared to their counterparts. In rural areas, maternal delivery at

home was significantly associated over time with religious affiliation, media exposure, parity,

and the educational attainment of the household head. However, these associations were not

consistently observed among urban women across survey periods (**Table 3**).

Table 3: Determinants of maternal delivery at home in Bangladesh by place of residence, 2012-2022

| | 2012 | 2019 | 2022 | 2012 | 2019 | 2022 |
| | Urban | Urban | Urban | Rural | Rural | Rural |
Variables	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)	AOR (95% CI)
Age of women						
15-19y	1	1	1	1	1	1
20-24y	1.51[0.73,3.1]	1.16 [0.7-1.93]	1.7[0.88,3.29]	1.03 [0.79,1.34]	0.93 [0.76-1.14]	0.75 [0.55,1.03]
25-29y	0.8[0.32,1.99]	1.08 [0.56-2.07]	0.95 [0.35,2.56]	1 [0.64,1.55]	1.02 [0.78-1.34]	0.86 [0.57,1.29]
30-34y	0.47[(0.12,1.8]	0.41 [0.17-1.01]	1.19 [0.38,3.75]	0.6 [0.33,1.08]	0.92 [0.65-1.32]	0.81 [0.45,1.48]
35-49y	0.64[0.13,3.02]	0.66 [0.22-2.13]	5.63 [1.59,19.92]**	0.52 [0.24,1.09]	0.79 [0.5-1.26]	1.2 [0.58,2.49]
Education of women						
No education	1	1	1	1	1	1
Primary	0.18 [0.07,0.46]***	1.26 [0.69,2.32]	1.58 [0.7,3.54]	1.11 [0.72,1.72]	0.79 [0.61,1.02]	1.2[0.71,2.04]
Secondary	0.29 [(0.12,0.7]**	0.67 [0.38,1.2]	1.37 [0.57,3.28]	0.81 [0.53,1.23]	0.61[0.47,0.79]***	1.03[0.6,1.76]
Higher	0.11 [0.04,0.27]***	0.3 [0.14,0.64]**	1.03 [0.34,3.11]	0.47 [0.29,0.77]**	0.41[0.3,0.56]***	0.65[0.34,1.24]
Religion						
Non-muslim		1	1	1	1	1
Muslim	1.34[0.65,2.77	1.90 [1.1-3.26]*	2.16[0.84,5.56]	1.43[1.02,2.01]*	1.98 [1.56-2.52]***	3.29[1.92,5.64]*
Media exposure						
No	1	1	1	1	1	1
yes	1.72[0.83,3.56]	0.89 [0.62-1.28]	0.85[0.51,1.43]	0.71[0.54,0.93]*	0.78 [0.67-0.88]***	0.73[0.59,0.91]**
Number of ANC visits						
No visit	1	1	1	1	1	1
1 visit	0.42[0.18,0.97]*	0.91 [0.49-1.7]	0.27[0.11,0.66]**	0.4[0.27,0.6]***	0.59 [0.47-0.75]***	0.57[0.37,0.88]*

2 visits	0.69[0.32,1.5]	0.38 [0.22-0.65]***	0.36[0.15,0.9]*	0.37[0.28,0.5]***	0.39 [0.32-0.48]***	0.32[0.21,0.48]***
3 visits	0.38[0.19,0.76]**	0.23 [0.13-0.41]***	0.25[0.09,0.68]**	0.29[0.21,0.4]***	0.29 [(0.23-0.35]***	0.22[0.13,0.36]***
4+ visits	0.12[0.06,0.24]***	0.14 [0.08-0.24]***	0.11[0.05,0.27]***	0.16[0.12,0.22]***	0.18 [0.15-0.22]***	0.17[0.11,0.27]***
Parity						
1-2	1	1	1	1	1	1
3 or more	1.1[0.54,2.23]	1.95 [1.25-3.02]**	1.37[0.73,2.57]	1.86[1.22,2.84]**	1.67 [1.39-2]***	2.04[1.49,2.78]***
Age of husband (in year)						
14-29	1	1	1	1	1	1
30-34	2.59[1.4,4.77]**	0.71 [0.46-1.11]	0.76[0.37,1.55]	1.19[0.88,1.61]	0.93 [0.76-1.14]	1.31[0.94,1.82]
35-39	2.13[0.89,5.08]	0.87 [0.46-1.62]	0.54[0.22,1.32]	1.06[0.66,1.7]	0.95 [0.72-1.26]	0.99[0.64,1.54]
40-44	4.59[1.41,14.98]*	0.87 [0.35-2.16]	0.68[0.25,1.89]	1.08[0.56,2.05]	1.02 [0.7-1.5]	0.83[0.45,1.53]
45-49	1[0.17,5.93]	1.17 [0.39-3.5]	0.63[0.15,2.61]	1.31[0.57,2.99]	1.01 [0.62-1.65]	0.64[0.27,1.53]
50-90	4.43[0.62,31.86]	0.56 [0.14-2.22]	0.22[0.05,1.03]	1.28[0.46,3.53]	1.45 [0.79-2.65]	0.87[0.35,2.12]
Spousal age gap (in year)						
<= 5 year	1	1	1	1	1	1
6-9 year	1.27[0.79,2.02]	0.93 [0.64-1.34]	1.07[0.59,1.91]	1.02[0.8,1.3]	1.07 [0.91-1.26]	0.94[0.68,1.3]
>9 year	0.53[0.25,1.12]	0.73 [0.42-1.25]	1.59[0.78,3.22]	0.94[0.65,1.35]	0.85 [0.67-1.08]	0.98[0.65,1.48]
Education of household head						
No education	1	1	1	1	1	1
Primary	1.05[0.53,2.08]	0.79[0.53,1.19]	0.88[0.47,1.65]	0.83[0.61,1.13]	0.94[0.8,1.09]	0.9[0.67,1.2]
Secondary	0.68[0.38,1.23]	0.64[0.44,0.93]*	0.56[0.32,0.99]*	0.85[0.66,1.11]	0.78[0.67,0.92]**	0.67[0.5,0.91]**
Higher	0.57[0.31,1.07]	0.63[0.36,1.1]	0.35[0.17,0.72]**	0.68[0.5,0.92]*	0.66[0.5,0.86]**	0.54[0.36,0.8]**

Wealth index						
Richest	1	1	1	1	1	1
Fourth	1.82[1.07,3.09]*	1.38 [0.95-2]	1.95[0.9,4.25]	1.4[1.07,1.81]*	1.78 [1.37-2.31]***	1.29[0.82,2.03]
Middle	3.17[1.48,6.82]**	1.39[(0.89-2.17]	2.83[1.48,5.41]**	2.31[1.69,3.17]***	2.23 [1.71-2.9]***	1.22[0.81,1.84]
Second	4.66[1.53,14.22]**	1.98 [1.17-3.34]*	4.29[1.96,9.4]***	2.18[1.51,3.14]***	2.77 [2.09-3.67]***	1.82[1.15,2.88]*
Poorest	4.44[1.84,10.74]**	2.17 [1.13-4.17]*	6.29[2.52,15.74]***	2.35[1.55,3.56]***	4.10 [3.04-5.53]***	2.11[1.31,3.41]**
Household size						
≤4	1	1	1	1	1	1
5+	1.2[0.73,1.96]	0.90 [0.66-1.22]	0.65[0.39,1.07]	1.06[0.83,1.35]	1.00 [0.87-1.15]	0.98[0.78,1.23]
Division						
Dhaka	1	1	1	1	1	1
Barishal	0.92[0.44,1.9]	1.20[0.67,2.15]	0.49[0.23,1.01]	1.83[1.26,2.66]**	1.9[1.45,2.48]***	1.56[0.97,2.51]
Chattogram	0.79[0.48,1.31]	0.96[0.67,1.38]	1.55[0.9,2.68]	1.65[1.14,2.38]**	1.32[1.04,1.68]**	0.94[0.62,1.42]
Khulna	0.37[0.21,0.67]**	0.33[0.17,0.64]**	0.32[0.12,0.9]*	0.74[0.54,1.01]	0.68[0.53,0.87]**	0.39[0.25,0.61]***
Mymensingh		0.42[0.28,0.72]**	1.33(0.71,2.47)		2.1[1.56,2.69]***	1.16[0.78,1.74]
Rajshahi	0.46[0.21,1.03]	0.60[0.36,0.99]*	0.44[0.19,1]*	0.76[0.54,1.07]	0.93[0.74,1.18]	0.75[0.49,1.17]
Rangpur	1.77[0.81,3.9]	1.42[0.77,3.62]	0.66[0.26,1.65]	1.67[1.18,2.37]**	1.24[0.95,1.54]	1.01[0.65,1.57]
Sylhet	1.05[0.45,2.44]	1.49[0.85,2.63]	0.76[0.4,1.47]	1.37[0.91,2.05]	1.55[1.17,2.06]**	1.18[0.77,1.8]

296   **Note:** AOR= Adjusted Odds Ratio, CI= Confidence interval, * p-value<0.05, ** p-value<0.01, *** p-value<0.001

Discussion

This study provides important insights into the changing landscape and persistent disparities of maternal delivery at home in Bangladesh over the past decade. Although the national prevalence of maternal delivery at home has declined markedly from 68.0% in 2012 to 34.9% in 2022, our findings reveal that this progress is not uniform across the country. Maternal delivery at home varied substantially at the district level, with the hill track districts-such as Rangamati, Bandarban, and Khagrachari-continuing to report rates exceeding 60% in 2022, nearly double the national average. These geographic inequities highlight the limited reach of facility-based delivery interventions in remote and marginalized areas, where barriers such as poor infrastructure, entrenched cultural preferences, and restricted access to healthcare persist despite national improvements.

Our analysis also identified several key factors associated with maternal delivery at home, including religion, number of antenatal care (ANC) visits, parity, education of the household head, household wealth, and administrative division. Muslim women were more likely to have maternal delivery at home compared to non-Muslims. Women with fewer than four ANC visits, those with higher parity, and those from the poorest households were significantly more likely to experience maternal delivery at home. Importantly, our findings show that the determinants of maternal delivery at home are not consistent between urban and rural areas, suggesting that context-specific, modifiable factors must be considered when designing interventions. While household wealth, ANC visits, and parity were the strongest predictors in rural settings, in urban areas, religion and the education level of the household head played a more prominent role. This urban-rural divergence indicates that interventions must be tailored to the unique drivers of maternal delivery at home in each setting-

prioritizing economic empowerment and improved ANC access in rural areas, and focusing on

education and cultural engagement in urban contexts.

Our findings reveal substantial division- and district-level differences in home delivery rates. Home

delivery remained particularly prevalent in the Barisal, Sylhet, and Mymensingh divisions, where

approximately half of the deliveries still occurred at home. These regions have historically faced

challenges in healthcare infrastructure, literacy rates, and accessibility, which may contribute to the

sustained high prevalence of home deliveries [30]. Previous studies have similarly highlighted that

healthcare services in the Mymensingh and Barisal divisions are inadequate, which may further

explain the persistence of home deliveries in these areas [31,32]. Similar findings were also reported

in different countries, which reported that home delivery rates were typically higher in less

developed areas with poor transportation and access to healthcare [33,34]. At the district level,

home deliveries were most common in peripheral and remote areas such as Bandarban, Rangamati,

Khagrachari, Bhola, and Habiganj. This pattern aligns with findings from Nepal, where home

delivery rates were notably higher in mountainous regions due to limited access to healthcare

facilities [35].

One important factor influencing home delivery was found to be educational attainment. Compared

with women without formal education, those with greater levels of education had a much lower

likelihood of giving birth at home. This result is in line with other research from several nations,

indicating that education is crucial in influencing maternal healthcare choices [36,37]. Awareness

and knowledge might be the main reasons for this type of scenario, as women with lower levels of

education may possess limited awareness regarding maternal healthcare facilities and the

advantages of in-facility deliveries [37–39]. Education empowers women to efficiently navigate

healthcare institutions and make informed decisions about their health [37]. While the gap in home

delivery rates between educated and less educated women has narrowed over time, further efforts

are required to ensure universal access to facility-based deliveries, particularly among

disadvantaged groups.

Home delivery rates are strongly impacted by the household head's educational attainment in

addition to the mother's education. The more educated the household head was, the less likely the

woman was to give birth at home. According to a Nigerian study, households with educated heads

spent more money on healthcare, which increased the rates of institutional delivery [40]. Another

study in Bangladesh revealed that a lower education level of household heads decreases maternal

healthcare utilization in Bangladesh [41]. These findings emphasize the broader role of education

not only in individual decision-making but also in shaping household-level health-seeking

behaviors.

Media exposure was identified as a protective factor against home delivery. Increasing exposure to

media can alter perceptions positively and encourage expectant mothers to choose facility-based

deliveries over home deliveries, as public awareness and health-related behaviors are strongly

influenced by media exposure. Owing to a lack of knowledge, cultural customs, and restricted

access to medical facilities, home deliveries continue in many areas. To promote institutional

deliveries and enhance the health of mothers and newborns, increasing media exposure can be a

useful tactic [42,43]. In Nigeria, similar findings have been reported, where increased usage of

maternal healthcare services was linked to media exposure [44]. Expanding media-based health

awareness programs could be an effective strategy to reduce home delivery rates further.

Religious beliefs also play a role in home delivery choices. Our research suggests that Muslim

women exhibited a greater tendency for home delivery than their non-Muslim peers did. Cultural

norms, traditional beliefs, and socioeconomic factors may contribute to this trend [45]. A similar

pattern has been observed in India, where women of the Muslim faith exhibited a greater propensity

for home births [36]. However, further research is necessary to investigate the underlying causes of

this difference and to develop culturally relevant treatments that encourage institutional deliveries

within various religious communities.

The utilization of antenatal care (ANC) is a crucial factor influencing home delivery, as women

who visit four or more ANC appointments are much less inclined to deliver at home. This aligns

with findings from Afghanistan, where ANC visits were strongly associated with lower home

deliveries [46]. High-quality ANC services provide essential counseling, health screenings, and

education on the benefits of skilled birth attendance, which collectively influence women's

decisions to seek facility-based delivery over home delivery [47–51]. The consistent relationship

between ANC utilization and a lower likelihood of home delivery highlights the need to strengthen

ANC services and ensure equitable access for all pregnant women.

Parity, or the number of previous childbirths, also influenced home delivery rates. Home deliveries

were more common among women with three or more children. This finding is similar to those of

studies conducted in Bangladesh and Ethiopia, which reported that women who are first-time

mothers or have fewer pregnancies are more likely to choose facility-based births [30,37]. Likewise,

multiparous women in rural Zambia are more inclined to give birth at home [52]. Women with

multiple previous births may feel more confident about delivering at home, particularly if they have

had uncomplicated deliveries in the past. Furthermore, multiparous women may be more likely to

opt for delivery at home because of logistical difficulties, financial limitations, and a dependence

on traditional birth attendants. Addressing these barriers through community-based interventions

and financial support programs could help encourage facility-based deliveries among high-parity

women.

Our findings highlight a strong wealth disparity in home delivery rates. Women from poorer

households were significantly more likely to give birth at home than those from affluent households

were. Similar findings have been reported in several countries, including Nigeria, Nepal, and India

[53–56]. Financial constraints, out-of-pocket fees, and indirect costs such as transportation and lost

wages may deter women from seeking facility-based deliveries. Studies from Kenya and West

Africa similarly report persistent economic disparities in maternal healthcare utilization,

emphasizing the need for financial support mechanisms such as conditional cash transfers and

subsidies to promote institutional deliveries among economically disadvantaged groups [47,48,57].

Urban–rural differences in home delivery rates were also evident. Home deliveries were more

common among women in rural areas. According to some studies conducted in Africa and

Indonesia, urban women are significantly more likely than their rural counterparts are to give birth

in a medical facility [58–60]. Women living in rural areas confront unique obstacles to facility

delivery, such as a lack of qualified healthcare personnel, restricted availability of medical services,

and poor healthcare infrastructure [44,49,61,62]. Bridging the urban–rural gap in maternal

healthcare requires strategic investments in rural healthcare facilities, improved transportation

networks, and community-based healthcare interventions.

Bangladesh has implemented a range of interventions over the past decade-including the expansion

of community clinics, maternal health voucher schemes, and awareness campaigns-that have

contributed to the overall reduction in maternal delivery at home and increased skilled birth

attendance [63]. The Maternal Health Voucher Scheme (MHVS), launched in 2007, has been

particularly impactful in improving access to antenatal care, skilled assistance at delivery, and

postnatal care among disadvantaged women, though administrative and coverage challenges remain

[64,65]. Our study affirms the success of these efforts at the national level but also reveals persistent

gaps: maternal delivery at home remains disproportionately high among marginalized groups such

as women in the hill tracts, high-parity mothers, and the poorest households. Additionally, our

findings highlight that determinants of maternal delivery at home differ between urban and rural

settings, with factors like religion and household head's education being more influential in urban

areas, while wealth, parity, and ANC visits are more critical in rural regions. These context-specific

disparities suggest that, despite overall progress, current interventions have not fully addressed the

unique barriers faced by these populations. To further reduce maternal delivery at home,

particularly in high-prevalence areas, our study recommends mobile and culturally sensitive

outreach in remote districts, greater engagement of religious and community leaders,

incentivization of complete ANC utilization, and adaptation of policies to the distinct socio-

demographic realities of different regions.

Strengths and Limitations

428 A key strength of this study is its ability to provide valuable insights at both the national and

429 subnational levels using data from the MICS 2012–13 and 2019, as well as the BDHS 2022. Despite

430 the widespread use of spatial analysis in disease transmission research, few studies in Bangladesh

431 have examined spatial and temporal trends in maternal delivery at home. The use of the most recent

BDHS 2022 dataset, which employed rigorous sampling techniques, enhances the generalizability of the findings to women of reproductive age (15–49 years) across the country. Additionally, the large sample size and application of appropriate statistical techniques allowed for robust identification of significant predictors and assessment of inequality gradients across multiple subgroups. The findings can inform policies aimed at reducing maternal delivery at home clusters and identifying areas with limited access to facility-based childbirth services.

However, the study has certain limitations. First, as with most MICS and DHS-based studies, data are self-reported and thus subject to recall bias and social desirability bias, which may affect the accuracy of reported reproductive behaviors. Second, due to the cross-sectional nature of the survey, the study cannot establish causal relationships between maternal delivery at home and its determinants. Third, some relevant covariates—such as maternal comorbidities, nutritional status, dietary patterns, physical activity, and psychosocial factors—were not included in the BDHS and thus could not be examined, even though they may have significant implications for maternal delivery at home. Finally, socioeconomic characteristics were measured at the time of the interview, which may not accurately reflect the woman's status at the time of pregnancy or childbirth.

Conclusion

Over the last decade, Bangladesh has made great strides in lowering the country's home delivery rates. However, one-third of births still occur at home, highlighting ongoing challenges in ensuring universal access to institutional deliveries. Geographic, socioeconomic, and demographic disparities persist, underscoring the need for targeted interventions. Strengthening healthcare infrastructure, particularly in remote and hilly areas, improving maternal health education, and increasing ANC utilization can further reduce home deliveries. Additionally, addressing financial barriers through subsidies and incentive programs may encourage greater use of facility-based care.

Culturally sensitive approaches, media-based awareness campaigns, and community engagement

initiatives should be prioritized to achieve equitable maternal healthcare access. To develop

inclusive and successful policies, future studies should look at how religious, cultural, and

economic issues affect decisions about maternal delivery at home. Bangladesh may achieve

universal facility-based birthing by resolving these inequities, which would ultimately improve the

health of expectant mothers and newborns.

Declarations

Acknowledgements: We are grateful to the MICS and DHS team for allowing us to conduct the

analysis of this study using the MICS 2012 and 2019 data set and the BDHS 2022 data set.

Author Contributions: Conceptualization, M.L.K, and R.B.I; Data curation, R.B.I, and M.L.K;

Formal analysis, R.B.I, and M.L.K; Methodology, R.B.I, and S.T.A.N; Visualization, R.B.I, and

S.T.A.N; Investigation, S.M., and A.A; Supervision; Validation, M.B.L; Writing—original draft,

R.B.I, S.T.A.N and M.L.K; Writing - review and editing, S.M., A.A., and M.B.L; All authors have

read and agreed to the published version of the manuscript.

Funding: The authors received no specific grant from funding agencies in the public, commercial,

or not-for-profit sectors.

Consent for publication: Not applicable.

Competing interests: There are no potential conflicts (financial, professional, or personal) for any

of the authors to disclose.

Data availability: All data relevant to the study are included in the article or uploaded as

supplemental information. All relevant data are within the manuscript. Raw MICS

(https://mics.unicef.org/) and BDHS (https://dhsprogram. com/) data are publicly available that can

be accessed upon request.

References

1. Trends in maternal mortality 2000 to 2023: estimates by WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division. [cited 13 May 2025]. Available: https://www.who.int/publications/i/item/9789240108462

2. El-Saharty S, Ohno N. South Asia's quest for reduced maternal mortality: What the data show. In: World Bank Blog: Investing in Health. 2015.

3. Hossain AT, Siddique AB, Jabeen S, Khan S, Haider MM, Ameen S, et al. Maternal mortality in Bangladesh: Who, when, why, and where? A national survey-based analysis. J Glob Health. 2023;13. doi:10.7189/JOGH.13.07002

4. The DHS Program - Bangladesh: DHS, 2022 - Final Report (English). [cited 13 May 2025]. Available: https://www.dhsprogram.com/publications/publication-fr386-dhs-final-reports.cfm?cssearch=1551986_1

5. Document Viewer. [cited 13 May 2025]. Available: https://docs.un.org/en/A/RES/70/1

6. UNISEF. Maternal and newborn health. 2023.

7. Kitila SB, Feyissa GT, Olika AK, Wordofa MA. Maternal Healthcare in Low- and Middle-Income Countries: A Scoping Review. Health Serv Insights. 2022;15. doi:10.1177/11786329221100310

8. Oladipo IA, Akinwaare MO. Trends and patterns of maternal deaths from 2015 to 2019, associated factors and pregnancy outcomes in rural Lagos, Nigeria: a cross-sectional study. Pan African Medical Journal. 2023;44.

9. Hernández-Vásquez A, Chacón-Torrico H, Bendezu-Quispe G. Prevalence of home birth among 880,345 women in 67 low- and middle-income countries: A meta-analysis of Demographic and Health Surveys. SSM Popul Health. 2021;16. doi:10.1016/j.ssmph.2021.100955

10. Shrestha SK, Banu B, Khanom K, Ali L, Thapa N, Stray-Pedersen B, et al. Changing trends on the place of delivery: Why do Nepali women give birth at home? Reprod Health. 2012;9: 1–8. doi:10.1186/1742-4755-9-25

11. Sahoo J, Singh SV, Gupta VK, Garg S, Kishore J. Do socio-demographic factors still predict the choice of place of delivery: A cross-sectional study in rural North India. J Epidemiol Glob Health. 2015;5: S27–S34.

12. Shrestha SK, Banu B, Khanom K, Ali L, Thapa N, Stray-Pedersen B, et al. Changing trends on the place of delivery: Why do Nepali women give birth at home? Reprod Health. 2012;9: 1–8. doi:10.1186/1742-4755-9-25

13. Tiruneh SA, Lakew AM, Yigizaw ST, Sisay MM, Tessema ZT. Trends and determinants of home delivery in Ethiopia: Further multivariate decomposition analysis of 2005-2016 Ethiopian Demographic Health Surveys. BMJ Open. 2020;10: 1–10. doi:10.1136/bmjopen-2019-034786

14. Caulfield T, Onyo P, Byrne A, Nduba J, Nyagero J, Morgan A, et al. Factors influencing place of
delivery for pastoralist women in Kenya: A qualitative study. BMC Womens Health. 2016;16: 1–11.
doi:10.1186/s12905-016-0333-3

15. Sarker BK, Rahman M, Rahman T, Hossain J, Reichenbach L, Mitra DK. Reasons for preference of
home delivery with traditional birth attendants (TBAs) in Rural Bangladesh: A qualitative
exploration. PLoS One. 2016;11. doi:10.1371/journal.pone.0146161

16. Islam MM, Shahjahan M. Exploring the reasons and factors influencing the choice of home
delivery of births in rural Bangladesh: a community-based cross-sectional study. J Health Res.
2022;36. doi:10.1108/JHR-07-2020-0284

17. World Health Organization. WORLD HEALTH STATISTICS - MONITORING HEALTH FOR THE SDGs.
World Health Organization. 2016; 1.121.

18. Dey R, Dey SR, Haque M, Rahman AB, Kundu S, Setu SP, et al. Mapping the prevalence and
covariates associated with home delivery in Bangladesh: A multilevel regression analysis. PLoS
One. 2024;19: e0313606.

19. Talukder A, Anik BH, Hossain MI, Haq I, Habib MJ. Socioeconomic and demographic factors for
mothers' delivery at home: a comparative study among BDHS 2007, 2011 and 2014. Asian Journal
of Social Health and Behavior. 2022;5: 10–17.

20. Ahmed KT, Karimuzzaman M, Mahmud S, Rahman L, Hossain MM, Rahman A. Influencing factors
associated with maternal delivery at home in urban areas: a cross-sectional analysis of the
Bangladesh Demographic and Health Survey 2017–2018 data. J Health Popul Nutr. 2023;42: 83.

21. BANGLADESH 2012-13 MICS FINAL REPORT RELEASED | UNICEF MICS. [cited 13 May 2025].
Available: https://mics.unicef.org/news/bangladesh-2012-13-mics-final-report-released

22. Bangladesh 2019 MICS Report - English | PDF | Drinking Water | Sanitation. [cited 13 May 2025].
Available: https://www.scribd.com/document/537704949/Bangladesh-2019-MICS-Report-English

23. Pathey P. Bangladesh multiple indicator cluster survey 2012–2013 Key findings. Bangladesh Bur
Stat UNICEF Bangladesh. 2014;2014: 2014.

24. Noor STA, Shil P, Talukdar A, Aktar S, Uddin MJ. Exploring Factors Influencing Wealth-Related
Disparities in Institutional Delivery: A Decomposition Analysis Using Bangladesh Multiple Indicator
Cluster Survey (MICS) 2019. Public Health Challenges. 2025;4: e70066.

25. National Institute of Population Research and Training (NIPORT), ICF. Bangladesh Demographic
and Health Survey 2022 Final Report. 2024. Available:
https://www.dhsprogram.com/publications/publication-FR386-DHS-Final-Reports.cfm

26. Ahmed KT, Karimuzzaman Md, Mahmud S, Rahman L, Hossain MdM, Rahman A. Influencing
factors associated with maternal delivery at home in urban areas: a cross-sectional analysis of the
Bangladesh Demographic and Health Survey 2017–2018 data. J Health Popul Nutr. 2023;42: 83.
doi:10.1186/s41043-023-00428-9

27.   Enuameh YAK, Okawa S, Asante KP, Kikuchi K, Mahama E, Ansah E, et al. Factors influencing health
facility delivery in predominantly rural communities across the three ecological zones in Ghana: A
cross-sectional study. PLoS One. 2016;11: 1–16. doi:10.1371/journal.pone.0152235

28.   Dey D, Haque MdS, Islam MdM, Aishi UI, Shammy SS, Mayen MdSA, et al. The proper application
of logistic regression model in complex survey data: a systematic review. BMC Med Res Methodol.
2025;25: 15. doi:10.1186/s12874-024-02454-5

29.   von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the
Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for reporting
observational studies. International Journal of Surgery. 2014;12: 1495–1499.
doi:https://doi.org/10.1016/j.ijsu.2014.07.013

30.   Ahmed KT, Karimuzzaman Md, Mahmud S, Rahman L, Hossain MdM, Rahman A. Influencing
factors associated with maternal delivery at home in urban areas: a cross-sectional analysis of the
Bangladesh Demographic and Health Survey 2017–2018 data. J Health Popul Nutr. 2023;42: 83.
doi:10.1186/s41043-023-00428-9

31.   Huda TM, Chowdhury M, El Arifeen S, Dibley MJ. Individual and community level factors
associated with health facility delivery: A cross sectional multilevel analysis in Bangladesh. PLoS
One. 2019;14: e0211113.

32.   Badiuzzaman M, Murshed SM, Rieger M. Improving maternal health care in a post conflict setting:
evidence from Chittagong Hill tracts of Bangladesh. J Dev Stud. 2020;56: 384–400.

33.   Teshale AB, Alem AZ, Yeshaw Y, Kebede SA, Liyew AM, Tesema GA, et al. Exploring spatial
variations and factors associated with skilled birth attendant delivery in Ethiopia: geographically
weighted regression and multilevel analysis. BMC Public Health. 2020;20: 1–19.

34.   Moshi FV, Lymo G, Gibore NS, Kibusi SM. Prevalence and factors associated with home childbirth
with unskilled birth assistance in Dodoma-Tanzania: a cross sectional study. East Afr Health Res J.
2020;4: 92.

35.   Shahabuddin ASM, De Brouwere V, Adhikari R, Delamou A, Bardaj A, Delvaux T. Determinants of
institutional delivery among young married women in Nepal: Evidence from the Nepal
Demographic and Health Survey, 2011. BMJ Open. 2017;7: e012446.

36.   Sahoo J, Singh SV, Gupta VK, Garg S, Kishore J. Do socio-demographic factors still predict the
choice of place of delivery: A cross-sectional study in rural North India. J Epidemiol Glob Health.
2015;5. doi:10.1016/j.jegh.2015.05.002

37.   Tiruneh SA, Lakew AM, Yigizaw ST, Sisay MM, Tessema ZT. Trends and determinants of home
delivery in Ethiopia: Further multivariate decomposition analysis of 2005-2016 Ethiopian
Demographic Health Surveys. BMJ Open. 2020;10: 1–10. doi:10.1136/bmjopen-2019-034786

38.   Straneo M, Fogliati P, Azzimonti G, Mangi S, Kisika F. Where do the rural poor deliver when high
coverage of health facility delivery is achieved? Findings from a community and hospital survey in
Tanzania. PLoS One. 2014;9: 1–17. doi:10.1371/journal.pone.0113995

39. Kruk ME, Hermosilla S, Larson E, Vail D, Chen Q, Mazuguni F, et al. Who is left behind on the road
to universal facility delivery? A cross-sectional multilevel analysis in rural Tanzania. Tropical
Medicine and International Health. 2015;20: 1057–1066. doi:10.1111/tmi.12518

40. Ogundari K, Abdulai A. Determinants of household's education and healthcare spending in
Nigeria: Evidence from survey data. African Development Review. 2014;26: 1–14.

41. Rahman MM, Amin MT, Ferdous Z, Patwary H, Haider MM. Education of household head and
maternal healthcare utilization: the case of Bangladesh. BMC Public Health. 2024;24: 3439.

42. Mills S, Williams JE, Adjuik M, Hodgson A. Use of health professionals for delivery following the
availability of free obstetric care in Northern Ghana. Matern Child Health J. 2008;12: 509–518.
doi:10.1007/s10995-007-0288-y

43. Agha S, Carton TW. Determinants of institutional delivery in rural Jhang, Pakistan. Int J Equity
Health. 2011;10: 1–12. doi:10.1186/1475-9276-10-31

44. Babalola S, Fatusi A. Determinants of use of maternal health services in Nigeria - Looking beyond
individual and household factors. BMC Pregnancy Childbirth. 2009;9: 43. doi:10.1186/1471-2393-
9-43

45. Kamal SMM, Hassan CH, Alam GM. Determinants of institutional delivery among women in
Bangladesh. Asia Pac J Public Health. 2015;27: NP1372–NP1388. doi:10.1177/1010539513486178

46. Rahman M, Saha P, Uddin J. Associations of antenatal care visit with utilization of institutional
delivery care services in Afghanistan: intersections of education, wealth, and household decision-
making autonomy. BMC Pregnancy Childbirth. 2022;22: 255.

47. Colombara D V., Hernández B, Schaefer A, Zyznieuski N, Bryant MF, Desai SS, et al. Institutional
delivery and satisfaction among indigenous and poor women in Guatemala, Mexico, and Panama.
PLoS One. 2016;11: 1–17. doi:10.1371/journal.pone.0154388

48. Tekelab T, Yadecha B, Melka AS. Antenatal care and women's decision making power as
determinants of institutional delivery in rural area of Western Ethiopia. BMC Res Notes. 2015;8:
1–8. doi:10.1186/s13104-015-1708-5

49. Séraphin MN, Ngnie-Teta I, Ayoya MA, Khan MR, Striley CW, Boldon E, et al. Determinants of
Institutional Delivery Among Women of Childbearing Age in Rural Haiti. Matern Child Health J.
2015;19: 1400–1407. doi:10.1007/s10995-014-1646-1

50. Agha S, Williams E. Quality of antenatal care and household wealth as determinants of
institutional delivery in Pakistan: Results of a cross-sectional household survey. Reprod Health.
2016;13: 1–8. doi:10.1186/s12978-016-0201-5

51. Feyissa TR, Genemo GA. Determinants of institutional delivery among childbearing age women in
Western Ethiopia, 2013: Unmatched case control study. PLoS One. 2014;9: 1–7.
doi:10.1371/journal.pone.0097194

52.   Scott NA, Henry EG, Kaiser JL, Mataka K, Rockers PC, Fong RM, et al. Factors affecting home
delivery among women living in remote areas of rural Zambia: a cross-sectional, mixed-methods
analysis. Int J Womens Health. 2018; 589–601.

53.   Newell R, Spillman I, Newell M-L. The use of facilities for labor and delivery: the views of women in
rural Uganda. J Public Health Afr. 2017;8: 592.

54.   Olorunsaiye CZ, Huber LB, Laditka SB, Kulkarni S, Boyd AS. Factors associated with health facility
delivery in West and Central Africa: a multilevel analysis of seven countries. Health Care Women
Int. 2020;41: 3–21.

55.   Singh PK, Rai RK, Singh L. Examining the effect of household wealth and migration status on safe
delivery care in urban India, 1992–2006. 2012.

56.   Devkota B, Maskey J, Pandey AR, Karki D, Godwin P, Gartoulla P, et al. Determinants of home
delivery in Nepal–A disaggregated analysis of marginalised and non-marginalised women from the
2016 Nepal Demographic and Health Survey. PLoS One. 2020;15: e0228440.

57.   Kinuthia J, Kohler P, Okanda J, Otieno G, Odhiambo F, John-Stewart G. A community-based
assessment of correlates of facility delivery among HIV-infected women in western Kenya. BMC
Pregnancy Childbirth. 2015;15: 1–9. doi:10.1186/s12884-015-0467-6

58.   Idris H, Budiastuti A, Razak R, Hasyim H. Delivery Services Utilization Based on Urban Rural Status
in Indonesia. Open Access Maced J Med Sci. 2022;10: 1147–1152.

59.   Newell R, Spillman I, Newell M-L. The use of facilities for labor and delivery: the views of women in
rural Uganda. J Public Health Afr. 2017;8: 592.

60.   Moyer CA, Mustafa A. Drivers and deterrents of facility delivery in sub-Saharan Africa: a
systematic review. Reprod Health. 2013;10: 1–14.

61.   Dahiru T, Oche OM. Determinants of antenatal care, institutional delivery and postnatal care
services utilization in Nigeria. Pan African Medical Journal. 2015;21: 1–17.
doi:10.11604/pamj.2015.21.321.6527

62.   Amano A, Gebeyehu A, Birhanu Z. Institutional delivery service utilization in Munisa Woreda,
South East Ethiopia: a community based cross-sectional study. BMC Pregnancy Childbirth.
2012;12: 1–6. doi:10.1186/1471-2393-12-105

63.   Mahmood SS, Amos M, Hoque S, Mia MN, Chowdhury AH, Hanifi SMA, et al. Does healthcare
voucher provision improve utilisation in the continuum of maternal care for poor pregnant
women? Experience from Bangladesh. Glob Health Action. 2019;12.
doi:10.1080/16549716.2019.1701324,

64.   Chowdhury AH, Hanifi SMA, Iqbal M, Hossain A, Stones W, Amos M, et al. Does maternal health
voucher scheme have association with distance inequality in maternal and newborn care
utilization? Evidence from rural Bangladesh. PLoS One. 2023;18: e0295306.

65.   Talukder MN, Rob U, Musa SAJM, Bajracharya A, Keya KT, Noor FR, et al. Evaluation of the impact
of the voucher program for improving maternal health behavior and status in Bangladesh. 2014.

Figure_1

Figure_2

Figure_3