Reduced risk of household secondary invasive Group A Streptococcal infections after a prophylaxis policy change, the Netherlands, 2022-2024

Brechje de Gier[1], Bart J.M. Vlaminckx[2], Annika van Roon[1], Mirjam J. Knol[1], Margreet J.M. te Wierik[1], Daan W. Notermans[1], Hester E. de Melker[1], Nina M. van Sorge[3,4]

1.Center for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands.
2. Department of Medical Microbiology, University Medical Center Utrecht, Utrecht, the Netherlands.
3. Department of Medical Microbiology and Infection Prevention, Amsterdam University Medical Center location AMC, University of Amsterdam, Amsterdam Institute for Infection and Immunity, Amsterdam, the Netherlands.
4. Netherlands Reference Laboratory for Bacterial Meningitis, Amsterdam University Medical Center location AMC, Amsterdam, the Netherlands.

Abstract

Importance: Household contacts of patients with invasive group A streptococcal (iGAS) disease have an increased risk of developing iGAS. In the Netherlands, the iGAS public health policy was changed in January 2023, offering antibiotic prophylaxis to household contacts of all iGAS patients rather than only those presenting with necrotising fasciitis or streptococcal toxic shock syndrome.

Objective: To estimate risk of iGAS in the general population and among household- and other contacts of primary iGAS patients, before and after the policy change.

Design: A nationwide cohort approach was used, linking population registry data with iGAS laboratory patient data.

Setting: Population-based.

Participants: All persons included in the Dutch population registry during the study period from 1 April 2022 to 31 December 2024. The iGAS case definition was an iGAS isolate submitted to the Netherlands Reference Laboratory for Bacterial Meningitis, with disease onset in the study period.

Exposure: For contacts of primary iGAS patients, exposure risk period was defined as the 30 days after culture date of the index patient. Exposure under the updated policy was defined as all person-time after 20 January 2023.

Main Outcomes and Measures: We estimated the incidence rate ratio (IRR) of iGAS during the 30-day risk period compared to unexposed person-time. Secondary attack rates among household contacts were estimated, calculating an odds ratio (OR) to compare attack rates before and after the policy change. Estimates were adjusted for age group, sex, household socioeconomic status and year-quarter.

Results: A total of 3,630 iGAS isolates were linked to the population registry, of which 14 were household secondary cases. Secondary attack rate among household contacts was 0.219% (n=7) before, and 0.047% (n=7) after the policy change, adjusted OR 0.17 (95%CI 0.05-0.53). The IRR for household contacts during the risk period was 235 (95%CI 94-587) before, and 74 (95%CI 35-156) after the policy change, compared to unexposed person-time. Among non-household relatives, colleagues, classmates and neighbours of primary iGAS cases, the number of secondary cases was too low to allow reliable estimates (<5).

Conclusions and relevance: We observed a reduction in secondary iGAS risk among household contacts after implementing an expanded antibiotic prophylaxis policy.

Introduction

Streptococcus pyogenes or Group A streptococcus (GAS) is a human-specific Gram-positive bacterium, that can be carried asymptomatically but also causes a large burden of non-invasive and invasive disease. Known risk factors for invasive group A streptococcus (iGAS) infections include young (0-5 years) or old (60+ years) age, having underlying medical conditions, the peripartum period, and co-infections such as influenza or chickenpox [1]. Although iGAS is uncommon, a surge of iGAS has been observed globally after lifting all societal restrictions related to the COVID-19 pandemic. iGAS has a high case fatality, estimated between 8 and 16% [1]. Close contacts of an iGAS patient have a greatly increased risk of developing iGAS, with estimates ranging between 800 to over 5000 per 100,000 person-years at risk, compared to background incidences of 2-4 per 100,000 person-years [1-3]. In the absence of effective vaccines, the only available public health measure to prevent iGAS is antibiotic prophylaxis for individuals after potential high-risk GAS exposure. There is currently no clear consensus how to most effectively prevent secondary iGAS.

In 2008, three clinical presentations of iGAS became notifiable by law in the Netherlands: streptococcal toxic shock syndrome (STSS), necrotising fasciitis and puerperal sepsis. Household contacts of patients with STSS or necrotising fasciitis were offered antibiotic prophylaxis to prevent secondary iGAS. In the Netherlands, a more than twofold increase in the annual number of notifiable iGAS infections was observed in 2022 compared to the annual average in the pre-COVID-19 pandemic years, leading to an estimated incidence of 10 per 100,000 per year [4, 5]. We have observed an absolute and relative increase of the globally disseminated *emm*1 clade M1$_{UK}$, as well as the emergence of new variants of concern such as *emm*1.134 in 2022-2023, followed by an expansion of a previously rare *emm* type 3.93 [5, 6]. A case-only analysis of the Dutch data showed that these three *emm* types more often clustered within households, suggesting an increased secondary attack rate [unpublished data]. These observations prompted a change in the Dutch public health policy regarding antibiotic prophylaxis in January 2023 where household contacts of all iGAS patients were eligible for antibiotic prophylaxis to prevent secondary iGAS [7]. To implement this policy, all iGAS infections (i.e. all iGAS compatible disease presentations with *S. pyogenes* detected in a normally sterile site) became notifiable by law. It was decided that a single daily dose of 500 mg azithromycin for three days was the preferred prophylaxis regimen to optimize acceptability and adherence. However, as macrolide resistance is not rare among *S. pyogenes* isolates [4], the prophylaxis can be changed based on the antibiotic susceptibility profile of the primary case. Alternative prophylactic regimes include either a 10-day course of clindamycin or oral penicillin in combination with four days of rifampicin. Both household- and other close contacts are informed about their increased risk and are advised to seek medical care without delay in case of any symptoms consistent with (i)GAS infection, to enable early treatment.

The main aim of this retrospective population-based cohort study was to assess the impact of the policy change on the risk of secondary iGAS among household- and other close contacts. We further describe the population incidence and risk factors for iGAS in the Netherlands in the post-pandemic years 2022-2024.

Methods

Data sources

The Netherlands Reference Laboratory for Bacterial Meningitis (NRLBM) has collected *S. pyogenes* isolates for nationwide bacteriological surveillance of iGAS since April 2022. All medical microbiological laboratories in the Netherlands are requested to submit iGAS isolates to the NRLBM, when cultured from a normally sterile site or from a non-sterile site in combination with a clinical presentation of iGAS. All iGAS isolates are *emm* (sub)typed according to the CDC protocol through PCR amplification and subsequent Sanger sequencing of the 180-bp domain that contains the 50

hypervariable codons [5]. Submitted isolates are accompanied by limited information regarding patient characteristics, including sex, birth date and postal code. Through the Statistics Netherlands (CBS) microdata system, isolates were linked to population registry data.

Data on birth dates and sex were retrieved from the population registry. A composite score of socioeconomic status, based on income, wealth and occupation status, was available at the household level. Household compositions, with start and end dates, were used to identify household contacts and socioeconomic status per participant. Non-household relatives, colleagues, classmates and neighbors were identified through the person-network datasets provided by Statistics Netherlands. As these datasets were available for the period up to and including 2022, we assumed the contacts in 2022 were still their contacts in 2024. For children born since 2022 and therefore not included in the person-network datasets, we assumed the non-household relatives of their household contacts were also the non-household relatives of the children. For clusters of co-primary cases, i.e. contacts with the same index date, the primary and secondary case labels were assigned at random and one of the two was included as outcome event in the main analysis.

Analytical methods

An open cohort approach was used, where persons were included from 1 April 2022 or their birth date (whichever came last) and followed until 31 December 2024, their first iGAS episode, or death, whichever came first. When multiple isolates were received from the same person within 30 days, these were considered as belonging to the same episode. As primary outcome, the first iGAS episode per person was used. Persons with multiple iGAS episodes could however contribute to multiple exposure periods for their contacts. The primary exposure of interest was being a contact of an iGAS patient, during the 30-day risk period after the primary patient's index date. The risk period was truncated in case of an event, therefore only the days leading up to the event of the contact are included in the person-time denominators. For secondary attack rate estimates, the number of contacts was the denominator.

Negative binomial regression was used to estimate IRR of contacts during the risk period compared to the population background incidence. IRRs were compared before and after the policy change, and an interaction term between period and the exposure was used to test for significance. Logistic regression was used to compare secondary attack rates among household contacts before and after the guideline change. All models were adjusted for age group, sex, calendar time (in year-quarters), and household socioeconomic status (SES) in quintiles (where quintile 1 represents the lowest SES).

Results are based on calculations by the RIVM, in project number 9892, using non-public microdata from Statistics Netherlands. As per the Statistics Netherlands policy, no numbers below 5 or estimates based on fewer than 5 events could be reported.

Ethics

The RIVM Centre for Clinical Expertise verified whether this study complies with the Dutch law for Medical Research Involving Human Subjects (WMO) or with the EU Clinical Trial Directive (2001/20/EC), and was of the opinion that review by an ethical research committee or institutional review board is not necessary by current national and European legislation (study number EPI-692).

Results

Figure 1 shows the number of *S. pyogenes* isolates from iGAS patients submitted to the NRLBM during the study period by *emm* type. During the 2022-2023 season, *emm*1.0 dominated, while *emm*3.93 clearly dominated during the 2023-2024 season, as described previously [5, 6]. A total of 4,215 iGAS isolate records were submitted to Statistics Netherlands, of which 3,729 (88.5%) could be deterministically linked to unique persons in the population registry (Supplementary Figure 1). Linkage was significantly more often possible for ages 46 years and older, females, and isolates of *emm*4.0, but did not differ by policy period (Supplementary Table 1). Eighteen isolates were excluded since they were collected before the start of the study period, and a further 67 isolates were excluded when de-duplicating isolates from the same person within 30 days. After exclusions, 3,644 iGAS isolates from 3,630 unique persons were included in the analysis.

Figure 1. Number of *S. pyogenes* isolates received and typed by the Netherlands Reference Laboratory for Bacterial Meningitis during the study period 1 April 2022-31 December 2024, by month and *emm* type (n=4,129).

Over the complete study period, 3,609/3,630 (99.4%) were primary cases, i.e. cases not occurring during the risk period of a known contact (Table 1), corresponding with a population background incidence of 7.07 per 100,000 person-years. The incidence peaked in the first quarter of 2023 at 14.15 per 100,000 person-years (Table 1). Age groups 0-5 and 66+ years had a significantly higher incidence compared to age group 45-65 years. Persons registered as male had a significantly lower incidence compared to persons of female or unknown sex. This is likely related to the risk of puerperal fever or -sepsis for women of childbearing age (see Supplementary Figure 1). Only the fourth quintile (second-to highest) of household SES had a significantly higher incidence compared to the first quintile. Incidences by year-quarter reflect the winter seasonality of iGAS.

In total, 14 secondary iGAS cases were identified among household contacts of primary iGAS patients during the 30-day risk period, corresponding to an incidence of 965 per 100,000 person-

years. The adjusted incidence rate ratio (IRR) for household contacts in the 30-day risk period compared to the population incidence was 100 (95%CI 55-181) over the entire study period. For non-household relatives, colleagues and classmates, and neighbors, the number of secondary cases was too low (<5) to present estimates according to the Statistics Netherlands policy.

Overall 30-day mortality was 10.2%, and significantly lower for cases with *emm* type 4.0 compared to *emm* type 1.0 (Table 2). Mortality was significantly lower for ages 0-45 years compared to 46-65 years, and significantly higher for 66+. No significant difference in mortality was observed between sexes, socioeconomic quintiles or year-quarters (Table 2).

Table 1. Number of person-years, iGAS events, incidence of iGAS per 100,000 person-years and by exposures of interest, 1 April 2022-31 December 2024, the Netherlands. Incidence rate ratios (IRR) with 95% confidence interval (CI), adjusted for age group, sex, household socioeconomic quintile and year-quarter.

		Person-years	iGAS events	Incidence per 100,000 person-years	IRR (95% CI)
Exposure	None	51,033,229	3,609	7.07	[ref]
	Household	1,451	14	965	**100 (55-181)**
	Non-household relatives	7,852	<5		
	Colleagues and classmates	11,178	<5		
	Neighbors	13,762	<5		
Age group (years)	0-5	3,023,746	355	11.74	**1.97 (1.70-2.28)**
	6-19	7,769,181	226	2.91	**0.49 (0.41-0.58)**
	20-45	17,488,349	1,127	6.44	1.03 (0.92-1.16)
	46-65	13,381,688	792	5.92	[ref]
	66+	9,404,508	1,130	12.02	**2.05 (1.82-2.32)**
Sex	Female or unknown	25,634,925	1,952	7.61	[ref]
	Male	25,432,548	1,678	6.60	**0.88 (0.81-0.96)**
Household socioeconomic status quintile	1 (lowest)	6,009,519	406	6.76	[ref]
	2	7,342,759	603	8.21	1.09 (0.94-1.26)
	3	8,947,470	691	7.72	1.07 (0.93-1.24)
	4	12,422,833	900	7.24	**1.16 (1.01-1.33)**
	5	14,961,044	979	6.54	1.13 (0.98-1.29)
	unknown	1,383,848	51	3.69	
Year-quarter	2022-Q2	4,670,115	278	5.95	[ref]
	2022-Q3	4,671,332	279	5.97	0.99 (0.81-1.21)
	2022-Q4	5,635,833	524	9.30	**1.49 (1.25-1.79)**
	2023-Q1	3,604,976	510	14.15	**2.15 (1.79-2.58)**
	2023-Q2	4,618,814	440	9.53	**1.47 (1.22-1.77)**

	2023-Q3	4,670,822	236	5.05	0.83 (0.67-1.02)
	2023-Q4	4,671,239	248	5.31	0.87 (0.71-1.07)
	2024-Q1	4,618,659	556	12.04	**1.87 (1.56-2.24)**
	2024-Q2	4,617,783	316	6.84	1.06 (0.87-1.29)
	2024-Q3	4,669,218	153	3.28	**0.53 (0.42-0.66)**
	2024-Q4	4,618,682	90	1.95	**0.32 (0.25-0.42)**

Table 2. 30-day mortality of iGAS cases by emm type and odds ratios (OR) with 95% confidence interval (CI), adjusted for agegroup, sex, household socioeconomic quintile and year-quarter.

		iGAS cases	Died within 30 days	Mortality percentage	OR (95% CI)
	Overall	3,630	369	10.2	
emm type	1.0	1,030	128	12.4	[ref]
	12.0	192	25	13	0.96 (0.56-1.60)
	22.0	266	21	7.9	0.73 (0.42-1.21)
	3.93	523	72	13.8	1.05 (0.66-1.67)
	4.0	362	17	4.7	**0.47 (0.26-0.78)**
	87.0	173	12	6.9	0.61 (0.30-1.16)
	89.0	187	18	9.6	0.74 (0.40-1.31)
	Other *emm* types	897	76	8.5	**0.65 (0.46-0.92)**
Age group	0-5	355	16	4.5	**0.48 (0.26-0.83)**
	6-19	226	7	3.1	**0.33 (0.14-0.68)**
	20-45	1,127	27	2.4	**0.29 (0.18-0.45)**
	46-65	792	71	9	[ref]
	66+	1,130	248	21.9	**2.64 (1.98-3.58)**
Sex	Female or unknown	1,952	165	8.5	[ref]
	Male	1,678	204	12.2	1.16 (0.92-1.46)
Household socioeconomic status	1 (lowest)	406	45	11.1	[ref]
	2	603	104	17.2	1.44 (0.98-2.15)
	3	691	82	11.9	1.02 (0.69-1.54)
	4	900	83	9.2	1.06 (0.71-1.60)
	5	979	49	5	0.89 (0.56-1.39)
	unknown	51	6	11.8	
Year-quarter	2022-Q2	278	20	7.2	[ref]
	2022-Q3	279	30	10.8	1.60 (0.86-3.02)
	2022-Q4	524	66	12.6	1.53 (0.88-2.76)
	2023-Q1	510	51	10	1.16 (0.65-2.12)
	2023-Q2	440	36	8.2	0.97 (0.53-1.81)
	2023-Q3	236	17	7.2	1.15 (0.56-2.36)
	2023-Q4	248	23	9.3	1.24 (0.63-2.47)
	2024-Q1	556	70	12.6	1.44 (0.78-2.72)

	2024-Q2	316	36	11.4	1.21 (0.63-2.35)
	2024-Q3	153	13	8.5	1.10 (0.49-2.39)
	2024-Q4	90	7	7.8	1.03 (0.37-2.57)

Impact of policy change

In the period before the policy change (1 April 2022- 19 January 2023), 7 secondary cases were identified among 3,195 household contacts, resulting in an IRR of 235 compared to the population incidence (95%CI 94-587; Table 2). After the policy change on 20 January 2023, there was a significant reduction in the IRR to 74 (95%CI 35-156; p for interaction 0.016), also based on 7 secondary cases among 14,974 household contacts. The population background incidence and the median number of household contacts per primary case did not differ between the two periods. Secondary attack rates (SAR) among household contacts were 0.219% before, and 0.047% after the guideline change (aOR 0.17, 95%CI 0.05-0.53).

Table 3. Secondary iGAS in households, stratified by periods with the old and the new policy in place. [1] Incidence rate ratio (IRR) adjusted for age group, gender, household socioeconomic quintile and year-quarter.

	Old policy (1 April 2022- 19 January 2023)	New policy (20 Jan 2023- 31 Dec 2024)
Primary iGAS cases in multi-person households	927	2,213
Household contacts	3,195	14,974
Median household contacts per case (IQR)	2 (1-3)	2 (1-3)
Secondary iGAS cases among household contacts	7	7
Secondary attack rate	0.219%	0.047%
Person-years household contacts	234	1217
Incidence among household contacts per 100,000 person-years	2995	575
Background population incidence per 100,000 person-years	7.17	7.03
IRR compared to unexposed [1]	235 (95% CI 94-587)	74 (95% CI 35-156)

Discussion

We observed a significant reduction in risk of secondary iGAS for household contacts after a policy change, which extended antibiotic prophylaxis eligibility to household contacts of all iGAS patients rather than only household contacts of primary cases with STSS or necrotizing fasciitis. This reduction was visible both as a 4.5-fold reduction in SAR, as well as a three-fold difference in IRR, where the person-time denominator was truncated in case of an event. Data was not available on the actual prescription or uptake of antibiotic prophylaxis, therefore we could not estimate direct effectiveness of the prophylaxis against secondary iGAS.

Based on a 2023 study in the Netherlands, we know that there was a high overall acceptance of antibiotic prophylaxis with an uptake of 95% among eligible contacts. Of these contacts, 4.4% developed symptoms consistent with mild GAS infection, but no secondary iGAS was observed [8]. A recent study by Birck *et al* linked prophylaxis prescription data, and found similar GAS infection rates among family contacts of primary iGAS cases who did and did not receive antibiotic prophylaxis [9].

According to a recent systematic review, there is one previous study reporting GAS risk reduction after antibiotic prophylaxis, but this was during an outbreak among persons experiencing homelessness. Since there is currently no general consensus as how to best prevent secondary iGAS in the general population, we believe our study contributes valuable insights on the benefits of antibiotic prophylaxis to prevent iGAS in high-exposure settings.

Adebanjo *et al* in 2020 summarized the available evidence on iGAS incidence among household members of primary iGAS cases [2]. The two largest previous studies found 1,240 and 4,520 iGAS cases per 100,000 person-years in the 30-day risk period among close contacts [2, 3]. Our incidence estimates for household contacts in this risk period, of 2,995 before and 575 per 100,000 person-years after the policy change, are in line with this order of magnitude.

After the COVID-19 pandemic, many countries reported an increase in iGAS incidence in 2022-2023, including the Netherlands [4]. We here report a population iGAS incidence of 7.07 per 100,000 person-years during 2022-2024, peaking in early 2023 at 14.15, which is roughly in line with published population incidences covering this period [10, 11]. A Danish study reported an incidence of 6.6 per 100,000 in children 0-17 years of age in 2022-2023, with the majority under the age of 5, consistent with the incidence in children in our study (Table 1) [12].

Surprisingly, we did not observe an association between iGAS risk and low household SES. Concordantly, 30-day mortality did not differ significantly by SES. Deprivation has been repeatedly shown to be an important risk factor for iGAS [1, 10, 13, 14]. Likely, our categorization of SES is not sensitive enough to capture the vulnerable populations most at risk for iGAS, such as people who inject drugs and people experiencing homelessness.

Our study has limitations, mainly affecting the estimated impact of the policy change on secondary iGAS incidence. The patient data submitted with the iGAS isolate did not allow us to distinguish between iGAS disease presentations for which prophylaxis was already indicated in 2022 (i.e. STSS and necrotizing fasciitis) versus newly notifiable iGAS. Especially for clusters with isolates cultured just around the date of the policy change, it is not clear if the contacts were already offered prophylaxis. One cluster had an index date before the policy change but secondary iGAS after. We decided to categorize this as a cluster under the old policy. Due to the low number of secondary cases, this decision impacted our results, increasing the SAR under the old policy. Also, the proportion of secondary cases occurring too soon after the primary case to receive prophylaxis may have differed between the two periods, which impacts the preventable fraction of secondary iGAS. The low number of (secondary) cases limited the power of our study, and precluded analyses among other types of close contacts such as non-household family members. Further, as dominant *emm* types also differed between the periods, it is not possible to fully disentangle effects of the policy change and differences in transmissibility or virulence of circulating *emm* types, despite adjusting estimates for calendar time. Lastly, the fact that 11.5% of submitted isolates could not be deterministically linked to a unique person in the population registry presents an important limitation, resulting in a lower incidence estimate. As linkage was based on postal code, birth date and sex, it is possible that twins and persons in large institutional households are overrepresented in the non-linked isolates. This might have resulted in an underestimation of the number of secondary household cases for both periods. While the median and IQR of household contacts per case were identical between the periods, the averages are very different, indicating a skewedness due to large institutional households included in the period with the new policy. Despite the adjustment for age group, this may have resulted in a more fragile population at risk included in the new policy period. On the other hand, the closeness of the contacts is likely to be less in such institutional households.

In conclusion, our results show a substantial reduction in the risk of secondary iGAS among household contacts after the Dutch iGAS antibiotic prophylaxis policy change. However, the overwhelming majority of iGAS occurs as sporadic cases. Therefore, while antibiotic prophylaxis likely prevents a significant disease burden in contacts at high risk, the impact on overall iGAS incidence can only be very modest.

References

1.	Watts, V., et al., *Antibiotic chemoprophylaxis for close contacts of invasive group A streptococcus in community settings: Evidence review.* J Infect, 2025. **90**(4): p. 106468.
2.	Adebanjo, T., et al., *Evaluating Household Transmission of Invasive Group A Streptococcus Disease in the United States Using Population-based Surveillance Data, 2013-2016.* Clin Infect Dis, 2020. **70**(7): p. 1478-1481.
3.	Mearkle, R., et al., *Household transmission of invasive group A Streptococcus infections in England: a population-based study, 2009, 2011 to 2013.* Euro Surveill, 2017. **22**(19).
4.	de Gier, B., et al., *Increase in invasive group A streptococcal (Streptococcus pyogenes) infections (iGAS) in young children in the Netherlands, 2022.* Euro Surveill, 2023. **28**(1).
5.	Rumke, L.W., et al., *Nationwide upsurge in invasive disease in the context of longitudinal surveillance of carriage and invasive Streptococcus pyogenes 2009-2023, the Netherlands: a molecular epidemiological study.* J Clin Microbiol, 2024. **62**(10): p. e0076624.
6.	Davies, M.A., et al., *Streptococcus pyogenes emm Type 3.93 Emergence, the Netherlands and England.* Emerg Infect Dis, 2025. **31**(2): p. 229-236.
7.	*RIVM - LCI-richtlijn Groep A-streptokokkeninfectie*. Available from: https://lci.rivm.nl/richtlijnen/groep-streptokokkeninfectie.
8.	Mujakovic, S., H.L.G. Ter Waarbeek, and J. Van den Boogaard, *Attitude towards antibiotic prophylaxis among household contacts of patients with invasive Group A Streptococcal disease and its impact on contacts' iGAS incidence in the Netherlands, February-June 2023*

in *ESCAIDE*. 2024: Stockholm.
9.	Birck, M.G., et al., *Choosing Wisely: Effectiveness and Safety of Antibiotic Prophylaxis in Close Contacts of Invasive Group A Streptococci Infection.* Clin Infect Dis, 2025. **80**(2): p. 314-315.
10.	Gregory, C.J., et al., *Invasive Group A Streptococcal Infections in 10 US States.* JAMA, 2025.
11.	Valcarcel Salamanca, B., et al., *Increase in invasive group A streptococcal infections (iGAS) in children and older adults, Norway, 2022 to 2024.* Euro Surveill, 2024. **29**(20).
12.	Nygaard, U., et al., *Invasive group A streptococcal infections in children and adolescents in Denmark during 2022-23 compared with 2016-17 to 2021-22: a nationwide, multicentre, population-based cohort study.* Lancet Child Adolesc Health, 2024. **8**(2): p. 112-121.
13.	Ammar, S., et al., *Post-pandemic increase in invasive group A strep infections in New Zealand.* J Infect Public Health, 2024. **17**(11): p. 102545.
14.	Zangarini, L., et al., *Incidence and clinical and microbiological features of invasive and probable invasive streptococcal group A infections in children and adults in the Brussels-Capital Region, 2005-2020.* Eur J Clin Microbiol Infect Dis, 2023. **42**(5): p. 555-567.