Cortical neural state topography reveals mesoscale heterogeneity in autism

Daniele Marcotulli[1,2,3]*, Valentina Francesca Cudia[1], Martina Vacchetti[3], Luca Berta[4], Samuela Morano[3], Marta Giacobbi[3], Carlotta Canavese[3], Barbara Svevi[3], Federica Silvia Ricci[1,3], Federico Amianto[3,5], Benedetto Vitiello[1,6], Andrea Martinuzzi[1,3], Tobias Banaschewski[7], Argyris Stringaris[2], Chiara Davico[1,3]

1. Department of Sciences of Public Health and Pediatrics, University of Turin, Turin, Italy.
2. Divisions of Psychiatry and Psychology and Language Science, University College London, London, United Kingdom.
3. Department of Child Care, Regina Margherita Hospital, Turin, Italy.
4. Department of Medical Physics, AOU Città della Salute e della Scienza, Turin, Italy.
5. Department of Neuroscience, University of Turin, Turin, Italy.
6. Department of Mental Health, School of Public Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA.
7. Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany.

* Corresponding author: daniele.marcotulli@unito.it

Abstract

Characterizing macroscopic brain organization in neuropsychiatric conditions relies on averaging neural activity within discrete regions, yet this approach collapses spatial information that likely carries distinct biological meaning. Here, we introduce a spatial autocorrelation framework to quantify the continuous topographical organization of local states and demonstrate its utility in autism spectrum conditions (ASC). Mapping the spatial heterogeneity of an electroencephalographic (EEG) excitability marker —the aperiodic exponent — across three independent datasets (n = 3767), we show that the autistic cortex exhibits a more heterogeneous topography, as preregistered. This pattern was specific to the mesoscale (~6–9 cm), replicated across cohorts, and persisted across wakefulness and sleep. This spatial metric outperformed both the global mean and regional variability in predicting ASC status, indicating that topographical arrangement captures biological variance not recovered by conventional approaches. Structural MRI analysis (n = 1198) revealed that local macroanatomy mirrors this functional heterogeneity, with stronger structure–function coupling in ASC, suggesting an anatomical basis for the observed topographical differences. By recovering spatial information typically collapsed through averaging, this framework provides a complementary axis for characterizing macroscopic brain organization across neuropsychiatric conditions.

Introduction

In any complex biological system, from the synchronization of cardiac cells to the collective motion of animal groups, the spatial arrangement of its constituents determines the system's dynamic properties. In the human cortex, the smoothness of the excitability landscape, how local neural states are arranged across the cortical surface, regulates how signals propagate across space[1–4]. Here, we test the hypothesis that Autism Spectrum Conditions (ASC), a common neurodevelopmental condition[5], are characterized by a distinct topological organization, specifically, increased heterogeneity in neural excitability[6].

Despite its importance for individuals and society, the neurobiological basis of ASC remains elusive[6, 7]. In the search for a neurophysiological mechanism, research has traditionally been divided into two dominant but often disconnected frameworks: the excitation/inhibition (E/I) imbalance hypothesis and the different brain connectivity hypothesis[8–11], both of which have received substantial independent support but lack integration and failed to provide reliable biomarkers[12].

The E/I hypothesis suggests that ASC results from altered E/I balance, yet findings remain heterogeneous, likely because studies relied on global averages or isolated regions, obscuring spatial topography[2,13–16]. Similarly, evidence for altered connectivity, such as the "local hyperconnectivity with global hypoconnectivity" hypothesis[12,17,18] remains vast yet inconsistent.

Here, we argue that these traditionally disconnected frameworks might conceptually converge at the mesoscale, the finest spatial resolution reliably accessible via non-invasive scalp electrophysiology. If local E/I states are heterogeneously distributed across the cortical surface, the resulting spatial gradients could drive a divergent organization of global network

dynamics, potentially accounting for findings from both theories within a single spatial

mechanism[13,19,20].

Specifically, we hypothesized that the ASC cortex is characterized by "patches" of relatively

distinct excitability states. These patches could arise from region-specific developmental

divergences or genetic determinants, reinforcing local connectivity and manifesting as the

enhanced local feature focus frequently observed in ASC[11, 17].

To test this, we quantified the spatial autocorrelation of the aperiodic exponent (slope). While

influenced by additional factors including neuromodulatory tone, the aperiodic exponent is

sensitive to shifts in excitation-inhibition dynamics[21] and serves here as a practical

electrophysiological index of local neural state. The spatial autocorrelation captures how

local neural dynamics at one location correlate with values at other locations, in relation to

the distance between electrodes. Therefore, lower spatial autocorrelation indicates the

presence of excitability "islands" that contrast sharply with their surroundings (**Fig. 1**).

Where local E/I states differ significantly from their neighbors, the resulting gradient would

impose a dynamic barrier, constraining the integration between adjacent circuits to follow a

divergent trajectory[1,3].

Crucially, this spatial autocorrelation framework diverges from traditional functional

connectivity. Rather than modeling communication between distinct brain regions, our

approach maps the continuous spatial distribution of local neural states. In highly complex

conditions like ASC, standard regional averaging obscures local excitation/inhibition

gradients, washing out crucial neurobiological variance.

By explicitly quantifying spatial heterogeneity rather than averaging it away, Moran's I

transforms cortical distribution into a single metric that still accommodates the possibility

that varying neural topographies underlie diverse clinical phenotypes.

To assess whether this heterogeneity is a stable trait rather than a spectral feature driven by

sensory processing or active cognition, we extended our analysis to sleep, a state where E/I

balance is strictly regulated and external confounds are minimized[22, 23].

Crucially, if this functional mesoscale heterogeneity is a stable developmental property, it

might be plausibly anchored in structural differences at the same spatial scale, offering a

multimodal framework to cross-validate these neurophysiological gradients. The local

morphological composite (encompassing gray matter, superficial white matter, and their

transitional boundary, represented by T1-weighted MRI intensity) served as a natural

structural candidate[24–27].

This study combines pre-registered confirmatory analyses with exploratory extensions. We

tested three pre-registered hypotheses:

- The spatial autocorrelation of E/I[21] is significantly lower in ASC individuals compared to

non-autistic controls.

- Increased E/I spatial variability can be observed in autistic individuals also during sleep.

- Greater spatial heterogeneity (i.e., lower spatial autocorrelation) of E/I is associated with

more pronounced autism-related traits (Autism Diagnostic Observation Scale-II, ADOS-2).

The pre-registration was conducted in our discovery sample and was later validated in two

independent cohorts. See Methods for any deviation.

Additionally, we deployed propensity-score matching to strengthen quasi-causal inferences

and triangulated our findings using MRI structure–function correlation. These exploratory

(non-pre-registered) analyses, which included the spatial sweep and independent-cohort

validation, were designed to assess the robustness and refine the primary hypotheses.

Results

The spatial heterogeneity of the aperiodic exponent is greater in ASC individuals compared to non-autistic individuals.

In total, 248 autistic subjects were recruited via self-referral or clinician-referral to a specialized tertiary care clinic (**Table 1**). A separate control group of 105 non-autistic individuals was selected from Emergency Room admissions. These patients had no history of epilepsy, acute brain injury, or neurodevelopmental or chronic neurological diagnoses. Clinical characteristics of the sample are presented in **Extended Data Table 1**.

During wakefulness, autistic individuals exhibited significantly greater spatial heterogeneity of the aperiodic exponent compared to non-autistic individuals (estimate = -0.55, 95% CI: -0.88 to -0.23; ASC n = 130, non-autistic n = 105; **Table 2**). This topographic alteration was independent of age and sex.

During sleep, we also observed a negative association between spatial autocorrelation of the aperiodic exponent and ASC, with an estimate of -0.85 (ASC EEGs n=211; non-autistic EEGs n=94; 95% CI: -1.17 to -0.52, **Extended Data Table 2**). This indicates that the higher spatial heterogeneity in ASC persists across wakefulness and sleep.

The reduced spatial autocorrelation persisted across the following sensitivity analyses: (i) balancing age and sex distributions via optimal propensity-score matching (**Extended Data Table 3**); (ii) excluding participants administered acute sleep aids (melatonin or dimetindene maleate) during recording[28, 29] (**Extended Data Tables 4a-b**); and (iii) excluding individuals with intellectual disability, global developmental delay, or epileptiform discharges (**Extended Data Tables 5-6**).

Independent-cohort validation.

141   After testing our primary hypothesis in the discovery cohort, we validated the findings in two

142   independent, publicly available datasets: 1) the Healthy Brain Network (HBN) dataset[30]; 2) a

143   dataset from the National Institute of Mental Health (NIMH) data archive (NIH), previously

144   published by Dede and colleagues[31].

145   *1) Healthy Brain Network validation analysis.* In the HBN dataset, we replicated the

146   association between ASC and lower aperiodic exponent spatial autocorrelation against

147   a community-referred non-autistic cohort heavily enriched for ADHD (ASC n = 416,

148   non-autistic n=2171, estimate = -0.34, 95% CI: -0.62 to -0.06, **Tables 1-2**). Similar

149   results were observed after adjusting for other neurodevelopmental conditions and in

150   age and sex matched samples (**Extended Data Table 7a and b**).

151   *2) NIH datasets validation analysis*[31]. Here too, we found that spatial autocorrelation of

152   aperiodic exponent was lower in the strictly defined autistic (AD) population, in

153   accordance with our hypothesis (**Tables 1-2**) (AD n=390, non-autistic n=555;

154   estimate: -0.25, 95% CI -0.43 to -0.08). This topographic alteration remained

155   significant after accounting for eye closure in a mixed-effects model (**Extended Data**

156   **Table 8a**)**.** Similarly, when using propensity-score matching for sex and age, we

157   observed an association between autism and higher spatial heterogeneity of the

158   aperiodic exponent (**Extended Data Table 8b**). The association persisted when

159   combining the AD and the lower traits penetrance ASD subgroup (**Extended Data**

160   **Table 8c**). After propensity score-matching, in this lower trait penetrance subgroup

161   the association did not reach statistical significance (**Extended Data Table 8d**).

Other spectral features are not consistently more heterogeneous in ASC.

Unlike the aperiodic exponent, peak alpha frequency and the aperiodic offset features did not

exhibit consistent heterogeneity across datasets. Peak alpha frequency was significantly more

heterogeneous in the ASC group within the HBN dataset but failed to replicate both in the

discovery sample and in the NIH dataset (**Supplementary Table 1**). Similarly, the aperiodic

offset showed increased heterogeneity only in the discovery dataset, and not in HBN

(**Supplementary Table 2**).

ASC is characterized by an increased heterogeneity of the aperiodic exponent specifically in

the mesoscale range.

To assess the scale specificity of our findings, we systematically mapped the aperiodic

exponent variability across a broad spatial range (30°–120°) in the discovery and (20°–120°)

HBN dataset. This analysis confirmed that the ASC-associated increase in heterogeneity is

not a global phenomenon but predominantly impacts the mesoscale network organization

(**Fig. 2A**).

To test whether mesoscale spatial organization carries diagnostic information beyond the

mean or variability of the aperiodic exponent, we compared nested logistic models predicting

ASC status (**Extended Data Table 9**). In both cohorts, mesoscale autocorrelation

outperformed competing summaries by AIC.

The mesoscale functional heterogeneity is mirrored at the structural level.

Using structural MRI from HBN, we tested whether the spatial profile of T1w intensity

mirrors the functional pattern.

The ASC coefficient on T1w spatial autocorrelation across region of interest (ROI) distances

showed a profile qualitatively similar to that of the EEG aperiodic exponent: negative at

mesoscale distances, returning toward positive values at larger scales (**Fig. 2B**).

A within-subject model revealed a significant EEG × ASC interaction on T1w spatial

autocorrelation (estimate = 0.16, CI 0.02 to 0.31), indicating different structure-function

coupling in ASC. Specifically, the positive association between functional and structural

heterogeneity was stronger in the ASC group. ASC status was also associated with lower

T1w spatial autocorrelation (estimate = -0.45, CI -0.86 to -0.03) (**Table 3**). The observed

cross-modal coupling remained robust when adjusting for global scan quality, and could not

be explained by the mean or variance of gray matter probability within the sampled ROIs (see

Methods).

Greater spatial heterogeneity of the aperiodic exponent is not associated with more

pronounced ASC traits.

In the discovery sample, we investigated the association between aperiodic exponent spatial

variability and ASC traits in autistic individuals. The ADOS total comparative score had an

estimate of 0.00 (EEGs n=184; 95% CI: -0.16 to 0.16), indicating no evidence for the pre-

registered hypothesis (**Extended Data Table 10**).

Discussion

Effective information processing relies on the spatial scale at which local units synchronize. Our findings indicate that in ASC, this arrangement has a divergent organization: the autistic cortex exhibits a more heterogeneous topography at the mesoscale.

While the aperiodic exponent is sensitive to shifts in E/I dynamics, it is not a pure measure of E/I ratio[32]. Nonetheless, our finding, while stemming from the autism E/I hypothesis, does not depend on the exact correspondence between the aperiodic exponent and E/I balance. By mapping the topological organization of the aperiodic exponent across the cortex, we demonstrate that the autistic cortex organizes a well-characterized property of neural activity more heterogeneously across space. This finding was replicated across three independent, large-scale datasets comprising 3767 participants, persisted during sleep, and remained significant after controlling for key confounds such as signal quality, comorbidities, and demographics.

With the above limitations, these results also extend prior reports of altered regional E/I balance in ASC[10,33,34] by revealing that such imbalances are spatially organized.

This heterogeneity persisted during sleep, indicating a stable, state-independent property not driven by sensory processing or active cognition[22]. Moreover, because Moran's I was evaluated against spatially permuted nulls, the observed difference reflects spatial organization rather than the value distribution itself.

Volume conduction, which acts as a spatial low-pass filter, is a potential confound[35,36]. However, three lines of evidence suggest that our findings reflect true neural heterogeneity rather than artifacts. First, biophysical simulations confine volume conduction blurring to a radius of 3–4 cm, attenuating rapidly beyond ~6–7 cm[36]; yet the group difference here is weak at the shortest lags where conduction is strongest, peaks at mesoscale distances and reverses sign beyond ~65° in the discovery sample and ~85° in HBN, a profile unlikely to

arise from a uniform confound. Second, volume conduction predicts smoother topographies

in larger heads; because ASC is associated with macrocephaly in approximately 15% of the

population, a head-size artifact would theoretically result in increased spatial autocorrelation.

Our observation of the opposite pattern (increased spatial heterogeneity) indicates that the

neural signal dominates the physical conduction effect. Third, a generalized conduction

artifact would uniformly affect all spectral features, yet our findings were metric-specific:

while the aperiodic exponent exhibited robust spatial heterogeneity, the aperiodic offset

showed no such effect (e.g., in the HBN dataset, the ASC coefficient for offset heterogeneity

was null: estimate = -0.02, p = 0.88).

The spatial specificity of our findings, peaking at the mesoscale (~6–9 cm), may provide

novel insight into the nature of E/I dysregulation in ASC. This window exceeds the radius of

maximum passive volume conduction (~3–4 cm), making a simple signal-leakage account

unlikely, yet remains local enough to capture the integration of distinct functional territories

before global signal dominance takes over. The increased heterogeneity at this specific scale

suggests that the neurobiological deficit in ASC may not be a failure of local micro-scale

circuits nor a whole-brain state shift at the macro-scale, but rather an alteration in the

functional coordination of neighboring neural populations.   Formal model comparison

reinforced this interpretation: mesoscale autocorrelation outperformed both the channel-wise

mean and standard deviation of the aperiodic exponent in predicting ASC status. The

diagnostic information carried by Moran's I thus reflects the spatial arrangement of

excitability states, not their central tendency or dispersion, ruling out the possibility that the

spatial signal merely recapitulates increased scatter across channels or average shifts in neural

states.

The relevance and specificity of the mesoscale range was further evidenced at the structural

level, where we observed regional heterogeneity in bias-corrected T1w signal intensity. At

the ROI dimensions utilized, the T1w signal reflects a local morphological composite,

capturing regional folding geometry, superficial white matter, and cytoarchitectural tissue

organization[37]. While this composite extends beyond the synaptic architecture generating the

EEG signal, it serves as a macroscopic proxy for local neuroanatomy. The coupling between

structural and functional heterogeneity was significantly stronger in autistic individuals,

suggesting that these two dimensions share a spatial organization in ASC. Consistent with

this, Stoner et al. identified patch-like cortical architecture in autistic individuals[38].

The association between reduced spatial autocorrelation and ASC was replicated across all

three datasets. However, in the NIH-derived dataset the association was absent when

combining the autism disorder subgroup with broader ASC criteria. The attenuation pattern

observed in this dataset is consistent with a signal that is modulated by both diagnostic

stringency and spatial measurement resolution: the NIH dataset's coarse 8-ROI parcellation

may lack the granularity to resolve mesoscale structure. Furthermore, recent evidence

suggests that as diagnostic criteria for neurodevelopmental conditions expand, the underlying

genetic and biological coherence tends to decrease[39]. Accordingly, the inclusion of older

individuals with lower trait penetrance in the NIH cohort likely introduces a broader, more

heterogeneous biological landscape, blurring the distinct mesoscale E/I topography that

specifically characterizes the more stringently defined neurodevelopmental phenotype in our

primary analysis.

However, within the ASC group, aperiodic exponent spatial autocorrelation was not

associated with autism trait penetrance as indexed by the ADOS comparative total score. The

absence of association with ADOS score, combined with the robust neurobiological signal,

suggests that mesoscale E/I heterogeneity may index a dimension of neural organization that

does not map into current summative behavioral phenotyping[40].

Our results remained consistent even when controlling for co-occurring neurodevelopmental

diagnoses (such as ADHD and intellectual disability), supporting the robustness of this signal

within our ASC cohort. However, whether this mesoscale spatial decoupling is uniquely

specific to ASC or represents a more general neurodevelopmental feature remains an open

question. Much like the extensive genetic pleiotropy observed across psychiatric

boundaries[41], this topographic E/I signature may reflect a transdiagnostic biological

vulnerability. This potential pleiotropy could partly explain why the spatial signal does not

scale linearly with ASC-specific behavioral metrics (e.g., the ADOS); rather, this

neurophysiological consistency coexists with marked clinical heterogeneity, underscoring

that a shared biological feature does not dictate a singular functional outcome.

Future studies with more refined clinical characterization will be essential. The spatial

distribution of EEG spectral features could help cluster autistic individuals into

phenotypically meaningful subtypes[41], and mapping individual E/I topography could inform

spatially targeted neuromodulation.

More broadly, the analytic framework developed here, i.e. mapping the spatial

autocorrelation of spectral parameters across the cortical surface, is not specific to autism. E/I

dysregulation features in schizophrenia, ADHD, and epilepsy, yet these conditions are

typically studied through approaches blind to spatial topology. Mesoscale spatial organization

of neural excitability may represent a general principle whose divergence characterizes

multiple neurodevelopmental conditions. Notably, this can be detected using scalp EEG,

making this insight accessible through common clinical tools.

These results should be interpreted in light of a few limitations. First, in all evaluated cohorts,

ASC diagnoses and assessments were performed at multiple centers and by different

clinicians. While this introduces noise, the persistence of the E/I signature across these

unstandardized environments suggests a robust phenotype relevant to real-world practice.

Indeed, the effect was largest in the discovery sample, where the clinical control group may

have contributed to a more conservative baseline and attenuated but significant in both

community-recruited cohorts. Second, potential referral biases, unequal phenotypic

characteristics across the diverse cohorts, and incomplete clinical characterization represent

inherent limitations of our study design, which we attempted to account for through targeted

sensitivity analyses.

About mesoscale specificity, scalp EEG provides more reliable spatial estimates at mesoscale

and longer distances than at shorter ones, where fewer electrode pairs and greater positioning

uncertainty reduce statistical power. As a result, our correlogram can more confidently

distinguish the mesoscale effect from longer-distance patterns than from shorter-distance

ones.

Regarding MRI, while our ROIs effectively capture local mesoscale morphological

composites, they inherently encompass tissue boundaries and partial volume effects,

precluding strict layer-specific claims about pure intracortical myelination. Moreover, our

radial projection method identifies the anatomical center of each sensor's macroscopic lead

field, representing a heuristic approximation that cannot fully account for the spatial smearing

inherent to EEG volume conduction.

This study identifies spatially heterogeneous neural dynamics as a fundamental

neurobiological feature of ASC. By shifting focus from global averages to mesoscale

topology, our results indicate that the spatial arrangement of local neural states carries

mechanistic information. The analytic framework developed here is generalizable beyond

autism and beyond E/I: mapping the spatial autocorrelation of any neural parameter across

330 the cortical surface offers a scalable, spatially informed approach to characterize macroscopic

331 brain organization using clinical EEG.

Materials and methods

<u>Discovery Sample's Participants</u>

This retrospective study was based on clinical data collected from participants who attended

the neurological Day Hospital at the Regina Margherita Children's Hospital, University of

Turin, between January 30, 2018, and May 29, 2024.

Inclusion criteria for the case group were the following:

- Children (age<18 years) with a previous diagnosis of autism spectrum disorder made by an

expert child and adolescent neuropsychiatrist, using standard diagnostic instruments.

- Written informed consent for research purposes signed by parents or legal guardians.

- An EEG performed on the date of the admission.

To ensure our sample was representative of the broad clinical population, we adopted an

inclusive recruitment strategy without restrictive exclusion criteria. All individuals with a

confirmed diagnosis of ASC who attended the neurological day hospital and completed an

EEG recording were included in the analysis, preserving the natural heterogeneity of the

cohort.

The control group for this study was derived from hospital data of participants (children

under 18 years of age) admitted to the emergency department during the same period. These

participants were referred for neuropsychiatric evaluation due to symptoms such as headache,

loss of consciousness, alteration of consciousness or paroxysmal movements. Exclusion

criteria applied to the control group included a diagnosis of developmental disorder, epilepsy,

acute brain injury, or seizure.

248 participants were recruited in the autistic group, and 105 participants were recruited in

the non-autistic group.

All the procedures were approved by the Citta della Salute e delle Scienza ethics committee

and parents or legal tutors of ASC individuals gave informed consent. Control EEGs were

collected anonymously.

<u>Publicly available datasets</u>

Healthy Brain Network. EEG data and participant demographics from the Healthy Brain

Network (HBN) (Releases 1–11 and the NC Release with available BIDS-formatted data)

were obtained from the public repository (

https://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/MRI_EEG.html). The

HBN uses a community-referred recruitment model in the New York City area and collects,

stores, and openly shares de-identified data on psychiatric, behavioral, cognitive, and lifestyle

phenotypes (e.g., fitness, diet), along with multimodal brain imaging (MRI),

electroencephalography (EEG), genetics, and actigraphy for children and adolescents aged 5–

21 years. EEG acquisition protocols and diagnostic criteria for the HBN cohort are described

elsewhere[30].

Sample collected from Dede et al. (NIH). The sample combined by Dede and colleagues[31]

and derived from the National Institute of Mental Health (NIMH) data archive

(https://nda.nih.gov/edit_collection.html?id=2022) were obtained from

https://github.com/adede1988/SheffieldAutismBiomarkers . In this dataset, we used the

aperiodic exponent normalized to the power spectrum distribution as our E/I measure to

mitigate inter-dataset differences in overall signal amplitude. Since our hypothesis focused on

autism, we included the "autistic disorder" (AD) group (using the original datasets

definition), the lower trait penetrance according to the ADOS nomenclature "ASD"

subgroup[31] and the non-autistic controls, excluding the schizophrenia and bipolar cohorts

from the Bipolar and Schizophrenia Consortium for parsing intermediate phenotypes (**Table**

1). We first isolated the strictly defined AD group to test for spatial E/I divergence in the

cohort with the most pronounced trait penetrance. We then analyzed the combined AD and

broader "ASD" cohorts against controls to test whether the topographic heterogeneity

generalizes across the wider diagnostic spectrum.

Both the HBN and the NIH-derived cohorts provided only wakefulness EEGs. In the NIH

dataset, we prioritized eyes-open recordings to maximize sample size.

<u>Measures and EEG Recordings in the discovery sample</u>

Demographic and clinical data were collected through electronic medical records. For the

case group, the collected data included: age at the admission, sex, comorbidities, traits

penetrance based on ADOS score, and EEG description. For the control group, the collected

data included: age, reason of admission, sex, anamnestic data, comorbidities, EEG

description.

According to clinical and technical considerations, EEGs were recorded during wakefulness,

sleep or both states.

The sleep recordings also served to exclude that increased E/I variability is associated with

information and/or sensory processing that occurs during wakefulness, or wake-related

artifacts (e.g. muscle activity).

Children in the case group underwent baseline standard EEG recording as part of a

comprehensive diagnostic protocol. For children in the control group, EEG recordings were

performed during their diagnostic evaluation for reasons other than neurodevelopmental

conditions, epilepsy and acute brain injury.

Trained pediatric neurologists evaluated the EEGs for presence and localization of

epileptiform discharges.

<u>EEG Data Analysis</u>

Extraction of E/I measure

To derive E/I from EEG segments, we employed the spectral parameterization technique

described by Donoghue and colleagues[21]. We focused on the exponent of the aperiodic

component, a well characterized electrophysiological proxy for E/I balance. Power spectral

density (PSD) was computed using Welch's method (1–40 Hz range). We then employed the

spectral parameterization algorithm (*specparam*, formerly *FOOOF*) to decompose the PSD

into periodic oscillatory peaks and an aperiodic 1/f exponent background.

For the NIH dataset we used data already available at

https://github.com/adede1988/SheffieldAutismBiomarkers .

For the NIH dataset, in which each ROI represents an average across multiple electrodes

from heterogeneous recording setups, we used the aperiodic exponent normalized to the

power spectrum distribution to mitigate inter-study amplitude biases.

Moran's I extraction

To quantify the spatial organization of the E/I landscape, we computed Moran's I for the

aperiodic exponent across the scalp surface.

Spatial autocorrelation was assessed using geodesic metrics to account for cranial topology

and arc lengths (great-circle distances) on a unit sphere, targeting a critical mesoscale

window of ~42–64° (corresponding to ~6–9 cm on a 51 cm circumference cranium). This

scale exceeds the radius of maximum passive volume conduction (~3–4 cm) to capture

genuine intra-hemispheric functional coordination (e.g., fronto-parietal integration) rather

than conductivity artifacts. We applied an inverse-distance kernel with a constant weighted-

mean neighbor angle of ~53° (~7.5 cm) to emphasize mid-range interactions. To ensure

consistent graph connectivity across varying resolutions, bandwidths were adapted inversely

to electrode density (8° for 128-channel, 11° for 19-channel, and 14° for 8-ROI datasets). The

primary target distance of 53° was selected during analysis development to center the kernel

above the dominant volume conduction radius and centered on the mesoscale; the subsequent

scale sweep (30°–120° for the discovery sample; 20°–120° for the higher resolution HBN

dataset) confirmed that the effect is not contingent on this specific choice. Because geodesic

angles are scale-invariant, the spatial weights are unaffected by head size variation across the

age range; the centimeter equivalents cited throughout are approximate, based on a

representative 51 cm head circumference (50th percentile for a 6-year-old boy, close to the

mean age of the discovery sample).

To verify that our results were not an artifact of the chosen spatial weighting scheme, we

conducted a sensitivity analysis specifically utilizing the higher-resolution HBN dataset (128

electrodes). In addition to the primary inversely weighted kernel, we re-computed the spatial

autocorrelations for both the EEG and MRI data using a continuous Gaussian spatial weight

matrix. The weight w_{ij} between electrodes I and j was defined as:

$$w_{ij} = \exp(-0.5 \times ((D_{ij} - d) / \sigma)^2)$$

where D_{ij} is the angular distance (in degrees), d is the target center distance of the spatial

filter, and σ is the bandwidth (fixed at 16°, corresponding to an approximate cortical distance

of 23 mm).

We applied this kernel to the HBN dataset only because a similar finely tuned spatial filtering

is methodologically robust only for high-density arrays (e.g., 128 channels), where the dense

spatial sampling provides sufficient pairs at varying distances.

To ensure consistent graph connectivity across the vastly different spatial resolutions of the

three datasets, we adapted the kernel bandwidth (δ) inversely to electrode density:

-       HBN (128-ch): Bandwidth = $\pm 8°$.

-       Discovery sample (19-ch): Bandwidth = $\pm 11°$.

-       NIH (8 ROIs): Bandwidth = $\pm 14°$.

Consequently, coarser arrays needed to integrate over a wider spatial band, which may dilute

scale specificity. Furthermore, for the HBN array, spatial weights were computed using a

regularized inverse-distance kernel with an offset = $5°$. This offset corresponds to the order-

of-magnitude positional uncertainty inherent to high-density electrode caps: cap fit variability

and inter-individual head geometry differences introduce electrode displacement errors of

approximately $3–7°$ in geodesic angle. Below this threshold, nominal distance differences

carry no reliable spatial information. The offset therefore acts as a nugget correction that

prevents the weight function from making spurious distinctions between pairs whose

geodesic separations differ by less than the measurement precision. For the 19-channel and 8-

ROI arrays, the wider kernel bandwidths ($\pm 11°$ and $\pm 14°$) already incorporate this positional

uncertainty, making the offset unnecessary.

This adaptive approach ensured that the spatial graph remained connected (reducing the

probability of isolated nodes) while targeting the same physical mesoscale distance in all

cohorts. Weights were row-standardized ("R" transform). Moran's I was computed against a

reference distribution generated via 499 random permutations. Lower (negative) Moran's I

values indicate greater spatial heterogeneity.

The spatial sweep analysis (30°–120° in the discovery sample; 20°–120° in the HBN) was

designed to characterize the distance-dependent profile of the ASC-E/I association, not to test

significance at each individual distance. We therefore report the full coefficient curve with

95% confidence intervals across all distances and interpret the overall shape of the profile

rather than applying pointwise significance thresholds. This approach avoids the inferential

distortions of multiple pointwise tests while preserving sensitivity to the scale-specificity of

the effect.

<u>T1-weighted MRI Preprocessing Electrode-matched MRI extraction</u>

Structural MRI data were obtained from the Healthy Brain Network (HBN) biobank. We

utilized the preprocessed T1-weighted (T1w) volumes generated by the Configurable Pipeline

for the Analysis of Connectomes (C-PAC)[42]. These data, available via the AWS Open Data

Registry (s3://fcp-indi/data/Projects/HBN/CPAC_preprocessed/), were processed using

standard anatomical registration workflows, resulting in skull-stripped, N4 bias-corrected

images non-linearly registered to the MNI152 template space at 2 mm isotropic resolution

(transform_Warped.nii.gz). To map functional EEG sensors to underlying cortical structure,

we utilized the standard GSN-HydroCel-129 montage geometry (EGI, Eugene, OR).

Electrode positions were mathematically transformed from the MNE-Python "Head"

coordinate frame (Neuromag convention: +X= Nasion, +Y= Left, +Z= Superior) to the

MNI152 RAS coordinate system (+X= Right, +Y= Anterior, +Z= Superior) via a 90-degree

axial rotation. For each electrode, a search vector was defined originating from the montage

center (0,0,0) and extending radially outward. We sampled the MNI152 2mm Gray Matter

(GM) probability template along this vector at depths ranging from 55 to 85 mm (2 mm

steps). The cortical coordinate for each electrode was defined as the depth maximizing GM

probability. To ensure valid cortical sampling, electrodes projecting to non-cortical tissue

(e.g., orbits, jaw, sinuses) were identified and excluded. Electrodes were retained only if their

projected 3 mm sphere captured a mean GM probability ≥ 0.30 (or ≥ 0.50 in a sensitivity

analysis). This procedure resulted in a final set of 119 validated (117 in the sensitivity

analysis with more stringent GM probability inclusion criteria) cortical electrodes for

downstream analysis.

Structural morphology was quantified by extracting T1w intensity values at each validated

cortical coordinate. For every subject (n = 1198), we defined spherical Regions of Interest

(ROIs) centered on the 119 (117 in the sensitivity analyses) projected coordinates. The

primary analysis utilized a 4 mm radius sphere, selected to capture the mesoscale

"morphological composite" (local gyrification and tissue contrast) of a single cortical gyrus

while minimizing overlap between adjacent sensors. Sensitivity analyses were performed

using radii of 3 mm and 6 mm. The mean T1w intensity was calculated within each sphere

using Python's nilearn. While raw T1w intensities are typically arbitrary units dependent on

scanner gain, our downstream use of Moran's I (a ratio of covariances) renders the analysis

scale-invariant, effectively normalizing global intensity differences between subjects.

Coordinate validation confirmed no systematic between-group bias in ROI placement, as

neither the mean nor the coefficient of variation of GM probability within the spherical ROIs

differed by diagnosis. Furthermore, the cross-modal dissociation test yielded a null

interaction (EEG_Moran_I x ASC) when evaluating GM probability ($p > 0.05$), suggesting

that the topological decoupling observed in the ASC group is specifically driven by T1w

signal rather than macroscopic differences in partial-volume tissue fractions.

While the spherical volumetric sampling offers lower cytoarchitectural specificity and gyral

homology than surface-based parcellations, it was specifically chosen to accommodate the

spatial constraints of the EEG array. To accurately compare spatial autocorrelation across

modalities, it is mathematically required that the structural and functional spatial weight

matrices (W) are identical. Employing spherical ROIs based on electrode projections

guarantees the topological equivalence.

Furthermore, volume conductor models demonstrate that EEG scalp signals are influenced by

the complex geometry of cortical folding and the distinct conductivity of the underlying

white matter.

<u>Statistical Analysis</u>

To test the hypothesis that greater spatial heterogeneity in the E/I balance is associated with

ASC features, we used Moran's I to quantify spatial heterogeneity and the aperiodic exponent

as an electrophysiological proxy for E/I[21,32,43]. Moran's I is a measure of spatial variability

approximately centered on zero, such that positive values indicate clustering and negative

values indicate dispersion (**Fig. 1**).

For analyses spanning multiple behavioral states (sleep, resting-state), we fit linear mixed-

effects models with subject-specific random intercepts. Associations between clinical

variables and E/I Moran's I, where no grouping structure was required, were tested with

linear regression. All continuous variables were standardized. In matched samples, matching

weights were used as regression weights.

Within-subject EEG–MRI spatial coupling in ASC and non-ASC participants was tested

using the linear model:

$\qquad$ MRI_Moran_I ~ EEG_Moran_I + ASC + EEG_Moran_I × ASC + age + sex

where EEG_Moran_I is the electrode-level Moran's I of the aperiodic exponent (eyes-closed),

and MRI_Moran_I is the T1w Moran's I at a given distance. The EEG_Moran_I × ASC

interaction term tests whether the strength of structure–function spatial coupling differs

between groups. We swept the MRI distance from 20° to 120° in 10° steps to assess the shape

of the association across spatial scales.

To verify robustness against potential confounds, we performed parallel control analyses

including: (i) T1w image quality metrics (coefficient of variation, entropy), (ii) head motion

(mean framewise displacement), and (iii) site-specific scanner performance metrics (from

MRIQC), demonstrating that the observed effects were not driven by data quality or

acquisition artifacts.

All reported p-values are two-sided.

Claude Code (Anthropic, Claude Opus), integrated in the Visual Studio Code IDE, was used

to assist with code linting and commenting. The authors reviewed all outputs and take full

responsibility for the accuracy and integrity of all code, analyses, and results.

<u>Study registration, open science and code availability.</u>

The three main hypotheses and their analyses were registered on the Open Science

Framework before analysing the data (https://osf.io/4q83m). Python codes for data cleaning

and   analysis   are   available   in   GitHub   at

https://github.com/danielemarcotulli/mesoscale_architecture_autism.git .

<u>Deviation from the pre-registered analyses.</u>

While the study adhered to the core pre-registered protocol (https://osf.io/4q83m/) the

following deviations were made. First, we expanded the spatial analysis from a single

distance to a sweep across 30°–120° / 20°–120° to characterize scale-specificity. Second, we

added propensity-score weighting and medication exclusions as robustness checks. Third, we

extended validation to two independent datasets (HBN and NIH). Fourth, we analyzed MRIs

from the HBN dataset to evaluate structure-function coupling. These analyses are exploratory

and are distinguished from confirmatory results throughout.

Acknowledgments: We thank the UCL MEG group and Dr. Daniel Rautio for offering constructive

feedback and discussion. Claude Code (Anthropic, Claude Opus) was used for code linting and

commenting. The authors take full responsibility for the accuracy and integrity of all code, analyses,

and results.

Funding: This work received no specific funding.

Author contributions:

Conceptualization: DM

Methodology: DM

Software: DM

Investigation: DM, VFC

Data Curation: VFC, DM

Validation: DM

Formal analysis: DM

Resources: CD, MV, BS, MG, SM, FSR, CC

Visualization: DM, LB

Project administration: DM

Supervision: AS, CD

Writing – original draft: DM, VFC

Writing – review & editing: All Authors

Competing Interests:

In the last 2 years, DM has received consultant honoraria from Angelini (Ethos).

In the last 2 years, BV has received consultant honoraria from Simon-Kucher, Medice, Angelini

(Ethos), and Alkermes Pharmaceuticals.

In the last 2 years, TB has received consultant honoraria from AGB pharma, Infectopharm, Medice,

Neuraxpharm, Neurim Pharmaceuticals, Oberberg GmbH and Takeda, and speaker's fee by AGB

pharma, Janssen-Cilag, Medice, Oberberg GmbH and Takeda.

Data availability:

Data used for results validation from the dataset of Dede et al.[31] are available at

https://github.com/adede1988/SheffieldAutismBiomarkers.

Use, sharing, adaptation, distribution, and reproduction are permitted under the Creative Commons

Attribution 4.0 International License (CC BY 4.0), provided appropriate credit is given to the original

author(s) and the source, a link to the license is provided, and any changes are indicated. Images or

other third-party material are included under the article's Creative Commons license unless otherwise

stated; for uses not covered by the license or permitted by law, permission may be required from the

rights holder. License: http://creativecommons.org/licenses/by/4.0/.

EEG data from the Healthy Brain Network (HBN) are freely available from the HBN portal:

https://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/ . Access to associated

phenotypic data is granted upon request via the same website, following the procedures described

there.

Discovery sample's individual-level EEG and clinical data contain protected health information from

minors and cannot be made openly available under the terms of the ethical approval granted by the

Citta della Salute e delle Scienza ethics committee. Fully de-identified data are available from the

corresponding author upon reasonable request. Access requests will be reviewed within 6 weeks and

granted subject to a data sharing agreement compliant with applicable privacy regulations (EU

General Data Protection Regulation).

References:

1. Shew, W. L., Yang, H., Yu, S., Roy, R. & Plenz, D. Information capacity and transmission are maximized in balanced cortical networks with neuronal avalanches. *J. Neurosci.* **31**, 55–63 (2011).

2. Yizhar, O., Fenno, L. E., Prigge, M., Schneider, F., Davidson, T. J., O'Shea, D. J., Sohal, V. S., Goshen, I., Finkelstein, J., Paz, J. T., Stehfest, K., Fudim, R., Ramakrishnan, C., Huguenard, J. R., Hegemann, P. & Deisseroth, K. Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature* **477**, 171–178 (2011).

3. Fries, P. Rhythms for cognition: communication through coherence. *Neuron* **88**, 220–235 (2015).

4. Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C., Laumann, T. O., Miezin, F. M., Schlaggar, B. L. & Petersen, S. E. Functional network organization of the human brain. *Neuron* **72**, 665–678 (2011).

5. Hirota, T. & King, B. H. Autism spectrum disorder: a review. *JAMA* **329**, 157–168 (2023).

6. Lord, C., Brugha, T. S., Charman, T., Cusack, J., Dumas, G., Frazier, T., Jones, E. J. H., Jones, R. M., Pickles, A., State, M. W., Taylor, J. L. & Veenstra-VanderWeele, J. Autism spectrum disorder. *Nat. Rev. Dis. Primer* **6** (2020).

7. Kamp-Becker, I. Autism spectrum disorder in ICD-11 — a critical reflection of its possible impact on clinical practice and research. *Mol. Psychiatry* **29**, 633–638 (2024).

8. Rubenstein, J. L. R. & Merzenich, M. M. Model of autism: increased ratio of excitation/inhibition in key neural systems. *Genes Brain Behav.* **2**, 255–267 (2003).

9. Chao, H. T., Chen, H., Samaco, R. C., Xue, M., Chahrour, M., Yoo, J., Neul, J. L., Gong, S., Lu, H. C., Heintz, N., Ekker, M., Rubenstein, J. L. R., Noebels, J. L., Rosenmund, C. & Zoghbi, H. Y. GABAergic dysfunction mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature* **468**, 263–269 (2010).

10. Lee, E., Lee, J. & Kim, E. Excitation/inhibition imbalance in animal models of autism spectrum disorders. *Biol. Psychiatry* **81**, 838–847 (2017).

651   11.   Belmonte, M. K., Cook, E. H., Anderson, G. M., Rubenstein, J. L. R., Greenough, W. T.,

652   Beckel-Mitchener, A., Courchesne, E., Boulanger, L. M., Powell, S. B., Levitt, P. R., Perry, E.

653   K., Jiang, Y. H., DeLorey, T. M. & Tierney, E. Autism as a disorder of neural information

654   processing: directions for research and targets for therapy. *Mol. Psychiatry* **9**, 646–663 (2004).

655   12.   Duffy, F. H. & Als, H. A stable pattern of EEG spectral coherence distinguishes children with

656   autism from neuro-typical controls — a large case control study. *BMC Med.* **10**, 64 (2012).

657   13.   Markicevic, M., Fulcher, B. D., Lewis, C., Helmchen, F., Rudin, M., Zerbi, V. & Wenderoth,

658   N. Cortical excitation:inhibition imbalance causes abnormal brain network dynamics as

659   observed in neurodevelopmental disorders. *Cereb. Cortex* **30**, 4922–4937 (2020).

660   14.   Nelson, S. B. & Valakh, V. Excitatory/inhibitory balance and circuit homeostasis in autism

661   spectrum disorders. *Neuron* **87**, 684–698 (2015).

662   15.   Antoine, M. W., Langberg, T., Schnepel, P. & Feldman, D. E. Increased excitation–inhibition

663   ratio stabilizes synapse and circuit excitability in four autism mouse models. *Neuron* **101**, 648–

664   661.e4 (2019).

665   16.   Cornew, L., Roberts, T. P. L., Blaskey, L. & Edgar, C. Resting-state oscillatory activity in

666   autism spectrum disorders. *J. Autism Dev. Disord.* **42**, 1884–1894 (2012).

667   17.   Happé, F. & Frith, U. The weak coherence account: detail-focused cognitive style in autism

668   spectrum disorders. *J. Autism Dev. Disord.* **36**, 5–25 (2006).

669   18.   Schwartz, S., Kessler, R., Gaughan, T. & Buckley, A. W. Electroencephalogram coherence

670   patterns in autism: an updated review. *Pediatr. Neurol.* **67**, 7–22 (2017).

671   19.   Deco, G., Ponce-Alvarez, A., Hagmann, P., Romani, G. L., Mantini, D. & Corbetta, M. How

672   local excitation–inhibition ratio impacts the whole brain dynamics. *J. Neurosci.* **34**, 7886–7898

673   (2014).

674   20.   Zhang, J., Cheng, W., Liu, Z., Zhang, K., Lei, X., Yao, Y., Becker, B., Liu, Y., Kendrick, K.

675   M., Lu, G. & Feng, J. Neural, electrophysiological and anatomical basis of brain-network

676   variability and its characteristic changes in mental disorders. *Brain* **139**, 2307–2321 (2016).

21.   Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A.

H., Wallis, J. D., Knight, R. T., Shestyuk, A. & Voytek, B. Parameterizing neural power spectra

into periodic and aperiodic components. *Nat. Neurosci.* **23**, 1655–1665 (2020).

22.   Peyrache, A., Dehghani, N., Eskandar, E. N., Madsen, J. R., Anderson, W. S., Donoghue, J. A.,

Hochberg, L. R., Halgren, E., Cash, S. S. & Destexhe, A. Spatiotemporal dynamics of

neocortical excitation and inhibition during human sleep. *Proc. Natl Acad. Sci. USA* **109**, 1731–

1736 (2012).

23.   Niethard, N., Hasegawa, M., Itokazu, T., Oyanedel, C. N., Born, J. & Sato, T. R. Sleep-stage-

specific regulation of cortical excitation and inhibition. *Curr. Biol.* **26**, 2739–2749 (2016).

24.   Koessler, L., Maillard, L., Benhadid, A., Vignal, J. P., Felblinger, J., Vespignani, H. & Braun,

M. Automated cortical projection of EEG sensors: anatomical correlation via the international

10–10 system. *NeuroImage* **46**, 64–72 (2009).

25.   Burt, J. B., Demirtaş, M., Eckner, W. J., Navejar, N. M., Ji, J. L., Martin, W. J., Bernacchia, A.,

Anticevic, A. & Murray, J. D. Hierarchy of transcriptomic specialization across human cortex

captured by structural neuroimaging topography. *Nat. Neurosci.* **21**, 1251–1259 (2018).

26.   Glasser, M. F. & Van Essen, D. C. Mapping human cortical areas in vivo based on myelin

content as revealed by T1- and T2-weighted MRI. *J. Neurosci.* **31**, 11597–11616 (2011).

27.   Andrews, D. S., Avino, T. A., Gudbrandsen, M., Daly, E., Marquand, A., Murphy, C. M., Lai,

M.-C., Lombardo, M. V., Ruigrok, A. N. V., Williams, S. C., Bullmore, E. T., The MRC AIMS

Consortium, Suckling, J., Baron-Cohen, S., Craig, M. C., Murphy, D. G. M. & Ecker, C. In

vivo evidence of reduced integrity of the gray–white matter boundary in autism spectrum

disorder. *Cereb. Cortex* **27**, 877–887 (2017).

28.   Kim, J. H., Ha, E. K., Han, B., Han, T., Shin, J., Chae, K. Y., Rhie, S. & Han, M. Y. First-

generation antihistamines and seizures in young children. *JAMA Netw. Open* **7**, e2429654

(2024).

29.   Dell'Isola, G. B., Tascini, G., Vinti, V., Tulli, E., Dini, G., Mencaroni, E., Ferrara, P., Di Cara,

G., Striano, P. & Verrotti, A. Effect of melatonin on sleep quality and EEG features in

childhood epilepsy: a possible non-conventional treatment. *Front. Neurol.* **14**, 1243917 (2023).

30.   Alexander, L. M., Escalera, J., Ai, L., Andreotti, C., Febre, K., Mangone, A., Vega-Potler, N.,

Langer, N., Alexander, A., Kovacs, M., Litke, S., O'Hagan, B., Andersen, J., Bronstein, B., Bui,

A., Bushey, M., Butler, H., Castagna, V., Camacho, N., Chan, E., Citera, D., Clucas, J., Cohen,

S., Dufek, S., Eaves, M., Fradera, B., Gardner, J., Grant-Villegas, N., Green, G., Gregory, C.,

Hart, E., Harris, B., Horton, J., Kahn, A., Kabotyanski, K., Karmel, B., Kelly, M. P., Kleinman,

B., Koo, E., Kramer, S., Lennon, E., Lord, A., Mantello, Y., Margolis, A., Merzon, G. R.,

Minshew, B. M., Moed, E. H., Naparstek, M. C., Panchal, A., Parra, M., Pugh, K. R.,

Racanello, C. R., Restrepo, A., Saltzman, R., Septimus, E., Tobe, R., Waltzer, K., Williams, A.,

Yeo, A., Castellanos, F. X. & Milham, M. P. An open resource for transdiagnostic research in

pediatric mental health and learning disorders. *Sci. Data* **4**, 170181 (2017).

31.   Dede, A. J. O., Xiao, W., Vaci, N., Cohen, M. X. & Milne, E. Exploring EEG resting state

differences in autism: sparse findings from a large cohort. *Mol. Autism* **16**, 13 (2025).

32.   Brake, N., Duc, F., Rokos, A., Arseneau, F., Shahiri, S., Khadra, A. & Plourde, G. A

neurophysiological basis for aperiodic EEG and the background spectral trend. *Nat. Commun.*

15, 1514 (2024).

33.   Bruining, H., Hardstone, R., Juarez-Martinez, E. L., Sprengers, J., Avramiea, A.-E., Simpraga,

S., Houtman, S. J., Poil, S.-S., Dallares, E., Palva, S., Oranje, B., Palva, J. M., Mansvelder, H.

D. & Linkenkaer-Hansen, K. Measurement of excitation–inhibition ratio in autism spectrum

disorder using critical brain dynamics. *Sci. Rep.* **10**, 9195 (2020).

34.   Huang, Q., Chen, D., Pereira, A. C., Leonard, A., Ellis, C. L., Velthuis, H., Dimitrov, M.,

Ponteduro, F. M., Wong, N. M. L., Kowalewski, L., Pretzsch, C. M., Daly, E., Murphy, D. G.

M. & McAlonan, G. M. Differential GABA dynamics across brain functional networks in

autism. *Commun. Biol.* **9**, 283 (2026).

35.   Grieve, P. G., Emerson, R. G., Isler, J. R. & Stark, R. I. Quantitative analysis of spatial

sampling error in the infant and adult electroencephalogram. *NeuroImage* **21**, 1260–1274

(2004).

36.   Srinivasan, R., Tucker, D. M. & Murias, M. Estimating the spatial Nyquist of the human EEG.

Behav. Res. Methods Instrum. Comput. **30**, 8–19 (1998).

37. Lerch, J. P., van der Kouwe, A. J. W., Raznahan, A., Paus, T., Johansen-Berg, H., Miller, K. L.,

Smith, S. M., Fischl, B. R. & Sotiropoulos, S. N. Studying neuroanatomy using MRI. *Nat.*

Neurosci. **20**, 314–326 (2017).

38. Stoner, R., Chow, M. L., Boyle, M. P., Sunkin, S. M., Mouton, P. R., Roy, S., Wynshaw-Boris,

A., Colamarino, S. A., Lein, E. S. & Courchesne, E. Patches of disorganization in the neocortex

of children with autism. *N. Engl. J. Med.* **370**, 1209–1219 (2014).

39. LaBianca, S., Lousdal, M. L., Krebs, M. D., Hansen, O. S., Georgii Hellberg, K.-L. L. A.,

Lundberg, M., Sørensen, J. Ø., Gaadin, J. R., Ohlsson, H., iPSYCH Study Consortium,

Borglum, A., Agerbo, E., Werge, T., Albinana, C., Vilhjalmsson, B. J., Kendler, K. S., Plana-

Ripoll, O. & Schork, A. J. As rates of ASD and ADHD rise, genetic contributions fall: evidence

for widening diagnostic criteria. Preprint at *medRxiv*

https://doi.org/10.1101/2025.11.10.25339891 (2025).

40. Lombardo, M. V., Lai, M.-C. & Baron-Cohen, S. Big data approaches to decomposing

heterogeneity across the autism spectrum. *Mol. Psychiatry* **24**, 1435–1450 (2019).

41. The Brainstorm Consortium. Analysis of shared heritability in common disorders of the brain.

Science **360**, eaap8757 (2018).

42. Craddock, C., Sikka, S., Cheung, B., Khanuja, R., Ghosh, S. S., Yan, C., Li, Q., Lurie, D.,

Vogelstein, J., Burns, R., Colcombe, S., Mennes, M., Kelly, C., Di Martino, A., Castellanos, F.

X. & Milham, M. Towards automated analysis of connectomes: the Configurable Pipeline for

the Analysis of Connectomes (C-PAC). *Front. Neuroinform.* Conference Abstract:

Neuroinformatics 2013 (2013).

43. McKeown, D. J., Finley, A. J., Kelley, N. J., Cavanagh, J. F., Keage, H. A. D., Baumann, O.,

Schinazi, V. R., Moustafa, A. A. & Angus, D. J. Test–retest reliability of spectral

parameterization by 1/f characterization using SpecParam. *Cereb. Cortex* **34**, bhad482 (2024).

Figure 1. Visual demonstration of increased spatial variability at the mesoscale.

Schematic representation of spatial autocorrelation in the aperiodic exponent. The plots illustrate two

contrasting spatial organizations of the EEG aperiodic exponent. Each dot (or star) represents a

sensor. For a given reference sensor (central star), spatial autocorrelation is calculated based on the

similarity of signals within a specific, geometrically defined spatial neighborhood or "lag" (here, the

mesoscale band of ~6–9 cm, denoted by dashed concentric circles). Sensor color reflects the local

*aperiodic exponent value. (**Left**) A highly coordinated spatial organization (high spatial*

autocorrelation). Sensors within the mesoscale neighborhood exhibit similar aperiodic exponent

values to the reference node and to each other, reflecting a smooth, continuous neural states

*architecture. (**Right**) A heterogeneous spatial topology (low spatial autocorrelation). Sensors within*

the exact same mesoscale boundary display highly variable aperiodic values, indicating a "patchy"

landscape characterized by abrupt local shifts in neural states.

Table 1. Demographic characteristics of the sample.

Dataset	Group	N, % of total	Age, years, mean (SD)	Males, n (%)
Discovery sample				
	ASC	248 (70.3)	5.5 (3.4)	196 (79)
	Non-autistic	105 (29.7)	8.5 (4.8)	82 (78)
Healthy Brain Network (HBN)[a]				
	ASC	416 (16.1)	10.6 (3.7)	341 (82.0)
	ADHD	1600 (61.8)	10.2 (3.3)	1154 (72.1)
	Intellectual Disability	66 (2.5)	12.2 (3.5)	40 (60.6)
	Non-autistic	585 (22.6)	10.9 (3.9)	281 (48.0)
NIH				
	AD	390 (37.6)	9.9 (8.1)	293 (74.9)
	ASD	93 (8.9)	16.2 (15.1)	54 (58.1)
	Non-autistic	555 (53.5)	20.4 (16.6)	296 (53.3)

778   *a. Diagnostic categories are not mutually exclusive*

779   **Table 2. Spatial autocorrelation of the aperiodic exponent during wake across the three**

780   **datasets.**

Dataset	Regressors	Estimates	95% CI	p-value
Discovery				
sample	ASC	-0.55	-0.88 – -0.23	**0.001**
n = 235	Sex (M)	-0.24	-0.63 – 0.14	0.207
	Age	0.13	-0.02 – 0.30	0.097
HBN				
n = 2587	ASC	-0.34	-0.62 – -0.06	**0.016**
	Sex (M)	-0.17	-0.39 – 0.04	0.110
	Age	0.25	0.15 – 0.35	**<0.001**
NIH				
n = 945	ASC (AD)	-0.25	-0.43 – -0.08	**0.005**
	Sex (M)	0.15	-0.02 – 0.32	0.089
	Age	0.12	0.04 – 0.21	**0.005**

783   *Bold p-values indicate significance at α = 0.05.*

784   *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

Table 3. T1w spatial autocorrelation (at 53°) in the HBN dataset.

Regressors	Estimates	95% CI	p-value
ASC	-0.45	-0.86 – -0.03	**0.03**
Age	-0.08	-0.21 – 0.05	0.23
Sex (M)	0.22	-0.05 – 0.50	0.11
EEG's Moran I	-0.05	-0.11 – 0.01	0.11
EEG's Moran I x ASC	0.16	0.02 – 0.31	**0.02**

788   Subjects with both MRIs and EEGs, n = 1198

789   *Bold p-values indicate significance at α = 0.05.*

790   *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

 Figure 2. Association between ASC and E/I spatial heterogeneity.

(A) Standardized regression coefficients (estimate) for the association between ASC diagnosis and

spatial autocorrelation (Moran's I) of the aperiodic exponent, computed across geodesic distances

from 20° to 120°. Negative coefficients indicate greater spatial heterogeneity in the ASC group.

Shaded bands represent 95% confidence intervals. The ASC-associated reduction in spatial

autocorrelation is concentrated in the mesoscale range (~42°–64°, corresponding to ~6–9 cm),

converging toward zero at shorter and longer distances. Results are shown as separate lines for the

discovery sample (red) and HBN (blue). At short distances (<30° for the 19-channel array), the

number of electrode pairs within the kernel bandwidth falls below the threshold for reliable Moran's I

estimation. The NIH dataset was excluded from the scale sweep because its 8 ROIs yield too few

spatial pairs per distance band for a stable correlogram. Shaded areas indicate confidence interval.

Across spatial lags, the ASD–control difference in spatial autocorrelation showed significant

heterogeneity in both datasets, indicating that the effect was scale-specific rather than uniform across

distances. In HBN, Cochran's Q confirmed significant heterogeneity across the 11 tested scales (20°–

120°; Q = 37.54, p < 0.001; I² = 73.4%), and meta-regression revealed a significant linear trend

toward less negative coefficients with increasing distance (estimate = 0.006 per degree, z = 4.76, p <

0.001), consistent with a group effect that peaks at mesoscale and reverses at longer lags. A similar

pattern was observed in the discovery sample across 10 scales (30°–120°; Q = 43.35, p < 0.001; I² =

79.2%; meta-regression estimate = 0.012 per degree, z = 5.90, p < 0.001). Together, these analyses

indicate that the ASC-related difference in spatial autocorrelation is not constant across the scalp but

is concentrated at mesoscale distances.

(B) Standardized regression coefficients for the association between ASC diagnosis and T1w spatial

autocorrelation (Moran's I) across electrode distances (20°–120°). The T1w profile paralleled the

EEG pattern: negative at mesoscale distances, returning toward positive values at larger scales.

Shaded areas indicate confidence interval.

 EXTENDED DATA

 Extended Data Table 1. Demographic, clinical and EEG-related characteristics of the discovery

 cohort.

Variable	*ASC (N=248)*	*Non-autistic (n=105)*
Sex, n (% males)	196 (79)	82 (78)
Age, months, mean (SD)	66.1 (41.3)	101.5 (57.2)
Age range, months	20-202	15-215
Intellectual Disability, n (%)	155 (62)	0 (0)
Comorbidities, n (%)	88 (35)	31 (29.5)
EEG anomalies reported, %	153 (61.7)	48 (45.7)
EEG state (n)		
Wake	130	105
Sleep	211	94
Medication during EEG (n)		
Melatonin	152	53
Dimetindene Maleate	60	-
Reason for admission non-autistic only, n (%)		
Loss of consciousness	-	38 (36.2)

Non-epileptic paroxysmal events	-	28 (26.7)
Headache	-	25 (23.8)
Altered consciousness	-	6 (5.7)
Vertigo	-	6 (5.7)
Other	-	2 (1.9)

 Extended Data Table 2. Aperiodic exponent (slope) spatial autocorrelation during sleep in the

827 **discovery cohort.**

Regressor	Estimate	95% CI	p-value
ASC	-0.85	-1.17 – -0.52	**<0.001**
Age	0.12	-0.04 – 0.27	0.139
Sex	-0.09	-0.46 – 0.28	0.630

828 EEGs n = 305

829 *Bold p-values indicate significance at α = 0.05.*

830 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

833 **Extended Data Table 3. Aperiodic exponent spatial autocorrelation during wake in the age and**

834 **sex matched samples (discovery cohort).**

Regressor	Estimate	95% CI	p
ASC	-0.62	-0.96 – -0.29	**<0.001**
Age	0.14	-0.03 – 0.31	0.098
Sex	0.28	-0.13 – 0.69	0.175

836 EEGs n = 210

837 *Bold p-values indicate significance at α = 0.05.*

838 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

840 **Extended Data Table 4a. Aperiodic exponent spatial autocorrelation excluding participants who**

841 **received dimetindene maleate in both sleep and wakefulness EEG (discovery sample).**

842

Regressor	Estimate	95% CI	p-value
ASC	-0.65	-0.84 – -0.45	**<0.001**
Age	0.14	0.04 – 0.25	**0.009**
Sex	-0.07	-0.33 – 0.19	0.595
Wake	0.18	0.00 – 0.36	0.050

843 EEGs n = 398

844 Subjects n = 207

845

846 **Extended Data Table 4b. Aperiodic exponent spatial autocorrelation excluding participants who**

847 **received melatonin as a sleep aid for their sleep EEG (discovery sample).**

Regressor	Estimate	95% CI	p-value
ASC	-0.55	-0.82 – -0.29	**<0.001**
Age	0.12	-0.03 – 0.28	0.121
Sex	-0.07	-0.48 – 0.35	0.763
Wake	0.13	-0.11 – 0.38	0.282

EEGs n = 213,

Subjects n = 114

848 *Bold p-values indicate significance at α = 0.05.*

849   *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

Extended Data Table 5. Aperiodic exponent spatial autocorrelation excluding ASC participants with intellectual disability/global developmental delay (discovery sample).

Regressor	Estimate	95% CI	p-value
ASC	-0.59	-0.86 – -0.33	**<0.001**
Age	0.15	-0.02 – 0.31	0.084
Sex	-0.13	-0.59 – 0.34	0.592
Wake	0.18	-0.05 – 0.40	0.132

EEGs n = 217

Subjects n = 99

Bold p-values indicate significance at α = 0.05.

Spatial autocorrelation was evaluated at a geodesic distance of 53°.

860 **Extended Data Table 6. Aperiodic exponent spatial autocorrelation excluding ASC participants**

861 **with epileptiform or slow wave anomalies (discovery sample).**

Regressor	Estimate	95% CI	p-value
ASC	-0.70	-0.95 – -0.45	**<0.001**
Age	0.17	0.04 – 0.31	**0.011**
Sex	-0.10	-0.42 – 0.21	0.530
Wake	0.05	-0.03 – 0.39	0.644

EEGs n = 232

Subjects n = 111

Bold p-values indicate significance at α = 0.05.

Spatial autocorrelation was evaluated at a geodesic distance of 53°.

Extended Data Table 7a. Aperiodic exponent spatial autocorrelation in age and sex matched

samples from the HBN dataset.

Regressor	Estimate	95% CI	p-value
ASC	-0.43	-0.79 – -0.06	**0.022**
Age	0.04	-0.01 – 0.09	0.09
Sex	0.12	-0.35 – 0.59	0.62

871 EEGs n = 832

873 **Extended Data Table 7b. Aperiodic exponent spatial autocorrelation adjusting for coexisting**

874 **conditions in the HBN dataset.**

Regressor	Estimate	95% CI	p-value
ASC	-0.35	-0.63 – -0.07	**0.014**
ADHD	0.03	-0.18 – 0.24	0.778
Intellectual Disability	0.16	-0.48 – 0.81	0.619
Age	0.25	0.15 – 0.35	**<0.001**
Sex	-0.18	-0.40 – 0.04	0.107

876 EEGs n = 2587

877 *Bold p-values indicate significance at α = 0.05.*

878 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

880 **Extended Data Table 8a. Aperiodic exponent spatial autocorrelation in the NIH dataset**

881 **adjusting for eyes closure (mixed-effects model).**

Regressor	Estimate	95% CI	p-value
ASC (AD)	-0.18	-0.33 – -0.03	**0.020**
Sex (M)	0.07	-0.07 – 0.21	0.343
Age	0.09	0.02 – 0.16	**0.013**
Eyes (open)	-0.46	-0.60 – -0.31	**<0.001**

883 EEGs n = 1396

885 **Extended Data Table 8b. Aperiodic exponent spatial autocorrelation in the NIH dataset in age**

886 **and sex matched samples.**

Regressor	Estimate	95% CI	p-value
ASC (AD)	-0.19	-0.36 – -0.02	**0.044**
Sex (M)	0.29	0.09 – 0.49	**0.004**
Age	0.02	-0.07 – 0.11	0.672

888 EEGs n = 702

889 **Extended Data Table 8c. Aperiodic exponent spatial autocorrelation in the NIH dataset**

890 **(including the lower penetrance category "ASD" from Dede et al.).**

Regressor	Estimate	95% CI	p-value
ASC (AD + ASD)	-0.19	-0.35 – -0.02	**0.025**
Sex (M)	0.10	-0.06 – 0.27	0.208
Age	0.14	0.06 - 0.22	**0.001**

892 EEGs n = 1038

894 **Extended Data Table 8d. Aperiodic exponent spatial autocorrelation in the NIH dataset**
895 **(including the lower penetrance category "ASD" from Dede et al.) in age and sex matched**
896 **samples.**

Regressor	Estimate	95% CI	p-value
ASC (ASD + AD)	-0.15	-0.32 – 0.02	0.081
Sex (M)	0.22	0.05 – 0.40	**0.013**
Age	0.12	0.03 – 0.20	**0.007**

898 EEGs n = 834

899 *Bold p-values indicate significance at α = 0.05.*

900 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

902   **Extended Data Table 9. Nested logistic regression model comparison for ASC classification.**

	Discovery (19ch, n = 235)		HBN (128ch, n = 2587)	
Model	**AIC**	**LR χ^2 vs A**	**AIC**	**LR χ^2 vs A**
A: age + sex	326.8	–	2212.5	–
B: A + mean exponent	323.8	5.00 (p = 0.025)	2209.7	4.83 (p = 0.028)
C: A + SD exponent	328.8	0.00 (p = 0.950)	2214.5	0.01 (p = 0.910)
D: A + mesoscale zI	**317.3**	11.53 (p < 0.001)	2208.5	6.00 (p = 0.014)
E: A + mean + SD + zI	318.0	–	**2206.9**	–

Contrast	**Discovery**	**HBN**
E vs B (spatial measures add to mean?)	$\chi^2(2) = 9.84$, **p = 0.007**	$\chi^2(2) = 6.80$, **p = 0.03**
E vs D (level measures add to zI?)	$\chi^2(2) = 3.31$, p = 0.19	$\chi^2(2) = 5.63$, p = 0.06

904   *Best AIC per cohort in bold. Bold p-values indicate significance at α = 0.05.*

908 **Extended Data Table 10. E/I spatial autocorrelation and ASC traits penetrance in the discovery**

909 **sample.**

911 **Aperiodic exponent spatial autocorrelation.**

Regressor	Estimate	95% CI	p-value
ADOS CSS	0.00	-0.16 – 0.16	0.990
Age	0.26	-0.01 – 0.53	0.061
Sex	-0.01	-0.40 – 0.39	0.976
Wake	0.31	0.04 – 0.59	**0.026**

912 EEGs n = 184

913 Subjects n = 102

914 *Bold p-values indicate significance at α = 0.05.*

915 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

918 **Supplementary Table 1. Peak Alpha Frequency spatial autocorrelation.**

Dataset	Regressor	Estimate	95% CI	p-value
Discovery sample **n = 204**	ASC	0.01	-0.29 – 0.31	0.948
	Sex (M)	-0.16	-0.54 – 0.21	0.387
	Age	-0.03	-0.18 – 0.12	0.676
HBN **n = 2545**	ASC	-0.70	-1.10 – -0.30	**0.001**
	Sex (M)	0.07	-0.23 – 0.38	0.630
	Age	0.11	0.07 – 0.15	**<0.001**
NIH **n = 932**	ASC	0.05	-0.11 – 0.21	0.544
	Sex (M)	-0.04	-0.18 – 0.13	0.746
	Age	0.02	-0.06 – 0.10	0.605

920 *Bold p-values indicate significance at $\alpha = 0.05$.*

921 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*

927 **Supplementary Table 2. Aperiodic offset spatial autocorrelation.**

Dataset	Regressors	Estimates	95% CI	p-value
Discovery sample **n = 235**	ASC	-0.65	-0.95 – -0.34	**<0.001**
	Sex (M)	-0.08	-0.44 – 0.29	0.679
	Age	0.05	-0.10 – 0.20	0.511
HBN **n = 2587**	ASC	-0.02	-0.36 – 0.31	0.882
	Sex (M)	0.33	0.07 – 0.59	**0.011**
	Age	0.15	0.11 – 0.18	**<0.001**

930 *Bold p-values indicate significance at α = 0.05.*

931 *Spatial autocorrelation was evaluated at a geodesic distance of 53°.*