Seroprevalence of Chagas Disease in Mothers and Children in Postervalle: A

Cross-Sectional Analysis in a Low-Endemic Setting

Authors and affiliations:

Bratriz Amparo Rodríguez-Olguin[1], Daniel F. Lozano Beltrán[2], Isabel Mariscal
Sejas[3], Brandon N. Mercado-Saavedra[4]*

[1] Facultad de Ciencias Farmacéuticas y Bioquímicas, Universidad Autónoma Gabriel
Rene Moreno. Santa Cruz-Bolivia.

[2] Fundación CEADES, Universidad Mayor de San Simón UMSS. Cochabamba,
Bolivia

[3] Hospital de Postervalle. Postervalle, Santa Cruz, Bolivia.

[4] Instituto de Investigación de Medicina, Universidad Católica Boliviana Santa Cruz,
Santa Cruz, Bolivia

*Corresponding author. Email: bmercado.s@ucb.edu.bo

Authors contribution:

BARO contributed to the conceptualization, study design, manuscript drafting, and

coordination of fieldwork. DFLB was responsible for the methodological adaptation

and refinement of the study design. IMS coordinated the sample collection team and

supervised field procedures. BNMS conducted the data analysis, interpreted the

statistical findings, and manuscript drafting.

ABSTRACT:

Chagas disease (CD), caused by *Trypanosoma cruzi*, remains a major public health concern in Latin America, particularly in Bolivia, which reports one of the highest infection burdens worldwide. While congenital transmission has gained importance in many countries, vectorial transmission persists in rural and peri-urban areas. This study aims to assess the epidemiological characteristics and seroprevalence of CD among mothers and their children in Postervalle, a Bolivian community officially classified as having low vector infestation. **Methods:** A cross-sectional study was conducted in July 2023, including 58 mothers and 104 children aged 4 to 17 years. Participants were recruited from the local school and underwent serological screening using three different diagnostic tests. Individuals testing positive on at least two assays were classified as seropositive. Sociodemographic data and exposure history were collected through structured questionnaires. Bivariate analysis was performed to identify associations between seropositivity and risk factors. **Results:** Seroprevalence was 15.8% among mothers and 3.8% among children. All seropositive children were born to seronegative mothers, suggesting likely vectorial rather than congenital transmission. Children who tested positive were more likely to have lived in rural areas and in houses with mud walls (p = 0.046 and p = 0.012, respectively). Among mothers, being bitten by kissing bugs was significantly associated with seropositivity (p = 0.004). No significant differences were found in age or number of children between groups. **Conclusion:** Despite its

classification as a low-infestation area, Postervalle presents evidence of ongoing

vectorial transmission of *T. cruzi*. These findings underscore the need for continued

entomological surveillance and integrated vector control strategies, especially in

vulnerable communities. The absence of congenital cases further highlights the

importance of local vector-related risk factors.

AUTHORS' SUMMARY

Chagas disease is a tropical infection caused by a parasite, known as *Trypanosoma cruzi*. This disease is mostly transmitted by vectors known as kissing bugs, but it can also be transmitted congenitally. In many Latin American countries, including Bolivia, this disease continues to affect large numbers of people, especially in rural areas where the vector is endemic. While some countries have made progress in controlling the vector that carries the parasite, transmission still occurs, often silently. In this study, we worked with mothers and their children in a small Bolivian town called Postervalle, which has been categorized as a low-risk area for the insect that spreads Chagas disease. We tested both the mothers and their children to see if they had been infected by the parasite. Surprisingly, none of the infected children had mothers who tested positive, suggesting they may have been infected by kissing bugs, not through birth. We also found that children who lived in homes with mud walls or rural areas were more likely to be infected. Our findings show that even in places thought to be low-risk, people—especially children—are still getting infected. This highlights the need for ongoing efforts to monitor and prevent Chagas disease, even in areas where it is assumed to be under control.

INTRODUCCION

Chagas disease (CD) is caused by the protozoan *Trypanosoma cruzi* and is prevalent in most countries of Latin America [1–3]. It is estimated that there are currently over 6 million people infected with the parasite and more than 70 million people at risk of getting the infection [1,4]. CD has different modes of transmission, including the vectorial, congenital, and by blood transfusion [1,3]. It is estimated that 1 out of 3 people infected with the parasite might progress to Chagas Cardiomyopathy, one of the chronic manifestations of CD [5–7]. Among these countries, Bolivia is particularly affected, presenting both the vector and congenital transmission, making it a critical context for the study of CD.

CD is a neglected infectious disease in several countries, including Bolivia, which is estimated to have the highest prevalence of the infection in the world [8]. The high prevalence of CD in Bolivia might be due to the presence of the vector in several areas of its territory, occupying over 40% of the communities in the country according to the Bolivian National Program of Chagas disease (BNPCD) [9]. A research study evaluated the prevalence of CD in a rural community in the Gran Chaco, a region highly endemic of the vector in Bolivia, presenting prevalence that ranged from 40-80% [10]. A community-based study performed in Spain in people from Latin American countries described that 40% of their participants came from Bolivia, and 36% of them were seropositives for CD [11].

Recent studies estimate that the prevalence of CD is approximately 12%

among pregnant women in South America as of 2020 [12]. In Bolivia, data from the

BNPCD reported a prevalence of 2.4% in children aged 5 to 15 years and a

congenital transmission rate of 1.4% in 2019 [13]. By 2022, the National Program of

Vector-Borne Diseases (ETVs) estimated a chronic infection prevalence of 19.2%

among individuals over 15 years of age [14]. While vector control programs and

improvements in maternal and child health services contributed to a reported

reduction in Chagas-related mortality to 2% by 2001 [15], subsequent reports

suggesting a decrease in vectorial transmission have not been consistently

supported by updated surveillance data. A 2017 study conducted in Camiri, a

municipality in the Gran Chaco region of Bolivia, confirmed a seroprevalence of

0.22% among children aged 5 to 21 years, with rates increasing with age and a mean

age of seropositivity of 13 years [16]. Similarly, studies in the urban areas of

Cochabamba reported prevalence rates ranging from 19% to 25% among children

aged 5 to 13 years, reinforcing that Chagas disease remains not only a rural but also

an urban public health concern [17].

This study aims to describe the epidemiological characteristics of Chagas disease

among mothers and their children in the community of Postervalle, a locality officially

classified as having a low vector infestation rate by the Bolivian National Chagas

Program [9] . By analyzing seroprevalence patterns and associated risk factors, this

study seeks to provide insight into whether transmission in this community is

primarily vectorial or congenital.

METHODS

Study Population

The study population included 58 mothers with their children (104 in total) ranging from 4 to 17 years old. The recruitment took place in July 2023 in Postervalle municipality, located in the southwest of the department of Santa Cruz, Bolivia. Postervalle has a population of 240000 inhabitants. The selection of the study population began with students from the public school of Postervalle, and the mothers of the students were also recruited to the study. All participants underwent an epidemiological questionnaire.

The study included children aged 4 to 17 and their mothers who gave their informed consent. Participants were from the local area and nearby towns and attended school regularly. Children who declined to participate or were unable to give oral consent were excluded.

Specimen collection

Blood samples were collected from all consenting students and their mothers. The samples were allowed to coagulate and centrifuged at 3500 rpm for clot and serum separation. The serum was used to perform three different serological tests: OnSite Chagas Ab Combo Rapid Test—CTK BIOTECH (USA), Chagatest HAI—Wiener lab (Rosario, Argentina), and Chagatest ELISA recombinant v.3.0—Wiener lab (Rosario, Argentina). Participants who had at least two positive serological tests were considered Chagas seropositive. All Chagas-positive participants (students

132 and their mothers) were referred to the National Program of Chagas in the

133 municipality for follow-up and treatment according to the national regulations.

Figure 1. Location of Postervalle in Santa Cruz, Bolivia

The map of Bolivia shows the location of Postervalle in the department of Santa Cruz,

Statistical Analysis

The data was analyzed using Stata 18.0 (StataCorp, College Station, US). Bivariate

analysis was done using Fisher's exact, Student's T-test, and the Mann-Whitney

test. The bi-variate analysis was performed using the Chagas serological status,

either mothers or children, with other epidemiological variables.

Ethics statement

The research protocol was revised and approved by the Institutional Review Board

(IRB) of the Bolivian Catholic University in Santa Cruz. Each participant provided

written informed consent for themselves and their children, and the children provided

oral informed consent to participate in the project.

RESULTS

Demographic characteristics of mothers

The study included 58 mothers, of whom 84.2% were seronegative for Chagas, and 15.8% were seropositive (Table 1). The mean age of the mothers was 37.39 years (SD±7.31), with no significant difference between seronegative and seropositive groups (p=0.475). The average number of children was slightly higher among seronegative mothers (2.08 ± 0.92) compared to seropositive mothers (1.78±0.44), but the difference was not statistically significant (p=0.335). While 86% of mothers had undergone a Chagas test, a more significant proportion of seropositive mothers (88.9%) had been tested, though this difference was insignificant (p=0.783). A significantly higher percentage of seropositive mothers (44.4%) reported being bitten by kissing bugs than seronegative mothers (8.3%; p=0.004). Additionally, seropositive mothers were more likely to receive Chagas treatment (22.2% vs. 2.1%, p=0.014).

Table 1. Demographic and Exposure Characteristics of Mothers Stratified by Chagas Serology Status (N=58)

| Variables | Serology status for Chagas in mothers | | | |
	Negative (84.2%)	Positive (15.8%)	Total (100%)	p value
Mother's age	37.083 (7.128)	39.000 (8.471)	37.386 (7.306)	0.475!
Number of children	2.083 (0.919)	1.778 (0.441)	2.035 (0.865)	0.335†

Chagas test (yes)*	41 (85.4%)	8 (88.9%)	49 (86.0%)	0.783
Received blood transfusion (yes)*	5 (10.4%)	0 (0.0%)	5 (8.8%)	0.311
Seen kissing bugs (yes)*	7 (14.6%)	3 (33.3%)	10 (17.5%)	0.175
Lived in rural areas (yes)*	18 (37.5%)	1 (11.1%)	19 (33.3%)	0.123
Been bitten by a kissing bug (yes)*	4 (8.3%)	4 (44.4%)	8 (14.0%)	0.004
Received treatment for Chagas (yes)*	1 (2.1%)	2 (22.2%)	3 (5.4%)	0.014
House received fumigation*	31 (64.6%)	8 (88.9%)	39 (68.4%)	0.150
Mud wall (yes)*	13 (27.1%)	2 (22.2%)	15 (26.3%)	0.761

During any pregnancy
! Mann-Whitney test
† Student t-test
All other tests were performed using X2 Fisher's exact test

Chagas Serology in Mothers and Children

Chagas serology testing was performed on 58 mother-child groups. Each mother

was paired with all their children included in the study. As mentioned in Table 1, the

prevalence of Chagas seropositivity among mothers was 15.8%. Table 2 describes

that 6.9% of mothers had seropositive children. All seropositive children are born to

seronegative mothers. Among children of seronegative mothers, 8.2% were

seropositive, though this difference was not statistically significant.

Table 2. Chagas Serology Results of Mothers and Their Children (N=58)

	Mothers' Chagas serology		
	Negative (84.4%)	Positive (15.6%)	Total
Children with Chagas disease			
No	45 (91.8%)	9 (100.0%)	54 (93.1%)
Si	4 (8.2%)	0 (0.0%)	4 (6.9%)

Characteristics of children

A total of 104 children were included in the analysis, of whom 96.2% were seronegative, and 3.8% were seropositive for Chagas (Table 3). Seropositive mothers were more likely to have lived in rural areas (75.0% vs. 28.3%, p=0.046) and to have mud walls in their homes (75.0% vs. 21.0%, p=0.012). While a larger proportion of seropositive mothers reported having received treatment for Chagas (25.0% vs. 1.0%, p<0.001), no significant differences were observed in other variables, including blood transfusion history (p=0.097) and fumigation of houses (p=0.206). The median age of seropositive children was 11 years (IQR: 8–14), compared to 15 years (IQR: 15.5 - 16.5) for seronegative children (Figure 1). Although there was a trend toward younger age in seropositive children, the difference was not statistically significant (Mann-Whitney U test, p=0.083).

Table 3. Characteristics of Children Stratified by Chagas Serology Status (N=104)

	Chagas serology			
	Negative (96.2%)	Positive (3.8%)	Total (100%)	p value
Mother received Chagas test (yes)*	72 (75.8%)	4 (100.0%)	76 (76.8%)	0.261
Mother received blood transfusion (yes)*	5 (5.1%)	1 (25.0%)	6 (5.9%)	0.097

Presence of kissing bugs in the house (yes)*	13 (13.8%)	1 (33.3%)	14 (14.4%)	0.344
Mother lived in a rural area (yes)*	28 (28.3%)	3 (75.0%)	31 (30.1%)	0.046
Mother bitten by a kissing bug (yes)*	9 (9.8%)	1 (33.3%)	10 (10.5%)	0.191
Mother received treatment for Chagas (yes)*	1 (1.0%)	1 (25.0%)	2 (1.9%)	<0.001
Hose received fumigation (yes)*	64 (71.1%)	4 (100.0%)	68 (72.3%)	0.206
House had mud walls (yes)*	21 (21.0%)	3 (75.0%)	24 (23.1%)	0.012

During pregnancy
All tests were performed using X2 Fisher's exact test

Graph 1. Age Distribution of Seropositive and Seronegative Children (N=104)

Mann Whitney test p=0.083

DISCUSSION

Latin American countries have several neglected diseases, such as Chagas disease, with approximately 70 million people at risk of infection. Congenital transmission is the main source of infection in most countries in the region [18,19] .

Bolivia is a country with one of the highest burdens of Chagas disease, where

multiple rural communities in the Chaco region are endemic to the vector [8,9,20] .

The unpublished official data presented by the Bolivian National Chagas program

reports that Postervalle is a low infestation community due to the low infestation rate

of vectors (1.32%) (unpublished data). In this study, we describe the situation of

Chagas disease in mothers and children in Postervalle, with a prevalence of mothers

of 15.8%. This prevalence is similar to that reported by the local authorities, which

estimate that the prevalence is 14.5% (unpublished data). However, a recent meta-

analysis described that the prevalence of pregnant women in Bolivia ranged from

17-65%, with lower rates in the past 5 years [1]. This is similar to a study that reports

that the prevalence among pregnant women at the public maternity in Santa Cruz is

22% [2]. Nevertheless, these results are mainly hospital-based, with women from all

regions (rural and urban) of the department, increasing the estimated prevalence of

each community. One factor that was significantly more present among Chagas

seropositive mothers was being bitten by the vector, where 44% of infected women

reported having been bitten by a kissing bug. However, having lived in rural areas,

lived in houses with mud walls was not significantly more prevalent in women with

Chagas disease. These results are not entirely consistent with other studies

performed in similar settings [3,4], which might be due to the small sample size or

the fact that most of the people included in the study have lived in Postervalle for

their entire lives, presenting similar epidemiological characteristics.

Bolivia joined the Southern Cone Initiative, a 1991 collaborative program with

Argentina, Brazil, Chile, Paraguay, and Uruguay, which aimed at interrupting

Chagas transmission by eliminating *Triatoma infestans* and ensuring safe blood

supplies [5]. While countries like Uruguay, Chile, and Brazil successfully halted

vectorial transmission by the late 1990s and early 2000s, Bolivia achieved

interruption in parts of La Paz and Potosí only by around 2011–2013, though

persistent bug populations and insecticide resistance continue to challenge

complete eradication in regions such as the Gran Chaco [6]. Interestingly, none of

the mothers in the study who tested positive for Chagas disease had seropositive

children, while all seropositive children in the study were born to mothers who tested

negative. This finding suggests that, in the community of Post

Postervalle, congenital transmission may be absent or minimal, and instead,

vectorial transmission is the likely source of infection among children. Although

maternal screening was conducted at the time of this study, it is possible that some

mothers may have seroconverted after giving birth or were false negatives due to

limitations in the sensitivity of serological tests during acute or early chronic stages

of infection. Alternatively, transmission could have occurred postnatally—after the

neonatal screening window—through direct vector contact during early childhood,

as all seropositive children were older than 8 years. This epidemiological pattern

aligns with other studies conducted in rural or peri-urban areas of Bolivia with similar

vector infestation classifications. For instance, Hopkins et al. (2019) reported

persistent *T. cruzi* transmission in school-aged children in the Bolivian Chaco, even

in areas under surveillance, with infection rates increasing with age and suggesting

ongoing vector exposure in domestic settings [16]. In another low-endemic setting in

Cochabamba, Medrano-Mercado et al. (2008) found seroprevalence rates in

children as high as 25%, reinforcing that low infestation classifications may not fully

reflect the true risk of transmission, particularly in settings with structural

vulnerabilities such as adobe walls and a lack of sustained vector control [17].

Furthermore, this result is consistent with the data reported by the local authorities

that estimate that the vertical transmission rate in Postervalle is 0% (unpublished

data). An additional concern is the potential impact of insecticide resistance in

Triatoma infestans, the principal vector species in Bolivia. Studies have documented

emerging resistance to pyrethroid insecticides in multiple regions of the Gran Chaco,

compromising the long-term effectiveness of vector control programs [21]. If such

resistance is present in Postervalle, even at low levels of infestation, vectorial

transmission may persist or resurge despite formal classification as a "low-risk" area.

This highlights the importance of integrating entomological surveillance with

serological screening and community-level epidemiological studies. Finally, the

characteristics associated with seropositivity in children included whether their

mothers had seen or been bitten by kissing bugs, and whether they had lived in

houses with mud walls—environments that likely reflect the domestic setting in which

the children were born and spent their early childhood. These conditions represent

key risk factors that facilitate vectorial transmission of *Trypanosoma cruzi* [4,6,7].

This study has several limitations, including a relatively small sample size, the

use of non-probabilistic sampling, and the fact that not all children of the participating

mothers were included. Participant recruitment was conducted primarily through the

local school, where most of the mothers were actively engaged. Although mothers

whose children did not attend the school may have been underrepresented, it is

unlikely that this introduced a significant selection bias, as there are no known

sociodemographic or epidemiological differences between mothers with school-

attending children and those whose children do not attend. While the overall sample

size maintained a statistical power above 80%, the limited number of individuals

within certain subcategories reduced the ability to detect statistically significant

associations. Finally, not all children of the 58 participating mothers were enrolled,

primarily due to the absence of verbal consent. Although this may influence the

observed prevalence of Chagas disease in children, it is unlikely to have significantly

impacted the general epidemiological patterns, as non-participating children are

presumed to share similar risk profiles with their siblings.

These findings highlight that, even in areas officially classified as low

infestation zones, vectorial transmission of *Trypanosoma cruzi* persists as a route of

infection among children. While Bolivia has made substantial progress under the

Southern Cone Initiative, particularly in selected regions, the continued presence of

infected vectors and active transmission in communities like Postervalle

underscores the fragility of these gains. Moreover, the overuse of insecticides—

potentially leading to vector resistance—and the broader ecological impacts of

climate change may further hinder vector control efforts, threatening to reverse

recent advances and sustain transmission in vulnerable rural populations [8].

ACKNOWLEDGEMENTS

To Drs. Freddy A. Tinajeros and Nayrha Villazón, Lizeth Gil, Mayra Morales for their

contribution during different steps of this project. To Drs Benjamin Quiroga, Célida

Montaño Gutiérrez, Judith Aguilar from the Chagas Program of Santa Cruz for their

support during this project. We also want to thank to the community of Postervalle

who received us, and all the participants of the study.

303
304
305   **REFERENCES**
306
307

1. Hochberg NS, Montgomery SP. Chagas Disease. Ann Intern Med. 2023;176:

ITC17–ITC32. doi:10.7326/aitc202302210

2. Rassi A, Rassi A, Marin-Neto JA. Chagas disease. Lancet (London, England).

2010;375: 1388–402. doi:10.1016/s0140-6736(10)60061-x

3. Bern C. Chagas' Disease. The New England journal of medicine. 2015;373:

456–66. doi:10.1056/nejmra1410150

4. WHO. Chagas disease (American trypanosomiasis). Available:

https://www.who.int/health-topics/chagas-disease#tab=tab_1

5. Saraiva RM, Meymandi S. Chagas Disease, A Neglected Tropical Disease.

2020; 145–162. doi:10.1007/978-3-030-44054-1_9

6. Echavarría NG, Echeverría LE, Stewart M, Gallego C, Saldarriaga C. Chagas

Disease: Chronic Chagas Cardiomyopathy. Current Problems in Cardiology.

2019;46: 100507. doi:10.1016/j.cpcardiol.2019.100507

7. Jr AR, Neto JAM, Rassi A. Chronic Chagas cardiomyopathy: a review of the

main pathogenic mechanisms and the efficacy of aetiological treatment

following the BENznidazole Evaluation for Interrupting Trypanosomiasis

(BENEFIT) trial. Memórias do Instituto Oswaldo Cruz. 2017;112: 224–235.

doi:10.1590/0074-02760160334

8. Pinazo M-J, Pinto J, Ortiz L, Sánchez J, García W, Saravia R, et al. A strategy

for scaling up access to comprehensive care in adults with Chagas disease

in endemic countries: The Bolivian Chagas Platform. PLoS Neglected Trop

Dis. 2017;11: e0005770. doi:10.1371/journal.pntd.0005770

9. Deportes M de S y. Tasa de infestación de Chagas en Bolivia 2013. Available:

geo.gob.bo

10. Chippaux J-P, Postigo JR, Santalla JA, Schneider D, Brutus L.

Epidemiological evaluation of Chagas disease in a rural area of southern

Bolivia. Transactions of the Royal Society of Tropical Medicine and Hygiene.

2008;102: 578–584. doi:10.1016/j.trstmh.2008.03.008

11. Ramos-Sesma V, Navarro M, Llenas-García J, Gil-Anguita C, Torrus-

Tendero D, Wikman-Jorgensen P, et al. Community-based screening of

Chagas disease among Latin American migrants in a non-endemic country:

an   observational   study.   Infect   Dis   Poverty.   2021;10:   117.

doi:10.1186/s40249-021-00897-2

12. Santana KH, Oliveira LGR, Castro DB de, Pereira M. Epidemiology of Chagas

disease in pregnant women and congenital transmission of Trypanosoma

cruzi in the Americas: systematic review and meta-analysis. Trop Med Int

Heal. 2020;25: 752–763. doi:10.1111/tmi.13398

13. PAHO. Evaluación Internacional de la situación epidemiológica y de control

de la enfermedad de Chagas en 44 municipios de los departamentos La Paz

y Potosí, previamente validados; y en 19 municipios de los departamentos

Cochabamba, Chuquisaca y Tarija, candidatos a validación. 2019. Available:

https://www.paho.org/sites/default/files/2020-01/Informe-evaluacion-

chagasBolivia-Julio2019.pdf

14. Deportes M de S y. PLAN NACIONAL DE ACCIÓN DE ENTOMOLOGÍA Y

MANEJOINTEGRADO DE VECTORESBOLIVIA 2023-2025. 2023.

Available: https://www.minsalud.gob.bo/8022-bolivia-presenta-el-plan-

nacional-de-accion-de-entomologia-y-manejo-integrado-de-vectores

15. Carlier Y, Truyens C. Congenital Chagas disease as an ecological model of

interactions between Trypanosoma cruzi parasites, pregnant women,

placenta and fetuses. Acta Trop. 2015;151: 103–115.

doi:10.1016/j.actatropica.2015.07.016

16. Hopkins T, Gonçalves R, Mamani J, Courtenay O, Bern C. Chagas disease

in the Bolivian Chaco: Persistent transmission indicated by childhood

seroscreening study. Int J Infect Dis. 2019;86: 175–177.

doi:10.1016/j.ijid.2019.07.020

17. Medrano-Mercado N, Ugarte-Fernandez R, Butrón V, Uber-Busek S, Guerra

H, Araújo-Jorge TC de, et al. Urban transmission of Chagas disease in

Cochabamba, Bolivia. Memórias Inst Oswaldo Cruz. 2008;103: 423–430.

doi:10.1590/s0074-02762008000500003

18. Buekens P, Cafferata ML, Alger J, Althabe F, Belizán JM, Bustamante N, et

al. Congenital Transmission of Trypanosoma cruzi in Argentina, Honduras,

and Mexico: An Observational Prospective Study. Am J Trop Med Hyg.

2018;98: 478–485. doi:10.4269/ajtmh.17-0516

19. Danesi E, Codebó MO, Sosa-Estani S. [Congenital transmission of

Trypanosoma cruzi. Argentina 2002-2014]. Medicina. 2019;79: 81–89.

20. Torrico RA. Conocimientos actuales sobre la epidemiología de la enfermedad

de Chagas en Bolivia. Available: https://iris.paho.org/handle/10665.2/11775

21. Mougabure-Cueto G, Picollo MI. Insecticide resistance in vector Chagas

disease: Evolution, mechanisms and management. Acta Trop. 2015;149: 70–

85. doi:10.1016/j.actatropica.2015.05.014

Figure 1

Figure 1