Machine learning-enhanced behavioural approach to detecting reactions to sound in infants and toddlers: proof-of-concept study

Chelsea M. Blankenship[1,2], Josef Schlittenlacher[3,4], Iain R. Jackson[3], Anisa S. Visram[3],

Kevin J. Munro[3,5], Lisa L. Hunter[1,2], David R. Moore[1,2,3]

[1] Division of Patient Services Research, Cincinnati Children's Hospital Medical Center,

Cincinnati, Ohio, USA

[2] Departments of Otolaryngology and Pediatrics, University of Cincinnati College of

Medicine, Cincinnati, Ohio, USA.

[3] Manchester Centre for Audiology and Deafness, University of Manchester, Manchester, UK

[4] Department of Speech, Hearing and Phonetic Sciences, University College London,

London, UK

[5] Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health

Science Centre, UK

Institutional Review Board Approval: This study was approved by the Cincinnati

Children's Hospital Medical Center Institutional Review Board (PI: Parikh, IRB#:2016-

4483).

Funding: Supported by NIH/NIDCD grant R01DC018734 (MPI: Hunter and Vannest),

NIHR Manchester Biomedical Research Centre (NIHR BRC-20007), William Demant

Foundation, and the Translational Science Award Program Grant 5UL1TR001425-04 (Center

for Clinical and Translational Science Training at the University of Cincinnati). Munro is

supported by an NIHR Senior Investigator award.

Corresponding Author:

Email addresses: JS: j.schlittenlacher@ucl.ac.uk

<u>Keywords:</u>

artificial intelligence, machine learning, infant, hearing, visual reinforcement audiometry,

behavioural response, behavioural observation, proof-of-concept, sound detection

Abstract

Objective: Show that a basic unsupervised machine learning (ML) algorithm can give

information on the direction of child and infant reactions to sound using non-identifiable

video-recorded facial features.

Design: Infants and toddlers were presented warble tones or single-syllable utterances 45

degrees to the left or right. A camera recorded their reactions, from which features like head

turns or eye gaze were extracted with OpenFace. Three clusters were formed using

Expectation Maximization on 80% of the toddler data. The remaining 20% and all infant data

were used to verify if the clusters represent groups for sound presentations to the left, to the

right, and both directions.

Study Sample: 28 infants (2-5 months) and 30 toddlers (2-4 years), born preterm (<32 weeks

gestational age) were presented ten sounds each.

Results: The largest cluster comprised 90% of the trials with sound presentations in both

directions, indicating "no decision." The remaining two clusters could be interpreted to

represent reactions to the left and the right, respectively, and average sensitivities of 96% for

the toddlers and 68% for the infants.

Conclusions: A simple machine learning algorithm demonstrated that it can form correct

decisions on the direction of sound presentation using non-identifiable facial behavioural

data.

Abbreviations

Artificial Intelligence (AI)

Behavioural Observation Audiometry (BOA)

Conditioned Play Audiometry (CPA)

Expectation Maximization (EM)

Machine Learning (ML)

Neonatal Intensive Care Unit (NICU)

Observer Psychoacoustic Procedure (OPP)

Principal Component Analysis (PCA)

Visual Reinforcement Audiometry (VRA)

1. Introduction

Congenital hearing loss is among the most common conditions at birth that result in

adverse neurodevelopmental outcomes, including language, academic, and social skill

deficits (Moeller et al., 2007; Yoshinaga-Itano et al., 2021). Infants at highest risk for hearing

loss are those born very prematurely who are cared for in neonatal intensive care units

(NICU) (Beswick et al., 2012; Hirvonen et al., 2018; Hunter et al., 2023). Fortunately,

congenital hearing loss is effectively addressed through universal newborn hearing screening

programs, if diagnosis and intervention are provided within the first few months after birth

(Ching et al., 2017; Sininger et al., 2009; Tomblin et al., 2015; Yoshinaga-Itano et al., 1998).

1.1 Paediatric Behavioural Hearing Assessment Methods

Behavioural assessment of infants' responses to sound is crucial to validate

electrophysiologic hearing assessments and determine the need for, and effectiveness of,

auditory intervention. Based on a large study of developmental milestones, infants alert to

sound by the age of 1 week, and orient towards sounds by 4 months (Capute et al., 1986).

Behavioural observation of subtle infant responses to sounds have long been used for

behavioural observation audiometry (BOA) (Bench et al., 1976a, 1976b, 1976c). BOA is an

unconditioned procedure that relies on behavioural response from the infant, such as changes

in sucking, eye shifts, forehead wrinkling or head orientation, in response to sound. However,

this behavioural observation method requires a highly trained observer, and success is

dependent on the child's age, developmental level, attention span, and cooperation (Widen &

Keener, 2003). BOA has high inter- and intrasubject variability in responses and rapid

response habituation (Hicks et al., 2000) and it is difficult to control for observer bias in a

clinical setting (Widen, 1993). Therefore responses obtained with BOA are not considered

suitable or reliable enough for clinical use (Ruth et al., 1983).

For research purposes, an observer-based psychoacoustic procedure (OPP) was

developed that requires highly trained human observers and specialized software (Olsho et

al., 1987; Werner & Marean, 1991). OPP controls observer bias by providing auditory

masking of the observer, and a two-alternative forced choice response paradigm with

psychoacoustic tracking, allowing testing in very young infants that are not yet

developmentally ready for VRA testing (Olsho et al., 1988). However, despite the use of

trained observers, many very young infants do not respond with adequate reliability, and they

have poorer thresholds than children and adults, either due to inattention or different listening

strategies. Thus, OPP is generally unsuitable for clinical application.

From around a developmental age of 6-7 months, infants start to reliably respond to

sound input with reinforced head turns. At this age, visual reinforcement audiometry (VRA)

is commonly used to diagnose hearing loss and track intervention outcomes (Widen &

Keener, 2003). VRA involves conditioning the child to make a head turn response following

the presentation of an auditory stimulus (detection not localization task). Most commonly,

this involves the infant looking towards a visual target such as a toy or video monitor, but

immature or subtle responses can also include eye shifts/movements and stilling or quieting.

VRA is generally recommended for infants and young children between 6 months to 2.5

years of age, since some older children may lose interest in the reinforcement or are too

active and need a task to maintain attention. Despite its widespread use, VRA has limitations:

responses can be influenced by the child's developmental level, fatigue, or distractibility.

Additionally, false positives may occur if the child anticipates reinforcement rather than

responding to the auditory stimulus, and testing can be challenging in children with motor

delays, limited visual attention, or visual impairments. Lastly, VRA typically involves two

trained individuals making it more resource-intensive which could present as a logistical

challenge in busy clinical settings or small clinics with limited personnel.

Conditioned Play Audiometry (CPA) is a behavioural hearing test typically used for

children between approximately 2.5 and 5 years of age, when they are developmentally ready

to engage in structured play tasks. CPA involves conditioning the child to perform simple

task like stacking rings, building towers, etc., every time they hear an auditory stimulus.

Transforming the hearing test into a game improves attention and motivation resulting in

more reliable threshold information across a wider range of test frequencies compared to

VRA (Thompson & Weber, 1974; Thompson & Thompson, 1972). However, successful CPA

requires that the child understands the task, has the control to wait for the stimulus before

responding, which can be challenging for children with developmental delays, attention

difficulties, or limited motor skills.

In summary, while there are several behavioural methods that are used clinically to

assess hearing in infants and young children, they each have limitations and challenges, and

success is highly dependent on the child's temperament, attention, and ability to engage with

the task. Additionally, many children with hearing loss have comorbid conditions such as

developmental delays or neurocognitive disorders, which make it difficult to complete

behavioural assessments. Consequently, assessing hearing in very young infants and in

difficult-to-test populations remains a significant clinical challenge. To address these gaps

and improve diagnostic accuracy, it is essential to explore objective approaches for hearing

assessments that don't rely on observers' subjective interpretation of behavioural responses.

1.2 Artificial Intelligence and Auditory Research

The use of artificial intelligence (AI) in medical settings is becoming increasingly

common and is rapidly transforming healthcare delivery and medical research. AI is a broad

term that encompasses several subsets, including machine learning (ML; system learns from

data and makes predictions without explicit programming), deep learning (a subset of ML

that uses multi-layered artificial neural networks) and, more recently, generative AI (develops

new content such as text, audio, pictures., etc.) and large language models. Furthermore, it's

well established that AI has the potential to improve medical care by improving diagnostic

accuracy, increasing treatment efficiency, and ultimately optimizing patient outcomes (Chinta

et al., 2025).

Despite the advantages of AI, its use in healthcare and research presents inherent

challenges related. First, many ML techniques require large datasets—often tens of thousands

of measures for basic algorithms and millions for complex image recognition tasks (Deng et

al., 2009; Krizhevsky, 2009). In paediatric hearing research, such datasets are unavailable and

would require extensive manual labelling by human observers. Second, ethical considerations

around data privacy are significant. High-performing, end-to-end deep learning models

typically rely on raw images, which cannot be fully anonymized because facial features and

other identifiers remain embedded in the data. This raises concerns about compliance with

privacy regulations and participant confidentiality. To mitigate these risks, studies should

prioritize storing only low-dimensional, non-identifiable representations of responses rather

than raw images.

OpenFace (Baltrusaitis et al., 2019; Baltrušaitis et al., 2015) is an open-source

software toolkit for analysing video data. It can be used to derive standardised behavioural

features from faces detected in video recordings, including three-dimensional head position,

head rotation across three different axes, eye gaze direction, and activation of facial action

units (e.g., eyebrow raising). In contrast to facial landmarks that encode where specific parts

of the face are located, these variables are less likely to allow identification of a child.

Additional processes, such as reducing the data with a principal component analysis (PCA),

can further ensure privacy. Future work should focus on approaches that use a limited set of

anonymized behavioural indicators, while building justification for collecting large, clinically

relevant datasets that could later support more sophisticated ML solutions, including for

example deep neural networks, for clinical implementation.

Within the field of audiology, there has been a recent influx of publications using AI

approaches (Frosolini et al., 2024). In a systematic review of AI in audiology, Frosolini and

colleagues identified 104 publications since 1990, with the majority (87.5%) published after

2020. Most publications focused on therapeutic and prognostic applications (34.6%), hearing

assessment (33.7%), unique applications and exploratory research (21.1%), temporal bone

radiology (5.8%) and vestibular diagnosis (5.8%). With regard to hearing assessments,

manuscripts were published on a variety of topics ranging from hearing screening and

diagnostic assessment (Gathman et al., 2023; Liu et al., 2021; Xie et al., 2025), measurement

of middle ear function and diagnosis of otitis media (Peng et al., 2025; Zeng et al., 2022),

hearing preservation (Zeitler et al., 2024) and post-implantation programming (Wathour,

Govaerts, Derue, et al., 2023; Wathour, Govaerts, Lacroix, et al., 2023). However, most of

the auditory AI research focusses on adults and older children. Infant AI literature primarily

focuses on classifying middle ear function and analysis of electrophysiologic assessments

including interpretation of auditory brainstem responses (McKearney et al., 2022),

categorisation of neural responses in infants to drumbeats versus repeated syllables (Gibbon

et al., 2021) and to pitch processing in neonates (Hart & Jeng, 2021). Currently, there are

few publications using AI approaches to detect behavioural responses to sound in young

children (Xie et al., 2025), none of which are in infants. Behavioural hearing assessments in

this population have inherent challenges, creating a critical need for alternative approaches.

Leveraging AI and machine learning (ML) may offer a solution for objectively measuring

sound detection in infants. If successful, it has the potential to improve infant assessment

methods which would ultimately impact diagnosis, guide treatment and improve outcomes.

1.3 Study Purpose and Research Questions

This paper presents an overview of our initial efforts to measure sound detection in

infants and toddlers using ML to identify eye, facial and body movements in response to

tonal and speech sounds. In this approach, a video camera replaces the human observer, and

an algorithm trained on validated data determines whether observed movements indicate a

reaction to sound. Specifically, we examine optical scanning of the infant's face and torso

combined with ML to monitor responses to sounds, including subtle cues a human observer

might not reliably detect. Therefore, the overall purpose of this study was to investigate

whether a behavioural observation-like auditory task could be trained with ML methods to

learn when an infant shifted eye gaze, turned their head, or reacted with facial expressions,

such as eye widening or eyebrow raising, in response to sounds. The study included a cohort

of infants (2-5 months) and toddlers (2-4 years) who were born preterm and underwent

hearing assessments on the same day as the ML protocol. In this proof-of-concept study, the

primary research questions were: (1) can ML be used to cluster responses based on head and

facial changes of infants and toddlers to presented sound stimuli? (2) can the trained

algorithm accurately detect the direction of sound presentation when infants and toddlers

respond to sound?

2. Method

2.1. Study Design and Participants

Participants were infants and toddlers born very or extremely preterm (< 32 weeks

gestational age or less) recruited soon after birth from NICUs in Cincinnati, Ohio, United

States and were enrolled in a longitudinal project to improve early detection of speech and

language delays. This population is of research interest due to their higher risk for hearing

loss and known difficulties with behavioural assessment due to more prevalent

developmental delay. Infants with known chromosomal abnormalities, or congenital

conditions affecting the central nervous system were excluded. All families were primarily

English-speaking and agreed to have their child video and audio recorded.

Participants included 58 children split into an infant group (n = 28, mean = 3.0 mos.,

range = 2.1 - 4.6 mos.) and a toddler group (n = 30, mean = 29.8 mos., range = 23.9 - 41.7

mos.). All ages are reported corrected for gestational age. All participants were seen once for

the ML study in conjunction with auditory brainstem response testing (infants) or age-

appropriate behavioural audiometry (toddlers). Infants had normal hearing in both ears as

determined with 1 kHz chirp-evoked auditory brainstem response thresholds (≤ 20 dB

eHL)(Sininger et al., 2020). For toddlers, hearing status was based on VRA or conditioned

play audiometry (CPA) thresholds, using earphones or sound field speakers with live speech

and pure tone stimuli (1, 2, 4, and 8 kHz)(Hunter et al., 2023). If only sound field testing was

completed, ear specific hearing status was determined through a combination of 226-Hz

tympanometry, distortion product otoacoustic emissions (2-10 kHz) measured at the same

visit along with repeated behavioural hearing assessment at follow-up study visits. Toddlers

had normal hearing in both ears (n=29, ≤ 20 dB HL), except one toddler with mild

conductive hearing loss (thresholds at 30 to 35 dB HL). The study was approved by the

Institutional Review Board of Cincinnati Children's Hospital Medical Center. Informed

consent was obtained from the caregiver of all children prior to enrolment.

2.2. Apparatus

All testing was completed in a sound-treated booth using a custom Python program.

The auditory stimuli were amplified (Crown D-75A, Harman) and presented through one of

two sound field speakers (Control 1Xtreme, JBL). Sound field speakers were positioned at 45

degrees to the left and right at ear level at approximately 1 meter from the infant/toddler. A

computer monitor was positioned directly in front of the child, at the midpoint between the

two sound field speakers. To record the child's responses, a video camera (AF/FHD1080P,

FUVISION) was positioned underneath the monitor (see Figure 1).

273   --- Insert Figure 1 here ---

275   *2.3. Stimuli*

276   The children were presented five different auditory stimuli including frequency-

modulated warble tones (duration of 500 msec, 5 Hz modulation rate, 5 % modulation depth,

carrier frequencies: 0.5, 1, 2 kHz), and utterances ("ba" and "da") through the right or left

sound field speakers at a level of 60 dB SPL, approximating conversational speech levels.

Each stimulus was presented twice in a randomized order, from either the left or right

speaker. The stimuli were calibrated using a Larson Davis System 824 sound level meter

(Depew, New York) and Brüel & Kjær ½" microphone.

2.4. Procedure

285   Toddlers sat in a highchair or, if restless, on their caregiver's lap while infants were

always positioned in their caregiver's lap (see Fig. 1). If the caregiver was in the video

recording space, they wore a face mask covering their nose and mouth to ensure the analysis

software did not detect and include their face in the collected data. Each child was tested in a

single session that started with an auditory visual primer: a black-and-white smiling face

displayed on the computer monitor while the phrase "Hello Baby" was played through both

sound field speakers to centre the child's gaze forward. For infants, a slowly rotating shape

with primary colours was presented throughout the test session to help maintain the infants'

attention. For the toddlers, a slowly moving colourful scene was presented throughout after

the auditory primer. The test assistant (seated inside the booth next to the computer monitor)

re-centred the child if needed between stimuli, ensured the child was upright, and could see

the computer monitor. When the child was centred and quiet, the test assistant manually

triggered the presentation of each sound (speech or tone) using a keypress. Responses to

sounds were not rewarded, as would be done in a typical clinical reinforced procedure. This

simplified approach was used to determine whether ML could detect spontaneous infant and

toddler responses to sound, rather than trained responses and reinforced head turns. This was

necessary for the proof-of-concept that naturalistic responses other than head turns could be

detected. The inter-stimulus interval was set to a minimum of 6 seconds. In some cases, a

longer interval was needed to re-centre the child between presentations. In this protocol, test

sessions lasted 114 seconds on average (SD = 27 sec; range = 81 - 287). All participants

included in the analysis successfully completed the test protocol in one session.

2.5. Data preprocessing with OpenFace

To protect participant privacy and avoid sharing raw images or videos for machine

learning analysis, all video data were stored internally. Each video image was processed

using OpenFace (Baltrušaitis et al., 2019), resulting in 55 standardised features per frame of

video. These features included estimates on the degree of activation of 35 facial action units

(Baltrušaitis et al., 2015), for example whether a part of an eyebrow is raised and by how

much. The remaining features encoded the gaze (8 features: x, y, and z directions for each

eye separately, and x and y angles each averaged across the two eyes (Wood et al., 2015), the

location and rotation of the head (6 features: 3 coordinates and 3 angles, one for each spatial

dimension/axis) and 6 features for the head shape. After an analysis of the reaction time to

stimulus onsets (see section Results), these 55 features were stored for a single frame of the

video, 2 seconds after stimulus onset. For this dataset, a maximum of 580 cases were

obtained (58 participants x 10 stimulus presentations per participant). However, for 20 cases

no data were available for 2 seconds after stimulus onset (2 infant cases, 18 toddler cases;

i.e., no behavioural response following sound presentation). Therefore, a total of 560 cases or

sets of features (278 infant cases, 282 toddler cases) were included in the analysis.

2.6. Clustering with Expectation Maximization (EM)

325   The need for annotation of the data for the training of an algorithm was avoided by

326 using unsupervised ML and verifying it via the direction of sound presentations.

327 Unsupervised clustering ML algorithms learn structures in the data, without the need for

328 labels, and allocate each measure to a cluster. The algorithm was trained to recognize three

329 clusters with facial features as the defining measure. The meaning of the clusters thus formed

330 was verified by matching the three predicted clusters to reactions to sound presentations from

331 the left, the right, or no decision for either direction.

332   Expectation maximization (EM, (Dempster et al., 1977) was used to model the data

333 by multivariate Gaussian distributions. Thus, each cluster is represented by its mean and

334 spread. EM not only learns the spread of the data in each dimension (e.g., head turn and eye

335 gaze across the vertical axis), but also covariances between these dimensions. For example,

336 turning the head to the right/left is likely to be correlated with a gaze to the right/left, and thus

337 would not give much additional information. EM updates its belief about the parameters

338 iteratively and learns on its own to match these distributions to the collected data. In brief, it

339 starts with initial means and covariances for each cluster and then assigns all data points to a

340 cluster (the expectation step). Thereafter, it updates the means and covariances of each cluster

341 based on the data that are presently predicted to belong to that cluster (the maximization

342 step). These two steps are repeated until the parameters converge.

343   The algorithm was trained on 80% (n=226) of the participants from the toddler group.

344 The remaining 20% (n=56) of the toddler group and all observations of the infant group

345 (n=278) were used as validation sets. The reason for using only toddler data for training was

346 because we expected more reliable responses from them. In particular, the infants might only

347 be able to perform a subset of the actions that toddlers can or not turn their heads at all.

348   EM was applied to four different sets of two features each: (1) Rotation of the head

349 along the vertical axis (yaw, turning left or right) and x-axis (pitch, nodding), (2) Gaze

350 direction in world coordinates of the left eye to the left/right and up/down, (3) Gaze angle

351 averaged across both eyes to the left/right and up/down, (4) the first two components of a

352 principal component analysis of all 55 features. For analyses 1 to 3, it could be expected that

353 the first variable that encodes looking left and right is most descriptive on reactions to the

sound presentations from the left or right, but the second variable could potentially give

general information about a reaction in any direction as well.

EM was performed using the R EMCluster package (Chen & Maitra, 2015). Since the

results of the EM algorithm depend on a random initialization, the following procedure was

followed to ensure replicable and consistent results: First, the algorithm was performed five

times, and the best performing solution was chosen (using the function "exhaust.EM") for the

initialization of an EM using the function "shortemcluster" and a convergent tolerance of

0.0001. Except for the analysis on the PCA, this was repeated 100 times or until the

maximum of all mean coordinates was at least half as large as the minimum and of opposite

sign, and vice versa. This was done to encourage the algorithm to have two clusters that are

of opposite sign, even though we did not enforce this to be in a particular or the same

dimension. After this, the function "emcluster" was run with default parameters to determine

the three clusters.

3. Results

3.1. Reaction Time and Absolute Head Angle Analysis

To determine if and when the children reacted to stimuli, the absolute angle of the

head looking left or right was analysed as a function of time after stimulus onset. Figure 2

shows that immediately before stimulus onset, both toddler and infant head angle was about

10 degrees off-centre on average (mean of absolute values). The toddler group showed a

maximum reaction 2 to 3 sec after stimulus onset, with an average head turn of 17 degrees

(0.30 radians), while the infant group only slightly or rarely turned their head, to about 12

degrees (0.21 rad) on average, also at around 2 seconds. In light of these results, we decided

to analyse the data from a single frame 2 seconds after stimulus onset and to use the toddler

group, with their stronger head reactions, for training the EM algorithm.

380   --- Insert Figure 2 here ---

3.2. Expectation Maximization Analysis

EM was applied to four different sets of two features each: (1) Rotation of the head

along the vertical axis (yaw, turning left or right) and x-axis (pitch, nodding), (2) Gaze

direction in world coordinates of the left eye to the left/right and up/down, (3) Gaze angle

averaged across both eyes to the left/right and up/down, (4) the first two components of a

principal component analysis computed from all 55 features. Head turn results for all trials

(Fig. 3) showed that most trials did not result in an observable head turn response. Most data

points were around the origin, indicating lack of head turns, no gaze in any particular

direction, nor any clear reaction that is reflected in the first two principal components. The

gaze for both eyes is very similar to the results for the left eye only, however, not all data

points are identical, indicating some gaze movements where left and right were not the same.

About 90% of all cases are close to the origin, leaving 10% where the algorithm makes a

decision. However, these data reflect a single frame only (2 seconds after stimulus onset), and

for proof-of-concept it was more important to show that a decision was correct when the

algorithm made one.

398   --- Insert Figure 3 here ---

Table 1 shows the number of predictions for each feature set, class (i.e., predicted

direction) and data set (training vs. validation) separated on whether a sound was presented

from the left or right speaker. This enabled us to calculate the specificity for each feature set.

For the head rotation feature set, the predictive decisions were made for six out of 56 cases

(ie., left or right predictive class, not centre) for the toddler validation group. Out of the six

cases where a decision was made, five of these head turns were in the expected direction (i.e.,

sound presented from left speaker and the predictive class was left and vice versa). For the

infant validation group, 13 of 24 head turns to the left followed a sound presentation from the

left speaker and 31 of 52 head turns to the right followed a sound presentation from the right

speaker. This results in a specificity of 58%. Using a one-sided binomial test of statistical

significance (i.e., assessing the probability that at least 44 of 76 reactions are correct when

guessing at 50%), it was not statistically significantly greater than a guessing rate of 50%, $p =$

0.07. The validation set of the toddlers contains too little data for meaningful statistical tests,

though 5 of 6 decisions (83%) were in the expected direction.

For gaze direction of the left eye, the toddler validation set showed a decision was

made in 4 of the 56 cases and all 4 of those predictions were in the expected direction. For

the infant group, a decision was made in 20 out of 278 cases of which 16 predictions were in

the expected direction, resulting in a specificity of 80% and statistical significance, $p = 0.006$.

Very similar results were found for gaze averaged across both eyes (Figure 3, bottom left

panel). All 6 reactions of the toddler validation set were in the expected direction (100%) and

so were 32 of the 48 reactions of the infant group, constituting a specificity of 67% and

statistical significance ($p < 0.001$).

PCA captured features, or combinations of features other than head turns and gaze.

Table 1 shows the performance for EM based on the first two components. For the toddler

validation set, only 6 predictions were made, though all of them were in the expected

direction (100%). For the infant validation group, 25 of 38 predictions were in the expected

direction resulting in a specificity of 66% ($p = 0.04$).

428   --- Insert Table 1 here ---

A human observer would base a decision on large or easily observed behavioural

responses like head turns and less reliably so on subtle features like eye movements.

Combining features can be a strength of more complex ML algorithms. We investigated the

potential to increase sensitivity by comparing the agreement between predictions based on

head movement with predictions based on the gaze averaged across both eyes. For the

toddlers, in four cases a decision was made by both head movement and gaze, and they

agreed in direction. There were two cases for each head movements and gaze where only one

of them made a decision. Thus, there were eight cases in total where a decision was based on

head movement or gaze, compared to six cases each when it was based on only one of them.

For the infants, combining the 76 reactions based on head turns and 48 reactions based on

gaze direction resulted only in a modest increase to 82 decisions. Of the 42 cases where both

features led to a decision, 39 predictions agreed in direction. The three cases where head

movement and gaze differed were correctly predicted by the gaze, and all three followed

sound presentations to the left.

4. Discussion

The ML approach used here was naïve to the nature and direction of the stimuli (left

vs. right speaker) and type of response yet was able to train an algorithm based on toddler

responses to detect gaze direction and angle. In infants, head turns were detected in 27% of

cases, but only 58% of those turns were toward the direction of the stimulus presentation. It

was somewhat higher, 67% to 80%, when predictions were based on eye gaze, however a

decision could be made in only 7% to 17% of cases. Because young infants do not generally

make clear head turns, an ML approach based on a variety of facial features may be helpful

during the first few months when early hearing diagnosis and initiation of rehabilitation is

critically important.

The EM algorithm formed clusters without knowledge to which side the stimuli were

presented. Nonetheless, three clusters always formed, one central, and one to each of the left

and right, with the left and right clusters showing about equal probability, as expected. This

was also true for the PCA data, which included additional information about facial

movement. It therefore appears possible to obtain meaningful lateralization of the

behavioural response to sound from the data alone.

For PCA analysis, a decision was made for 11% of the toddler validation group and

14% of the infant validation group. In these cases, the PCA results showed high specificity,

with all six predictions correct in the toddler group and 66% for the infant group. However,

this does not constitute an improvement compared to using gaze alone. PCA is a rather

simple and linear approach to combine features. It remains to be seen if more complex

approaches based on bigger data sets, based on 1,000 or more participants, and supervised

combination of features may offer improved performance. The high specificity of only two

components of the PCA, which were formed from all 55 features in an unsupervised way, is

promising for such an approach. The lowest accuracy for the infant group was achieved when

analysing head turns, which indicates that infants show fewer head turns than the toddlers, on

whom the algorithm was trained.

A significant drawback in the present analysis is the substantial proportion of data for

which no decision was made on whether the child looked to the left or to the right. This was

the case for about 90% of the toddler validation data, and 73% to 93% for the infant

validation data, depending on feature set. This may be partly explained by the child not

reacting to the stimulus, or the reaction being finished before or starting later than 2 seconds

after stimulus onset. Figure 3 shows most of the head rotation and gaze data to be focused on

the computer screen or camera rather than one of the two sound field speakers. This is

consistent with previous publications that showed a 32% head turn response rate for

unreinforced complex noise in infants between 12 to 18 months of age (Moore et al., 1975)

and only 25% of infants (6-30 months) made a reliable head turn to warble tones (Shaw &

Nikolopoulos, 2004). Another possibility is that the test assistant, whose task was to keep the

child looking ahead and centring the child if they were looking away, may have interfered

with the response. Furthermore, a considerable number of head turns, and eye gazes were

made in the direction opposite to that expected, as is evident by blue and red symbols on both

sides of the figures. Theoretically, the algorithm could be improved by counting head turns in

either direction, and by considering a wider window of time in which responses may occur

(as opposed to a single frame 2 seconds after stimulus onset), which would allow for

improved decision rates. This would require control trial (no sound) data and a comparison

between control and sound trials as opposed to the left/right comparisons made in the present

study. This would be consistent with traditional VRA approaches that require only detection

of a sound by means of a conditioned head turn response, and do not require the head turn to

be in the direction of the sound source. In other words, VRA is more sensitive when used as a

detection task rather than a localization task (Widen et al., 2000). Furthermore, such

techniques could be used in a VRA procedure in which a child's head turns are reinforced

which would significantly increase the turning behaviour observed and therefore improve the

likelihood of appropriate response classification. Real observers can learn about the child's

response time through observation and are able to include responses that are slow but

consistent. An effective AI approach would similarly be able to learn and adapt to the

individual's reaction time.

Altogether, the present analyses with simple machine learning techniques and few

data demonstrate proof-of-concept and categorize the head turn direction with reasonable

specificity. More data that is anonymized via preprocessing with OpenFace will allow us to

employ more complex and possibly more accurate machine learning approaches, like

variational autoencoders (Kingma & Welling, 2013) to replace the PCA, neural networks to

capture possible non-linear interactions between the features, and a more complex AI that

does not analyse a single frame but the whole-time span after stimulus onset. These

approaches could improve the sensitivity of detection, enabling diagnostic and rehabilitative

audiometric applications. Further improvement may be realized with data that is manually

labelled with a human observer's judgement of whether a child showed a facial reaction or

not. This is likely to show higher accuracies since a detection task is easier than the present

localization paradigm.

Lastly, this proof-of-concept study was conducted in infants and toddlers with normal

hearing that demonstrated high completion rates on behavioural hearing assessments (60% of

toddlers completed full assessments, defined as 2 speech thresholds and 4 pure-tone

thresholds in each ear), which may partly explain the strong classification performance

observed for the infant and toddler validation groups when a decision was made (7 to 11%

toddler cases; 7 to 17% infant cases). Future work should evaluate this approach in more

clinically complex populations, such as children with hearing loss and those with additional

neurological or developmental conditions—groups that are often the least testable using

conventional behavioural methods. Examining the clinical implications of this method,

including comparisons between machine learning–based classifications and responses judged

by trained clinicians, will provide valuable insight. Ongoing data collection includes full-term

infants as well as a subset of infants and toddlers with hearing loss and developmental

conditions, which will allow us to explore these questions in future studies.

5. Conclusions

(1) In this proof-of-concept study, an unsupervised ML algorithm was tasked to form three

meaningful clusters. The cluster representing "no decision by the algorithm" encompassed

90% of the data. When a decision was made about the direction of sound presentation

("right" and "left' clusters), it was mostly correct for the toddlers and significantly above

chance correct for the infants, despite the algorithm being trained on toddlers' reactions.

(2) Automatic VRA could be possible with unsupervised machine learning techniques and

few data, as demonstrated in this study using expectation maximization. Performance level

demonstrated here may be used to provide a human rater with an additional opinion.

(3) The approach was applied to anonymous data, achieved by preprocessing the images to

result in actions performed by the head and face rather than a description of the face. This

allows us to collect and share data across clinics and apply more powerful AI techniques like

deep learning to larger data sets.

Acknowledgements

References

Baltrusaitis, T., Ahuja, C., & Morency, L. P. (2019). Multimodal Machine Learning: A
    Survey and Taxonomy. *IEEE Trans Pattern Anal Mach Intell*, *41*(2), 423–443.
    https://doi.org/10.1109/tpami.2018.2798607

Baltrušaitis, T., Mahmoud, M., & Robinson, P. (2015). Cross-dataset learning and person-
    specific normalisation for automatic Action Unit detection. *11th IEEE International
    Conference and Workshops on Automatic Face and Gesture Recognition*, *6*, pp 1–6.

Bench, J., Collyer, Y., Mentz, L., & Wilson, I. (1976a). Studies in infant behavioural
    audiometry. I. Neonates. *Audiology*, *15*(2), 85–105.
    https://doi.org/10.3109/00206097609071766

Bench, J., Collyer, Y., Mentz, L., & Wilson, I. (1976b). Studies in infant behavioural
    audiometry. II. Six-week-old infants. *Audiology*, *15*(4), 302–314.
    https://doi.org/10.3109/00206097609071790

Bench, J., Collyer, Y., Mentz, L., & Wilson, I. (1976c). Studies in infant behavioural
    audiometry. III. Six-month-old infants. *Audiology*, *15*(5), 384–394.
    https://doi.org/10.3109/00206097609071798

Beswick, R., Driscoll, C., & Kei, J. (2012). Monitoring for postnatal hearing loss using risk
    factors: a systematic literature review. *Ear and Hearing*, *33*(6), 745–756.
    https://doi.org/10.1097/AUD.0b013e31825b1cd9

Capute, A. J., Palmer, F. B., Shapiro, B. K., Wachtel, R. C., Schmidt, S., & Ross, A. (1986).
    Clinical linguistic and auditory milestone scale: prediction of cognition in infancy.
    Developmental Medicine and Child Neurology, *28*(6), 762–771.
    https://doi.org/10.1111/j.1469-8749.1986.tb03930.x

Chen, W. C., & Maitra, R. (2015). *EMCluster: EM algorithm for model-based clustering of*
finite mixture Gaussian distribution, R Package. URL: https://cran.r-
project.org/package=EMCluster.
Ching, T. Y. C., Dillon, H., Button, L., Seeto, M., Van Buynder, P., Marnane, V., Cupples,
L., & Leigh, G. (2017). Age at Intervention for Permanent Hearing Loss and 5-Year
Language Outcomes. *Pediatrics*, *140*(3). https://doi.org/10.1542/peds.2016-4274
Chinta, S. V., Wang, Z., Palikhe, A., Zhang, X., Kashif, A., Smith, M. A., Liu, J., & Zhang,
W. (2025). AI-driven healthcare: Fairness in AI healthcare: A survey. *PLOS Digit*
Health, *4*(5), e0000864. https://doi.org/10.1371/journal.pdig.0000864
Dempster, A. P., Laird, N. M., & Rubin, D. B. (1977). Maximum likelihood from incomplete
data via the EM algorithm. *Journal of the Royal Statistical Society: Series B*
(methodological), *39*(1), 1–22.
Deng, J., Dong, W., Socher, R., Li, L. J., Li, K., & Fei-Fei, L. (2009). Imagenet: A large-
scale hierarchical image database. . IEEE conference on computer vision and pattern
recognition,
Frosolini, A., Franz, L., Caragli, V., Genovese, E., de Filippis, C., & Marioni, G. (2024).
Artificial Intelligence in Audiology: A Scoping Review of Current Applications and
Future Directions. *Sensors (Basel)*, *24*(22). https://doi.org/10.3390/s24227126
Gathman, T. J., Choi, J. S., Vasdev, R. M. S., Schoephoerster, J. A., & Adams, M. E. (2023).
Machine Learning Prediction of Objective Hearing Loss With Demographics, Clinical
Factors, and Subjective Hearing Status. *Otolaryngology and Head and Neck Surgery*,
169(3), 504–513. https://doi.org/10.1002/ohn.288
Gibbon, S., Attaheri, A., Ni Choisdealbha, A., Rocha, S., Brusini, P., Mead, N., Boutris, P.,
Olawole-Scott, H., Ahmed, H., Flanagan, S., Mandke, K., Keshavarzi, M., &
Goswami, U. (2021). Machine learning accurately classifies neural responses to
rhythmic speech vs. non-speech from 8-week-old infant EEG. *Brain and Language*,
220, 104968. https://doi.org/10.1016/j.bandl.2021.104968
Hart, B. N., & Jeng, F. C. (2021). A Demonstration of Machine Learning in Detecting
Frequency Following Responses in American Neonates. *Perceptual and Motor Skills*,
128(1), 48–58. https://doi.org/10.1177/0031512520960390
Hicks, C. B., Tharpe, A. M., & Ashmead, D. H. (2000). Behavioral auditory assessment of
young infants: methodological limitations or natural lack of auditory responsiveness?
Am J Audiol, *9*(2), 124–130. https://doi.org/10.1044/1059-0889(2000/015)
Hirvonen, M., Ojala, R., Korhonen, P., Haataja, P., Eriksson, K., Gissler, M., Luukkaala, T.,
& Tammela, O. (2018). Visual and Hearing Impairments After Preterm Birth.
Pediatrics, *142*(2). https://doi.org/10.1542/peds.2017-3888
Hunter, L. L., Vannest, J., Moore, D. R., Barnes-Davis, M., Blankenship, C., Prather, L.,
Caldwell-Kurtzman, J., & Parikh, N. (2023). Hearing, Speech, and Language in
Infants and Toddlers Born Prematurely. *Volta Rev*, *123*(1), 1–20.
Kingma, D. P., & Welling, M. (2013). Auto-encoding variational bayes. *arXiv*.
https://doi.org/arXiv:1312.6114v11[stat.ML]
Krizhevsky, A. (2009). *Learning Multiple Layers of Features from Tiny Images*. (Technical
Report, University of Toronto, Issue. https://www.cs.toronto.edu/~kriz/learning-
features-2009-TR.pdf
Liu, Y., Xu, R., & Gong, Q. (2021). Maximising the ability of stimulus-frequency
otoacoustic emissions to predict hearing status and thresholds using machine-learning
models. *International Journal of Audiology*, *60*(4), 263–273.
https://doi.org/10.1080/14992027.2020.1821252
McKearney, R. M., Bell, S. L., Chesnaye, M. A., & Simpson, D. M. (2022). Auditory
Brainstem Response Detection Using Machine Learning: A Comparison With
Statistical Detection Methods. *Ear and Hearing*, *43*(3), 949–960.
https://doi.org/10.1097/aud.0000000000001151
Moeller, M. P., Tomblin, J. B., Yoshinaga-Itano, C., Connor, C. M., & Jerger, S. (2007).
Current state of knowledge: language and literacy of children with hearing
impairment. *Ear and Hearing*, *28*(6), 740–753.
https://doi.org/10.1097/AUD.0b013e318157f07f

Moore, J. M., Thompson, G., & Thompson, M. (1975). Auditory localization of infants as a
function of reinforcement conditions. *Journal of Speech and Hearing Disorders*,
40(1), 29–34. https://doi.org/10.1044/jshd.4001.29
Olsho, L. W., Koch, E. G., Carter, E. A., Halpin, C. F., & Spetner, N. B. (1988). Pure-tone
sensitivity of human infants. *Journal of the Acoustical Society of America*, *84*(4),
1316–1324. https://doi.org/10.1121/1.396630
Olsho, L. W., Koch, E. G., Halpin, C. F., & Carter, E. A. (1987). An observer-based
psychoacoustic procedure for use with young infants. . *Developmental Psychology*,
23(5), 627.
Peng, S., Zhao, Y., Yao, X., Yin, H., Ma, B., Liu, K., Li, G., & Cao, Y. (2025). Using
Machine Learning for Analysis of Wideband Acoustic Immittance and Assessment of
Middle Ear Function in Infants. *Audiol Res*, *15*(2).
https://doi.org/10.3390/audiolres15020035
Ruth, R. A., Horner, J. S., McCoy, G. S., & Chandler, C. R. (1983). Comparison of auditory
brainstem response and behavioral audiometry in infants. *Scandinavian Audiology.*
Supplementum, *17*, 94–98.
Shaw, P., & Nikolopoulos, T. (2004). The effect of initial stimulus type for visual
reinforcement audiometry. *International Journal of Audiology*, *43*(4), 193–197.
https://doi.org/10.1080/14992020400050027
Sininger, Y. S., Hunter, L. L., Roush, P. A., Windmill, S., Hayes, D., & Uhler, K. M. (2020).
Protocol for Rapid, Accurate, Electrophysiologic, Auditory Assessment of Infants and
Toddlers. *Journal of the American Academy of Audiology*, *31*(6), 455–468.
https://doi.org/10.3766/jaaa.19046
Sininger, Y. S., Martinez, A., Eisenberg, L., Christensen, E., Grimes, A., & Hu, J. (2009).
Newborn hearing screening speeds diagnosis and access to intervention by 20-25
months. *Journal of the American Academy of Audiology*, *20*(1), 49–57.
https://doi.org/10.3766/jaaa.20.1.5
Thompson, G., & Weber, B. A. (1974). Responses of infants and young children to behavior
observation audiometry (BOA). *Journal of Speech and Hearing Disorders*, *39*(2),
140–147. https://doi.org/10.1044/jshd.3902.140
Thompson, M., & Thompson, G. (1972). Response of infants and young children as a
function of auditory stimuli and test methods. *Journal of Speech and Hearing*
Research, *15*(4), 699–707. https://doi.org/10.1044/jshr.1504.699
Tomblin, J. B., Harrison, M., Ambrose, S. E., Walker, E. A., Oleson, J. J., & Moeller, M. P.
(2015). Language Outcomes in Young Children with Mild to Severe Hearing Loss.
Ear Hear, *36 Suppl 1*, 76S–91S. https://doi.org/10.1097/AUD.0000000000000219
Wathour, J., Govaerts, P. J., Derue, L., Vanderbemden, S., Huaux, H., Lacroix, E., &
Deggouj, N. (2023). Prospective Comparison Between Manual and Computer-
Assisted (FOX) Cochlear Implant Fitting in Newly Implanted Patients. *Ear and*
Hearing, *44*(3), 494–505. https://doi.org/10.1097/aud.0000000000001314
Wathour, J., Govaerts, P. J., Lacroix, E., & Naïma, D. (2023). Effect of a CI Programming
Fitting Tool with Artificial Intelligence in Experienced Cochlear Implant Patients.
Otology & Neurotology, *44*(3), 209–215.
https://doi.org/10.1097/mao.0000000000003810
Werner, L. A., & Marean, G. C. (1991). Methods for estimating infant thresholds. *Journal of*
the Acoustical Society of America, *90*(4 Pt 1), 1867–1875.
https://doi.org/10.1121/1.401666
Widen, J. E. (1993). Adding objectivity to infant behavioral audiometry. *Ear and Hearing*,
14(1), 49–57. https://doi.org/10.1097/00003446-199302000-00007
Widen, J. E., Folsom, R. C., Cone-Wesson, B., Carty, L., Dunnell, J. J., Koebsell, K., Levi,
A., Mancl, L., Ohlrich, B., Trouba, S., Gorga, M. P., Sininger, Y. S., Vohr, B. R., &
Norton, S. J. (2000). Identification of neonatal hearing impairment: hearing status at 8
to 12 months corrected age using a visual reinforcement audiometry protocol. *Ear and*
Hearing, *21*(5), 471–487.
Widen, J. E., & Keener, S. K. (2003). Diagnostic testing for hearing loss in infants and young
children. *Ment Retard Dev Disabil Res Rev*, *9*(4), 220–224.
https://doi.org/10.1002/mrdd.10083

Wood, E., Baltrušaitis, T., Zhang, X., Sugano, Y., Robinson, P., & Bulling, A. (2015).
Rendering of Eyes for Eye-Shape Registration and Gaze Estimation IEEE
International Conference on Computer Vision,
Xie, W., Li, C., Peng, H., Liu, Y., Zhang, Z., Cheng, X., & Liu, J. (2025). Behavior
recognition technology based on deep learning used in pediatric behavioral
audiometry. *Scientific Reports*, *15*(1), 15648. https://doi.org/10.1038/s41598-025-
97519-w
Yoshinaga-Itano, C., Manchaiah, V., & Hunnicutt, C. (2021). Outcomes of Universal
Newborn Screening Programs: Systematic Review. *J Clin Med*, *10*(13).
https://doi.org/10.3390/jcm10132784
Yoshinaga-Itano, C., Sedey, A. L., Coulter, D. K., & Mehl, A. L. (1998). Language of early-
and later-identified children with hearing loss. *Pediatrics*, *102*(5), 1161–1171.
https://doi.org/10.1542/peds.102.5.1161
Zeitler, D. M., Buchlak, Q. D., Ramasundara, S., Farrokhi, F., & Esmaili, N. (2024).
Predicting Acoustic Hearing Preservation Following Cochlear Implant Surgery Using
Machine Learning. *Laryngoscope*, *134*(2), 926–936.
https://doi.org/10.1002/lary.30894
Zeng, J., Kang, W., Chen, S., Lin, Y., Deng, W., Wang, Y., Chen, G., Ma, K., Zhao, F.,
Zheng, Y., Liang, M., Zeng, L., Ye, W., Li, P., Chen, Y., Chen, G., Gao, J., Wu, M.,
Su, Y.,…Cai, Y. (2022). A Deep Learning Approach to Predict Conductive Hearing
Loss in Patients With Otitis Media With Effusion Using Otoscopic Images. *JAMA*
Otolaryngol Head Neck Surg, *148*(7), 612–620.
https://doi.org/10.1001/jamaoto.2022.0900

Figure Legends

Figure 1. Experimental setup for toddlers who were seated in the high chair with sound field

speakers positioned to the right and left at 45° azimuth relative to the ears at the same height.

A computer monitor was straight ahead and played a centring video continuously throughout

the test. The video camera was positioned underneath the computer monitor.

Figure 2. Average absolute angle of the head turns to the left or right (yaw), relative to the

position of the camera, as a function of time after stimulus onset for the toddler group (purple

line) and infant group (green line). Red line indicates the analysis window (2 seconds) after

stimulus onset.

Figure 3. Multivariate Gaussian distributions for three clusters predicted by Expectation

Maximization. Black crosses and ellipses show means and covariances for the distributions.

Red symbols show reactions where a sound was presented to the right, blue symbols where a

sound was presented to the left. Crosses (x) show the training data taken from the toddler

group. Validation data is shown in triangles (infants) and circles (toddler).

Table 1. Number of predictions for each of the four feature sets and three predictive classes for the training and validation sets

Feature Sets	Predictive Classes	Training Set (Toddler)		Validation (Toddler)		Validation (Infant)		p-value
		Left Speaker	Right Speaker	Left Speaker	Right Speaker	Left Speaker	Right Speaker	
Head Rotation	Left	48	4	4	1	13	11	0.07
	Centre	65	72	23	27	102	100	
	Right	9	28	0	1	21	31	
Gaze Direction (Left Eye)	Left	39	4	4	0	7	2	*0.006*
	Centre	78	84	23	29	127	131	
	Right	5	16	0	0	2	9	
Gaze Angle (Both Eyes)	Left	43	5	5	0	14	10	*<0.001*
	Centre	67	78	22	28	116	114	
	Right	12	21	0	1	6	18	
Principal Component Analysis	Left	45	4	5	0	15	10	*0.04*
	Centre	72	82	22	28	118	122	
	Right	5	18	0	1	3	10	

* PCA analysis only includes the first two components

Figure 1.

Figure 2.

Figure 3.