1

Pathophysiological mechanisms of uterine niches (cesarean scar defects) associated with

abnormal bleeding and infertility: a systematic review

Running title: Pathophysiological mechanisms of uterine niches

Gabriel M. Lino[1], Pauliana V. M. Galvão[2]*, Maria L. F. da Silva[1], Priscila Maria de Barros

Rodrigues[1], Valda L. M. Luna[1], Marcos Cezar Feitosa de Paula Machado[1], George A. M.

Conrado[1]

[1]University of Pernambuco, Serra Talhada, Brazil

[2]Faculty of Medical Sciences, University of Pernambuco, Recife, Brazil

Correspondence: Pauliana V. M. Galvão; pauliana.galvao@upe.br

ABSTRACT

Aim: To review the pathological findings of uterine niches in women presenting with

infertility and abnormal uterine bleeding.

Methods: The Medline, Embase, and Cochrane Central databases were searched until 1 May

2025. We included cohorts, case-controls, and case series. The risk of bias was evaluated with

ROBINS-I, and the certainty of evidence was presented according to GRADE.

Results: We included fourteen studies (637 women). Chronic inflammation was observed in

54% (95% CI 25 to 81%; 5 studies, 280 women; $I^2 = 94.3\%$); fibrosis in 83% (95% CI 51 to

96%; 6 studies, 266 women; $I^2 = 92.1\%$); endometrial defects in 90% (95% CI 75 to 96%; 4

studies, 177 women; $I^2 = 64.6\%$); ectopic endometrium in 26% (95% CI 18 to 36%; 9 studies,

411 women; $I^2 = 74.2\%$); and vascular remodeling in 81% (95% CI 75 to 86%; 6 studies, 205

women; $I^2 = 0.0\%$). There was an increased risk of endometrial defects in women with niches

compared to those with prior cesarean deliveries (RR 2.13, 95% CI 1.27 to 3.55; 2 studies,

124 women; $I^2 = 0.0\%$), similarly for ectopic endometrium (RR 2.86, 95% CI 1.03 to 7.96; 2

studies, 97 women; $I^2 = 0.0\%$). Additionally, niches exhibited a pro-inflammatory phenotype,

characterized by elevated expression of CD138 and TNF-α.

Conclusions: The observed pathological features suggest an underlying process driven by the

interplay of chronic inflammation, altered immune cell recruitment, fibrosis, and vascular

remodeling within the regenerating endometrium.

Registration number: CRD420251049667.

Keywords: Cesarean scar defect; Uterine niche; Isthmocele; Infertility; Abnormal uterine

bleeding.

INTRODUCTION

Long-term morbidity following cesarean delivery has become a major concern in

obstetrics [1]. Reduced fecundability, increased risks of placenta accreta, placenta previa, and

abnormal uterine bleeding have all been associated with prior cesarean [1,2]. However, the

mechanisms underlying these associations are extremely unclear. Over the past decade,

growing attention has been directed toward anatomic defects at the site of the uterine scar,

commonly referred to as uterine niches or cesarean scar defects, as key contributors to these

adverse outcomes.

Uterine niches are present in 50 to 60% of women who have had a cesarean delivery [3],

classically associated with abnormal uterine bleeding and infertility, although dysmenorrhea

and dyspareunia are also often reported [4]. Crucially, these symptoms can persist for years after

the surgery.

In theory, damage to the endometrium and endometrial-myometrial interface is one of

the promoters of niche pathogenesis [5]. Thereafter, local dysfunctional hormonal, metabolic, and

inflammatory mechanisms lead to reproductive disorders [6]. It is assumed that the presence of

extensive fibrosis and ectopic endometrium promotes symptom onset. However, the exact

mechanisms are yet unknown. Such an understanding could provide valuable insights into both

clinical management and prevention, aimed at reducing long-term adverse reproductive

outcomes following cesarean delivery. Therefore, this review aimed to systematically review

pathological findings in symptomatic uterine niches.

METHODS

This systematic review was registered prospectively in the PROSPERO database

(CRD420251049667). We followed the PRISMA 2020 guidelines for the reporting of

systematic reviews and meta-analyses [7].

Criteria for considering studies for this review

We considered randomized controlled trials, cohorts, case controls, and case series

reporting pathological findings of women diagnosed with uterine niches after cesarean delivery

and presenting with abnormal uterine bleeding, infertility, or related disorders. The following

outcomes were sought in these studies: inflammation (and inflammatory markers), fibrosis,

endometrial defects, presence of ectopic endometrium, asynchrony of the menstrual cycle in

the scar, and angiogenesis or changes in the surrounding vasculatures (including dilated

capillaries and thick-walled vessels).

Search methods for identification of studies

We searched the Medical Literature Analysis and Retrievel System Online (Medline),

Embase, and the Cochrane Central Register of Controlled Trials (Central) databases until 1 May

2025. A combination of the following terms, and variations, composed the search query:

"cesarean scar defect", "cesarean scar", "uterine scar", "uterine defect", "isthmocele", "uterine

niche", and "histology". The search was adapted for each database. There were no language or

publication date restrictions.

Data collection and analysis

Two investigators were responsible for independently screening the title and abstract of

the identified records. In this stage, if one of the authors considered the record eligible, both

assessed the full text. Subsequently, they proceeded with the selection and extraction of key

information from the included studies into a previously cross-validated file. Disagreements on

which studies should be included in the review were resolved by discussing with a third

investigator. Extracted information included authorship, conflict of interest, funding, country,

setting, population, inclusion and exclusion criteria, information on baseline characteristics,

adjustment methods for baseline imbalances, outcomes reported and how they were assessed,

follow-up of participants, information on risk of bias, suture techniques in uterine closure,

histological observations, and immunohistochemistry markers of inflammation, proliferation,

and hormone expression (CD138, CD16, CD3, CD20, CD56, CD68, CD163, myeloperoxidase

[MPO], tryptase, tumor necrosis factor-α [TNF-α], α-smooth muscle actin [α-SMA], Ki67,

CD10, CD31, estrogen receptor [ER], and progesterone receptor [PR]).

Specific definitions for the outcomes were not required for inclusion in the review, as

long as they were reported in the histological analyses. Uterine niches were considered

appropriately defined when described as defects in the cesarean scar measuring more than 2

mm. When clear definitions for uterine niche diagnosis or outcome classification were lacking,

the available information was recorded and used to inform the risk of bias assessment.

Risk of bias assessment in included studies

Two investigators independently used the Cochrane risk-of-bias tool for nonrandomized

studies of intervention (ROBINS), version 2024, to assess the risk of bias/quality of the included

studies [8]. When different classifications were reached, the final risk assessment was reached by

discussion.

Synthesis methods

We used an inverse-variance random effects model, reporting dichotomous data as

Relative Risk (RR) with 95% Confidence intervals (CI) and continuous outcomes as mean

differences (MD) and 95% CI. We also presented the baseline risk and risk difference (RD) for

the critical outcomes [9]. Statistical heterogeneity was assessed by calculating tau, tau^2, and I^2.

No specific threshold of I^2 was considered a significant indicator of heterogeneity; rather, a

contextualized I^2 was considered based on visual inspection and acceptable between-study

estimate variation [10]. We investigated publication bias with funnel plots.

The analyses were conducted in R, with the package metafor [11].

Certainty of the evidence assessment

The overall certainty of the evidence for each outcome was assessed by two

investigators using the GRADE approach, formulating a summary of findings and certainty of

evidence tables containing the critical outcomes investigated. There were no disagreements;

this process was conducted by deliberation.

The Grading of Recommendations, Assessment, Development and Evaluation

(GRADE) handbook was followed to determine the quality of the evidence of the analyses

conducted [12]. For the evaluation of imprecision, the update provided in 2022 was followed [13].

Inconsistency was evaluated by investigating contextualized heterogeneity, based on whether

reasonable explanations for study variations could be provided [10].

125   The tables were generated using the GRADEpro GDT software available online.

127 RESULTS

128 **Results of the search**

129   We identified 447 records in three databases, Medline, Embase, and Central (Figure 1).

130 After excluding duplicates and records marked as ineligible (case reports and reviews), we

131 assessed the title and abstract of 318 records. Subsequently, we excluded 292 records for being

132 outside the review scope. One additional study was excluded because the digital record was not

133 available. The full text was extracted from 25 records. Finally, we excluded eight studies that

134 did not report any histological analysis and three that did not investigate uterine niches.

136 **Included studies**

137   Fourteen studies met the inclusion criteria for this review. Their main characteristics are

138 summarized in Table 1, with additional details provided in Table S1. Additionally, the most

139 common symptoms reported were abnormal uterine bleeding and infertility. However, in some

140 cases, these symptoms could not be clearly attributed to the uterine niche due to coexisting

141 comorbidities – an issue further addressed in the risk of bias assessment.

142   Histological samples were obtained either through hysteroscopic resection or

143 hysterectomy, with varying time intervals between symptom onset and surgical management.

144   Commonly reported histological features included chronic inflammation, fibrosis,

145 endometrial defects, ectopic endometrium, asynchronous menstrual cycle, vascular remodeling,

146 and immunohistochemical markers of inflammation and cell proliferation. A visual summary

147 of these findings across studies is provided in Figure S1.

148   Most studies employed single-arm, retrospective, or prospective designs, typically

149 involving nonconsecutive patient series.

Risk of bias

The overall risk of bias is presented in Table 1, while a more detailed description of

each domain is provided in Table S2. In general, the quality of most studies was limited by

selection bias and unclear reporting of pathological findings (reporting bias). Except for Wang

2025, Raimondo 2015, and Gubbini 2011, inadequate adjustment for baseline imbalances and

nonconsecutive patient recruitment led to substantial residual confounding, resulting in at least

a serious risk of bias for all other studies.

Analyses of histopathological observations

Chronic inflammation

Six studies investigated chronic inflammation, resulting in a pooled prevalence of 48%

(95% CI 23 to 74%; 6 studies, 318 women; $I^2 = 93.5\%$; very low-certainty evidence; Figure

2). Excluding the only study at critical risk of bias, the prevalence changes to 54% (95% CI 25

to 81%; 5 studies, 280 women; $I^2 = 94.3\%$; very low-certainty evidence; Figure S2). Raimondo

2015 had a conflicting estimate with the remaining studies; the pooled prevalence further

changed to 66% when it was excluded (95% CI 54 to 76%; 4 studies, 160 women; $I^2 = 48.9\%$;

very low-certainty evidence).

Fibrosis

Six studies investigated fibrosis, with a pooled prevalence of 83% (95% CI 51 to 96%;

6 studies, 266 women; $I^2 = 92.1\%$; very low-certainty evidence; Figures 3 and S3). As in the

previous outcome, Gubbini 2011 and Raimondo 2015 had conflicting results. Excluding

Gubbini from the fibrosis analysis results in a prevalence of 88% (95% CI 81 to 93%; 5 studies,

225 women; $I^2 = 24,3\%$; low-certainty evidence).

Endometrial defects

Four studies reported endometrial defects, resulting in a prevalence of 90% (95% CI 75

to 96%; 4 studies, 177 women; $I^2 = 64.6\%$; very low-certainty evidence; Figure 4). Additionally,

Huang 2023 and Higuchi 2021 observed an increased risk of endometrial defects in women

with niches compared to those with prior cesarean deliveries but without niches (RR 2.13, 95%

CI 1.27 to 3.55; 2 studies, 124 women; $I^2 = 0.0\%$; low-certainty evidence; Figure S4). Further,

Wang 2025 also observed an increased risk of endometrial defects in women with symptomatic

niches compared to asymptomatic niches (RR 2.71, 95% CI 2.00 to 3.67; 1 study, 155 women;

low-certainty evidence).

Ectopic endometrium

Ten studies reported the presence of ectopic endometrium within the scar, with a pooled

prevalence of 31% (95% CI 20 to 44%; 10 studies, 449 women; $I^2 = 84.8\%$; very low-certainty

evidence; Figure S5). Excluding studies at critical risk of bias resulted in a prevalence of 26%

(95% CI 18 to 36%; 9 studies, 411 women; $I^2 = 74.2\%$; very low-certainty evidence).

Additionally, two studies compared women with uterine niches to those with prior cesarean

deliveries but without niches. The presence of ectopic endometrium was associated with an

increased risk of niche (RR 2.86, 95% CI 1.03 to 7.96; 2 studies, 97 women; $I^2 = 0.0\%$; low-

certainty evidence; Figure S6). Further, Wang 2025 compared the risk of symptoms in niches

with and without ectopic endometrium, finding a large increase in risk (RR 7.48, 95% CI 3.06

to 18.25; 1 study, 155 women; low-certainty evidence).

Asynchronous menstrual cycle

Three studies investigated asynchrony of the menstrual cycle, comparing cells in the

niche to specimens of the uterine cavity. The estimated pooled prevalence was 30% (95% CI

13 to 55%; 3 studies, 157 women; $I^2 = 87.4\%$; very low-certainty evidence; Figure S7).

Restricting the analysis to studies not at critical risk of bias yields a slightly higher estimate of

42% (95% CI 23 to 63%; 2 studies, 119 women; $I^2 = 80.9\%$; very low-certainty evidence).

Vascular remodeling

Vessels with prominent wall proliferation, referred to as thick-walled vessels, were

investigated by six studies, observing a pooled prevalence of 81% (95% CI 75 to 86%; 6 studies,

205 women; $I^2 = 0.0\%$; low-certainty evidence; Figure S8).

Immunohistochemistry markers (inflammation and cell proliferation)

Four studies conducted immunohistochemistry analyses on markers of inflammation,

inflammatory cells, cell proliferation, or hormonal expression (Tables S3 and S4). All studies,

except for Wang 2025, compared these markers in women with niches with those without

niches but with prior cesareans. Three studies assessed inflammatory markers in niche

specimens, including CD138, CD16, CD3, CD20, CD56, CD68, CD163, myeloperoxidase

(MPO), tryptase, and tumor necrosis factor-α (TNF-α). They reported increased expression of

CD138, CD16, and TNF-α; decreased expression of CD3, CD20, CD68, CD163, and tryptase;

and no significant change in CD56 and MPO (Table S3).

Regarding cell proliferation and hormonal receptors, three studies assessed markers

such as α-smooth muscle actin (α-SMA), Ki67, CD10, CD31, estrogen receptor (ER), and

progesterone receptor (PR). They reported significantly increased expression of CD31; α-SMA

was either similar or increased; and Ki67, CD10, ER, and PR showed no significant change

(Table S4).

Certainty of evidence

As most of the publications consisted of retrospective observational studies, the

certainty of our observations was low to very low. The domains in which we downgraded the

quality of evidence were inconsistency (one level), as studies generally reported non-

standardized findings, and risk of bias extracted from ROBINS-I (two levels). Therefore, the

certainty started at high and was downgraded by three levels (very low-certainty) in some cases

and by two levels when inconsistency was not substantial (low-certainty).

DISCUSSION

Summary of findings

In this systematic review of women diagnosed with uterine niches in the cesarean scar

and presenting chiefly with infertility and abnormal uterine bleeding, the most prevalent

histological findings were fibrosis in 83% (51% to 96%), endometrial defects in 88% (81% to

93%), thick-walled vessels in 81% (75% to 86%), and chronic inflammation in 54% (25% to

81%). Furthermore, compared to women with prior cesarean deliveries but without niches, there

is higher endometritis (CD138), increased neutrophils (CD16), and TNF-α expression,

especially around thick-walled vessels, while T lymphocytes (CD3), B lymphocytes (CD20),

macrophages (CD68), M2 macrophages (CD163), and tryptase expression are comparatively

reduced.

Although ectopic endometrium was not highly prevalent, women with uterine niches

had nearly a threefold higher risk of ectopic endometrium compared to those with prior cesarean

deliveries. A similar increase in risk was observed for endometrial defects. A less frequent but

notable observation was menstrual cycle asynchrony between the niche and surrounding cells.

Limitations of the findings

249   A key limitation was the unclear reporting of pathological findings. Certain histological

250 features were described in some studies but not in others, raising uncertainty about whether

251 these abnormalities were truly absent or simply not assessed. This ambiguity also raises

252 concerns about the consistency of specimen evaluation, specifically, whether all samples were

253 analyzed using the same criteria and by the same group of pathologists.

254   In addition, the high heterogeneity observed in our analyses represents an important

255 limitation. While some variability was anticipated, given the unstandardized patient selection

256 and inherent design limitations of the included studies, the extent of heterogeneity may also

257 reflect true differences in patient factors (age and comorbidities), the inflammatory stage, and

258 the degree of tissue remodeling. However, it remains unclear whether these variations stem

259 from between-study bias or genuine biological differences in the inflammatory response across

260 study populations.

261   One observation that supports the latter hypothesis is the apparent association between

262 the prevalence of chronic inflammation and extensive fibrosis across studies. In some cases,

263 fibrosis was highly prevalent while chronic inflammation was not reported, whereas in others,

264 chronic inflammation was common and extensive fibrosis was less frequently observed.

266 *Immune-mediated inflammation*

267   The extent of damage to the uterine wall, particularly the endometrium, may influence

268 both the magnitude of the inflammatory response and the likelihood of its chronic persistence

269 [28,29]. Other factors, such as genetic predisposition, are also likely to play a role, although these

270 remain poorly understood in uterine physiology.

271   Altogether, there is a plausible indication that immune-mediated inflammation is central

272 to the pathogenesis of niches. Higuchi 2021 [20] observed a distinct phenotype of immune cell

273 recruitment in patients with niches compared to those with previous cesarean deliveries. There

is reduced expression of anti-inflammatory macrophages and increased markers of chronic

inflammatory responses.

In comparison, a similar immune environment is observed in eutopic endometrial cells

of women with adenomyosis, with upregulation of pro-inflammatory markers[30]. Persistence of

inflammation is attributed to both altered hormonal expression and signaling from ectopic cells.

The initial disturbance leading to an altered immune microenvironment in uterine niches

could be hypoxia caused by surgical injury or suture techniques. Increased expression of

hypoxia-inducible factor 1α and consequential classic epithelial-mesenchymal changes has

been reported in patients with endometriosis [31]. Although reversal of these conditions can

ultimately lead to the resolution of abnormal signaling, chronic expression of transforming

growth factor (TGF)-β1 and long-term effects on tissue organization are possible [32]. Resistance

to cell death, in a dysfunctional state, for example, is evidenced by increased expression of

TNF-α in these patients.

Endometrial dysfunction

Structural defects in the eutopic endometrium were present in nearly all women, which

might explain the 'chronic wound' behavior. After impaired cell-cell communication in the

eutopic endometrium, there is an increased probability of disorganization and growth into the

myometrium. Further, it has been proposed that the presence of ectopic endometrium and

activation of TGF-β signaling increase local metabolic demand, trigger hypoxia signaling [33],

and pro-angiogenic factors associated with inflammation [34]. This could contribute to the

development of remodeled vessels and increased α-SMA deposition observed in the defect area.

Also, perivascular fibroblasts, activated by endothelial-derived signals, are prone to adopt a

pro-inflammatory phenotype [35].

In healthy women, endometrial stromal cells' responsiveness to estrogen promotes

angiogenesis and increased capillary diameter and growth from the basalis layer. Injury to the

endometrium causes an intracellular increase in COX-2 activity, and consequently of

prostaglandins, of note, prostaglandin E2 (PGE2) [36]. Thereafter, PGE2 increases cAMP activity,

which mediates aromatase conversion of androgens. In persistent inflammation, estrogen

interacts with estrogen alpha receptors on the adjacent myometrium, promoting muscle

contraction, tissue auto-traumatization, and subverts conventional regenerative pathways [37].

Furthermore, altered vasculature, angiogenesis, and asynchrony could increase the

propensity for bleeding within uterine niches. These areas of frequent bleeding lead to

activation of the coagulation cascade and pro-inflammatory responses [39].

Secondary infertility is also common. A recent systematic review by Vitagliano and

collaborators [40] demonstrated that infertility is more closely associated with the presence of

uterine niches than with cesarean delivery itself. Such observation reinforces the view that

adequately healed cesarean scars are generally not problematic and may only confer a very

modest increase in risk. In contrast, poor scar healing, characterized by the formation of uterine

niches, creates the necessary conditions for adverse reproductive outcomes.

Plausible mechanisms for niche-associated infertility include reduced endometrial

receptivity during implantation due to dysfunctional signaling and embryotoxic effects of

chronic inflammation [6]. In particular, fibrosis may be associated with a defective or poorly

responsive endometrium. This mirrors observations in other gynecological conditions, such as

endometriosis, where extensive fibrosis is linked to impaired endometrial function and reduced

implantation potential [41].

For both infertility and abnormal uterine bleeding, hysteroscopic resection results in

restoration of normal functional endometrium and resolution or improvement of symptoms in

approximately 70% of women [25,26,42]. The underlying mechanism may involve re-

epithelialization driven by healthy progenitor cells [29].

One study reported better bleeding control with levonorgestrel-releasing intrauterine

devices in patients showing more clinical signs of vascular proliferation, suggesting that certain

histological patterns might influence treatment efficacy [43]. In theory, pronounced fibrosis

results in decreased PGE2 activity, whereas substantial vascular remodeling could indicate

higher underlying inflammation and hormonal imbalance.

DISCLOSURE

Conflicts of interest

The authors declare no competing interests, nor was funding received for this project.

REFERENCES

1. Sandall J, Tribe RM, Avery L, et al. Short-term and long-term effects of caesarean section on the health of women and children. *Lancet*. 2018;392(10155):1349-1357. doi:10.1016/S0140-6736(18)31930-5

2. Murji A, Sanders AP, Monteiro I, et al. Cesarean scar defects and abnormal uterine bleeding: a systematic review and meta-analysis. *Fertil Steril*. 2022;118(4):758-766. doi:10.1016/j.fertnstert.2022.06.031

3. Vissers J, Hehenkamp W, Lambalk CB, Huirne JA. Post-Caesarean section niche-related impaired fertility: hypothetical mechanisms. *Hum Reprod*. 2020;35(7):1484-1494. doi:10.1093/humrep/deaa094

4. Donnez O. Cesarean scar defects: management of an iatrogenic pathology whose prevalence has dramatically increased. *Fertil Steril*. 2020;113(4):704-716. doi:10.1016/j.fertnstert.2020.01.037

5. Leyendecker G, Wildt L, Mall G. The pathophysiology of endometriosis and adenomyosis: tissue injury and repair. *Arch Gynecol Obstet*. 2009;280(4):529-538. doi:10.1007/s00404-009-1191-0

6. Vannuccini S, Clifton VL, Fraser IS, et al. Infertility and reproductive disorders: impact of hormonal and inflammatory mechanisms on pregnancy outcome. *Human Reproduction Update*. 2016;22(1):104-115. doi:10.1093/humupd/dmv044

7. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *PLOS Medicine*. 2021;18(3):e1003583. doi:10.1371/journal.pmed.1003583

8. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919. doi:10.1136/bmj.i4919

9. Deeks JJ, Higgins JP, Altman DG, McKenzie J, Veroniki A. Chapter 10: Analysing data and undertaking meta-analyses [last updated November 2024]. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Published online 2024. Accessed April 5, 2025. https://training.cochrane.org/handbook/current/chapter-10

10. Guyatt G, Zhao Y, Mayer M, et al. GRADE guidance 36: updates to GRADE's approach to addressing inconsistency. *J Clin Epidemiol*. 2023;158:70-83. doi:10.1016/j.jclinepi.2023.03.003

11. Viechtbauer W. Conducting meta-analyses in {R} with the {metafor} package. *Journal of Statistical Software*. 2010;36(3).

12. Schünemann H, Brożek J, Guyatt G, Oxman A, editor(s). Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach (updated October 2013). Published online 2013. https://gdt.gradepro.org/app/handbook/handbook.html

13. Zeng L, Brignardello-Petersen R, Hultcrantz M, et al. GRADE Guidance 34: update on rating imprecision using a minimally contextualized approach. *Journal of Clinical Epidemiology*. 2022;150:216-224. doi:10.1016/j.jclinepi.2022.07.014

14. Wang Y, Han Y, Guo X, et al. Caesarean Scar Endometrial Defects Contribute to Post-Caesarean Abnormal Uterine Bleeding and Chronic Endometritis: A Retrospective Case-Control Study. *BJOG*. 2025;132 Suppl 2(Suppl 2):132-139. doi:10.1111/1471-0528.18089

15. Maudot C, Debras E, Laurent-Bellue A, et al. Caesarean section scar: Histological analysis on hysterectomy specimen. A pilot study. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2024;303:236-243. doi:10.1016/j.ejogrb.2024.10.048

16. Amro B, Ramirez M, Farhan R, et al. Isthmoceles - Accuracy of imaging diagnosis and clinical correlation with histology: A prospective cohort study. *Facts Views Vis Obgyn*. 2024;16(2):173-183. doi:10.52054/FVVO.16.2.021

17. Hladchuk ZI, Rozhkovska NM, Sytnikova VO, Syvyi SM. Morphofunctional features of isthmocele complicated by abnormal uterine bleeding. *REPRODUCTIVE ENDOCRINOLOGY*. 2024;(73):30-34. doi:10.18370/2309-4117.2024.73.30-34

18. Huang J, Liu X, Hou Y, et al. Macrophage polarisation in caesarean scar diverticulum. *J Clin Pathol*. 2023;76(6):379-383. doi:10.1136/jclinpath-2021-207926

19. AbdullGaffar B, Almulla A. A Histopathologic Approach to Uterine Niche: What to Expect and to Report in Hysteroscopy-Resected Isthmocele Specimens. *Int J Surg Pathol*. 2022;30(3):240-250. doi:10.1177/10668969211039415

20. Higuchi A, Tsuji S, Nobuta Y, et al. Histopathological evaluation of cesarean scar defect in women with cesarean scar syndrome. *Reprod Med Biol*. 2021;21(1):e12431. doi:10.1002/rmb2.12431

21. Karpathiou G, Chauleur C, Dridi M, et al. Histologic Findings of Uterine Niches. *Am J Clin Pathol*. 2020;154(5):645-655. doi:10.1093/ajcp/aqaa080

22. Shapira M, Mashiach R, Meller N, et al. Clinical Success Rate of Extensive Hysteroscopic Cesarean Scar Defect Excision and Correlation to Histologic Findings. *J Minim Invasive Gynecol*. 2020;27(1):129-134. doi:10.1016/j.jmig.2019.03.001

23. Donnez O, Donnez J, Orellana R, Dolmans MM. Gynecological and obstetrical outcomes after laparoscopic repair of a cesarean scar defect in a series of 38 women. *Fertil Steril*. 2017;107(1):289-296.e2. doi:10.1016/j.fertnstert.2016.09.033

24. Tanimura S, Funamoto H, Hosono T, et al. New diagnostic criteria and operative strategy for cesarean scar syndrome: Endoscopic repair for secondary infertility caused by cesarean scar defect. *J Obstet Gynaecol Res*. 2015;41(9):1363-1369. doi:10.1111/jog.12738

25. Raimondo G, Grifone G, Raimondo D, Seracchioli R, Scambia G, Masciullo V. Hysteroscopic treatment of symptomatic cesarean-induced isthmocele: a prospective study. *J Minim Invasive Gynecol*. 2015;22(2):297-301. doi:10.1016/j.jmig.2014.09.011

26. Gubbini G, Centini G, Nascetti D, et al. Surgical hysteroscopic treatment of cesarean-induced isthmocele in restoring fertility: prospective study. *J Minim Invasive Gynecol*. 2011;18(2):234-237. doi:10.1016/j.jmig.2010.10.011

27. Morris H. Surgical pathology of the lower uterine segment caesarean section scar: is the scar a source of clinical symptoms? *Int J Gynecol Pathol*. 1995;14(1):16-20. doi:10.1097/00004347-199501000-00004

28. Ben-Nagi J, Walker A, Jurkovic D, Yazbek J, Aplin JD. Effect of cesarean delivery on the endometrium. *Int J Gynaecol Obstet*. 2009;106(1):30-34. doi:10.1016/j.ijgo.2009.02.019

29. Ang CJ, Skokan TD, McKinley KL. Mechanisms of Regeneration and Fibrosis in the Endometrium. *Annu Rev Cell Dev Biol*. 2023;39:197-221. doi:10.1146/annurev-cellbio-011723-021442

30. Bourdon M, Santulli P, Jeljeli M, et al. Immunological changes associated with adenomyosis: a
systematic review. *Hum Reprod Update*. 2021;27(1):108-129. doi:10.1093/humupd/dmaa038

31. Xiong Y, Liu Y, Xiong W, et al. Hypoxia-inducible factor 1α-induced epithelial-mesenchymal
transition of endometrial epithelial cells may contribute to the development of endometriosis.
Hum Reprod. 2016;31(6):1327-1338. doi:10.1093/humrep/dew081

32. Lin X, Dai Y, Xu W, et al. Hypoxia Promotes Ectopic Adhesion Ability of Endometrial Stromal
Cells via TGF-β1/Smad Signaling in Endometriosis. *Endocrinology*. 2018;159(4):1630-1641.
doi:10.1210/en.2017-03227

33. Wu MH, Chen KF, Lin SC, Lgu CW, Tsai SJ. Aberrant expression of leptin in human
endometriotic stromal cells is induced by elevated levels of hypoxia inducible factor-1alpha. *Am J*
Pathol. 2007;170(2):590-598. doi:10.2353/ajpath.2007.060477

34. Laschke MW, Menger MD. In vitro and in vivo approaches to study angiogenesis in the
pathophysiology and therapy of endometriosis. *Hum Reprod Update*. 2007;13(4):331-342.
doi:10.1093/humupd/dmm006

35. Di Carlo SE, Peduto L. The perivascular origin of pathological fibroblasts. *J Clin Invest*.
2018;128(1):54-63. doi:10.1172/JCI93558

36. Banu SK, Lee J, Speights VO, Starzinski-Powitz A, Arosh JA. Cyclooxygenase-2 regulates
survival, migration, and invasion of human endometriotic cells through multiple mechanisms.
Endocrinology. 2008;149(3):1180-1189. doi:10.1210/en.2007-1168

37. García-Solares J, Donnez J, Donnez O, Dolmans MM. Pathogenesis of uterine adenomyosis:
invagination or metaplasia? *Fertil Steril*. 2018;109(3):371-379.
doi:10.1016/j.fertnstert.2017.12.030

38. Jabbour HN, Sales KJ. Prostaglandin receptor signalling and function in human endometrial
pathology. *Trends Endocrinol Metab*. 2004;15(8):398-404. doi:10.1016/j.tem.2004.08.006

39. Yu L, Shen H, Ren X, et al. Multi-omics analysis reveals the interaction between the complement
system and the coagulation cascade in the development of endometriosis. *Sci Rep*.
2021;11(1):11926. doi:10.1038/s41598-021-90112-x

40. Vitagliano A, Cicinelli E, Viganò P, et al. Isthmocele, not cesarean section per se, reduces in vitro
fertilization success: a systematic review and meta-analysis of over 10,000 embryo transfer cycles.
Fertility and Sterility. 2024;121(2):299-313. doi:10.1016/j.fertnstert.2023.11.007

41. Vissers G, Giacomozzi M, Verdurmen W, Peek R, Nap A. The role of fibrosis in endometriosis: a
systematic review. *Hum Reprod Update*. 2024;30(6):706-750. doi:10.1093/humupd/dmae023

42. Zhang J, Zhu C, Yan L, et al. Comparing levonorgestrel intrauterine system with hysteroscopic
niche resection in women with postmenstrual spotting related to a niche in the uterine cesarean
scar: a randomized, open-label, controlled trial. *Am J Obstet Gynecol*. 2023;228(6):712.e1-
712.e16. doi:10.1016/j.ajog.2023.03.020

43. Huang SY, Chen YY, Tsai CC, Lin H, Ou YC. Evaluating resectoscopy and the levonorgestrel
intrauterine system for intermenstrual bleeding associated with cesarean scar defects: A
retrospective study of treatment outcomes. *J Obstet Gynaecol Res*. 2024;50(2):253-261.
doi:10.1111/jog.15840

FIGURE LEGENDS

Figure 1 Legend PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-

Analyses)

2020 flow diagram of study selection.

Figure 2 Legend Proportion of chronic inflammation in histopathological observations in

women with symptomatic uterine niches. IV, inverse variance.

Figure 3 Legend Proportion of fibrosis in histopathological observations in women with

symptomatic uterine niches. IV, inverse variance.

Figure 4 Legend Proportion of endometrial defects in histopathological observations in women

with symptomatic uterine niches. IV, inverse variance.

Figure S1 Legend Prevalence of histological findings per study.

Figure S2 Legend Subgroup analysis based on the risk of bias of the proportion of chronic

inflammation in histopathological observations in women with symptomatic uterine niches. IV,

inverse variance.

Figure S3 Legend Subgroup analysis based on the risk of bias of the proportion of fibrosis in

histopathological observations in women with symptomatic uterine niches. IV, inverse

variance.

Figure S4 Legend Relative risk of endometrial defects in histopathological observations in

women with symptomatic uterine niches compared to women with previous cesarean deliveries

without uterine niches. IV, inverse variance.

Figure S5 Legend Proportion of ectopic endometrium in histopathological observations in

women with symptomatic uterine niches. IV, inverse variance.

Figure S6 Legend Relative risk of ectopic endometrium in histopathological observations in

women with symptomatic uterine niches compared to women with previous cesarean deliveries

without uterine niches. IV, inverse variance.

Figure S7 Legend Proportion of asynchronous menstrual cycle in histopathological

observations in women with symptomatic uterine niches. IV, inverse variance.

Figure S8 Legend Proportion of thick-walled vessels in histopathological observations in

women with symptomatic uterine niches. IV, inverse variance.

Study	Events	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Amro 2024	33	60	17.7%	0.55 [0.42, 0.68]	
Karpathiou 2020	6	8	13.3%	0.75 [0.35, 0.97]	
Mirre 1995	3	51	17.5%	0.06 [0.01, 0.16]	
Hladchuk 2024	9	38	17.0%	0.24 [0.11, 0.40]	
Raimondo 2015	12	120	17.5%	0.10 [0.05, 0.17]	
Gubbini 2011	32	41	17.0%	0.78 [0.62, 0.89]	
Total (95% CI)		**318**	**100.0%**	**0.48 [0.23, 0.74]**	

Heterogeneity: Tau² = 1.72; Chi² = 77.17, df = 5 (P < 0.01); I² = 93.5%

Study	Events	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Donnez 2017	30	38	19.2%	0.79 [0.63, 0.90]	
AbdullGaffar 2022	22	22	11.9%	1.00 [0.85, 1.00]	
Huang 2023	33	37	18.5%	0.89 [0.75, 0.96]	
Karpathiou 2020	8	11	11.7%	1.00 [0.63, 1.00]	
Raimondo 2018	106	120	19.6%	0.88 [0.81, 0.93]	
Gubbini 2011	7	41	19.1%	0.83 [0.07, 0.32]	
Total (95% CI)		**266**	**100.0%**	**0.83 [0.51, 0.96]**	

Heterogeneity: Tau² = 2.91; Chi² = 63.23, df = 5 (P < 0.01); I² = 92.1%

Study	Events	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Higuchi 2021	49	63	41.3%	0.78 [0.66, 0.87]	
Wang 2025	63	69	36.7%	0.91 [0.82, 0.97]	
Huang 2023	37	37	11.2%	1.00 [0.91, 1.00]	
Karpathiou 2020	8	8	10.8%	1.00 [0.69, 1.00]	
Total (95% CI)		177	100.0%	0.90 [0.75, 0.96]	

Heterogeneity: Tau² = 2.74; Chi² = 8.47, df = 3 (P = 0.04); I² = 64.6%

Identification of new studies via databases and registers