Prevalence and Risk Factors of Relapse in Pulmonary Sarcoidosis: First Systematic Review and Meta-Analysis

Zia Hashim[1], Naresh Kumar Tripathy[2], Alok Nath[1], Mansi Gupta[1], Ajmal Khan[1], Ajit Kumar[1], Anup Kumar[3]

[1]Department of Pulmonary Medicine; [2]Department of Hematology, [3]Department of Biostatistics and Health Informatics; Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India

ABSTRACT

Background and Aim: Relapses are known to adversely affect the prognosis of patients with pulmonary sarcoidosis (PS). However, a systematic review or meta-analysis on the subject is not yet available in the literature.

Research question: What are the global prevalence and risk factors for relapse in PS patients?

Methods: We systematically searched PubMed, Google Scholar, Scopus, and Embase electronic databases to collect eligible PS studies published until 31 January 2025. Eligible studies were peer-reviewed, English-language studies reporting relapse and risk factors in PS patients. Prevalence and risk factors were analyzed using proportions and log odds ratios (LORs) with 95% confidence intervals (CIs). Quality was assessed by Hoy et al tool for prevalence studies

Results: A total of 50 studies with 5978 patients were included in this meta-analysis. Among these patients, 3648 were followed up, and PS relapse was analyzed. Relapse prevalence heterogeneity was significant (Q test, p<0.01). The pooled prevalence of relapse was 0.40 (95% CI: 0.34, 0.46) in all included studies. Sensitivity analyses of the pooled prevalence estimate showed minimal variation, with estimates ranging from 35% to 41% across analyses removing outliers, performing leave-one-out, adjusting for study size, excluding lower-quality studies, and excluding studies without relapse definitions. The pooled prevalence of relapse in prospective vs retrospective studies was 0.32 (95% CI: 0.26, 0.38) vs. 0.44 (95% CI: 0.36, 0.54), with a meta-regression coefficient of 0.14 (p = 0.02). The pooled LOR of relapse was greater for patients who were black [0.91 (95% CI: 0.44, 1.37); p<0.001], had stage II & III disease [0.55 (95% CI: 0.07, 1.03); p=0.02], and had stage IV disease [1.04 (95% CI: 0.25, 1.83); p=0.01], whereas it was comparable for age, sex, treatment type, and PS type (isolated or systemic PS).

Conclusion: The pooled prevalence of PS relapse in all studies was 40%, but it was lower in prospective studies. A black race and higher-grade radiographic stages were the underlying risk factors for disease relapse.

KEYWORDS: Pulmonary Sarcoidosis, Relapse, Risk Factors, Meta-analysis

[1] Prof. Alok Nath, DM
Department of Pulmonary Medicine
Sanjay Gandhi Postgraduate Institute of Medical Sciences
Raebareli Road, Lucknow, Uttar Pradesh 226014, India

<div align="center">**Take-Home Message**</div>

Research Question: What is the global occurrence and etiology of relapse among patients with pulmonary sarcoidosis illness?

Interpretation:
- Our first meta-analysis of 3,646 follow-up PS patients from 50 studies revealed that the pooled prevalence of PS relapse was 40% (95% CI: 0.34–0.46).
- The relapse prevalence was lower in prospective studies than in retrospective studies (0.32 [95% CI: 0.2 0.38] vs. 0.44 [95% CI: 0.36, 0.54], with a meta-regression coefficient=0.14, p= 0.02), indicating the potential impact of study design on the occurrence or detection of relapses in patients with this disease.
- Black race (LOR: 0.91, p<0.001) and stage II & III PS (LOR: 0.55, p=0.02) and IV PS (LOR 1.04, p=0.0 were the underlying risk factors for relapse, whereas age, sex, treatment type, and type of PS (isolated systemic) had no associations.
- This research emphasizes the critical need for a consensus definition of PS relapse and prospective larg cohort studies identifying new biomarkers and treatments of relapse to improve the routine clinic practice and outcome of this disease.

INTRODUCTION

Pulmonary sarcoidosis (PS) is a nonnecrotizing granulomatous inflammatory disease that primarily affects the lungs and/or thoracic lymph nodes. It can be categorized into two types, viz. Isolated PS affects only the lungs (isolated PS) or systemic PS, which, along with the lungs, can affect several other organs, including the eyes, skin, liver, spleen, kidneys, heart, central nervous system, musculoskeletal system, bone marrow, and peripheral lymph nodes.[1][2] Pathologically, this disease may have a self-limiting phenotype with a good prognosis or a progressive phenotype with a poor prognosis, including a risk of severe organ dysfunction and increased morbidity and mortality.[3][4] Progressive disease often requires optimal treatment to prevent disease severity and mortality and improve quality of life.[5][6]

Despite tremendous advancements in PS diagnosis and treatment, unpredictable and frequent relapses remain a major challenge in the clinical management of this disease. They commonly occur at various time points after withdrawal of treatment, including corticosteroids (CSs) or steroid-sparing agents (SSAs), and may persist for years, adversely affecting the prognosis of the disease.[7-10] Thus, comprehensive data on the prevalence of relapse and underlying risk factor relapse are essential for developing strategies for the early detection and prevention of relapses and improving clinical outcomes.

Currently, there is no standard definition of relapse, and different studies have reported that patients relapse in different ways without clear documentation of the underlying risk factors. Furthermore, the data available in the literature on the prevalence and risk factors for PS have not yet been meta-analyzed. Therefore, this study aimed to evaluate the global pooled prevalence and risk factors for relapse through a comprehensive systematic review and meta-analysis of the data of the PS studies available in the literature to date. We hope that this analysis will provide new insights into relapse in PS patients and lead to the development of a consensus definition of PS relapse, as well as the formulation of new guidelines to improve the outcome of relapse-affected PS patients.

METHODS

Study Protocol and Registration

We performed a systematic review and meta-analysis following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines (www.prisma-statement.org). The study protocol was registered with PROSPERO, number CRD420251075384.
https://www.crd.york.ac.uk/PROSPERO/view/CRD420251075384.

Definitions of sarcoidosis states

Sarcoidosis Relapse:
Relapse: (i) worsening of pulmonary/extrapulmonary symptoms and radiology postremission requiring retreatment; (ii) >20% worsening of functional/biological features; (iii) exclusion of alternative causes.
Progression: Worsening of clinical, radiological, functional, or biological markers from baseline, including new organ involvement.
Exacerbation: Acute disease flare-up during treatment tapering/withdrawal.
Remission: Sustained (1–3 months) resolution of symptoms, stabilized/normalized radiology, and normalized pulmonary functions/serum biomarkers.
Anti-sarcoid treatment withdrawal time:
Time when the disease is stable with no clinical, radiological, or biological signs of activity.

Search strategy:

We searched PubMed, Google Scholar, Scopus, and Embase for studies on pulmonary sarcoidosis and relapse published up to 31 January 2025. Two authors (NKT and ZH) independently used Boolean combinations of MeSH terms: (("Sarcoidosis, Pulmonary"[Mesh] OR "pulmonary sarcoidosis"[Title/Abstract] OR "lung

sarcoidosis"[Title/Abstract] OR "sarcoidosis"[Title/Abstract]) AND ("Recurrence"[Mesh] OR "relapse"[Title/Abstract] OR "recurrence"[Title/Abstract] OR "flare-up"[Title/Abstract] OR "exacerbation"[Title/Abstract]) AND ("Prevalence"[Mesh] OR "prevalence"[Title/Abstract] OR "incidence"[Title/Abstract] OR "frequency"[Title/Abstract] OR "rate"[Title/Abstract] OR "Risk Factors"[Mesh] OR "risk factors"[Title/Abstract] OR "predictors"[Title/Abstract] OR "associations"[Title/Abstract] OR "determinants"[Title/Abstract])). Reference lists and citations were manually searched for additional studies.

Inclusion and Exclusion Criteria:
Eligible studies met the following criteria: (i) full-text observational or clinical studies in English, peer-reviewed journals; (ii) studies reporting relapse in PS patients treated with corticosteroids (CSs) and/or steroid-sparing agents (SSAs); and (iii) studies reporting risk factors for relapse. The exclusion criteria included non-English publications, non-peer-reviewed publications, reviews, case reports, studies with missing data, and studies of single-organ nonpulmonary sarcoidosis.

Study Selection:
We retrieved 1432 records through electronic databases and manual searches. We imported the search results into EndNote (version 25.0, Clarivate Analytics, NY, USA) to remove duplicates. After 112 duplicates were removed, 1320 records were screened by title and abstract. Among these, 79 studies were selected for full-text review, and 1241 were excluded. Of the 79 full-text articles assessed, 50 met the inclusion criteria, while 25 studies were excluded because of missing clear relapse data, and 4 studies were excluded because full-text articles were not available (Figure 1). We used a two-level screening process to select eligible studies. First, two authors (NKT and ZH) independently screened titles and abstracts to identify relevant full-text articles. Second, they independently reviewed the full texts against the eligibility criteria to select studies for inclusion. Disagreements were resolved through discussion, with unresolved issues adjudicated by a third author (AN).

Data Extraction:
The authors NKT and ZH extracted data from the study texts and supplementary materials via Excel 365, including author details, publication year, country, study design, patient demographics, sarcoidosis organ involvement, treatment, remission/relapse timing, and PS diagnosis criteria. The study authors were contacted for additional data. The extracted variables included age (continuous), sex (binary), and ethnicity (categorical: Black, White, other). Radiographic stages followed scoring criteria (I–IV). Acute worsening during treatment changes was classified as relapse if reported as such, although this variability was noted as a limitation. (Supplementary Table 2)

Subgrouping of Studies:
The studies were categorized into the following subgroups: (i) study type, geographical location, and time period; (ii) patient age, sex, and ethnicity; (iii) radiographic stages of PS[11]; (iv) treatment types (CS, SSA, or combined CS+SSA); and (v) isolated PS and multiorgan PS.

Assessment of Quality:
We selected the Hoy et al. (2012) risk of bias tool[12] over the nine-item JBI Checklist[13] for its focus on prevalence-specific biases and validated, high-interrater-reliability framework, making it ideal for assessing prevalence studies in this meta-analysis. The tool employs a four-point response system ("yes," "no," "unclear," "not applicable") to evaluate 10 characteristics: (i) representativeness of the target population; (ii) appropriateness of the sampling method; (iii) adequacy of the sample size; (iv) description of the study setting and participants; (v) coverage of the identified sample; (vi) validity of measurement methods; (vii) reliability of measurement methods; (viii) appropriateness of statistical analyses; (ix) adequacy of the response rate; and (x) management of nonresponse bias. Studies earn one point per "yes" response, with scores ranging from 0 to 10. Scores ≥

validated as indicating good quality, denote a lower risk of bias suitable for meta-analyses, although contextu
factors may classify some studies as moderate risk.

Statistical analysis:
We used STATA 18.0 (StataCorp, College Station, TX) for the statistical analysis. Prevalence was calculated
proportions with 95% confidence intervals (CIs) via the DerSimonian–Laird random effects model[14]. Th
restricted-maximum likelihood method was used for dichotomous outcomes because of its robustness with spar
data and heterogeneity. Cochrane's Q was used to assess subgroup differences in pooled proportions across stud
designs. Pooled LORs with 95% CIs were calculated via a random effects model with the restricted-maximu
likelihood method.

Subgroup analysis: Continuous variables were evaluated via the empirical Bayes method and are reported as th
mean differences with 95% CIs. Continuous variables (e.g., age) were analysed as continuous variables to avo
loss of information from categorization, with mean differences reported for age comparisons. For subgrou
comparisons, the Z test was used for two groups, and the ANOVA-equivalent Q test was used for more than tw
groups.

Meta-regression: To account for multiple hypotheses across risk factors (e.g., age, sex, race, radiographic stag
and treatment type), p values were interpreted cautiously, and meta-regression was used to confirm the robustne
of the significant findings.[15]

Heterogeneity: The amount of heterogeneity, τ^2, was estimated via the restricted maximum likelihoo
estimator.[16] In addition, the Q test for heterogeneity and the I^2 statistic were reported.[17] If any amount
heterogeneity was detected (i.e., $\tau^2 > 0$, regardless of the results of the Q test), a prediction interval for th
true outcomes was also calculated. Proportions were compared via LORs; if the 95% CI included zero, th
difference was considered nonsignificant. Studentized residuals and Cook's distances are used to examin
whether studies may be outliers or influential in the context of the model.[18] Outliers were identified using
Baujat plot, which identifies studies with high contributions to heterogeneity and significant influence on th
overall effect size. (Supplementary Figure 10)

Publication Bias:
We assessed publication bias visually via a funnel plot and objectively with Egger's and Begg's tests.[19 20]

Sensitivity analysis: To assess the robustness of the pooled relapse prevalence estimate, we conducted sever
sensitivity analyses. First, we removed outliers from the analysis. Second, we performed a leave-one-out analys
by sequentially excluding each study and recalculating the pooled prevalence estimate. Third, we excluded studi
that did not specify explicit relapse criteria and repeated the analysis on the remaining studies. Fourth, to evalua
the influence of sample size, we excluded studies with fewer than 50 patients in follow-up and recalculated th
pooled prevalence. Fifth, we excluded studies that were assessed as having a moderate risk of bias. Finally,
address potential issues with variance instability, we applied the Freeman–Tukey double arcsine transformation
an alternative to the DerSimonian–Laird method. These sensitivity analyses were conducted to confirm th
consistency and reliability of our pooled prevalence estimate under different methodological variations.

RESULTS
Study characteristics:
Of the 50 studies, 18 were from Europe, 18 from Asia, 12 from North America, and 2 from Africa. Twenty-ni
studies were retrospective, and 21 were prospective. Studies were also classified on the basis of the availability

a relapse definition (Table 1). A consistent definition of relapse was utilized in 22 studies, whereas other autho relied on clinical observations. (Supplementary Table 1)

Quality:
A review of study quality revealed that 62% (31 of 50 studies) presented a low risk of bias, 38% (19 of 50 studie demonstrated a moderate risk of bias, and no studies were classified as having a high risk of bias. These variation stem from a lack of proper definitions for relapse or improper reporting of various parameters and their retrospective design. (Table 2)

Relapse:
Among the 5,978 patients across 50 studies, 3,646 were followed up posttreatment, with 1,437 experiencing relapse. The pooled prevalence of relapse was 0.40 (95% CI: 0.34, 0.46), ranging from 18% to 88% (Figure 2). Most relapses (54%) were clinical, radiological, functional, or based on biomarkers, and 30% were undefined (Figure 3a). The timing of relapse was mentioned in only 58% of the studies. The overall time to relapse ranged from one month to 60 months. The median time to relapse ranged from 3 (IQR 1.8–6.5) to 13 (IQR 9.5–31.5) months (Figure 3b).

Heterogeneity: Substantial heterogeneity was observed in the prevalence of relapse, as evidenced by the Cochran's Q test (Q(49) = 1029, p < 0.001). The estimated variance of the true effect sizes was $\tau^2 = 0.04$, with a heterogeneity index of $I^2 = 95\%$. The 95% prediction interval for the true outcomes ranged from 0.008-0.787. Th positive range of this interval suggests that despite the heterogeneity, the true outcomes of the studies generally align with the direction of the estimated average outcome. We identified study number 34, 48, 41, 45, and 24 wei outliers. (Table 1, Supplementary Figure 1)

Sensitivity analysis

We assessed the robustness of the pooled relapse prevalence estimate in pulmonary sarcoidosis patients using a random-effects REML model across 50 studies. Sensitivity analyses demonstrated the following:
- After removing outliers, the pooled prevalence was 0.35 (95% CI: 0.30, 0.40) (Supplementary Figure 2).
- For studies with a relapse definition (N=22), the proportion was 0.38 (95% CI: 0.30, 0.45), and for studie without a relapse definition (N=28), it was 0.41 (95% CI: 0.33, 0.50) (Supplementary Figure 4).
- There were no significant differences in relapse proportions between studies with ≥50 patients (N=24) an those with <50 patients (N=26). Additionally, for studies with low bias risk (N=31), the proportion was 0.38 (95% CI: 0.32, 0.45), and for studies with moderate bias risk (N=19), it was 0.42 (95% CI: 0.31, 0.5 (Supplementary Figure 5).
- The Freeman–Tukey transformation yielded a proportion of 0.40 (95% CI: 0.33, 0.46) (Supplementary Figure 6).
- Leave-one-out analysis showed stable proportions ranging from 0.39 to 0.41 with a 95% CI of 0.33 to 0.4 (Supplementary Figure 9).

Despite high heterogeneity ($I^2=95\%$, $Tau^2=0.05$, $chi^2=1029$, p=0.0), even in prospective studies ($I^2=86\%$), the pooled relapse proportion remained stable at 40% (Figure 4, Supplementary Table 2).

Table 1: Characteristics of the Included Studies

Study No.	Study	Country	Design	Total Patients	Follow-up Patients	Relapsed Patients	Time to Relapse (Months)	Definition of Remission	Relapse definition
1	Bouchut et al. (2024)[21]	France	Retrospective	175	173	91	48-156	NA	Yes*
2	Guliyeva et al. (2024)[22]	Turkey	Retrospective	52	52	14	2-60	Clinical normalization	NA
3	Dhooria et al. (2023)[23]	India	Prospective	86	86	39	6-15	NA	Yes
4	Ozdemirel et al. (2023)[24]	Turkey	Retrospective	73	73	16	NA	NA	NA
5	Spyropoulos et al. (2022)[25]	Greece	Prospective	147	147	42	NA	NA	NA
6	Madan et al. (2022)[26]	India	Prospective	327	327	73	NA	NA	NA
7	Sakkat et al. (2022)[27]	Canada, UK	Retrospective	33	11	7	2-10	Clinical & biological normalization	NA
8	Morar et al. (2022)[28]	South Africa	Retrospective	102	89	54	NA	NA	NA
9	Feng et al. (2022)[29]	China	Retrospective	432	123	24	NA	NA	Yes
10	Gavrysyuk et al. (2021a)[30]	Ukraine	Prospective	143	84	28	6-12	Clinical & radiological normalization	Yes
11	Murata et al. (2021)[31 32]	Japan	Retrospective	103	103	23	NA	NA	Yes
12	Gavrysyuk et al. (2021b)[32]	Ukraine	Prospective	108	108	56	6-12	NA	Yes
13	Fang et al. (2019)[33]	China	Retrospective	60	11	2	1-30	NA	Yes
14	Nath et al. (2019)[34]	India	Retrospective	120	120	29	2-9	Clinical, radiological, & biological normalization	Yes
15	Zheng et al. (2019)[35]	China	Retrospective	203	96	30	1-36	Clinical & radiological normalization for at least 06 months after treatment withdrawal.	Yes
16	Ozyilmaz et al. (2018)[36]	Turkey	Prospective	91	29	13	NA	NA	NA

#	Author	Country	Study type						
17	Belhomme et al. (2018)[37]	France	Retrospective	80	34	10	NA	NA	Yes
18	Chen et al. (2018)[38]	China	Retrospective	23	23	12	NA	Clinical, & radiological normalization	Yes
19	Mañá et al. (2017)[39]	Spain	Prospective	640	266	30	NA	NA	Yes
20	Maturu et al. (2016)[40]	India	Prospective	27	27	10	3-12	Clinical, & radiological normalization	Yes
21	Liu et al. (2016)[41]	China	Prospective	42	42	20	NA	NA	Yes
22	Polverosi et al. (2014)[42]	Italy	Retrospective	56	44	9	NA	NA	NA
23	Vorselaars et al. (2014)[43]	Netherlands	Retrospective	47	47	29	8-14	NA	Yes
24	Sharma et al. (2003)[44]	India	Retrospective	56	53	40	On CS Taper	NA	NA
25	Kiper et al. (2001)[45]	Turkey	Retrospective	15	15	6	NA	Clinical, radiological, & biological normalization	Yes
26	Sharma et al. (2001)[46]	India	Prospective	106	85	21	NA	NA	Yes
27	Baculard et al. (2001)[47]	France	Retrospective	21	21	4	On CS Taper	Radiological, & PFT normalization	Yes
28	Muller-Quernheim et al. (1999)[48]	Germany	Prospective	11	11	3	8-22	NA	NA
29	Kim et al. (1999)[49]	Korea	Prospective	27	27	4	On CS withdrawal	NA	NA
30	Baltzan et al. (1999)[50]	Canada	Prospective	23	18	8	NA	NA	Yes
31	Rizzato et al. (1998)[51]	Italy	Retrospective	702	82	30	12-36	NA	NA
32	Wyser et al. (1997)[52]	South Africa	Prospective	37	16	7	4-48	NA	Yes
33	Gottlieb et al. (1997)[53]	Israel	Prospective	337	221	86	2-12	NA	Yes
34	Lower et al. (1995)[54]	USA	Retrospective	105	40	35	NA	NA	Yes
35	Takada et al. (1993)[55]	Japan	Retrospective	295	76	36	On CS withdrawal	NA	NA
36	Chadelat et al. (1993)[56]	France	Prospective	8	8	2	NA	NA	NA

37	Prior et al. (1991)[57]	UK	Prospective	25	25	6	NA	NA	NA
38	Rizzato et al. (1991)[58]	Italy	Retrospective	40	40	4	On CS withdrawal	NA	NA
39	Baumann et al. (1990)[59]	USA	Retrospective	15	15	10	3-6	NA	NA
40	Lower et al. (1990)[60]	USA	Retrospective	15	14	5	NA	NA	NA
41	Johns et al. (1986)[61]	USA	Retrospective	181	171	128	1-2	NA	NA
42	Wallaert et al. (1986)[62]	France	Prospective	12	12	8	12-18	NA	NA
43	Turner-Warwick et al. (1986)[63]	UK	Prospective	32	32	4	NA	NA	NA
44	Eule et al. (1986)[64]	Germany	Prospective	172	110	24	NA	NA	NA
45	Johns et al. (1983)[65]	USA	Retrospective	103	99	74	On CS taper or withdrawal	NA	NA
46	Rohatgi et al. (1981)[66]	USA	Prospective	35	20	6	On CS Taper	NA	NA
47	Johns et al. (1976)[67]	USA	Retrospective	192	146	102	On CS withdrawal	NA	NA
48	Johns et al. (1974)[68]	USA	Retrospective	152	145	113	On CS taper or withdrawal	NA	NA
49	Emirgil et al. (1969)[69]	USA	Retrospective	84	22	6	On CS Withdrawal	NA	NA
50	Morse et al. (1961)[70]	USA	Retrospective	7	7	4	1-3	NA	NA

Biological tests: Serological, histological or other in vitro laboratory tests for determining the sarcoid activity of involve organs. CS: Corticosteroids, NA: Not Available, PFT: Pulmonary Function Test, PS: Pulmonary Sarcoidosis, SSA: Steroi Sparing Agents, Relapse definition: Yes (available).

Geographic location and time period of the studies:
No significant difference was found in the pooled proportions of relapses between Asian (N=18: 0.34, 95% CI: 0.26, 0.41) and non-Asian studies (N=32: 0.43, 95% CI: 0.35, 0.51; p=0.08), or between studies conducted before 2000 CE (N=23: 0.36, 95% CI: 0.29, 0.42) and after 2000 CE (N=27: 0.44, 95% CI: 0.34, 0.54; p=0.16) (Figure 4 Supplementary Figures 7-8).

Meta-regression: Meta-regression revealed that study design (prospective vs. retrospective) significantly affected relapse rates (coefficient = 0.14, p = 0.02), with pooled estimates of 0.32 (95% CI: 0.26–0.38) for prospective studies (N = 21) and 0.44 (95% CI: 0.36–0.54) for retrospective studies (N = 29). However, there were no significant associations with relapse proportions for year (coefficient = 0.08, p = 0.31), Asian population

(coefficient = -0.07, p = 0.33), sample size greater than 50 years or quality of studies (coefficient = -0.03, p = 0.72). (Supplementary Table 5)

.

Patient mean age, sex and ethnicity:
In studies with data available for two groups, the pooled log odds ratios (LORs) for relapsed vs. nonrelapsed patients were not significantly different for mean age [-0.01 (95% CI: -1.47, 1.46); p=0.99] or sex [0.19 (95% CI: -0.12, 0.49); p=0.23]. The mean difference of age in the relapse group vs non-relapse -0.01 (95% CI: -1.47, 1.46); p=0.99 group was for black patients compared with white patients, 0.91 (95% CI: 0.44, 1.37) with a P value of <0.001 indicating statistical significance (Fig. 5).

Radiographic Staging of Disease
In studies with data available for two groups, the pooled LORs of relapse were greater for stage IV disease than for stage I disease [1.04 (95% CI: 0.25, 1.83); p=0.01] and for stage II and III disease than for stage I disease [0.5 (95% CI: 0.07, 1.03); p=0.02]. No significant difference was found for stage II and III vs. stage IV disease [-0.54 (95% CI: -1.39, 0.31); p=0.21] (Fig. 6).

Isolated PSs and Multiorgan PSs:
The pooled proportions of relapses in patients with isolated pulmonary sarcoidosis (PS) and multiorgan PS were not significantly different [0.35 (95% CI: 0.25, 0.46) vs. 0.41 (95% CI: 0.34, 0.48); p=0.35] (Fig. 7).

CS, SSA and combined treatment (CS+SSA):
Corticosteroids (CSs) were used in 70% of the studies, whereas steroid-sparing agent (SSA) and combined CS+SSA treatment were each used in 15% of the studies. The pooled proportions of relapse did not significantly differ across treatments: CS: 0.40 (95% CI: 0.33, 0.47); SSA: 0.47 (95% CI: 0.27, 0.66); combined CS+SSA: 0.2 (95% CI: 0.19, 0.39); p=0.11. Compared with the CS (0.40) and SSA (0.47) treatments, the combined CS+SSA treatment had the lowest pooled proportion of relapses (0.29, 95% CI: 0.19, 0.39) (Fig. 8).

Publication bias: The funnel plot was symmetrical, indicating that there was no visual publication bias. Similarly the regression-based Egger's test [intercept=0.60 (95% CI: -2.07, 3.27), p=0.645] and nonparametric Begg's test (Kendall's tau=0.07, p=0.465) showed no evidence of publication bias (Fig. 9). There was no bias due to missing data, as we excluded studies with missing data.

DISCUSSION:
This study, the first systematic review and meta-analysis on the global incidence of pulmonary sarcoidosis (PS relapse, analysed 50 studies with 3,646 follow-up patients. The key findings included a 40% pooled relaps prevalence (range: 10%–87%). Sensitivity analyses confirmed robustness, with estimates ranging from 0.3 (outliers removed, lower heterogeneity) to 0.41, and most analyses were stable at approximately 0.38–0.40.

Some review articles in the literature have reported that the prevalence of PS relapse ranges from 13–80%, which is consistent with our findings.[7 10 71] The variability in relapse prevalence observed in our study, as well as in thes review articles, may stem from inconsistent definitions of relapse, remission, and PS diagnosis. Although th WASOG and ATS criteria for the diagnosis of sarcoidosis have been available since 1999.[11 53 72], most of th studies have used no standard criteria, such as histological confirmation of the sarcoid diagnosis, leading misdiagnosis of conditions resembling PS or PS relapse.

The timing of relapse after remission varies depending on disease severity and treatment efficacy. Our met[...] analysis revealed that the minimum and maximum median times to relapse were 3 and 13 months, respectivel[...] Most relapses occurred within the first three years, with approximately 10% occurring in the fourth or fifth yea[...] suggesting the need for active patient follow-up for five years. Similarly, several studies in the literature repo[...] relapse occurrence from 1–24 months after remission and, in certain cases, after 24 months of treatme[...] withdrawal.[6 10 53] In accordance with our data, one study reported no further relapses in patients who remaine[...] asymptomatic for three years after prednisone withdrawal.[51]

The type of relapse, characterized by clinical, radiological, pulmonary function, or biochemical parameters, [...] clinically important for improving the prognosis of patients with pulmonary sarcoidosis. Over 60% of patien[...] experienced symptomatic relapses with radiological, functional, or biochemical features, whereas approximate[...] 10% experienced asymptomatic relapses, with the remainder undefined or based on other criteria. In the literatur[...] most studies report clinical and/or radiological relapses, some report functional or biochemical relapses, and mar[...] do not specify relapse types.[5 73] On the basis of these findings, relapse diagnosis should not rely solely on clinic[...] parameters but should include radiological, functional, or biochemical parameters. Asymptomatic relapse[...] detected by biomarkers, may sometimes precede clinical and/or radiological signs, facilitating early detection ar[...] management of relapses.

Relapse-associated risk factors are crucial for determining treatment strategies for this disease but remain large[...] unknown. In this study, we identified several potential risk factors for PS relapse. Compared with white rac[...] black race is a well-documented risk factor for sarcoidosis and is associated with greater disease severity ar[...] mortality.[74 75] Our observation aligns with the high severity of sarcoidosis in the Black population,[74 75] which [...] supported by studies reporting high relapse rates in this group.[28 65 69] The increased relapse risk in Black patien[...] may reflect genetic predispositions (HLA-DRB1*14 allele) or disparities in healthcare, warranting furth[...] investigation.[76] There was no effect of age, gender, geographical region, or year of study—whether in Asia [...] elsewhere or conducted before or after 2000 CE—suggesting the cosmopolitan nature of this problem.

Additionally, we observed that higher radiographic scoring stages of the lungs (i.e., stages II, III, and IV) a[...] important risk factors for PS relapse. The lower probability of disease resolution ranging from 0% (stage IV) [...] 10–40% (stages II and III) than from 50–90% (stage I) supports our findings.[77] The higher relapse rate observed [...] stage IV sarcoidosis patients (pooled LOR: 1.04, 95% CI: 0.25, 1.83) may partly reflect chronic, nonresolvin[...] disease rather than true relapse, particularly given the retrospective nature of many included studies (58[...] retrospective). Some stage IV patients may remain stable without active biomarkers, suggesting that selection bia[...] could overestimate relapse rates in this group. This distinction warrants cautious interpretation of our finding[...] Furthermore, prior studies, including ours and others, have reported higher relapse rates in patients with advance[...] radiographic stages, supporting these findings.[25 34 59] Parenchymal damage or fibrosis in advanced stages ma[...] reflect persistent inflammation, increasing the likelihood of relapse.

To determine whether isolated or systemic PS could be a risk factor for relapse, we compared relapse rat[...] between these subgroups but found no significant difference. However, previous studies have shown a lir[...] between multiorgan involvement and relapse risk,[32 37] reporting high relapse rates in patients with lung,[24] eye[...] heart[31], musculoskeletal system[21] involvement central nervous system (CNS).[27 78]

We investigated whether treatments for pulmonary sarcoidosis (PS), including corticosteroids (CSs; e.g[...] prednisone, prednisolone, and methylprednisolone), immunosuppressive agents (SSAs; e.g., methotrexat[...] azathioprine, and infliximab), or combined regimens (CSs + SSAs), influence the incidence of relapse. Owing [...]

limited data (15% of studies), SSAs were grouped despite their heterogeneity (e.g., methotrexate vs. TNF-alph inhibitors), potentially obscuring specific relapse risks. Pooled relapse rates were similar across the CS, SSA, an combined treatments, indicating that there was no association between treatment type and relapse. Although n significant, the lower relapse rate with combined treatment (0.29) than with CS (0.40) or SSA monotherapy (0.4 suggests a possible synergistic effect, particularly in severe cases, warranting further exploration. In contra several studies have reported that CS monotherapy is a major risk factor for PS relapse.[10 31 53 79] A high relaps rate in corticosteroid (CS)-treated pulmonary sarcoidosis patients was observed when the CS-treated group wa not compared with the non-CS-treated group. Most CS-treated patients had severe disease at baseline, suggestin that high disease severity, not CS therapy, may drive the elevated relapse rate. A recent report showin comparable outcomes between patients with CS- versus MTX-treated sarcoidosis further supports this notion. Interestingly, some reports have revealed a significantly lower relapse rate and better disease outcomes in patien receiving combined treatment (CS + MTX) than in those receiving CS treatment alone.[81 82] Further studies on th combined treatment of PS with CS + MTX or other SSAs may provide new insights into this area.

Determining the factors contributing to the variability in relapse timing among patients is critically important. Relapse rates may differ between spontaneous remission and treatment-induced remission, with the latter potentially posing a higher risk, especially after corticosteroid cessation. However, this study focused solely on treated patients[53] We hypothesize that residual disease activity (subclinical posttreatment activity) drives relapse in pulmonary sarcoidosis patients. The literature supports this finding, showing that posttreatment residual sarcoi activity is significantly more common in patients who relapse than in those who do not relapse.[51] The likelihood of pulmonary sarcoidosis relapse may depend on residual sarcoid activity at treatment withdrawal: moderate activity may cause early relapses, mild activity may cause late relapses, and minimal/no activity may cause no relapses. Treatment type (70% CS, 15% SSA, 15% CS+SSA) and duration influence relapse timing (median: 3–1 months), with premature cessation linked to earlier relapses. Reliable methods to measure sarcoid activity during treatment and follow-up are needed to optimize treatment duration and timing for tapering/withdrawal to minimize relapses.

The studies exhibited high heterogeneity, slightly lower in prospective designs ($I^2 = 86\%$) than in retrospectiv ones ($I^2 = 94\%$). Despite this, the positive 95% prediction interval aligns with the estimated average outcom Prospective studies showed less heterogeneity, suggesting stricter follow-up reduces variability. To account f heterogeneity, a random effects model was applied across all analyses. No studies had a high risk of bias; 31 had low risk, and the rest had a moderate risk. Funnel plot inspection and Egger's/Begg's tests indicated n publication bias Multiple sensitivity analyses, including outlier removal, subgroup analyses stratified by relaps definition, sample size, and study quality, Freeman–Tukey transformation, and leave-one-out analysis, validate the robustness of the findings, which were unaffected by sample size (<50 patients), study quality, or the presenc or absence of a relapse definition. Further substantiation was achieved through meta-regression, confirmin prospective design as the sole significant contributor.

Our meta-analysis revealed a 40% pooled relapse incidence in pulmonary sarcoidosis (PS) patients (range: 18% 88%), which was lower in prospective (32%) studies than in retrospective (44%) studies, indicating that stud design influences relapse rates. A black race and higher-grade radiographic stages were key risk factors f relapse. Additionally, this study provides a solid foundation for developing a consensus definition of PS relaps and other sarcoidosis-related events, including remission, exacerbation, disease progression, and optimal treatme withdrawal timing. While this study offers the best available evidence on PS relapse and its risk factors, furth large-scale prospective cohort studies are essential to validate these findings, identify new biomarkers ar treatments for relapse, and improve routine clinical practice and patient outcomes.

Data Availability Statements: The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

References:

1. Partin M, Clebak KT, Chen R, et al. Sarcoidosis: Evaluation and Treatment. *American family physician* 2024;109(1):19–29. [published Online First: 2024/01/16]

2. Tavana S, Alizadeh M, Mohajerani SA, et al. Pulmonary and extra-pulmonary manifestations of sarcoidosis. *Nigerian medical journal : journal of the Nigeria Medical Association* 2015;56(4):258–62. doi: 10.4103/0300-1652.169702 [published Online First: 2016/01/14]

3. Judson MA. Extrapulmonary sarcoidosis. *Seminars in respiratory and critical care medicine* 2007;28(1):83–101. doi: 10.1055/s-2007-970335 [published Online First: 2007/03/03]

4. Gerke AK. Morbidity and mortality in sarcoidosis. *Current opinion in pulmonary medicine* 2014;20(5):472–8. doi: 10.1097/MCP.0000000000000080 [published Online First: 2014/07/17]

5. Gerke AK. Treatment of Granulomatous Inflammation in Pulmonary Sarcoidosis. *J Clin Med* 2024;13(3):738. doi: 10.3390/jcm13030738 [published Online First: 2024/02/10]

6. Baughman RP, Valeyre D, Korsten P, et al. ERS clinical practice guidelines on treatment of sarcoidosis. *The European respiratory journal* 2021;58(6) doi: 10.1183/13993003.04079-2020 [published Online First: 2021/06/19]

7. Thillai M, Atkins CP, Crawshaw A, et al. BTS Clinical Statement on pulmonary sarcoidosis. *Thorax* 2021;76(1):4–20. doi: 10.1136/thoraxjnl-2019-214348 [published Online First: 2020/12/04]

8. Johns CJ, Michele TM. The Clinical Management of Sarcoidosis A 50-Year Experience at the Johns Hopkins Hospital. *Medicine* 1999;78(2)

9. Shobha V. Pulmonary sarcoidosis – Predictors of relapse: Are we there yet? *Indian Journal of Rheumatology* 2019;14(4) doi: 10.4103/injr.injr_168_19

10. Baughman RP, Judson MA. Relapses of sarcoidosis: what are they and can we predict who will get them? *The European respiratory journal* 2014;43(2):337–9. doi: 10.1183/09031936.00138913 [published Online First: 2014/02/04]

11. Statement on sarcoidosis. Joint Statement of the American Thoracic Society (ATS), the European Respiratory Society (ERS) and the World Association of Sarcoidosis and Other Granulomatous Disorders (WASOG) adopted by the ATS Board of Directors and by the ERS Executive Committee, February 1999. *American journal of respiratory and critical care medicine* 1999;160(2):736–55. doi: 10.1164/ajrccm.160.2.ats4-99 [published Online First: 1999/08/03]

12. Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *Journal of clinical epidemiology* 2012;65(9):934–9. doi: 10.1016/j.jclinepi.2011.11.014 [published Online First: 20120627]

13. Munn Z, Moola S, Lisy K, et al. The Joanna Briggs Institute approach for conducting systematic reviews of prevalence and incidence data. *International Journal of Evidence-Based Healthcare* 2015;13(3):163–69. doi: 10.1097/XEB.0000000000000054

14. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled clinical trials* 1986;7(3):177–88. doi: 10.1016/0197-2456(86)90046-2 [published Online First: 1986/09/01]

15. Thompson SG, Higgins JP. How should meta-regression analyses be undertaken and interpreted? *Stat Med* 2002;21(11):1559–73. doi: 10.1002/sim.1187

16. Viechtbauer W. Bias and Efficiency of Meta-Analytic Variance Estimators in the Random-Effects Model. *Journal of Educational and Behavioral Statistics* 2005;30(3):261–93. doi: 10.3102/10769986030003261

17. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539–58. doi: 10.1002/sim.1186

18. Viechtbauer W, Cheung MW. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods* 2010;1(2):112–25. doi: 10.1002/jrsm.11 [published Online First: 20101004]

19. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50(4):1088–101. [published Online First: 1994/12/01]

20. Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *BMJ (Clinical research ed)* 1997;315(7109):629–34. doi: 10.1136/bmj.315.7109.629 [published Online First: 1997/10/06]

21. Bouchut A, Lhote R, Maksud P, et al. Prognostic value of hypermetabolic bone sarcoidosis observed by 18F-fluorodeoxyglucose positron emission tomography. *Rheumatology (Oxford, England)* 2024 doi: 10.1093/rheumatology/keae019 [published Online First: 2024/01/21]

22. Guliyeva V, Demirkan FG, Yigit RE, et al. A clinical overview of paediatric sarcoidosis: Multicentre experience from Turkey. *Modern rheumatology* 2024;34(3):639–45. doi: 10.1093/mr/road050 [published Online First: 2023/05/27]

23. Dhooria S, Sehgal IS, Agarwal R, et al. High-dose (40 mg) versus low-dose (20 mg) prednisolone for treating sarcoidosis: a randomised trial (SARCORT trial). *The European respiratory journal* 2023;62(3) doi: 10.1183/13993003.00198-2023 [published Online First: 2023/09/11]

24. Sahin Ozdemirel T, Akinci Ozyurek B, Tatci E, et al. Relationships Between Systemic Inflammatory Markers and 18F-FDG PET/CT Imaging and Clinical Findings in Pulmonary Sarcoidosis. *Cureus* 2023;15(3):e36521. doi: 10.7759/cureus.36521 [published Online First: 2023/04/24]

25. Spyropoulos G, Domvri K, Manika K, et al. Clinical, imaging and functional determinants of sarcoidosis phenotypes in a Greek population. *Journal of thoracic disease* 2022;14(6):1941–49. doi: 10.21037/jtd-21-1760 [published Online First: 2022/07/12]

26. Madan K, Sryma PB, Pattnaik B, et al. Clinical Profile of 327 patients with Sarcoidosis in India: An Ambispective Cohort Study in a Tuberculosis (TB) Endemic Population. *Lung India* 2022;39(1):51–57. doi: 10.4103/lungindia.lungindia_960_20 [published Online First: 2022/01/04]

27. Sakkat A, Cox G, Khalidi N, et al. Infliximab therapy in refractory sarcoidosis: a multicenter real-world analysis. *Respiratory research* 2022;23(1):54. doi: 10.1186/s12931-022-01971-5 [published Online First: 2022/03/11]

28. Morar R, Feldman C. Sarcoidosis in Johannesburg, South Africa: A retrospective study. *Afr J Thorac Crit Care Med* 2022;28(4):150–56. doi: 10.7196/AJTCCM.2022.v28i4.205 [published Online First: 2023/02/14]

29. Feng H, Yan L, Zhao Y, et al. Neutrophils in Bronchoalveolar Lavage Fluid Indicating the Severity and Relapse of Pulmonary Sarcoidosis. *Front Med (Lausanne)* 2021;8:787681. doi: 10.3389/fmed.2021.787681 [published Online First: 2022/02/22]

30. Gavrysyuk V, Merenkova I, Dziublyk Y, et al. Efficacy and Tolerability of Methotrexate and Methylprednisolone in a Comparative Assessment of the Primary and Long-Term Outcomes in Patients with Pulmonary Sarcoidosis. *Diagnostics* 2021; 11(7). https://mdpi-res.com/d_attachment/diagnostics/diagnostics-11-01289/article_deploy/diagnostics-11-01289.pdf?version=1626676520.

31. Murata O, Suzuki K, Takeuchi T, et al. Incidence and baseline characteristics of relapse or exacerbation in patients with pulmonary sarcoidosis in Japan. *Sarcoidosis, vasculitis, and diffuse lung diseases : official journal of WASOG* 2021;38(3):e2021026. doi: 10.36141/svdld.v38i3.11327 [published Online First: 2021/11/09]

32. Gavrysyuk V, Merenkova I, Vlasova N, et al. Clinical Factors Associated with the Risk of Pulmonary Sarcoidosis Relapse. *Georgian medical news* 2021(313):84–88. [published Online First: 2021/06/10]

33. Fang C, Zhang Q, Wang N, et al. Effectiveness and tolerability of methotrexate in pulmonary sarcoidosis: A single center real-world study. *Sarcoidosis, vasculitis, and diffuse lung diseases : official journal of WASOG* 2019;36(3):217–27. doi: 10.36141/svdld.v36i3.8449 [published Online First: 2020/06/02]

34. Nath A, Hashim Z, Khan A, et al. Experience of Sarcoidosis and Factors Predicting Relapse at a Tertiary Care Institute in North India. *Indian Journal of Rheumatology* 2019;14(4)

35. Zheng Y, Wang H, Xu Q, et al. Risk factors of relapse in pulmonary sarcoidosis treated with corticosteroids. *Clinical rheumatology* 2019;38(7):1993–99. doi: 10.1007/s10067-019-04507-3 [published Online First: 2019/03/17]

36. Ozyilmaz E, Ozturk OG, Durmaz A, et al. Early prediction of sarcoidosis prognosis with HLA typing: a 5 year follow-up study. *Sarcoidosis, vasculitis, and diffuse lung diseases : official journal of WASOG* 2018;35(3):184–91. doi: 10.36141/svdld.v35i3.6781 [published Online First: 2018/01/01]

37. Belhomme N, Jouneau S, Bouzille G, et al. Role of serum immunoglobulins for predicting sarcoidosis outcome: A cohort study. *PLoS One* 2018;13(4):e0193122. doi: 10.1371/journal.pone.0193122 [published Online First: 2018/04/12]

38. Chen H, Jin R, Wang Y, et al. The Utility of (18)F-FDG PET/CT for Monitoring Response and Predicting Prognosis after Glucocorticoids Therapy for Sarcoidosis. *Biomed Res Int* 2018;2018(1):1823710. doi: 10.1155/2018/1823710 [published Online First: 2018/04/25]

39. Mañá J, Rubio-Rivas M, Villalba N, et al. Multidisciplinary approach and long-term follow-up in a series of 640 consecutive patients with sarcoidosis: Cohort study of a 40-year clinical experience at a tertiary referral center in Barcelona, Spain. *Medicine* 2017;96(29)

40. Maturu VN, Rayamajhi SJ, Agarwal R, et al. Role of serial F-18 FDG PET/CT scans in assessing treatment response and predicting relapses in patients with symptomatic sarcoidosis. *Sarcoidosis, vasculitis, and diffuse lung diseases : official journal of WASOG* 2016;33(4):372–80. [published Online First: 2017/01/13]

41. Liu Y, Qiu L, Wang Y, et al. The Circulating Treg/Th17 Cell Ratio Is Correlated with Relapse and Treatment Response in Pulmonary Sarcoidosis Patients after Corticosteroid Withdrawal. *PLoS One* 2016;11(2):e0148207. doi: 10.1371/journal.pone.0148207 [published Online First: 2016/02/06]

42. Polverosi R, Russo R, Coran A, et al. Typical and atypical pattern of pulmonary sarcoidosis at high-resolution CT: relation to clinical evolution and therapeutic procedures. *Radiol Med* 2014;119(6):384–92. doi: 10.1007/s11547-013-0356-x [published Online First: 2013/12/04]

43. Vorselaars AD, Verwoerd A, van Moorsel CH, et al. Prediction of relapse after discontinuation of infliximab therapy in severe sarcoidosis. *The European respiratory journal* 2014;43(2):602–9. doi: 10.1183/09031936.00055213 [published Online First: 2013/08/31]

44. Sharma SK, Balamurugan A, Pandey RM, et al. Human leukocyte antigen-DR alleles influence the clinical course of pulmonary sarcoidosis in Asian Indians. *Am J Respir Cell Mol Biol* 2003;29(2):225–31. doi: 10.1165/rcmb.2003-0007OC [published Online First: 2003/02/26]

45. Kiper N, Anadol D, Ozcelik U, et al. Inhaled corticosteroids for maintenance treatment in childhood pulmonary sarcoidosis. *Acta paediatrica (Oslo, Norway : 1992)* 2001;90(8):953–6. [published Online First: 2001/09/01]

46. Sharma SK, Mohan A, Guleria JS. Clinical characteristics, pulmonary function abnormalities and outcome of prednisolone treatment in 106 patients with sarcoidosis. *The Journal of the Association of Physicians of India* 2001;49:697–704. [published Online First: 2001/09/28]

47. Baculard A, Blanc N, Boule M, et al. Pulmonary sarcoidosis in children: a follow-up study. *The European respiratory journal* 2001;17(4):628–35. doi: 10.1183/09031936.01.17406280 [published Online First: 2001/06/13]

48. Muller-Quernheim J, Kienast K, Held M, et al. Treatment of chronic sarcoidosis with an azathioprine/prednisolone regimen. *The European respiratory journal* 1999;14(5):1117–22. doi: 10.1183/09031936.99.14511179 [published Online First: 1999/12/22]

49. Kim DS, Paik SH, Lim CM, et al. Value of ICAM-1 expression and soluble ICAM-1 level as a marker of activity in sarcoidosis. *Chest* 1999;115(4):1059–65. doi: 10.1378/chest.115.4.1059 [published Online First: 1999/04/20]

50. Baltzan M, Mehta S, Kirkham TH, et al. Randomized trial of prolonged chloroquine therapy in advanced pulmonary sarcoidosis. *American journal of respiratory and critical care medicine* 1999;160(1):192–7. doi: 10.1164/ajrccm.160.1.9809024 [published Online First: 1999/07/03]

51. Rizzato G, Montemurro L, Colombo P. The late follow-up of chronic sarcoid patients previously treated with corticosteroids. *Sarcoidosis, vasculitis, and diffuse lung diseases : official journal of WASOG* 1998;15(1):52–8. [published Online First: 1998/05/08]

52. Wyser CP, van Schalkwyk EM, Alheit B, et al. Treatment of progressive pulmonary sarcoidosis with cyclosporin A. A randomized controlled trial. *American journal of respiratory and critical care medicine* 1997;156(5):1371–6. doi: 10.1164/ajrccm.156.5.9506031 [published Online First: 1997/12/31]

53. Gottlieb JE, Israel HL, Steiner RM, et al. Outcome in sarcoidosis. The relationship of relapse to corticosteroid therapy. *Chest* 1997;111(3):623–31. doi: 10.1378/chest.111.3.623 [published Online First: 1997/03/01]

54. Lower EE, Baughman RP. Prolonged use of methotrexate for sarcoidosis. *Archives of internal medicine* 1995;155(8):846–51. [published Online First: 1995/04/24]

55. Takada K, Ina Y, Noda M, et al. The clinical course and prognosis of patients with severe, moderate or mild sarcoidosis. *Journal of clinical epidemiology* 1993;46(4):359–66. doi: 10.1016/0895-4356(93)90150-y [published Online First: 1993/04/01]

56. Chadelat K, Baculard A, Grimfeld A, et al. Pulmonary sarcoidosis in children: serial evaluation of bronchoalveolar lavage cells during corticosteroid treatment. *Pediatr Pulmonol* 1993;16(1):41–7. doi: 10.1002/ppul.1950160109 [published Online First: 1993/07/01]

57. Prior C, Haslam PL. Increased levels of serum interferon-gamma in pulmonary sarcoidosis and relationship with response to corticosteroid therapy. *Am Rev Respir Dis* 1991;143(1):53–60. doi: 10.1164/ajrccm/143.1.53 [published Online First: 1991/01/01]

58. Rizzato G, Fraioli P, Montemurro L. Long-term therapy with deflazacort in chronic sarcoidosis. *Chest* 1991;99(2):301–9. doi: 10.1378/chest.99.2.301 [published Online First: 1991/02/01]

59. Baumann MH, Strange C, Sahn SA. Do chest radiographic findings reflect the clinical course of patients with sarcoidosis during corticosteroid withdrawal? *AJR Am J Roentgenol* 1990;154(3):481–5. doi: 10.2214/ajr.154.3.2106208 [published Online First: 1990/03/01]

60. Lower EE, Baughman RP. The use of low dose methotrexate in refractory sarcoidosis. *Am J Med Sci* 1990;299(3):153–7. doi: 10.1097/00000441-199003000-00002 [published Online First: 1990/03/01]

61. Johns CJ, Schonfeld SA, Scott PP, et al. Longitudinal study of chronic sarcoidosis with low-dose maintenance corticosteroid therapy. Outcome and complications. *Ann N Y Acad Sci* 1986;465(1):702–12. doi: 10.1111/j.1749-6632.1986.tb18549.x [published Online First: 1986/01/01]

62. Wallaert B, Ramon P, Fournier EC, et al. High-dose methylprednisolone pulse therapy in sarcoidosis. *European journal of respiratory diseases* 1986;68(4):256–62. [published Online First: 1986/04/01]

63. Turner-Warwick M, McAllister W, Lawrence R, et al. Corticosteroid treatment in pulmonary sarcoidosis: do serial lavage lymphocyte counts, serum angiotensin converting enzyme measurements, and gallium-67 scans help management? *Thorax* 1986;41(12):903–13. doi: 10.1136/thx.41.12.903 [published Online First: 1986/12/01]

64. Eule H, Weinecke A, Roth I, et al. The possible influence of corticosteroid therapy on the natural course of pulmonary sarcoidosis. Late results of a continuing clinical study. *Ann N Y Acad Sci* 1986;465(1):695–701. doi: 10.1111/j.1749-6632.1986.tb18548.x [published Online First: 1986/01/01]

65. Johns CJ, Zachary JB, MacGregor MI, et al. The longitudinal study of chronic sarcoidosis. *Transactions of the American Clinical and Climatological Association* 1983;94:173–81. [published Online First: 1982/01/01]

66. Rohatgi PK, Ryan JW, Lindeman P. Value of serial measurement of serum angiotensin converting enzyme in the management of sarcoidosis. *The American journal of medicine* 1981;70(1):44–50. doi: 10.1016/0002-9343(81)90410-1 [published Online First: 1981/01/01]

67. Johns CJ, Macgregor MI, Zachary JB, et al. Extended experience in the long-term corticosteroid treatment of pulmonary sarcoidosis. *Ann N Y Acad Sci* 1976;278(1):722–31. doi: 10.1111/j.1749-6632.1976.tb47087.x [published Online First: 1976/01/01]

68. Johns CJ, Zachary JB, Ball WC, Jr. A ten-year study of corticosteroid treatment of pulmonary sarcoidosis. *The Johns Hopkins medical journal* 1974;134(5):271–83. [published Online First: 1974/05/01]

69. Emirgil C, Sobol BJ, Williams MH, Jr. Long-term study of pulmonary sarcoidosis. The effect of steroid therapy as evaluated by pulmonary function studies. *J Chronic Dis* 1969;22(2):69–86. doi: 10.1016/0021-9681(69)90060-5 [published Online First: 1969/07/01]

70. Morse SI, Cohn ZA, Hirsch JG, et al. The treatment of sarcoidosis with chloroquine. *The American journal of medicine* 1961;30:779–84. doi: 10.1016/0002-9343(61)90213-3 [published Online First: 1961/05/01]

71. Nagai S, Handa T, Ito Y, et al. Outcome of sarcoidosis. *Clinics in chest medicine* 2008;29(3):565–74, x. doi: 10.1016/j.ccm.2008.03.006 [published Online First: 2008/06/10]

72. Hunninghake GW, Costabel U, Ando M, et al. ATS/ERS/WASOG statement on sarcoidosis. American Thoracic Society/European Respiratory Society/World Association of Sarcoidosis and other Granulomatous Disorders. *Sarcoidosis Vasc Diffuse Lung Dis* 1999;16(2):149–73. [published Online First: 1999/11/24]

73. Crouser ED, Maier LA, Wilson KC, et al. Diagnosis and Detection of Sarcoidosis. An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med* 2020;201(8):e26–e51. doi: 10.1164/rccm.202002-0251ST [published Online First: 2020/04/16]

74. Burke RR, Stone CH, Havstad S, et al. Racial differences in sarcoidosis granuloma density. *Lung* 2009;187(1):1–7. doi: 10.1007/s00408-008-9111-9 [published Online First: 2008/08/22]

75. Mirsaeidi M, Machado RF, Schraufnagel D, et al. Racial difference in sarcoidosis mortality in the United States. *Chest* 2015;147(2):438–49. doi: 10.1378/chest.14-1120 [published Online First: 2014/09/05]

76. da Costa CH, Silva VL, Fabricio-Silva GM, et al. HLA in a cohort of Brazilian patients with sarcoidosis. *Hum Immunol* 2013;74(10):1326–32. doi: 10.1016/j.humimm.2013.07.009 [published Online First: 20130801]

77. Scadding JG. Prognosis of intrathoracic sarcoidosis in England. A review of 136 cases after five years' observation. *British medical journal* 1961;2(5261):1165–72. doi: 10.1136/bmj.2.5261.1165 [published Online First: 1961/11/04]

78. Belperio JA, Fishbein MC, Abtin F, et al. Pulmonary sarcoidosis: A comprehensive review: Past to present. *Journal of autoimmunity* 2023:103107. doi: 10.1016/j.jaut.2023.103107 [published Online First: 2023/10/22]

79. Akram MJ, Khalid U, Abu Bakar M, et al. Sarcoidosis: epidemiology, characteristics, and outcomes over 10 years - a single-center study in Pakistan. *Expert review of respiratory medicine* 2022;16(1):133–43. doi: 10.1080/17476348.2021.1924062 [published Online First: 2021/08/18]

80. Vis R, Mathijssen H, Keijsers RGM, et al. Prednisone vs methotrexate in treatment naive cardiac sarcoidosis. *Journal of nuclear cardiology : official publication of the American Society of Nuclear Cardiology* 2023;30(4):1543–53. doi: 10.1007/s12350-022-03171-6 [published Online First: 2023/01/15]

81. Ballul T, Borie R, Crestani B, et al. Treatment of cardiac sarcoidosis: A comparative study of steroids and steroids plus immunosuppressive drugs. *International journal of cardiology* 2019;276:208–11. doi: 10.1016/j.ijcard.2018.11.131 [published Online First: 2018/12/12]

82. Nagai S, Yokomatsu T, Tanizawa K, et al. Treatment with methotrexate and low-dose corticosteroids in sarcoidosis patients with cardiac lesions. *Internal medicine (Tokyo, Japan)* 2014;53(5):427–33. doi: 10.2169/internalmedicine.53.0794 [published Online First: 2014/03/04]

Study No	Study (Year)	Frame	Method	Size	Description	Coverage	Validity	Reliability	Analysis	Response	Bias	Score	Risk
1	Bouchut et al. (2024)	Low	High	Low	Low	Low	Low	Low	Low	Low	Low	9	Low
2	Guliyeva et al. (2024)	Low	High	Low	Low	Low	Low	Low	Low	Low	Low	9	Low
3	Dhooria et al. (2023)	Low	Low	Low	Low	Low	Low	Low	Low	Low	Low	10	Low
4	Ozdemirel et al. (2023)	Low	High	Low	Low	Low	High	High	Low	Low	Low	7	Moderate
5	Spyropoulos et al. (2022)	Low	Low	Low	Low	Low	High	Low	Low	Low	Low	9	Low
6	Madan et al. (2022)	Low	Low	Low	Low	Low	Low	Low	Low	Low	Low	10	Low
7	Sakkat et al. (2022)	Low	High	High	High	Low	High	Low	Low	Low	Low	6	Moderate
8	Morar et al. (2022)	Low	High	Low	Low	Low	High	Low	Low	Low	Low	8	Low
9	Feng et al. (2022)	Low	High	Low	Low	Low	Low	Low	Low	Low	Low	9	Low
10	Gavrysyuk et al. (2021)	Low	Low	Low	Low	Low	Low	Low	Low	Low	Low	10	Low
11	Murata et al. (2021)	Low	High	Low	Low	Low	Low	Low	Low	Low	Low	9	Low
12	Gavrysyuk et al. (2021)	Low	Low	Low	Low	Low	Low	Low	Low	Low	Low	10	Low
13	Fang et al. (2019)	Low	High	High	Low	Low	Low	Low	Low	Low	Low	8	Low
14	Nath et al. (2019)	Low	High	Low	Low	Low	Low	Low	Low	Low	Low	9	Low
15	Zheng et al. (2019)	Low	High	Low	High	Low	Low	Low	Low	Low	Low	8	Low
16	Ozyilmaz et al. (2018)	Low	Low	High	High	Low	High	Low	Low	Low	Low	7	Moderate
17	Belhomme et al. (2018)	Low	High	High	High	Low	Low	Low	Low	Low	Low	7	Moderate
18	Chen et al. (2018)	Low	High	High	Low	Low	Low	Low	Low	Low	Low	8	Low
19	Mañá et al. (2017)	Low	Low	Low	High	Low	Low	Low	Low	Low	Low	9	Low
20	Maturu et al. (2016)	Low	Low	High	Low	Low	Low	Low	Low	Low	Low	9	Low
21	Liu et al. (2016)	Low	Low	High	Low	Low	Low	Low	Low	Low	Low	9	Low
22	Polverosi et al. (2014)	Low	High	High	Low	Low	High	Low	Low	Low	Low	7	Moderate
23	Vorselaars et al. (2014)	Low	High	High	Low	Low	Low	Low	Low	Low	Low	8	Low
24	Sharma et al. (2003)	Low	High	Low	Low	Low	High	Low	Low	Low	Low	8	Low
25	Kiper et al. (2001)	Low	High	High	Low	Low	Low	Low	Low	Low	Low	8	Low
26	Sharma et al. (2001)	Low	Low	Low	Low	Low	Low	Low	Low	Low	Low	10	Low
27	Baculard et al. (2001)	Low	High	High	Low	Low	Low	Low	Low	Low	Low	8	Low
28	Muller-Quernheim et al. (1999)	Low	Low	High	Low	Low	High	Low	Low	Low	Low	8	Low
29	Kim et al. (1999)	Low	Low	High	Low	Low	High	High	Low	Low	Low	7	Moderate
30	Baltzan et al. (1999)	Low	Low	High	Low	Low	Low	Low	Low	Low	Low	9	Low
31	Rizzato et al. (1998)	Low	High	Low	High	Low	High	Low	Low	Low	Low	7	Moderate
32	Wyser et al. (1997)	Low	Low	High	High	Low	Low	Low	Low	Low	Low	8	Low
33	Gottlieb et al. (1997)	Low	Low	Low	High	Low	Low	Low	Low	Low	Low	9	Low
34	Lower et al. (1995)	Low	High	High	High	Low	Low	Low	Low	Low	Low	7	Moderate
35	Takada et al. (1993)	Low	High	Low	High	Low	High	Low	Low	Low	Low	7	Moderate
36	Chadelat et al. (1993)	Low	Low	High	Low	Low	High	Low	Low	Low	Low	8	Low
37	Prior et al. (1991)	Low	Low	High	Low	Low	High	Low	Low	Low	Low	8	Low
38	Rizzato et al. (1991)	Low	High	High	Low	Low	High	Low	Low	Low	Low	7	Moderate
39	Baumann et al. (1990)	Low	High	High	Low	Low	High	Low	Low	Low	Low	7	Moderate
40	Lower et al. (1990)	Low	High	High	Low	Low	High	Low	Low	Low	Low	7	Moderate
41	Johns et al. (1986)	Low	High	Low	Low	Low	High	High	Low	Low	Low	7	Moderate
42	Wallaert et al. (1986)	Low	Low	High	Low	Low	High	High	Low	Low	Low	7	Moderate
43	Turner-Warwick et al. (1986)	Low	Low	High	Low	Low	High	High	Low	Low	Low	7	Moderate
44	Eule et al. (1986)	Low	Low	Low	High	Low	High	Low	Low	Low	Low	8	Low
45	Johns et al. (1983)	Low	High	Low	Low	Low	High	Low	Low	Low	Low	8	Low
46	Rohatgi et al. (1981)	Low	Low	High	High	Low	High	Low	Low	Low	Low	7	Moderate
47	Johns et al. (1976)	Low	High	Low	High	Low	High	Low	Low	Low	Low	7	Moderate
48	Johns et al. (1974)	Low	High	Low	Low	Low	High	Low	Low	Low	Low	8	Low
49	Emirgil et al. (1969)	Low	High	High	High	Low	High	Low	Low	Low	Low	6	Moderate
50	Morse et al. (1961)	Low	High	High	Low	Low	High	Low	Low	Low	Low	7	Moderate

Table 2: Quality assessment of studies using the Hey et al. (2012) risk of bias tool. Columns: Study No (study identifier), Study (Year) (author, publication year), Frame (sampling frame quality), Method (study methodology rigor), Size (sample size adequacy), Description (clarity of study description), Coverage (population coverage), Validity (measurement validity), Reliability (measurement reliability), Analysis (analytical rigor), Response (response rate quality), Bias (potential bias), Score (overall quality score), Risk (risk of bias: Low, Moderate).

Fig. 1

The PRISMA flow chart shows the selection of eligible studies

Study	Number of relapses	Followed-up		Proportion with 95% CI	Weight (%)
Bouchut et al. (2024)	91	173		0.53 [0.45, 0.60]	2.23
Guliyeva et al. (2024)	14	52		0.27 [0.15, 0.39]	2.11
Dhooria et al. (2023)	39	86		0.45 [0.35, 0.56]	2.15
Ozdemirel et al. (2023)	16	73		0.22 [0.12, 0.31]	2.18
Spyropoulos et al. (2022)	42	147		0.29 [0.21, 0.36]	2.23
Madan et al. (2022)	73	327		0.22 [0.18, 0.27]	2.28
Sakkat et al. (2022)	7	11		0.64 [0.35, 0.92]	1.50
Morar et al. (2022)	54	89		0.61 [0.51, 0.71]	2.16
Feng et al. (2022)	24	123		0.20 [0.13, 0.27]	2.24
Gavrysyuk et al. (2021)	28	84		0.33 [0.23, 0.43]	2.16
Murata et al. (2021)	23	103		0.22 [0.14, 0.30]	2.22
Gavrysyuk et al. (2021)	56	108		0.52 [0.42, 0.61]	2.18
Fang et al. (2019)	2	11		0.18 [0.00, 0.41]	1.71
Nath et al. (2019)	29	120		0.24 [0.17, 0.32]	2.22
Zheng et al. (2019)	30	96		0.31 [0.22, 0.41]	2.19
Ozyilmaz et al. (2018)	13	29		0.45 [0.27, 0.63]	1.89
Belhomme et al. (2018)	10	34		0.29 [0.14, 0.45]	2.00
Chen et al. (2018)	12	23		0.52 [0.32, 0.73]	1.81
Mañá et al. (2017)	30	266		0.11 [0.07, 0.15]	2.29
Maturu et al. (2016)	10	27		0.37 [0.19, 0.55]	1.89
Liu et al. (2016)	20	42		0.48 [0.33, 0.63]	2.00
Polverosi et al. (2014)	9	44		0.20 [0.09, 0.32]	2.11
Vorselaars et al. (2014)	29	47		0.62 [0.48, 0.76]	2.05
Sharma et al. (2003)	40	53		0.75 [0.64, 0.87]	2.12
Kiper et al. (2001)	6	15		0.40 [0.15, 0.65]	1.64
Sharma et al. (2001)	21	85		0.25 [0.16, 0.34]	2.19
Baculard et al. (2001)	4	21		0.19 [0.02, 0.36]	1.94
Muller-Quernheim et al. (1999)	3	11		0.27 [0.01, 0.54]	1.58
Kim et al. (1999)	4	27		0.15 [0.01, 0.28]	2.06
Baltzan et al. (1999)	8	18		0.44 [0.21, 0.67]	1.71
Rizzato et al. (1998)	30	82		0.37 [0.26, 0.47]	2.15
Wyser et al. (1997)	7	16		0.44 [0.19, 0.68]	1.65
Gottlieb et al. (1997)	86	221		0.39 [0.32, 0.45]	2.25
Lower et al. (1995)	35	40		0.88 [0.77, 0.98]	2.16
Takada et al. (1993)	36	76		0.47 [0.36, 0.59]	2.13
Chadelat et al. (1993)	2	8		0.25 [0.00, 0.55]	1.44
Prior et al. (1991)	6	25		0.24 [0.07, 0.41]	1.95
Rizzato et al. (1991)	4	40		0.10 [0.01, 0.19]	2.19
Baumann et al. (1990)	10	15		0.67 [0.43, 0.91]	1.67
Lower et al. (1990)	5	14		0.36 [0.11, 0.61]	1.62
Johns et al. (1986)	128	171		0.75 [0.68, 0.81]	2.25
Wallaert et al. (1986)	8	12		0.67 [0.40, 0.93]	1.56
Turner-Warwick et al. (1986)	4	32		0.12 [0.01, 0.24]	2.12
Eule et al. (1986)	24	110		0.22 [0.14, 0.30]	2.22
Johns et al. (1983)	74	99		0.75 [0.66, 0.83]	2.20
Rohatgi et al. (1981)	6	20		0.30 [0.10, 0.50]	1.82
Johns et al. (1976)	102	146		0.70 [0.62, 0.77]	2.23
Johns et al. (1974)	113	145		0.78 [0.71, 0.85]	2.24
Emirgil et al. (1969)	6	22		0.27 [0.09, 0.46]	1.88
Morse et al. (1961)	4	7		0.57 [0.20, 0.94]	1.21
Overall				0.40 [0.34, 0.46]	

Heterogeneity: τ^2 = 0.04, I^2 = 93.81%, H^2 = 16.16

Test of $\theta_i = \theta_j$: Q(49) = 1028.61, p = 0.00

Test of θ = 0: z = 13.30, p = 0.00

Random-effects REML model

Legend for Figure 2:
Forest plot showing the pooled proportion of the prevalence of sarcoidosis relapse in follow-up patients in the included studies. The blue square represents the prevalence, and the horizontal bars represent the 95% confidence intervals (CIs). The diamond at the bottom represents the pooled prevalence with a 95% CI.

(a)   (b)

Legend for Figure 3:
Relapse types and time in follow-up patients with pulmonary sarcoidosis. (a) Pie chart showing different types of relapses observed in the patients and (b) Box plots showing the distribution of median times to relapse, with the left plot depicting the minimum median time (3 months, range 1-12 months) and the right plot depicting the maximum median time (13 months, range 2-60 months), including interquartile ranges.

Study	K		Proportion with 95% CI
Design of study			
Prospective	21		0.32 [0.26, 0.38]
Retrospective	29		0.45 [0.36, 0.54]
Test of group differences: $Q_b(1) = 6.07$, p = 0.01			
Relapse definition			
Yes	22		0.38 [0.30, 0.45]
No	28		0.41 [0.33, 0.50]
Test of group differences: $Q_b(1) = 0.37$, p = 0.54			
Year of study			
After 2000	27		0.36 [0.29, 0.42]
Before 2000	23		0.44 [0.34, 0.54]
Test of group differences: $Q_b(1) = 1.95$, p = 0.16			
Asian vs Non-Asian			
Asian	18		0.34 [0.26, 0.41]
Non-Asian	32		0.43 [0.35, 0.51]
Test of group differences: $Q_b(1) = 2.98$, p = 0.08			
Studies with patients>50			
Yes	24		0.41 [0.33, 0.50]
No	26		0.38 [0.30, 0.46]
Test of group differences: $Q_b(1) = 0.32$, p = 0.57			
Quality of study: Risk of bias			
Low	31		0.38 [0.32, 0.45]
Moderate	19		0.42 [0.31, 0.53]
Test of group differences: $Q_b(1) = 0.36$, p = 0.55			
Overall			0.40 [0.34, 0.46]

Heterogeneity: $\tau^2 = 0.04$, $I^2 = 93.81\%$, $H^2 = 16.16$

Test of $\theta_i = \theta_j$: Q(49) = 1028.61, p = 0.00

Random-effects REML model

Legend for Figure 4: Forest plots showing pooled proportions of the prevalence of relapse in subgroups according to study design, definition of relapse, year of study, place of study, sample size and quality

Legend to Figure 5: Forest plot showing (a) pooled mean differences in the ages of patients with pulmonary sarcoidosis at relapse, (b) log odds ratios for the sex of patients with pulmonary sarcoidosis, and (c) log odds ratios for the race of patients with pulmonary sarcoidosis

Study	Relapse	No-relapse	Relapse	No-relapse		Log odds-ratio with 95% CI	Weight (%)
Stage IV vs Stage I							
Dhooria et al. (2023)	31	37	4	9		0.63 [-0.64, 1.90]	5.83
Ozdemirel et al. (2023)	13	27	2	18		1.47 [-0.14, 3.07]	4.59
Spyropoulos et al. (2022)	31	53	10	51		1.09 [0.28, 1.90]	8.01
Murata et al. (2021)	5	34	4	20		-0.31 [-1.73, 1.12]	5.21
Nath et al. (2019)	16	66	3	14		0.12 [-1.24, 1.48]	5.46
Zheng et al. (2019)	18	51	6	11		-0.44 [-1.57, 0.70]	6.45
Kiper et al. (2001)	6	5	0	4		2.36 [-0.77, 5.50]	1.75
Baculard et al. (2001)	3	5	1	12		1.97 [-0.52, 4.47]	2.53
Eule et al. (1986)	17	52	7	34		0.46 [-0.52, 1.44]	7.16
Emirgil et al. (1969)	5	21	0	1		-0.26 [-3.60, 3.07]	1.58
Heterogeneity: τ^2= 0.10, I^2= 17.01%, H^2= 1.21						0.55 [0.07, 1.03]	
Test of θ_i= θ_j: Q(9) = 10.48, p = 0.31							
Test of θ = 0: z = 2.24, p = 0.02							
Stage II, III vs Stage I							
Dhooria et al. (2023)	4	1	4	9		2.20 [-0.29, 4.68]	2.54
Ozdemirel et al. (2023)	1	12	2	18		-0.29 [-2.80, 2.22]	2.51
Spyropoulos et al. (2022)	1	1	10	51		1.63 [-1.22, 4.48]	2.05
Murata et al. (2021)	0	6	4	20		-1.05 [-4.10, 2.00]	1.83
Nath et al. (2019)	10	11	3	14		1.45 [-0.07, 2.96]	4.90
Zheng et al. (2019)	6	4	6	11		1.01 [-0.60, 2.62]	4.57
Kiper et al. (2001)	0	0	0	4		2.20 [-2.69, 7.09]	0.80
Baculard et al. (2001)	0	0	1	12		2.12 [-2.15, 6.39]	1.02
Eule et al. (1986)	0	0	7	34		1.53 [-2.47, 5.52]	1.15
Emirgil et al. (1969)	1	8	0	1		-0.64 [-4.28, 3.00]	1.36
Heterogeneity: τ^2= 0.00, I^2= 0.00%, H^2= 1.00						1.04 [0.25, 1.83]	
Test of θ_i= θ_j: Q(9) = 5.48, p = 0.79							
Test of θ = 0: z = 2.59, p = 0.01							
Stage II, III vs Stage IV							
Dhooria et al. (2023)	31	37	4	1		-1.56 [-3.81, 0.68]	2.96
Ozdemirel et al. (2023)	13	27	1	12		1.75 [-0.39, 3.90]	3.16
Spyropoulos et al. (2022)	31	53	1	1		-0.54 [-3.34, 2.27]	2.10
Murata et al. (2021)	5	34	0	6		0.73 [-2.29, 3.74]	1.87
Nath et al. (2019)	16	66	10	11		-1.32 [-2.34, -0.31]	6.99
Zheng et al. (2019)	18	51	6	4		-1.45 [-2.82, -0.07]	5.41
Kiper et al. (2001)	6	5	0	0		0.17 [-3.91, 4.25]	1.11
Baculard et al. (2001)	3	5	0	0		-0.45 [-4.59, 3.69]	1.08
Eule et al. (1986)	17	52	0	0		-1.10 [-5.06, 2.86]	1.17
Emirgil et al. (1969)	5	21	1	8		0.64 [-1.65, 2.94]	2.86
Heterogeneity: τ^2= 0.51, I^2= 30.14%, H^2= 1.43						-0.54 [-1.39, 0.31]	
Test of θ_i= θ_j: Q(9) = 10.55, p = 0.31							
Test of θ = 0: z = -1.25, p = 0.21							
Overall						0.35 [-0.11, 0.80]	
Heterogeneity: τ^2= 0.50, I^2= 37.64%, H^2= 1.60							
Test of θ_i= θ_j: Q(29) = 42.39, p = 0.05							
Test of θ = 0: z = 1.50, p = 0.13							
Test of group differences: Q_b(2) = 7.55, p = 0.02							

Random-effects REML model

Legend for Figure 6:
Forest plot showing pooled log odd ratios of the prevalence of relapse in pulmonary sarcoidosis patients for: Top: radiographic Stage IV vs Stage I Middle: Stage II & III vs Stage I Lower: Stage II & III combined vs Stage IV

Legend for Figure 7:

Forest plots showing pooled proportions of relapses in patients with isolated single-organ pulmonary sarcoidosis and patients with multiorgan systemic pulmonary sarcoidosis.

Legend for Figure 8:

Forest plot showing the pooled proportions of relapses in pulmonary sarcoidosis patients receiving steroids, steroid-sparing agents and mixed treatment (steroid+steroid-sparing agents)

Legend to Figure 9:
Symmetrical funnel plot depicting no visual publication bias. The numbers adjacent to the marker depict the study numbers given in Table 1. The box at the upper right corner of the plot shows the numerical values of Egger's test and the nonparametric rank correlation (Begg) test.

Fig 10: Figure 10: Line graphs showing trajectories of pulmonary sarcoidosis evolution, including (A) disease progression, (B) disease exacerbation, and (D) disease relapse.

Supplementary Figures

Supplementary Figure 1: Boujat plot with study numbers. Study number 34, 48, 41, 45 and 24 identified as outliers

Supplementary Forest plots (Figures 2-7)

Each forest plot displays the proportion of relapses with 95% confidence intervals (CIs) for individual studies, alongside an overall estimate using a random-effects REML model. Studies are listed with the number of relapses and total number of followed-up patients. The size of each study's square reflects its weight (%). Heterogeneity is assessed via τ^2, I^2, and H^2, with Q and p values for study differences and z and p values for the overall effect. The x-axis represents the proportion of relapses (0 to 1).

Supplementary Figure 2: Omitting outliers, model. Study numbers 34, 48, 41, 45 and 24 were identified as outliers and hence removed

Supplementary Figure 3: (a) Prospective studies and (b) retrospective studies

Supplementary Figure 4: (a) Studies with relapse definition (b) Studies without relapse definition

Supplementary Figure 5: (a) Studies with low risk of bias (b) Studies without moderate risk of bias

Supplementary Figure 6: (a) Studies with patient follow-up ≥50 (b) Studies with patient follow-up <50

Supplementary Figure 7: (a) Studies form Asian countries and (b) Studies form non-Asian countries

Supplementary Figure 8: (a) Studies after 2000 CE and (b) before 2000 CE

Supplementary Figure 9: Leave-one-out analysis by sequentially excluding each study and recalculating the pooled prevalence estimate

Supplementary Tables and Analysis

Supplementary Table 1: Sarcoidosis Relapse Definitions

Study No	Study (Year)	Definition
1	Bouchut et al. (2024)	Radiological changes (e.g., new nodules or lymphadenopathy on CT) or biological markers (e.g., elevated serum ACE levels) indicating disease recurrence, necessitating treatment.
2	Guliyeva et al. (2024)	Recurrence of clinical symptoms (e.g., cough, dyspnea) or signs post-remission, after at least 6 months of absent inflammatory activity, with or without ongoing treatment.
3	Dhooria et al. (2023)	Worsening of clinical symptoms (e.g., dyspnea, fatigue), pulmonary function tests (e.g., reduced FEV1 or FVC), or radiological findings (e.g., new infiltrates on chest imaging).
8	Morar et al. (2022)	Recurrence of sarcoidosis symptoms or signs (e.g., respiratory or systemic manifestations) during tapering or discontinuation of anti-sarcoidosis medications, such as corticosteroids.
9	Feng et al. (2022)	Recurrence of sarcoidosis symptoms or signs (e.g., cough, shortness of breath, or radiological changes) after complete disease control, indicating relapse.
10	Gavrysyuk et al. (2021)	Worsening of clinical symptoms (e.g., dyspnea, cough) and/or radiological findings (e.g., new lymphadenopathy or infiltrates on CT) post-normalization, requiring retreatment.
11	Murata et al. (2021)	Reappearance of symptoms (e.g., dyspnea, dry cough) or radiological signs (e.g., bilateral hilar lymphadenopathy or

Study No	Study (Year)	Definition
		parenchymal infiltration on chest X-ray/CT) post-remission, or sudden worsening of stable disease.
12	Gavrysyuk et al. (2021)	Appearance of new sarcoidosis areas on CT (e.g., pulmonary infiltrates) with or without worsening clinical symptoms (e.g., cough), necessitating retreatment.
13	Fang et al. (2019)	New pulmonary sarcoidosis symptoms (e.g., cough, shortness of breath) or exacerbation of existing symptoms, confirmed by clinical or radiological assessment.
14	Nath et al. (2019)	Worsening of clinical symptoms (e.g., respiratory distress), radiological findings (e.g., new nodules on CT), or biological markers (e.g., elevated serum ACE or inflammatory markers).
15	Zheng et al. (2019)	A chest HRCT showing radiographic progression in combination of worsening of clinical symptoms to warrant retreatment following a decrease in dose or discontinuation of corticosteroids, without alternative causes such as infections, heart failure, or pulmonary embolism.
17	Belhomme et al. (2018)	Progressive disease requiring escalation of corticosteroid therapy (e.g., increased dose) or resumption of corticosteroids after cessation, due to worsening symptoms or imaging findings.
18	Chen et al. (2018)	Progression of clinical symptoms (e.g., increased cough or dyspnea) and/or radiological changes (e.g., new lesions on imaging) post-clinical and radiological normalization.
19	Mañá et al. (2017)	The reappearance of former signs or symptoms associated with pulmonary sarcoidosis after remission, such as BHL or parenchymal infiltration on chest X-rays or chest CT scans, or appearance of dyspnea or dry cough; "exacerbation" was the sudden worsening of stable disease, such as BHL, parenchymal infiltration or fibrosis on chest X-rays or chest CT scans, or increased frequency of dyspnea or dry cough; "improved," was amelioration of signs or symptoms associated with pulmonary sarcoidosis; and "stable," was neither "relapse," "exacerbation" nor "improved."

Study No	Study (Year)	Definition
20	Maturu et al. (2016)	Worsening of pulmonary or extrapulmonary symptoms (e.g., dyspnea, skin lesions) and/or radiological progression (e.g., new infiltrates) without alternative causes, requiring reintroduction of steroids or steroid-sparing agents.
25	Kiper et al. (2001)	Presence of more than two of these criteria: Worsening of radiological findings and pulmonary function tests, occurrence of hepatosplenomegaly, new eye involvement, elevation of serum globulins, alkaline phosphatase and serum ACE, hypercalcemia and hypercalciuria..
27	Baculard et al. (2001)	Return of respiratory symptoms (e.g., cough, shortness of breath) after normalization of radiological findings and pulmonary function tests (e.g., normal FEV1, FVC).
30	Baltzan et al. (1999)	A decline in lung function from time 0 (the beginning of run in) in any value of FEV1, FVC, or DLCO-SB of 15% or more for two consecutive measurements. Radiography alone was not used to define relapse.
32	Wyser et al. (1997)	Clinical deterioration (e.g., worsening respiratory symptoms) at least 3 months after treatment cessation, or failure after $\geq$ 9 months of treatment, requiring intervention.
33	Gottlieb et al. (1997)	Return of severe clinical symptoms (e.g., dyspnea, fatigue) warranting treatment, after > 1 month of symptom-free remission without corticosteroids.
34	Lower et al. (1995)	Increased sarcoidosis symptoms (e.g., cough, shortness of breath) or worsening pulmonary function test results (e.g., reduced FEV1 or DLCO).

ACE: Angiotensin-Converting Enzyme, BHL: Bilateral Hilar Lymphadenopathy, CS: Corticosteroids, CT: Computed Tomography, DLCO: Diffusing Capacity of the Lung for Carbon Monoxide, DLCO-SB: Diffusing Capacity of the Lung for Carbon Monoxide – Single Breath, FEV1: Forced Expiratory Volume in One Second, FVC: Forced Vital Capacity, HRCT: High-Resolution Computed Tomography.

Supplementary Table 2: Sensitivity analysis

Analysis Type	Studies	Pooled ES (95% CI)	I² (%)	τ²
Leave-one-out	50	0.40 (0.33, 0.46)	95	0.05
Removing outliers	45	0.35 (0.30, 0.40)	89	0.02
Freeman-Tukey	50	0.40 (0.33, 0.46)	95	0.05
Prospective Studies	22	0.32 (0.30, 0.45)	88	0.03
Retrospective Studies	28	0.45 (0.33, 0.50)	97	0.04
Studies with relapse definition	22	0.38 (0.30, 0.45)	92	0.03
Follow-up < 50 studies	24	0.41 (0.33, 0.50)	97	0.04
High-quality studies	31	0.38 (0.32, 0.45)	97	0.04

Supplementary Table 3: Meta-regression

```
Effect-size label: Proportion
      Effect size: _meta_es
        Std. err.: _meta_se

Random-effects meta-regression          Number of obs  =        50
Method: REML                            Residual heterogeneity:
                                                      tau2 =   .03502
                                                    I2 (%) =    92.37
                                                        H2 =    13.10
                                             R-squared (%) =     9.39
                                            Wald chi2(6)   =     9.43
                                            Prob > chi2    =    0.1507

    _meta_es │  Coefficient  Std. err.      z    P>|z|     [95% conf. interval]
─────────────┼────────────────────────────────────────────────────────────────
     rel_def │   -.0219262    .0734148    -0.30   0.765    -.1658166     .1219642
        year │    .0948448    .0784528     1.21   0.227    -.0589198     .2486094
      design │    .1398559      .06235     2.24   0.025     .0176521     .2620596
       asian │    .0705943    .0690715     1.02   0.307    -.0647833     .2059718
   more_fifty │   -.0464423    .0646201    -0.72   0.472    -.1730954     .0802108
     quality │   -.0291071    .0814531    -0.36   0.721    -.1887521      .130538
       _cons │    .0662692    .1720054     0.39   0.700    -.2708552     .4033937

Test of residual homogeneity: Q_res = chi2(43) = 569.37  Prob > Q_res = 0.0000
```

Outlier analysis: A Baujat plot is depicted in Supplementary Figure 1. On the Baujat plot, the horizontal axis denotes each study's contribution to the overall heterogeneity (Cochran Q-test), while the vertical axis indicates each study's impact on the summary effect size. To facilitate understanding, we subdivided the Baujat plot into four quadrants. Studies located in Quadrants 2 and 3, representing the far right of the horizontal axis, substantially contribute to observed heterogeneity. Data from the upper quadrants (1 and 2) exerted a considerable influence on the overall meta-analysis result. The study number 34, 48, 41, 45 and 24 were found to be outliers. After the removal of these studies, the pooled prevalence of relapse from the rest of the studies was 0.35 (95% CI: 30, 40)

Sensitivity analysis using the Freeman–Tukey double arcsine transformation for relapse proportions across 50 studies shows a random pooled effect size (ES) of 0.40 (95% CI: 0.33 to 0.46)

```
.  metaprop relapse followed, random study(study) freeman ci

             Study    |    ES    [95% Conf. Interval]    % Weight
--------------------+----------------------------------------------------
1                     |    0.53      0.45      0.60    2.18
2                     |    0.27      0.17      0.40    2.09
3                     |    0.45      0.35      0.56    2.12
4                     |    0.22      0.14      0.33    2.15
5                     |    0.29      0.22      0.36    2.18
6                     |    0.22      0.18      0.27    2.22
7                     |    0.64      0.35      0.85    1.59
8                     |    0.61      0.50      0.70    2.13
9                     |    0.20      0.13      0.27    2.19
10                    |    0.33      0.24      0.44    2.13
11                    |    0.22      0.15      0.31    2.17
12                    |    0.52      0.43      0.61    2.15
13                    |    0.18      0.05      0.48    1.77
14                    |    0.24      0.17      0.33    2.18
15                    |    0.31      0.23      0.41    2.15
16                    |    0.45      0.28      0.62    1.92
17                    |    0.29      0.17      0.46    2.00
18                    |    0.52      0.33      0.71    1.85
19                    |    0.11      0.08      0.16    2.23
20                    |    0.37      0.22      0.56    1.92
21                    |    0.48      0.33      0.62    2.01
22                    |    0.20      0.11      0.35    2.09
23                    |    0.62      0.47      0.74    2.04
24                    |    0.75      0.62      0.85    2.10
25                    |    0.40      0.20      0.64    1.71
26                    |    0.25      0.17      0.35    2.15
27                    |    0.19      0.08      0.40    1.96
28                    |    0.27      0.10      0.57    1.66
29                    |    0.15      0.06      0.32    2.06
30                    |    0.44      0.25      0.66    1.77
31                    |    0.37      0.27      0.47    2.13
32                    |    0.44      0.23      0.67    1.72
33                    |    0.39      0.33      0.45    2.20
34                    |    0.88      0.74      0.95    2.13
35                    |    0.47      0.37      0.58    2.11
36                    |    0.25      0.07      0.59    1.54
37                    |    0.24      0.11      0.43    1.96
38                    |    0.10      0.04      0.23    2.15
39                    |    0.67      0.42      0.85    1.74
40                    |    0.36      0.16      0.61    1.70
41                    |    0.75      0.68      0.81    2.20
42                    |    0.67      0.39      0.86    1.65
43                    |    0.12      0.05      0.28    2.10
44                    |    0.22      0.15      0.30    2.18
45                    |    0.75      0.65      0.82    2.16
46                    |    0.30      0.15      0.52    1.86
47                    |    0.70      0.62      0.77    2.18
48                    |    0.78      0.71      0.84    2.19
49                    |    0.27      0.13      0.48    1.91
50                    |    0.57      0.25      0.84    1.33
--------------------+----------------------------------------------------
Random pooled  ES    |    0.40      0.33      0.46  100.00
--------------------+----------------------------------------------------

   Heterogeneity chi^2 =    1028.61 (d.f. = 49) p =        0.00
   I^2 (variation in ES attributable to heterogeneity) =    95.24%
   Estimate of between-study variance Tau^2 =        0.05

   Test of ES=0 : z=      11.77 p =        0.00
```

Random effect model

Effect-size label: Proportion
Effect size: _meta_es
Std. err.: _meta_se
Study label: study

Meta-analysis summary		Number of studies =	50	
Random-effects model		Heterogeneity:		
Method: REML		tau2 =	0.0387	
		I2 (%) =	93.81	
		H2 =	16.16	

Study	Proportion	[95% conf. interval]		% weight
Bouchut et al. (2024)	0.526	0.452	0.600	2.23
Guliyeva et al. (2024)	0.269	0.149	0.390	2.11
Dhooria et al. (2023)	0.453	0.348	0.559	2.15
Ozdemirel et al. (2023)	0.219	0.124	0.314	2.18
Spyropoulos et al. (2022)	0.286	0.213	0.359	2.23
Madan et al. (2022)	0.223	0.178	0.268	2.28
Sakkat et al. (2022)	0.636	0.352	0.921	1.50
Morar et al. (2022)	0.607	0.505	0.708	2.16
Feng et al. (2022)	0.195	0.125	0.265	2.24
Gavrysyuk et al. (2021)	0.333	0.233	0.434	2.16
Murata et al. (2021)	0.223	0.143	0.304	2.22
Gavrysyuk et al. (2021)	0.519	0.424	0.613	2.18
Fang et al. (2019)	0.182	0.000	0.410	1.71
Nath et al. (2019)	0.242	0.165	0.318	2.22
Zheng et al. (2019)	0.312	0.220	0.405	2.19
Ozyilmaz et al. (2018)	0.448	0.267	0.629	1.89
Belhomme et al. (2018)	0.294	0.141	0.447	2.00
Chen et al. (2018)	0.522	0.318	0.726	1.81
Mañá et al. (2017)	0.113	0.075	0.151	2.29
Maturu et al. (2016)	0.370	0.188	0.553	1.89
Liu et al. (2016)	0.476	0.325	0.627	2.00
Polverosi et al. (2014)	0.205	0.085	0.324	2.11
Vorselaars et al. (2014)	0.617	0.478	0.756	2.05
Sharma et al. (2003)	0.755	0.639	0.871	2.12
Kiper et al. (2001)	0.400	0.152	0.648	1.64
Sharma et al. (2001)	0.247	0.155	0.339	2.19
Baculard et al. (2001)	0.190	0.023	0.358	1.94
Muller-Quernheim et al. (1999)	0.273	0.010	0.536	1.58
Kim et al. (1999)	0.148	0.014	0.282	2.06
Baltzan et al. (1999)	0.444	0.215	0.674	1.71
Rizzato et al. (1998)	0.366	0.262	0.470	2.15
Wyser et al. (1997)	0.438	0.194	0.681	1.65
Gottlieb et al. (1997)	0.389	0.325	0.453	2.25
Lower et al. (1995)	0.875	0.773	0.977	2.16
Takada et al. (1993)	0.474	0.361	0.586	2.13
Chadelat et al. (1993)	0.250	0.000	0.550	1.44
Prior et al. (1991)	0.240	0.073	0.407	1.95
Rizzato et al. (1991)	0.100	0.007	0.193	2.19
Baumann et al. (1990)	0.667	0.428	0.905	1.67
Lower et al. (1990)	0.357	0.106	0.608	1.62
Johns et al. (1986)	0.749	0.684	0.814	2.25
Wallaert et al. (1986)	0.667	0.400	0.933	1.56
Turner-Warwick et al. (1986)	0.125	0.010	0.240	2.12
Eule et al. (1986)	0.218	0.141	0.295	2.22
Johns et al. (1983)	0.747	0.662	0.833	2.20
Rohatgi et al. (1981)	0.300	0.099	0.501	1.82
Johns et al. (1976)	0.699	0.624	0.773	2.23
Johns et al. (1974)	0.779	0.712	0.847	2.24
Emirgil et al. (1969)	0.273	0.087	0.459	1.88
Morse et al. (1961)	0.571	0.205	0.938	1.21
theta	0.398	0.339	0.456	

Test of theta = 0: z = 13.30   Prob > |z| = 0.0000
Test of homogeneity: Q = chi2(49) = 1028.61   Prob > Q = 0.0000

```
.  meta summarize, leaveoneout

   Effect-size label: Proportion
         Effect size: _meta_es
           Std. err.: _meta_se
         Study label: followed

Leave-one-out meta-analysis summary    Number of studies =       50
Random-effects model
Method: REML
```

Omitted study	Proportion	[95% conf. interval]		p-value
173	0.395	0.335	0.454	0.000
52	0.401	0.341	0.460	0.000
86	0.397	0.337	0.456	0.000
73	0.402	0.342	0.461	0.000
147	0.400	0.341	0.460	0.000
327	0.402	0.342	0.461	0.000
11	0.394	0.335	0.453	0.000
89	0.393	0.334	0.452	0.000
123	0.402	0.343	0.462	0.000
84	0.399	0.339	0.459	0.000
103	0.402	0.342	0.461	0.000
108	0.395	0.335	0.455	0.000
11	0.402	0.342	0.461	0.000
120	0.401	0.342	0.461	0.000
96	0.400	0.340	0.459	0.000
29	0.397	0.337	0.456	0.000
34	0.400	0.340	0.460	0.000
23	0.395	0.336	0.455	0.000
266	0.404	0.346	0.463	0.000
27	0.398	0.339	0.458	0.000
42	0.396	0.336	0.456	0.000
44	0.402	0.343	0.461	0.000
47	0.393	0.334	0.452	0.000
53	0.390	0.332	0.448	0.000
15	0.398	0.338	0.457	0.000
85	0.401	0.342	0.461	0.000
21	0.402	0.343	0.461	0.000
11	0.400	0.340	0.459	0.000
27	0.403	0.344	0.462	0.000
18	0.397	0.337	0.457	0.000
82	0.398	0.339	0.458	0.000
16	0.397	0.338	0.457	0.000
221	0.398	0.338	0.458	0.000
40	0.387	0.331	0.443	0.000
76	0.396	0.336	0.456	0.000
8	0.400	0.341	0.459	0.000
25	0.401	0.341	0.460	0.000
40	0.404	0.346	0.463	0.000
15	0.393	0.334	0.452	0.000
14	0.398	0.339	0.458	0.000
171	0.390	0.332	0.447	0.000
12	0.393	0.335	0.452	0.000
32	0.404	0.345	0.462	0.000
110	0.402	0.342	0.461	0.000
99	0.390	0.332	0.448	0.000
20	0.400	0.340	0.459	0.000
146	0.391	0.332	0.449	0.000
145	0.389	0.331	0.446	0.000
22	0.400	0.341	0.460	0.000
7	0.396	0.336	0.455	0.000
theta	0.398	0.339	0.456	0.000