Associations of plasma ergothioneine levels with cognitive function change in

non-demented older Chinese adults: A community-based longitudinal study

Author:

Danyang Zhao, PhD candidate[1,*], Chengwu Feng, PhD[1], Peiyu Li, PhD candidate[1],

Yunan Liu, PhD candidate[1], Long Lu, PhD[2] , Bin Jiang[3], Shifu Xiao, MD[4,#], Geng

Zong, PhD[5,#]

[#] Shifu Xiao and Geng Zong contributed equally as senior authors.

Author Affiliations:

[1]Shanghai Institute of Nutrition and Health, University of Chinese Academy of

Sciences, Chinese Academy of Sciences, Shanghai 200031, China

[2]Shanghai Institute of Organic Chemistry, University of Chinese Academy of

Sciences, Chinese Academy of Sciences, Shanghai 200032, China

[3]Xiamen Kingdomway Group Company, Xiamen, 361028, China

[4]Department of Geriatric Psychiatry, Shanghai Mental Health Center, Shanghai Jiao

Tong University School of Medicine, Shanghai 200032, China; Alzheimer's Disease

and Related Disorders Center, Shanghai Jiao Tong University, Shanghai 200032,

China

[5]Institute of Nutrition, Fudan University, Shanghai 200032, China.

Correspondence to:

Shifu Xiao, MD, Department of Geriatric Psychiatry, Shanghai Mental Health Center,

Shanghai Jiao Tong University School of Medicine, 600 South Wanping Road,

Shanghai 200032, China; Alzheimer's Disease and Related Disorders Center,

Shanghai Jiao Tong University, 600 South Wanping Road, Shanghai 200032, China.

Email, xiaoshifu@msn.com

Geng Zong, PhD, Institute of Nutrition, Fudan University, No.130 Dong-An Road,

Shanghai 200032, China.

Email, zonggeng@fudan.edu.cn

Abstract

Aim: Animal studies have reported that ergothioneine (ET) exert a protective effect on cognitive function, but evidence from humans remains limited. In this study, we explored the association between plasma ET levels and cognitive change in a prospective cohort study specifically focused on older adults without dementia in Shanghai.

Methods: This analysis was based on 1,131 community-dwelling elder Chinese (mean age 69 years) from the Shanghai Brain Aging Study. Ultra-high-performance liquid chromatography-mass spectrometry was used to measure plasma ET at baseline. Cognitive function was assessed at baseline and follow-up (mean 2.0 years) via the Montreal Cognitive Assessment (MoCA) by trained clinicians. Multivariable linear regression evaluated ET-cognition associations with confounder adjustment.

Results: Each 1-SD increase in ET was associated with a 0.23 higher point in MoCA changes (95% CI: 0.04–0.43, P=0.021) after adjustment for age, sex, smoking, alcohol, diet score, educational level, BMI, hypertension, hyperlipidemia, diabetes, cardiovascular disease, *APOE ε4* status and baseline MoCA scores. Sex-stratified analysis revealed a significant association in men (β=0.78 per SD, 95% CI: 0.35–1.21, $P < 0.001$), but not in women ($P_{\text{for interaction}} = 0.017$). Dose-response curves indicated plateauing effects above 1000 ng/mL. Domain-specific analyses identified significant protective associations for visuospatial/executive function (β=0.08, 95% CI: 0.02–

52   0.14) and delayed recall (β=0.11, 95% CI: 0.02–0.20).

53   Conclusions: Our findings indicate that higher plasma ET levels are significantly

54   associated with slower cognitive decline independent of confounders in non-demented

55   community-dwelling elderly participants, with such association observed in men but

56   not women.

Background

An estimated 57 million people worldwide currently suffer from dementia in 2019,

with projections suggesting a tripling of this number by 2050[1]. Given the central roles

of oxidative stress and neuroinflammation in dementia pathogenesis, dietary

antioxidants have emerged as promising candidates for cognitive preservation[2].

Ergothioneine (ET), a diet-derived sulfur-containing amino acid derivative and

potent thione-based antioxidant, is primarily obtained from mushrooms and fermented

foods[3]. Its absorption and tissue distribution in humans are mediated by the organic

cation transporter OCTN1 (encoded by *SLC22A4*), which facilitates pH-dependent

intestinal uptake[4] and selective accumulation in organs[5]. Notably, ET has been

detected in human parietal brain tissues[6] and medulla oblongata[7], confirming its

ability to cross the blood-brain barrier. Animal studies have shown that ET may

protect against neuronal damage induced by cisplatin[8], β-amyloid (Aβ)[9,10], and age-

related cognitive decline[11]. Emerging observational studies have found that lower

plasma levels of ET are significantly associated with higher risks of

neurodegenerative diseases, such as mild cognitive impairment (MCI)[12,13] and

dementia[14–18], however, several knowledge gaps remain. First, most of the above

studies were based on cross-sectional study design[12–17], therefore potential reverse

causation cannot be excluded. It has been suggested that plasma ET declines

concomitantly with the deterioration of cognitive function[14]. Second, since the impact

of a single dietary factor on health is mild, it is prone to be affected by the baseline

characteristics of subjects (such as sex, educational level, disease status and gene

polymorphism). However, no study has systematically evaluated potential effect

modifiers on the association between ET levels and cognitive function. Third, the

dose-response distribution between ET and cognitive function remains undetermined.

To fill these gaps, we utilized prospectively collected data of 1,131 community-

dwelling older adults from the Shanghai Brain Aging Study (SBAS) cohort, and

examined the association between baseline plasma ET levels and subsequent changes

in cognitive function. We particularly analyzed the potential effect modification of

baseline characteristics (e.g., sex, educational level, disease status and genetic factor)

on the associations and explore the dose-response associations between ET and

cognitive decline.

Methods

Study population

The Shanghai Brain Aging Study (SBAS) is an ongoing population-based prospective

cohort study established in 2016 to investigate the influencing factors related to

cognitive decline among community-dwelling adults aged ≥55 years in Shanghai,

China. Follow-up assessments have been conducted biennially since baseline, with

three waves completed as of 2024. At enrollment, a total of 2,430 eligible residents

were screened, of whom 1,627 participants: (1) completed standardized in-person

interviews, (2) provided fasting blood specimens at baseline, and (3) authorized health

insurance data linkage, thus included for follow-up. The study protocol was approved

by the Shanghai Clinical Research Ethics Committee (Approval No. 2018-10-16), and

all participants provided written informed consent.

For the current analysis, baseline and first follow-up data (mean follow-up duration: 2.0 ± 0.1 years; retention rate: 74.0%) were used to minimize sample reduction because of death and loss to follow-up. We excluded 88 participants with dementia at baseline and an additional 408 participants with missing data on the Montreal Cognitive Assessment (MoCA) at either baseline or first follow-up, resulting in a final sample of 1,131 participants (eFigure 1).

Cognitive assessments

Cognitive function was assessed using the MoCA (Beijing Version) following standardized protocols at baseline and during each follow-up visit[19,20]. All assessments were administered by trained clinicians. The MoCA domains include visuospatial/executive, naming, memory, attention, language, abstraction, delayed recall and orientation. The range of overall score was between 0 and 30 points. The primary outcome was defined as the absolute change in MoCA scores during follow-up ($\Delta MoCA = MoCA_{\text{follow-up1}} - MoCA_{\text{baseline}}$).

Plasma Ergothioneine Measurements

At each research site, fasting blood samples (10 mL) were collected in EDTA tubes from participants after a 12-hour overnight fast in the following morning. All samples were centrifuged at 4°C and stored at -80°C until analysis. Informed consent was signed by all participants prior to sample collection. An internal standard (IS) solution was prepared in methanol:acetonitrile (1:1, v/v) containing: L-glutamic acid-2,3,3,4,4-d5 (2.5 µg/mL). A stock solution (5 mg/mL) was prepared by dissolving ET standard

in 50% methanol.

A randomized sequence was employed for continuous sample analysis, with a

quality control sample added to each batch of samples (n = 10) during instrumental

analysis. The analysis was performed as described by Luo et al[21]. Samples were

separated using the Vanquish ultra-high-performance liquid chromatography system

(Agilent 1290 Infinity III) and then subjected to mass spectrometry with a Q Exactive

HF-X mass spectrometer (Thermo). Details of sample pretreatment and instrumental

parameters were provided in the **Supplementary Method.**

Chromatographic peak integration for ET and calibration curve construction

(prepared by gradient dilution of the stock solution) were performed using

TraceFinder software, with quantification achieved via IS method. The linear

calibration curve for ET was $y = 0.0001041x - 0.002337$ with an R^2 of 0.9951 and a

linear range of 50–25000 ng/mL, while the coefficients of variation for quality control

samples were 19.6%.

Ascertainment of Covariates

A structured questionnaire was administered by trained interviewers to collect

demographic characteristics including age, sex, and highest attained education.

Educational level was categorized as illiterate, primary school, middle school and

high school or above. Smoking status was dichotomized as "ever-smoker" (current or

former smoking) versus "never-smoker", while alcohol use was defined as "ever-

drinker" versus "never-drinker". Dietary quality was evaluated using a composite diet

score (range 0-4) based on four components: 1 point each for daily intake of fruits,

vegetables, or fish, and 1 point for abstaining from red meat consumption. Scores

were summed to create an ordinal diet quality metric.

Anthropometric measurements including body weight, height and blood pressure

were obtained following standardized protocols. Body mass index (BMI) was

calculated as weight (kg)/height squared (m^2). The history of diseases was ascertained

from clinical interviews and health insurance data, with conditions classified as

present or absent. Hypertension was defined as systolic blood pressure ≥140 mm Hg

and/or diastolic blood pressure ≥90 mm Hg, or history of antihypertensive medication

use. Diabetes mellitus was defined as fasting plasma glucose of 7.0 mmol/L or higher,

or hypoglycemic medication use. Hyperlipidemia was defined as meeting any of the

following criteria[22]: fasting plasma total cholesterol ≥6.2 mmol/L, LDL-C ≥4.1

mmol/L, triglycerides ≥2.3 mmol/L, HDL-C <1.0 mmol/L, or lipid-lowering

medication use. Cardiovascular disease (CVD), specifically including stroke and

coronary heart disease, was defined as clinical interviews or health insurance data.

CVD cases identified through health insurance data were defined using ICD-10 codes

I20-I25 and I60-I64. Apolipoprotein E (*APOE*) genotyping was performed as

described previously. Participants were categorized into three groups: *APOE ε4*

carriers (defined as *ε3/ε4* or *ε4/ε4*), *APOE ε4* non-carriers and *APOE ε2/ε4* carriers.[23].

Statistical analysis

Plasma ET was normalized using the SD as the scaling factor[24], when it was analyzed

as a continuous variable (ng/mL per 1 SD). Owing to gender differences in plasma ET

levels, sex-specific quartiles were derived for categorical analysis. When baseline

characters were presented, mean and SD were reported for continuous variables,

while n (percentages) was provided for categorical variables. We used one-way

ANOVA to assess baseline differences in continuous variables across sex-stratified

quartiles of ET, and χ^2 tests for categorical values.

The associations of plasma ET concentrations with MoCA score change were

explored using generalized linear models (GLM) with gaussian distribution and

identity link function. We imputed missing values using means or modes if

missingness is <5% of covariates, otherwise missing indicators were used. Plasma ET

concentration was analyzed as a continuous variable (z-scaled) and sex-stratified

quartiles. Three models of increasing complexity were designed. Model 1 was

adjusted for age, sex and follow-up years. Model 2 was further adjusted for smoking,

alcohol, diet score, educational level, BMI, *APOE ε4* status, baseline hypertension,

hyperlipidemia, diabetes and cardiovascular disease. Model 3 was additionally

adjusted for baseline MoCA scores. The same categories and adjustment models were

used to assess MoCA domain score as the outcome. The linear trend was assessed by

replacing categorical plasma ET levels with their median values in GLMs.

To assess the impact of covariates on the association between ET levels and

cognitive function, we conducted the following two analyses. We conducted the

partial correlation analysis between ET concentrations and all candidate covariates.

Second, we performed a multivariable linear regression where the ET levels were

treated as the outcome variable and regressed against all baseline phenotypes present.

We also used restricted cubic splines with three knots to explore the shape of the

dose–response relationship between baseline plasma ET and changes in MoCA score.

Additionally, the joint significance of the non-linear spline terms was evaluated

through the 'anova' function implemented in the 'rms' R package, which tests the null

hypothesis that all non-linear spline terms collectively equal zero. This approach

avoids manual model comparisons by directly evaluating the contribution of non-

linearity within the spline framework. These spline models were adjusted for the same

potential confounders as the main multivariable linear regression analyses.

We conducted stratified analyses by sex (male vs. female), baseline MoCA score

(median cutoff: <22 vs. ≥22), educational level (less than high school vs. high school

or above), *APOE ε4* status (*APOE-ε4* carriers vs. *APOE-ε4* non-carriers) and

hyperlipidemia (yes vs. no) as a priori defined and assessed potential interactions with

these variables using the likelihood ratio test comparing two nested models: a full

model containing the interaction term between the stratifying variable (e.g., sex) and

plasma ET, and a reduced model excluding the interaction term. A P-value < 0.05 for

the likelihood ratio chi-squared statistic indicated significant effect modification. All

analyses were performed using R software (version 4.2.2). A two-sided $P < 0.05$ was

considered statistically significant.

Results

Basic characteristics

Table 1 presents baseline characteristics of the 1,131 participants (65.7% women,

mean age 69 years) by sex-stratified plasma ET quartiles. The median (Q1–Q3) ET

concentrations for each quartile were 534.7 (430.6–613.0), 744.2 (693.3–819.6),

930.1 (861.1–1,007.2), and 1,200.0 (1,113.9–1,373.8) ng/mL, respectively.

Participants with higher ET levels were younger and demonstrated better educational

attainment, and higher prevalence of hyperlipidemia.

Baseline MoCA scores were higher across ET quartiles (Q1:21.1±4.6,

Q2:21.3±4.6, Q3:21.8±4.3, Q4:22.3±3.7; P = 0.002), and this pattern persisted at

follow-up (Q1:20.5±4.9, Q2:20.7±4.9, Q3:21.9±4.9, Q4:22.5±4.2; P < 0.001). Change

in total MoCA score (ΔMoCA) diverged significantly (P = 0.008), with declines

observed in Q1 (-0.6±3.7) and Q2 (-0.5±3.3), contrasting with marginal improvements

in Q3 (+0.1±3.2) and Q4 (+0.2±3.3).

Assessment of potential confounding

Partial correlation analysis showed weak associations between ET and covariates,

with all absolute values of partial correlation coefficients ≤0.15 (eFigure 2). In

multivariable linear regression with ET as the outcome, female gender was positively

associated with ET concentrations (β=0.39, 95% CI: 0.23–0.54; P < 0.001), as were

higher educational attainment (high school or above vs. illiterate: β=0.46, 95% CI:

0.07–0.84; P = 0.02) and hyperlipidemia (β=0.20, 95% CI: 0.08–0.33; P = 0.001)

(eFigure 3).

Associations of baseline plasma ET levels with change in total MoCA scores

Associations between baseline plasma ET concentrations and change in total MoCA

score are presented in Table 2. Each 1-SD increase in ET was positively associated

with a 0.23 higher point in MoCA changes (β=0.23, 95% CI: 0.04–0.43; P = 0.021)

after adjustment for age, sex, follow-up years, educational level, smoking, alcohol,

diet score, BMI, hypertension, hyperlipidemia, diabetes, cardiovascular disease,

APOE ε4 status and baseline MoCA scores. The mean changes (95% CIs) in total

MoCA score across ET quartiles were -0.58 (-0.74 to -0.43) for Q1, -0.52 (-0.68 to -

0.37) for Q2, 0.09 (-0.07 to 0.24) for Q3, and 0.18 (0.03 to 0.33; $P_{\text{for trend}}$ = 0.001) for

Q4.

When analyses were stratified by sex, the positive association between plasma ET

and MoCA score changes persisted only in men ($P_{\text{for interaction}}$ = 0.004 for continuous

ET; $P_{\text{for interaction}}$ = 0.017 for ET quartiles). Among them, each 1-SD increase in ET

was associated 0.78 higher total MoCA scores change (β=0.78, 95% CI: 0.35–1.21; P

< 0.001). The mean changes (95% CIs) in total MoCA score across ET quartiles were

-1.33 (-1.38 to -1.27) for Q1, -1.1 (-1.16 to -1.04) for Q2, 0.36 (0.31 to 0.41) for Q3

and 0.35 (0.29 to 0.40; $P_{\text{for trend}}$ < 0.001) for Q4 among men. In contrast, no

significant associations were observed in women.

In Figure 1, RCS showed no significant departure from linear associations overall

($P_{\text{non-linearity}}$ = 0.143) or within sex strata (men: $P_{\text{non-linearity}}$ = 0.577; women: $P_{\text{non-linearity}}$ = 0.758). However, visual inspection of the spline curves revealed a potential

plateauing effect at ET concentrations ≥1,000 ng/mL in the total population (Figure

1A). Baseline ET concentrations differed between men and women (eTable 1). Most

men (81.5%) had concentrations below 1,000 ng/mL (median 754.2, IQR 592.0–937.9

ng/mL), while 35.7% of women exceeded 1,000 ng/mL (median 890.1, IQR 709.7–

1,095.6 ng/mL, eFigure 4).

To facilitate clinical interpretation of effect sizes, we conducted an analysis

substituting raw plasma ET concentrations (per 200 ng/mL increment) for z-scores in

Model 3 (Figure 2). In the total cohort, each 200 ng/mL higher baseline ET was

marginally associated with a 0.13-point improvement in MoCA change (β=0.13, 95%

CI: 0.02–0.23), roughly equivalent to the association observed for 1.6 years younger

age (β=0.08 per year decrease, 95% CI: 0.06–0.11). Among men, each 200 ng/mL

higher ET was associated with a 0.43-point improvement in MoCA change (β=0.43,

95% CI: 0.19–0.66). This effect magnitude corresponds approximately to 4.3 years

younger age, based on the age association (β=0.10 per year decrease, 95% CI: 0.05–

0.15). No significant associations were observed in women.

Plasma ET and specific cognitive domains of the MoCA score

Table 3 shows the association between baseline plasma ET concentrations and

changes in each individual MoCA component. A significant positive association

between ET (per 1 SD) and change in visuospatial/executive score (β=0.08, 95% CI:

0.02–0.14; $P = 0.012$) and delayed recall score (β=0.11, 95% CI: 0.02–0.20; $P =$

0.014) were observed after adjusting for potential confounders. Supplementary

analyses characterizing the population and associations using the annualized change

of total MoCA score (ΔMoCA / follow-up years) and MoCA domain score (ΔMoCA

domain / follow-up years) are presented in eTables 2 and 3, which aligns with the

main findings.

Stratified analysis

No statistically significant interactions were detected across any subgroups (eFigure

5A-D; all $P_{\text{for interaction}} > 0.05$), yet the direction and magnitude of associations

between baseline plasma ET and MoCA score change exhibited heterogeneity in

specific strata.

Discussion

In this prospective cohort study of 1,131 community-dwelling older adults (mean age

69 years), higher baseline plasma ET levels were significantly associated with slower

cognitive decline, as assessed by MoCA scores, during a 2-year follow-up period.

More specifically, when the plasma concentration of ET exceeds 1,000 ng/mL, the

decline in cognitive function significantly slows down. However, this association has

only been observed in men. Domain-specific analysis found that the observed ET-

MoCA association was mainly driven by the temporary slowdown in the decline of

visuospatial/executive and delayed recall.

While cross-sectional studies have consistently shown that higher ET levels were

significantly associated with higher cognitive function[12,13] and lower risk of

dementia[14–17], longitudinal evidence remains scarce and methodologically limited. A

previous French study on the basis of a nested case-control study design (N=842)

screened the associations of over 10 food metabolites with cognitive function and

found that higher ET level was significantly associated with slower cognitive decline

[25]. Similarly, a study from Singapore[18] reported that lower ET level was associated

with accelerated cognitive decline among 470 elderly participants, yet the sample of

this study could not represent the general community-dwelling elderly, as 40% of the

participants had been diagnosed with dementia at baseline. From the perspective of

dementia prevention, exploring the association between plasma ET and cognitive

function in a community-dwelling elderly population is of greater public health

significance, and our study has filled this gap. Notably, determining the temporal

sequence between the decline in ET levels and cognitive decline is crucial, as it is

possible that cognitive impairment directly affects dietary intake, leading to reduced

consumption of ET-rich foods and a decline in plasma ET levels. Previous studies

have struggled to address this question as they did not exclude participants with

dementia at baseline[18]. In contrast, our study included only participants with normal

cognitive function, and the results remained robust even after excluding those with

baseline cognitive function at the lower end of the normal range. Collectively, our

findings support that low ET intake occurs prior to cognitive decline.

The observed associations between ET levels and specific cognitive domains,

namely delayed recall and visuospatial/executive function, warrant consideration

regarding ET's potential neuroprotective roles. Impaired delayed recall represents one

of the earliest and most sensitive cognitive markers of dementia progression,

predictive of conversion from MCI to dementia[26,27]. The preferential preservation of

this function by ET suggests targeted neuroprotective effects within the hippocampus,

the brain region orchestrating declarative memory consolidation and retrieval.

Experimental evidence shows that ET protects hippocampal neurons from oxidative

stress[28], reduces Aβ accumulation and lipid peroxidation[29] in the hippocampus, and

ET deficiency disrupts hippocampal neurogenesis[30] while supplementation promotes

dendritic spine formation[31]. Human neuroimaging studies corroborate this

hippocampal vulnerability, showing reduced hippocampal volume in MCI and

dementia patients with systemic ET deficiency[14,18]. Specifically,

visuospatial/executive function critically depends on the prefrontal cortex[32] and

fronto-parietal network modulation. The susceptibility of these regions to oxidative

stress and metabolic dysregulation suggests they might be sensitive to ET's

antioxidant properties, though the precise mechanisms require further exploration.

The men-specific cognitive changes associated with ET likely reflect a dose-

dependent association modulated by baseline sex differences. Women exhibited

substantially higher median plasma ET concentrations (890.1 ng/mL; IQR: 709.7–

1,095.6) than men (754.2 ng/mL; IQR: 592.0–937.9), with 35.7% of women

exceeding the 1,000 ng/mL saturation threshold where dose-response relationships

plateaued in our analysis. Conversely, most men (81.5%) operated below this

threshold, possibly enabling detectable dose-dependent neuroprotection. This aligns

with broader nutrient-sex interactions wherein antioxidants (e.g., vitamin C[33], fish-

oil[34]) show men-predominant associations for cognitive function, possibly due to

women's higher baseline antioxidant reserves. Human microbiome studies

demonstrated divergent bacterial communities between sexes[35], the sexual

dimorphism in microbiome could theoretically modulate ET bioavailability through

bacterial absorption[36–38] or degradation[39,40], although direct evidence is lacking.

Ultimately, the dose effect observed for ET provides the most plausible explanation

for this sex divergence, aligning with plateau effects observed for other

neuroprotective agents and consistent with our cohort's ET distribution. However, the

potential contribution of alternative mechanisms requires further investigation.

Our study design and analytical approaches robustly mitigate concerns regarding

reverse causation. First, the longitudinal analysis of baseline plasma ET levels against

subsequent changes in cognitive function inherently establishes temporal precedence.

Second, the positive association between ET and MoCA score changes strengthened

rather than attenuated after comprehensive adjustment for baseline MoCA scores in

multivariable Model 3, contradicting the expectation that confounding by pre-existing

cognitive status would diminish the effect. Third, weak correlations ($|r| \leq 0.15$)

between baseline ET levels and key covariates—including demographics, lifestyle

factors, and comorbidities—indicate minimal confounding potential, a conclusion

further reinforced by our inclusion of these variables in all primary models.

There are also limitations to our study. First, most variables, including

demographic data, health conditions and diet were self-reported, which could be

inherently challenging in elderly populations due to recall bias and cognitive

limitations. Second, the median 2-year follow-up period, though sufficient to detect

significant cognitive changes in this cohort, precluded the observation of long-term

associations. Extended surveillance would clarify ET's durability in dementia

prevention. Third, reliance ET was only measured at baseline therefore cannot capture

potential intraindividual fluctuations over time, although the observed plateau effect

above 1,000 ng/mL suggests stable biological activity beyond this threshold. Fourth,

the difference of standardized experimental and analytical procedures in

metabolomics (e.g., instrumentation, quantification methods, sample preparation)

means that the ET concentrations quantified in this study cannot be directly compared

with quantitative results from other metabolomic investigations. Finally, as

participants were exclusively community-dwelling Chinese older adults,

generalizability to younger populations, other ethnicities, or clinical cohorts requires

verification. Notwithstanding these limitations, our use of a quantitative biomarker,

prospective design, comprehensive confounder adjustment, and stratification analyses

provides robust evidence supporting a sex-specific protective role of ET against

cognitive decline in older adults.

In conclusion, our findings indicate that higher plasma ET levels are significantly

associated with slower cognitive decline independent of confounders in general

community elderly participants without dementia at baseline, whereas such

association was only observed in men but not in women. Further analysis with large

sample size and longer follow-up period is needed to verify our findings.

Article information

Author Contributions: Xiao and Zong had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Zhao, Xiao and Zong.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Zhao, Zong.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Zhao, Feng, Li and Liu.

Administrative, technical, or material support: Lu, Jiang and Xiao.

Supervision: Xiao and Zong.

Conflict of Interest Disclosures: Jiang serves as the Chairman of Board and Chief Executive Officer of Xiamen Kingdomway Group, a company engaged in the research, development, and production of pharmaceutical ingredients and dietary supplements. The other authors declare no competing interests.

Additional Contributions: This study was conducted using the Shanghai Brain Aging Study. All authors are thankful to the participants agreed to contribute to study. We also gratefully acknowledge Xiamen Kingdomway Group Company for their in-kind contribution of the ergothioneine reference standard used in this study.

References

1. GBD 2019 Dementia Forecasting Collaborators. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. *Lancet Public Health* **7**, e105–e125 (2022).

2. Andersen, J. K. Oxidative stress in neurodegeneration: cause or consequence? *Nat. Med.* **10**, S18–S25 (2004).

3. Halliwell, B. & Cheah, I. Are age-related neurodegenerative diseases caused by a lack of the diet-derived compound ergothioneine? *Free Radic. Biol. Med.* **217**, 60–67 (2024).

4. Harwood, M. D., Zhang, M., Pathak, S. M. & Neuhoff, S. The Regional-Specific Relative and Absolute Expression of Gut Transporters in Adult Caucasians: A Meta-Analysis. *Drug Metab. Dispos. Biol. Fate Chem.* **47**, 854–864 (2019).

5. Gründemann, D., Hartmann, L. & Flögel, S. The ergothioneine transporter (ETT): substrates and locations, an inventory. *FEBS Lett.* **596**, 1252–1269 (2022).

6. Novotny, B. C. *et al.* Profiling the metabolic landscape of AD. *Alzheimers Dement.* **17**, e050086 (2021).

7. Graham, S. F., Chevallier, O. P., Kumar, P., Türko Gcaron Lu, O. & Bahado-Singh, R. O. Metabolomic profiling of brain from infants who died from Sudden Infant Death Syndrome reveals novel predictive biomarkers. *J. Perinatol. Off. J. Calif. Perinat. Assoc.* **37**, 91–97 (2017).

8. Song, T.-Y., Chen, C.-L., Liao, J.-W., Ou, H.-C. & Tsai, M.-S. Ergothioneine protects against neuronal injury induced by cisplatin both in vitro and in vivo. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* **48**, 3492–3499 (2010).

9. Yang, N.-C. *et al.* Ergothioneine protects against neuronal injury induced by β-amyloid in mice. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* **50**, 3902–3911 (2012).

10. Cheah, I. K. *et al.* Inhibition of amyloid-induced toxicity by ergothioneine in a transgenic Caenorhabditis elegans model. *FEBS Lett.* **593**, 2139–2150 (2019).

11. Song, T.-Y. *et al.* Ergothioneine and melatonin attenuate oxidative stress and protect against learning and memory deficits in C57BL/6J mice treated with D-galactose. *Free Radic. Res.* **48**, 1049–1060 (2014).

12. Cheah, I. K., Feng, L., Tang, R. M. Y., Lim, K. H. C. & Halliwell, B. Ergothioneine levels in an elderly population decrease with age and incidence of cognitive decline; a risk factor for neurodegeneration? *Biochem. Biophys. Res. Commun.* **478**, 162–167 (2016).

13. Kondoh, H., Teruya, T., Kameda, M. & Yanagida, M. Decline of ergothioneine in frailty and cognition impairment. *FEBS Lett.* **596**, 1270–1278 (2022).

14. Wu, L.-Y. *et al.* Low plasma ergothioneine levels are associated with neurodegeneration and cerebrovascular disease in dementia. *Free Radic. Biol. Med.* **177**, 201–211 (2021).

15. Teruya, T., Chen, Y.-J., Kondoh, H., Fukuji, Y. & Yanagida, M. Whole-blood metabolomics of dementia patients reveal classes of disease-linked metabolites. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2022857118 (2021).

16. Novotny, B. C. *et al.* Profiling the metabolic landscape of AD. *Alzheimers Dement.* **17**, (2021).

17. Oka, T. *et al.* Multiomics analysis to explore blood metabolite biomarkers in an Alzheimer's Disease Neuroimaging Initiative cohort. *Sci. Rep.* **14**, 6797 (2024).

18. Wu, L.-Y. *et al.* Low Plasma Ergothioneine Predicts Cognitive and Functional Decline in an Elderly Cohort Attending Memory Clinics. *Antioxid. Basel Switz.* **11**, 1717 (2022).

19. Wen, H.-B. *et al.* The application of Montreal cognitive assessment in urban Chinese residents

of Beijing. *Zhonghua Nei Ke Za Zhi* **47**, 36–39 (2008).(in Chinese)

20. Nasreddine, Z. S. *et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J. Am. Geriatr. Soc.* **53**, 695–699 (2005).

21. Luo, D., Deng, T., Yuan, W., Deng, H. & Jin, M. Plasma metabolomic study in Chinese patients with wet age-related macular degeneration. *BMC Ophthalmol.* **17**, 165 (2017).

22. Joint Committee on the Chinese Guidelines for Lipid Management. Chinese guideline for lipid management (primary care version 2024). *Zhonghua Xin Xue Guan Bing Za Zhi* **52**, 330–337 (2024). (in Chinese)

23. Frick, E. A. *et al.* Serum proteomics reveal APOE-ε4-dependent and APOE-ε4-independent protein signatures in Alzheimer's disease. *Nat. Aging* **4**, 1446–1464 (2024).

24. van den Berg, R. A., Hoefsloot, H. C. J., Westerhuis, J. A., Smilde, A. K. & van der Werf, M. J. Centering, scaling, and transformations: improving the biological information content of metabolomics data. *BMC Genomics* **7**, 142 (2006).

25. González-Domínguez, R. *et al.* Food and Microbiota Metabolites Associate with Cognitive Decline in Older Subjects: A 12-Year Prospective Study. *Mol. Nutr. Food Res.* **65**, 2100606 (2021).

26. Caselli, R. J. *et al.* Neuropsychological decline up to 20 years before incident mild cognitive impairment. *Alzheimers Dement. J. Alzheimers Assoc.* **16**, 512–523 (2020).

27. Heister, D. *et al.* Predicting MCI outcome with clinically available MRI and CSF biomarkers. *Neurology* **77**, 1619–1628 (2011).

28. Kushairi, N., Phan, C. W., Sabaratnam, V., Naidu, M. & David, P. Dietary amino acid ergothioneine protects HT22 hippocampal neurons against H2O2-induced neurotoxicity via antioxidative mechanism. *PharmaNutrition* **13**, 100214 (2020).

29. Yang, N.-C. *et al.* Ergothioneine protects against neuronal injury induced by β-amyloid in mice. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* **50**, 3902–3911 (2012).

30. Ishimoto, T. *et al.* TrkB phosphorylation in serum extracellular vesicles correlates with cognitive function enhanced by ergothioneine in humans. *NPJ Sci. Food* **8**, 11 (2024).

31. Nakamichi, N. *et al.* Oral Administration of the Food-Derived Hydrophilic Antioxidant Ergothioneine Enhances Object Recognition Memory in Mice. *Curr. Mol. Pharmacol.* **14**, 220–233 (2021).

32. Friedman, N. P. & Robbins, T. W. The role of prefrontal cortex in cognitive control and executive function. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* **47**, 72–89 (2022).

33. Lee, L. *et al.* Relationships between dietary intake and cognitive function level in Korean elderly people. *Public Health* **115**, 133–138 (2001).

34. Kuszewski, J. C., Howe, P. R. C. & Wong, R. H. X. Evaluation of Cognitive Performance following Fish-Oil and Curcumin Supplementation in Middle-Aged and Older Adults with Overweight or Obesity. *J. Nutr.* **150**, 3190–3199 (2020).

35. Ma, Z. S. & Li, W. How and Why Men and Women Differ in Their Microbiomes: Medical Ecology and Network Analyses of the Microgenderome. *Adv. Sci. Weinh. Baden-Wurtt. Ger.* **6**, 1902054 (2019).

36. Zhang, Y. *et al.* Discovery and structure of a widespread bacterial ABC transporter specific for ergothioneine. *Nat. Commun.* **13**, 7586 (2022).

37. Dumitrescu, D. G. *et al.* A microbial transporter of the dietary antioxidant ergothioneine. *Cell* **185**, 4526-4540.e18 (2022).

488 38. Cheah, I. K., Lee, J. Z., Tang, R. M. Y., Koh, P. W. & Halliwell, B. Does Lactobacillus reuteri
489 influence ergothioneine levels in the human body? *FEBS Lett.* **596**, 1241–1251 (2022).
490 39. Wolff, J. B. Ergothionase from Escherichia coli. *J. Biol. Chem.* **237**, 874–881 (1962).
491 40. Maurer, A., Leisinger, F., Lim, D. & Seebeck, F. P. Structure and Mechanism of Ergothionase
492 from Treponema denticola. *Chem. Weinh. Bergstr. Ger.* **25**, 10298–10303 (2019).
493

Tables

Table 1. Baseline Characteristics of Participants Stratified by Sex-Specific Quartiles of Plasma ET Concentrations

	Overall (n = 1131)	Q1 (n = 271)	Q2 (n = 280)	Q3 (n = 288)	Q4 (n = 292)	P-value
Plasma ET, ng/mL*	842.2 (657.8-1052.5)	534.7 (430.6-613.0)	744.2 (693.3-819.6)	930.1 (861.1-1007.2)	1200.0 (1113.9-1373.8)	<0.001
Age, years, mean (SD)	69.0 (7.5)	69.7 (7.6)	69.7 (7.8)	68.3 (7.2)	68.5 (7.2)	0.028
Women, n (%)	743 (65.7)	179 (66.1)	187 (66.8)	188 (65.3)	189 (64.7)	0.959
Educational level, n (%)						0.001
Illiterate	30 (2.7)	15 (5.5)	8 (2.9)	6 (2.1)	1 (0.3)	
Primary school	92 (8.1)	29 (10.7)	25 (8.9)	17 (5.9)	21 (7.2)	
Middle school	376 (33.2)	90 (33.2)	99 (35.4)	106 (36.8)	81 (27.7)	
High school or above	633 (56.0)	137 (50.6)	148 (52.9)	159 (55.2)	189 (64.7)	
Ever smokers, n (%)	211 (18.7)	52 (19.2)	53 (18.9)	58 (20.1)	48 (16.4)	0.699
Ever drinkers, n (%)	160 (14.1)	30 (11.1)	43 (15.4)	55 (19.1)	32 (11.0)	0.013
Diet score, mean (SD)	3.0 (0.6)	3.0 (0.7)	3.0 (0.6)	3.0 (0.5)	3.0 (0.6)	0.671
BMI, kg/m^2, mean (SD)	24.0 (3.3)	23.8 (3.4)	24.2 (3.5)	24.3 (3.3)	23.9 (3.0)	0.186
Hypertension, n (%)	905 (80.0)	225 (83.0)	227 (81.1)	225 (78.1)	228 (78.1)	0.383
Hyperlipidemia, n (%)	787 (69.6)	172 (63.5)	193 (68.9)	206 (71.5)	216 (74.0)	0.046
Diabetes, n (%)	420 (37.1)	98 (36.2)	100 (35.7)	119 (41.3)	103 (35.3)	0.400
Cardiovascular disease, n (%)	740 (65.4)	185 (68.3)	177 (63.2)	186 (64.6)	192 (65.8)	0.642
APOE genotype[†], n (%)						0.325
ε4 carriers	200 (17.7)	56 (20.7)	40 (14.3)	54 (18.8)	50 (17.1)	
ε4 non-carriers	907 (80.2)	208 (76.8)	237 (84.6)	226 (78.5)	236 (80.8)	
ε2/ε4 carriers	24 (2.1)	7 (2.6)	3 (1.1)	8 (2.8)	6 (2.1)	
Baseline MoCA score, mean (SD)	21.6 (4.3)	21.1 (4.6)	21.3 (4.6)	21.8 (4.3)	22.3 (3.7)	0.002
First follow-up MoCA score, mean (SD)	21.4 (4.8)	20.5 (4.9)	20.7 (4.9)	21.9 (4.9)	22.5 (4.2)	<0.001
Δ MoCA score, mean (SD)	-0.2 (3.4)	-0.6 (3.7)	-0.5 (3.3)	0.1 (3.2)	0.2 (3.3)	0.008

Values are percentages for categorical variables and means ± SD for continuous variables. One-way ANOVA factor was used for continuous variables, and a χ2 test was used for categorical variables. $P < 0.05$ was considered significant.

*Plasma ET concentrations are presented with the interquartile range (Q1–Q3).

†APOE ε4 carriers were defined as carriers of ε3/ε4 or ε4/ε4 genotypes.

BMI, Body Mass Index; APOE, Apolipoprotein E; MoCA, Montreal Cognitive Assessment.

Table 2. Associations between Baseline Plasma ET Concentrations and Change in Total MoCA Score in the Total Population and Stratifying by Sex

	ET (ng/mL) per 1-SD increment		Q1	Q2	Q3	Q4	P for trend
	β (95% CI)	P-value	Δ MoCA score, mean (95% CI)	Δ MoCA score, mean (95% CI)	Δ MoCA score, mean (95% CI)	Δ MoCA score, mean (95% CI)	
Total							
Model 1	0.19 (-0.02, 0.40)	0.070	-0.58 (-0.63, -0.54)	-0.52 (-0.57, -0.48)	0.09 (0.04, 0.13)	0.18 (0.14, 0.23)	0.004
Model 2	0.17 (-0.04, 0.38)	0.114	-0.58 (-0.65, -0.51)	-0.52 (-0.59, -0.46)	0.09 (0.02, 0.15)	0.18 (0.12, 0.24)	0.006
Model 3	0.23 (0.04, 0.43)	0.021	-0.58 (-0.74, -0.43)	-0.52 (-0.68, -0.37)	0.09 (-0.07, 0.24)	0.18 (0.03, 0.33)	0.001
Men							
Model 1	0.75 (0.31, 1.19)	0.001	-1.33 (-1.63, -1.03)	-1.1 (-1.36, -0.84)	0.36 (0.08, 0.64)	0.35 (0.08, 0.61)	<0.001
Model 2	0.79 (0.33, 1.24)	<0.001	-1.33 (-1.52, -1.13)	-1.1 (-1.23, -0.96)	0.36 (0.22, 0.50)	0.35 (0.21, 0.49)	<0.001
Model 3	0.78 (0.35, 1.21)	<0.001	-1.33 (-1.38, -1.27)	-1.1 (-1.16, -1.04)	0.36 (0.31, 0.41)	0.35 (0.29, 0.40)	<0.001
Women							
Model 1	0.02 (-0.21, 0.25)	0.842	-0.20 (-0.40, 0)	-0.24 (-0.43, -0.05)	-0.06 (-0.25, 0.13)	0.09 (-0.10, 0.28)	0.468
Model 2	0.02 (-0.26, 0.21)	0.850	-0.20 (-0.30, -0.10)	-0.24 (-0.34, -0.14)	-0.06 (-0.15, 0.03)	0.09 (0.02, 0.16)	0.732
Model 3	0.06 (-0.16, 0.27)	0.618	-0.20 (-0.26, -0.14)	-0.24 (-0.30, -0.18)	-0.06 (-0.12, 0)	0.09 (0.04, 0.14)	0.346

Linear regression models were used: Model 1 was adjusted for age, sex and follow-up years. Model 2 was further adjusted for smoking, alcohol, diet score, educational level, BMI, APOE ε4 status, hypertension, hyperlipidemia, diabetes and cardiovascular disease. Model 3 was additionally adjusted for baseline MoCA scores.

The interaction between sex and continuous plasma ET was significant (LRT: $\chi^2 (1) = 8.39$, $P = 0.004$).

The interaction between sex and the four-category ET groups was also significant (LRT: $\chi^2 (4) = 10.24$, $P = 0.017$).

P for trend was tested in all models by treating ET categories as a continuous variable assigned the median value of each category.

Q1, quartile 1; Q2, quartile 2; Q3, quartile 3; Q4, quartile 4.

510 Table 3. Associations between Baseline Plasma ET Concentrations and Changes in MoCA Domain Scores in the Total Population

	ET (ng/mL) per 1-SD increment		Q1	Q2	Q3	Q4	P for trend
	β (95% CI)	P-value	Δ MoCA score, mean (95% CI)	Δ MoCA score, mean (95% CI)	Δ MoCA score, mean (95% CI)	Δ MoCA score, mean (95% CI)	
Visuospatial/executive							
Model 1	0.06 (-0.01, 0.13)	0.085	-0.17 (-0.18, -0.16)	-0.21 (-0.22, -0.20)	-0.13 (-0.14, -0.12)	0.01 (0, 0.02)	0.049
Model 2	0.06 (-0.01, 0.13)	0.115	-0.17 (-0.19, -0.16)	-0.21 (-0.23, -0.19)	-0.13 (-0.15, -0.11)	0.01 (-0.01, 0.03)	0.061
Model 3	0.08 (0.02, 0.14)	0.012	-0.17 (-0.24, -0.10)	-0.21 (-0.28, -0.15)	-0.13 (-0.20, -0.06)	0.01 (-0.06, 0.08)	0.016
Naming							
Model 1	0.01 (-0.03, 0.05)	0.565	0.07 (0.07, 0.08)	-0.02 (-0.02, -0.02)	0.03 (0.03, 0.03)	0.08 (0.08, 0.08)	0.622
Model 2	0.01 (-0.04, 0.05)	0.760	0.07 (0.06, 0.08)	-0.02 (-0.03, -0.01)	0.03 (0.02, 0.04)	0.08 (0.07, 0.09)	0.828
Model 3	0.02 (-0.02, 0.06)	0.303	0.07 (0.03, 0.12)	-0.02 (-0.06, 0.01)	0.03 (-0.01, 0.06)	0.08 (0.04, 0.12)	0.418
Attention							
Model 1	-0.02 (-0.09, 0.05)	0.621	-0.15 (-0.15, -0.15)	-0.17 (-0.17, -0.17)	-0.02 (-0.03, -0.02)	-0.08 (-0.08, -0.07)	0.277
Model 2	-0.03 (-0.10, 0.04)	0.467	-0.15 (-0.16, -0.13)	-0.17 (-0.18, -0.16)	-0.02 (-0.04, -0.01)	-0.08 (-0.09, -0.07)	0.388
Model 3	0 (-0.05, 0.06)	0.890	-0.15 (-0.22, -0.07)	-0.17 (-0.25, -0.09)	-0.02 (-0.09, 0.04)	-0.08 (-0.14, -0.01)	0.081
Language							
Model 1	0.03 (-0.03, 0.08)	0.310	-0.03 (-0.04, -0.03)	-0.08 (-0.08, -0.07)	-0.01 (-0.01, 0)	0.02 (0.02, 0.03)	0.342
Model 2	0.02 (-0.03, 0.08)	0.371	-0.03 (-0.05, -0.01)	-0.08 (-0.1, -0.06)	-0.01 (-0.02, 0.01)	0.02 (0.01, 0.04)	0.331
Model 3	0.04 (-0.01, 0.08)	0.120	-0.03 (-0.09, 0.02)	-0.08 (-0.13, -0.02)	-0.01 (-0.06, 0.05)	0.02 (-0.03, 0.08)	0.179
Abstraction							
Model 1	-0.02 (-0.08, 0.03)	0.375	-0.20 (-0.21, -0.19)	-0.03 (-0.04, -0.02)	-0.08 (-0.09, -0.07)	-0.09 (-0.10, -0.08)	0.358
Model 2	-0.03 (-0.08, 0.03)	0.356	-0.20 (-0.21, -0.19)	-0.03 (-0.04, -0.02)	-0.08 (-0.09, -0.07)	-0.09 (-0.01, -0.08)	0.364
Model 3	0.01 (-0.04, 0.05)	0.775	-0.20 (-0.26, -0.14)	-0.03 (-0.09, 0.03)	-0.08 (-0.14, -0.02)	-0.09 (-0.15, -0.03)	0.085
Delayed recall							
Model 1	0.10 (-0.01, 0.21)	0.064	0.06 (0.05, 0.08)	0.24 (0.22, 0.25)	0.43 (0.42, 0.44)	0.38 (0.36, 0.39)	0.032
Model 2	0.11 (0, 0.21)	0.056	0.06 (0.03, 0.09)	0.24 (0.21, 0.26)	0.43 (0.40, 0.46)	0.38 (0.35, 0.40)	0.029
Model 3	0.11 (0.02, 0.20)	0.014	0.06 (-0.05, 0.18)	0.24 (0.12, 0.35)	0.43 (0.31, 0.55)	0.38 (0.26, 0.49)	0.004
Orientation							
Model 1	0.03 (-0.02, 0.08)	0.250	-0.17 (-0.17, -0.16)	-0.25 (-0.25, -0.24)	-0.13 (-0.14, -0.13)	-0.14 (-0.15, -0.13)	0.479
Model 2	0.02 (-0.02, 0.07)	0.331	-0.17 (-0.18, -0.15)	-0.25 (-0.26, -0.24)	-0.13 (-0.14, -0.12)	-0.14 (-0.15, -0.13)	0.601
Model 3	0.02 (-0.02, 0.06)	0.396	-0.17 (-0.22, -0.11)	-0.25 (-0.29, -0.2)	-0.13 (-0.18, -0.09)	-0.14 (-0.18, -0.10)	0.412

511 Linear regression models were used. Adjustment for covariates in Models 1 and 2 was identical to Table 2. Model 3 was additionally adjusted for baseline MoCA domain scores.

512 P for trend was tested in all models by treating ET categories as a continuous variable assigned the median value of each category. Q, quartile.

Figures

Figure 1. Multivariate-adjusted Relation of Baseline Plasma ET with MoCA Score Changes for the Total Population (A) and Stratifying by Sex (B). Associations were evaluated with the use of restricted cubic splines. The solid lines represent the central β estimate and the shaded area represent the 95% confidence intervals (CIs).

519   A   B   C

521   Figure 2. Forest Plots of Multivariable Associations Between Covariates and MoCA Score Change (A) Total population, (B) Men, (C) Women. Age analyzed per 1 year decrement.

522   ET concentrations analyzed per 200 ng/mL increment.