Myocardial Native T1 Mapping in the German National Cohort (NAKO): Associations with Age, Sex, and Cardiometabolic Risk Factors

Clemens Ammann[1,2,3,4], Jan Gröschel[1,2,3,5], Hadil Saad[1,2,4], Susanne Rospleszcz[6], Christopher Schuppert[6], Thomas Hadler[1,2,3], Richard Hickstein[1,2,3], Thoralf Niendorf[7], Janis M. Nolde[8], Matthias B. Schulze[9,10], Karin H. Greiser[11], Josua A. Decker[12], Thomas Kröncke[12], Thomas Küstner[13], Konstantin Nikolaou[14], Stefan N. Willich[15], Thomas Keil[15,16,17], Marcus Dörr[18,19], Robin Bülow[19,20], Fabian Bamberg[6], Tobias Pischon[1,21,22], Christopher L. Schlett[6], Jeanette Schulz-Menger[1,2,3,4,*]

Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany
Working Group on CMR, Experimental and Clinical Research Center, a cooperation between Charité – Universitätsmedizin Berlin and the Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany
DZHK (German Centre for Cardiovascular Research), partner site Berlin, Berlin, Germany
Department of Cardiology and Nephrology, HELIOS Hospital Berlin-Buch, Berlin, Germany
Deutsches Herzzentrum der Charité – Medical Heart Center of Charité and German Heart Institute Berlin, Department of Cardiology, Angiology and Intensive Care, Berlin, Germany
Department of Diagnostic and Interventional Radiology, University of Freiburg, Freiburg, Germany
Berlin Ultrahigh Field Facility, Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany
Department of Nephrology, University Medical Centre Freiburg, Freiburg, Germany
Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany
Institute of Nutritional Science, University of Potsdam, Nuthetal, Germany
Division of Cancer Epidemiology, German Cancer Research Center, Heidelberg, Germany
Department of Diagnostic and Interventional Radiology, University Hospital Augsburg, Augsburg, Germany
Medical Image and Data Analysis (MIDAS.Lab), University Hospital of Tübingen, Tübingen, Germany
Department of Diagnostic and Interventional Radiology, University Hospital Tübingen, Tübingen, Germany
Institute of Social Medicine, Epidemiology and Health Economics, Charité – Universitätsmedizin Berlin, Berlin, Germany
Institute of Clinical Epidemiology and Biometry, University of Würzburg, Würzburg, Germany
State Institute of Health I, Bavarian Health and Food Safety Authority, Erlangen, Germany
Institute of Community Medicine (SHIP/KEF), University Medicine Greifswald, Greifswald, Germany
German Centre for Cardiovascular Research (DZHK), partner Site Greifswald, Greifswald, Germany
Institute of Diagnostic Radiology and Neuroradiology, University Medicine Greifswald, Greifswald, Germany
Molecular Epidemiology Research Group, Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany
Biobank Technology Platform, Max Delbrück Center for Molecular Medicine in the Helmholtz Association, Berlin, Germany
* Corresponding author, jeanette.schulz-menger@charite.de

Abstract

Background and Aims: In cardiovascular magnetic resonance (CMR), myocardial native T1 mapping enables quantitative, non-invasive tissue characterization and is sensitive to subclinical changes in myocardial structure and composition. We investigated how age, sex, and cardiometabolic risk factors are associated with myocardial T1 in a population-based analysis within the German National Cohort (NAKO).

Methods: This cross-sectional study included 29,573 prospectively enrolled participants who underwent CMR-based midventricular T1 mapping at 3.0 T, alongside clinical phenotyping. After artificial intelligence-assisted myocardial segmentation, a subset of 9,162 outliers was subjected to manual quality control according to clinical evaluation standards. Associations with cardiometabolic risk factors, identified through self-reported medical history, clinical chemistry, and blood pressure measurements, were evaluated using adjusted linear regression models.

Results: Women had higher T1 values than men, with sex differences progressively declining with age. T1 was significantly elevated in individuals with diabetes (β=3.91 ms; p<0.001), kidney disease (β=3.44 ms; p<0.001), and current smoking (β=6.67 ms; p<0.001). Conversely, hyperlipidaemia was significantly associated with lower T1 (β=−4.41 ms; p<0.001). Associations with hypertension showed a sex-specific pattern: T1 was lower in women but increased with hypertension severity in men.

Conclusions: Myocardial native T1 varies by sex and age and shows associations with major cardiometabolic risk factors. Notably, lower T1 times in participants with hyperlipidaemia may indicate a direct effect of blood lipids on the heart. Our findings support the utility of T1 mapping as a sensitive marker of early myocardial changes and highlight the sex-specific interplay between cardiometabolic health and myocardial tissue composition.

Keywords

German National Cohort (NAKO); T1 mapping; cardiovascular magnetic resonance; myocardium; cardiometabolic; cardiovascular disease

Graphical Abstract

Key Question

How are age, sex, and cardiometabolic risk factors associated with myocardial native T1, a quantitative magnetic resonance imaging marker of myocardial tissue composition, in a large-scale population-based evaluation within the German National Cohort (NAKO)?

Key Finding

T1 relaxation times were higher in women and gradually converged between sexes with age. Diabetes, kidney disease, smoking, and hypertension in men were associated with prolonged T1 times. Unexpectedly, hyperlipidaemia and hypertension in women showed a negative association with T1.

Take-Home Message

Native T1 mapping is sensitive to subclinical myocardial changes and reflects a close interplay between metabolic and myocardial health. It reveals marked age-dependent sex differences and sex-specific responses in myocardial tissue composition to cardiometabolic risk factors.

Abbreviations

AHA	American Heart Association
AI	artificial intelligence
BMI	body mass index
BP	blood pressure
CAD	coronary artery disease
CI	confidence interval
CMR	cardiovascular magnetic resonance
CVD	cardiovascular disease
ESC	European Society of Cardiology
HR	heart rate
HT	hypertension
LDL-c	LDL-cholesterol
MAP	mean arterial pressure
MOLLI	modified look-locker inversion recovery
MR	magnetic resonance
NAKO	German National Cohort
SD	standard deviation
TD	thyroid disease

Introduction

Myocardial native T1 mapping by cardiovascular magnetic resonance (CMR) is a quantitative, non-invasive imaging marker of myocardial tissue composition (1,2). The technique is sensitive to diffuse processes including inflammation, oedema, interstitial fibrosis or myocardial infiltration with iron or lipid deposits and has proven helpful in diagnosing and monitoring cardiovascular disease (3). Importantly, T1 mapping enables the detection of subclinical changes in myocardial tissue composition, potentially capturing early stages of cardiac pathology (4).

The German National Cohort (NAKO) is a population-based prospective study comprising over 100,000 women and 100,000 men aged 19 to 75 years with extensive phenotyping, aimed at investigating the causes of chronic diseases (5). Its whole-body magnetic resonance (MR) imaging sub-cohort comprises 30,868 individuals who underwent standardized 3.0 T MR imaging closely aligned in time with clinical and laboratory assessments (6).

Major cardiometabolic risk factors such as diabetes, hypertension, hyperlipidaemia, chronic kidney disease, and smoking are believed to promote adverse structural myocardial remodelling through mechanisms including low-grade inflammation and fibrosis (7–11). Myocardial native T1 may therefore serve as a sensitive, albeit non-specific, imaging marker of target-organ damage. In the UK Biobank, native T1 mapping at 1.5 T was found to be associated with prevalent cardiovascular disease and incident events in participants aged 44 to 84 based on hospital registry data (12). However, evidence gaps persist. Specifically, there are still limited data from younger individuals and large-scale population-based studies relating the aforementioned risk factors to myocardial T1. Moreover, while previous studies have reported a sex difference of myocardial native T1, the role of age remains inconsistent and insufficiently understood (12–14).

In this cross-sectional analysis of the NAKO MR cohort, we investigated how cardiometabolic risk factors, age, and sex are associated with myocardial native T1 as a marker of early changes in myocardial tissue composition.

Methods

Study Population
This study leverages data from the NAKO at baseline and included 29,573 participants (44.2% women) with a midventricular T1 map obtained by CMR and corresponding data from self-questionnaires and clinical chemistry. A modified look-locker inversion recovery (MOLLI) 5(3)3 T1 mapping technique was used to acquire a midventricular short-axis T1 pixel map.

Participants were scanned across five imaging centres, ensuring uniform protocols and equipment (3.0 T MAGNETOM Skyra, Siemens Healthineers, Forchheim, Germany). Imaging parameters are published in (15) and include a voxel size of 1.4 × 1.4 × 8.0 mm and a flip angle of 35°. The average interval between clinical examination and MR imaging was 28 to 42 days.

Analysis and Quality Control

Automatic segmentation of the myocardium and identification of the right ventricular insertion point were performed using a deep learning-based research algorithm (Siemens Healthineers, Forchheim, Germany) (16). Mean T1 values were calculated globally and for the six midventricular segments defined by the American Heart Association (AHA) (17) using dedicated research software (18).

After exclusion of repeated or failed T1 mapping acquisitions, image quality and artificial intelligence (AI)-generated segmentations were reviewed manually by an experienced physician if the mean T1 time of at least one myocardial segment differed by more than three standard deviations from age- (10-year groups) and sex-specific means. For this purpose, a custom application was developed in Python 3.11 that allows the inspection of T1 maps with overlayed contours alongside corresponding source images and metadata. In case of insufficient image quality due to inadequate breathing motion correction or co-registration of individual images, electrocardiographic trigger errors, incorrect orientation or other image artifacts not limited to individual myocardial segments, the entire examination was excluded from analysis. If at most three segments had to be excluded, the mean myocardial T1 time was calculated over the remaining segments weighted by their area. Manual quality control followed recommendations for clinical evaluation (1).

To validate the analysis pipeline, a representative sample of 100 cases was selected from the full dataset (matched for age, sex, height, weight, blood pressure as well as cardiovascular and metabolic condition prevalence). These cases were additionally analysed manually in CE-certified medical software and the results were quantitatively compared to those obtained from the automated segmentation followed by quality control.

Statistical Analysis

The statistical analysis was carried out in Python 3.12 using package statsmodels 0.14.4 and following the recommendations for analysing data from the NAKO cohort (19). Variables of interest included: age, sex, body mass index (BMI), blood pressure, heart rate, cardiovascular or metabolic self-reported diseases, current smoking, and laboratory parameters, including HbA1c, LDL-cholesterol and serum creatinine. Variables were primarily sourced from the clinical examination with only age extracted from the DICOM image metadata.

Mean T1 relaxation time and the averaged standard deviation in T1 were analysed by sex and myocardial segment. Since lateral segments generally exhibit more artifacts and lower T1 values, all statistical analyses were additionally performed for the two septal segments only and are reported in the supplementary information. A healthy subcohort was defined as follows: none of the diseases listed in Table 1, no current smoking, a BMI between 18.5 and 30.0 kg/m^2, blood pressure < 140/90 mmHg, LDL-cholesterol < 3.0 mmol/l (116 mg/dl), HbA1c < 48 mmol/mol (6.5%) and serum creatinine < 106 µmol/l (1.2 mg/dl) for men or < 88 µmol/l (1.0 mg/dl) for women. By combining reported conditions and clinical chemistry, we aimed to exclude both treated patients with controlled disease and undiagnosed diseased participants.

An ordinary least squares linear regression model was used to examine the associations of age, sex, and their interaction (age×sex) with myocardial T1. Heart rate was included in the analysis as a covariate since the MOLLI technique may be sensitive to heart rate, mainly due to variable intervals between inversions (1,20). Age was mean-centred to facilitate interpretation and reduce collinearity. We observed a strong sex dependency that varied by age. This relationship was visualized using linear regression, median and interquartile range in 2-year bins. Heart rate was linearly associated with T1 (β = 0.35 ms/bpm [0.31 – 0.38]; p < 0.001). Accordingly, subsequent analyses were adjusted for heart rate as well as age, sex, and their interaction; where appropriate, analyses were also stratified by sex.

To evaluate the association between cardiometabolic risk factors and myocardial T1 relaxation time, we conducted linear regression analyses, modelling native myocardial T1 as the dependent variable. Independent variables of interest comprised the presence of self-reported diabetes or elevated HbA1c, reported hyperlipidaemia or elevated LDL-cholesterol, reported hypertension or elevated measured blood pressure, reported kidney disease or elevated creatinine levels, and current smoking. Elevated blood parameters are based on the cut-off values used in the definition of the healthy subcohort. The European Society of Cardiology (ESC) and European Atherosclerosis Society define an LDL-cholesterol target value of less than 116 mg/dl (3.0 mmol/l) for low-risk populations (21). Other cardiovascular disease (myocardial infarction, coronary artery disease, heart failure, arrythmia, peripheral artery disease) as well as thyroid disease and cancer were included as covariates in the regression model.

In a sub-analysis, measured blood pressure was categorized according to the hypertension classification of the ESC guidelines (22) into grade I (≥ 140/90 mmHg), grade II (≥ 160/110 mmHg), grade III (≥ 180/110 mmHg), and isolated systolic or diastolic hypertension. We calculated sex-stratified linear regression models for myocardial T1 as the dependent variable, hypertension categories as independent variates of interest and included all other variables except hypertension from the previous analysis as covariates. In a further

linear regression analysis restricted to participants with self-reported hyperlipidaemia, the influence of lipid-lowering therapy was examined, adjusting for all covariates from the previous models except for hyperlipidaemia and elevated LDL-cholesterol.

We investigated nonlinear associations between BMI, mean arterial pressure (MAP) and heart rate with myocardial T1 using locally estimated scatterplot smoothing stratified by sex. Median and 95% confidence intervals were computed via bootstrapping for equal-width bins. Sex-stratified density plots are displayed along the same horizontal axis for each variable. BMI was excluded from the regression models as it may act as a mediator in the causal pathway from lifestyle or metabolic factors to cardiometabolic conditions such as hypertension and diabetes. Adjusting for BMI in this context could attenuate the observed effects of these conditions on myocardial T1. Variance inflation factors were calculated for all variates in the regression models and remained below two. Subjects with missing values were omitted from the respective regression analyses.

Ethics

This study was approved by the NAKO Use and Access Committee, based on the participants' informed consent and consistency with the objectives of the NAKO. Ethical approval for the NAKO study was obtained from the responsible local ethics committees of all study centres. The study was conducted in accordance with the Declaration of Helsinki and subsequent amendments.

Results

Quality Control

Of the 29,573 available T1 maps, 9,162 images with corresponding myocardial segmentation were selected for manual quality control, following the criteria outlined above. Among these, 1,779 subjects were excluded entirely from the analysis, while 4,805 cases involved the exclusion of one to three individual myocardial segments. The quality control procedure and results are illustrated in Figure 1. Artifacts were most frequently found in the inferolateral (3,634) and inferior (2,761) myocardial segments. The exclusions reduced the standard deviation of the overall mean T1 in the final dataset from 48 ms to 35 ms, while the mean itself remained unchanged (difference < 0.1 ms). Following quality control, a total of 27,794 (44.6% female) participants were included in the final analysis.

Comparison of the automated segmentation and manual quality control with a fully manual analysis in the validation subset revealed only a small mean deviation of -1.16 ± 10.49 ms (Supplementary Figure 1). Intraclass correlation for the two-way mixed, single-score model (ICC 3,1) was excellent, with a coefficient of 0.96 [0.94 – 0.97] (p < 0.001).

Figure 1: Manual Quality Control

A) T1 maps with abnormal values that deviate by more than three standard deviations from age- and sex-specific means in at least one myocardial segment are displayed together with the artificial intelligence-generated segmentation in a Python-based interface. Individual segments with artifacts can be excluded from the analysis by clicking on the respective area. B) Results from manual quality control.

Participant Characteristics

Participant characteristics and the prevalence of self-reported conditions are summarized in Table 1. Following the definition outlined above, 3,910 (14.1%) participants were assigned to the healthy subcohort. For the combination of self-reported medical history and elevated blood parameters, 1,341 (4.8%) participants reported diabetes or had elevated HbA1c, 18,263 (65.7%) reported hyperlipidaemia or had elevated LDL-cholesterol, 10,363 (37.3%) reported hypertension or had elevated blood pressure, and 1,147 (4.1%) reported kidney disease or had elevated serum creatinine levels.

Table 1: Basic Characteristics of Participants and Prevalence of Self-Reported Conditions

	Male (n = 15,393)	Female (n = 12,401)	Missing Values
Age [years]	48.3 ± 12.4 [20 – 75]	49.0 ± 12.2 [20 – 74]	0 (0.0%)
BMI [kg/m²]	26.8 ± 4.1 [15.6 – 52.3]	25.8 ± 5.2 [14.9 – 54.6]	352 (1.3%)
Heart Rate [bpm]	68.8 ± 10.8 [24 – 147]	70.8 ± 10.1 [28 – 120]	15 (0.1%)
Systolic BP [mmHg]	131.0 ± 14.6 [83 – 218]	122.9 ± 16.3 [81 – 234]	15 (0.1%)
Diastolic BP [mmHg]	80.4 ± 9.8 [49 – 136]	76.8 ± 9.7 [48 – 129]	15 (0.1%)
HbA1c [mmol/mol]	36.0 ± 6.5 [15 – 131]	35.7 ± 5.6 [6 – 127]	651 (2.3%)
LDL-cholesterol [mmol/l]	3.4 ± 0.9 [0.4 – 8.0]	3.2 ± 0.9 [0.7 – 9.9]	664 (2.4%)
Serum Creatinine [μmol/l]	80.8 ± 14.5 [7 – 641]	64.7 ± 10.8 [11 – 347]	544 (2.0%)
Diabetes	659 (4.3%)	505 (4.1%)	31 (0.1%)
Hyperlipidaemia	3,470 (22.5%)	2,608 (21.0%)	192 (0.7%)
Hypertension	3,903 (25.4%)	2,594 (20.9%)	57 (0.2%)
Kidney Disease	245 (1.6%)	261 (2.1%)	46 (0.2%)
Current Smoker	2,954 (19.2%)	2,169 (17.5%)	537 (1.9%)
Myocardial Infarction	67 (0.4%)	19 (0.2%)	24 (0.1%)
CAD or Angina Pectoris	157 (1.0%)	60 (0.5%)	68 (0.2%)
Arrhythmia	958 (6.2%)	940 (7.6%)	146 (0.5%)
Heart Failure	205 (1.3%)	168 (1.4%)	94 (0.3%)
Peripheral Artery Disease	152 (1.0%)	99 (0.8%)	61 (0.2%)
Thyroid Disease	1,326 (8.6%)	4,061 (32.7%)	95 (0.3%)
Cancer	654 (4.2%)	810 (6.5%)	61 (0.2%)

Values are mean ± standard deviation [min-max] or count (%). Missing values were omitted for the regression models. BMI: body mass index; BP: blood pressure; CAD: coronary artery disease

Age and Sex Dependence of Myocardial Native T1

The mean myocardial T1 relaxation time for 27,794 participants was 1,219 ± 35 ms at 3.0 T. Septal T1 (1,237 ± 40 ms) was significantly (p < 0.001 by two-sided paired t-test) higher than in lateral segments (1,200 ± 46 ms). Inferolateral and inferior segments had the highest variance in T1 values (Figure 2).

Figure 2: Polar Plots for Mean Myocardial T1 per AHA Segment

Values are given as the mean and average standard deviation of T1 within each segment (i.e., the mean of the standard deviations of individual values). AHA: American Heart Association

In the linear regression model, sex was significantly associated with myocardial T1 relaxation time (β = 20.92 ms [20.14 – 21.70] at the mean age of 48.6 years; p < 0.001). However, the effect of sex varied with age and showed a significant interaction (p < 0.001). While T1 increased with age in male participants (β = 0.21 ms/year [0.17 – 0.25]; p < 0.001), the age dependence was even stronger but inverse for women (β = −0.37 ms/year [−0.42 – −0.32]; p < 0.001) when stratified by sex (Figure 3). Visually, the age-dependency appeared less pronounced in the healthy subcohort, which was statistically confirmed by a significant three-way interaction term (p = 0.006). Mean values are shown in Table 2 for 10-year age groups (with the last bin covering ages 60 to 75). Corresponding values for septal segments only are provided in Supplementary Table 2.

Table 2: Mean Myocardial T1 for Age Groups by Sex

Age Group	Male mean ± std	n	Female mean ± std	n
20-29	1,205 ± 28	1,532	1,235 ± 33	1,167
30-39	1,207 ± 30	2,082	1,238 ± 34	1,427
40-49	1,208 ± 31	4,246	1,235 ± 32	3,445
50-59	1,210 ± 35	4,348	1,227 ± 31	3,643
60-75	1,213 ± 38	3,185	1,224 ± 32	2,719
Total	1,209 ± 34	15,393	1,231 ± 33	12,401

Two-way analysis of variance: age group p < 0.001; sex p < 0.001; age group-sex interaction p < 0.001. std: standard deviation; n: number of subjects

Figure 3: Age Dependence of Myocardial T1 in Men and Women

Linear regression (solid line), median (dotted line), and interquartile range (shaded area) shown in 2-year age bins, stratified by sex. The last bin is defined as ≥ 70 years for the whole cohort and ≥ 58 years for the healthy subcohort.

Associations of Cardiometabolic Risk Factors with Myocardial T1

Participants with self-reported diabetes or elevated HbA1c showed significantly higher T1 times in the overall cohort (p < 0.001, Figure 4). When stratified by sex, this association remained significant in men (p < 0.001), but not in women (p = 0.482). Similarly, kidney disease or elevated creatinine levels were linked to prolonged T1 relaxation times overall (p = 0.001), which was also significant in men (p < 0.001) but not women (p = 0.272).

Current smoking was consistently associated with elevated myocardial T1 across the entire cohort and in both sexes (p < 0.001 respectively). In contrast, hyperlipidaemia or elevated LDL-cholesterol was significantly associated with lower myocardial T1 values in the overall cohort and in both sexes (p < 0.001 respectively). Sex-specific effects were observed for reported hypertension or elevated blood pressure, with a significant positive β in men (p < 0.001) and a significant negative β in women (p < 0.001). In the entire cohort, the association was not significant (p = 0.855).

Corresponding regression coefficients and confidence intervals are provided in Supplementary Table 1. A confirmatory analysis restricted to septal segments rendered comparable results (Supplementary Figure 2). Among participants with hyperlipidaemia, lipid-lowering therapy was associated with a nonsignificant, small decrease in myocardial T1 (β = −0.82 ms [−2.89 – 1.26]; p = 0.440).

Figure 4: Associations of Cardiometabolic Risk Factors with Myocardial T1

Displayed are beta coefficients and 95% confidence intervals from linear regression models for the whole sample (left panel) and stratified by sex (right panel). HR: heart rate; CVD: cardiovascular disease; TD: thyroid disease

Body mass index exhibited a nonlinear, U-shaped relationship with myocardial T1, with higher T1 values observed in both underweight and overweight individuals (Figure 5). Consistent with the findings from the regression model, there was a pronounced and unexpected negative correlation at low blood pressure values (MAP < 90 mmHg) in female participants, while blood pressure was almost linearly positively correlated in male participants. The heart rate showed an approximately linear positive relationship to T1 for both sexes.

Figure 5: Local Regressions for Body Mass Index, Mean Arterial Pressure and Heart Rate

Locally estimated scatterplot smoothing, stratified by sex. Median and 95% confidence interval (bootstrapped with 1,000 iterations) are shown for nine equal-width bins. Values below the 0.1st or above the 99.9th percentile for heart rate were excluded. Two outliers were excluded for mean arterial pressure. Kernel density plots are displayed on a secondary y-axis along the same horizontal axis. CI: confidence interval

Impact of Hypertension Severity by ESC Classification

In the sex-stratified analysis based on the ESC classification of measured blood pressure (Figure 6), a progressive association between hypertension severity and myocardial native T1 time was observed in men. Compared to normotensive individuals, men with grade I hypertension had higher myocardial T1 (β = 3.14 ms [1.02 – 5.27]; p = 0.004), which further increased with grade II (β = 5.98 ms [3.12 – 8.83]; p < 0.001) and grade III hypertension (β = 8.44 ms [2.62 – 14.27]; p = 0.005). Isolated diastolic or systolic hypertension showed no significant trend (p = 0.422 and p = 0.303 respectively). Among women, isolated diastolic hypertension (β = −3.73 ms [−7.13 – −0.33]; p = 0.031) and isolated systolic hypertension (β = −3.62 ms [−5.89 – −1.34]; p = 0.002) were significantly associated with lower myocardial T1 time, while no significant associations were observed for grade I (p = 0.276), grade II (p = 0.277), or grade III hypertension (p = 0.778).

Figure 6: Sex-Stratified Associations Between ESC Hypertension Categories and Myocardial T1

Beta coefficients and 95% confidence intervals from multiple linear regression, stratified by sex. Covariates for diabetes, hyperlipidaemia and kidney disease are based on medical history and clinical chemistry. HT: hypertension; HR: heart rate; CVD: cardiovascular disease; TD: thyroid disease

Discussion

Summary of Key Findings

In this first, large-scale analysis of CMR-based T1 mapping in the NAKO study, we report associations between age, sex, and cardiometabolic risk factors with myocardial native T1 in 27,794 participants. Women exhibited overall significantly higher T1 values, with the sex difference progressively decreasing with age. Notably, we discovered a previously unreported inverse relationship between hyperlipidaemia and myocardial T1 and found sex-specific associations for cardiometabolic risk factors, especially hypertension.

Sex and Age Dependence of Myocardial Native T1

The observed sex differences in T1 are consistent with previous findings (12–14,23). While the influence of age remains controversial in the literature (13,23), our analysis aligns with results from healthy participants in the UK Biobank (12). The broader age distribution in the NAKO cohort allowed inclusion of younger adults and the capture of the menopausal transition, which may contribute to the observed sex-age interactions. Although thinner myocardial walls in women (24) might increase sensitivity to partial volume effects, this alone is unlikely to account for the marked difference. Results were comparable in the confirmatory septal analysis, reinforcing confidence in the quality control procedures and the AI-based segmentations. In the healthy subcohort, the age dependency was less pronounced and may therefore encompass differences in age-specific prevalence of diseases between sexes.

Hyperlipidaemia and Myocardial T1

The significant association of elevated blood lipids with lower myocardial T1 is a novel and unexpected finding. Hyperlipidaemia typically promotes fibrosis via atherosclerotic pathways (25) and would therefore be more consistent with prolonged T1 times (1). Our results suggest additional mechanisms, with possible explanations including altered myocardial lipid metabolism or early lipid infiltration. There is growing evidence that serum lipids have direct effects on the heart independent of atherosclerosis (11). Obesity is associated with adipocyte infiltration of the myocardium (26) and although the underlying mechanisms may differ, lipid overload as in Fabry disease presents with shortened T1 times (27). Another possible explanation would be a protective effect of statin therapy, which has been reported to attenuate extracellular volume expansion (28). Lipid-lowering therapy showed a modest negative association with myocardial T1, suggesting a potential moderating role. However, this association did not reach statistical significance.

Sex-Specific Associations with Hypertension

The observation that hypertension is associated with elevated myocardial T1 in men and lower values in women, indicates a sex-specific cardiac response to pressure overload. One possible explanation lies in sex-specific hormonal and genetic regulation of blood pressure and target organ damage (29). The observed difference also aligns with (30), where the authors reported greater strain impairment in hypertensive men, but contrasts with (31), where female sex was independently associated with left ventricular remodelling. Differing results may be related to women losing the cardioprotective effects of oestrogen in the context of obesity and diabetes, potentially influencing the myocardial response to hypertension (29,30). In men, the observed progressive increase in native T1 values across hypertension grades I to III suggests a dose-response relationship, where higher blood pressure levels are associated with greater myocardial remodelling.

Diabetes, Kidney Disease, and Smoking

Our findings of elevated myocardial T1 in individuals with diabetes, kidney disease or current smoking likely reflect known pathophysiological mechanisms: In recent studies, diabetes was independently associated with higher native T1 values, indicating potential subclinical myocardial dysfunction before structural remodelling occurs (12,32). Higher T1 values in individuals with diabetes may reflect expanded intracellular water volume in hypertrophied cardiomyocytes (32). Additionally, advanced glycation end-products increasingly form in the presence of chronic hyperglycaemia and are linked to oxidative stress and coronary artery disease (33). Diffuse myocardial fibrosis, as measured by myocardial native T1 and extracellular volume, has also been reported in early-stage chronic kidney disease (8) and is a recognized pathophysiological factor in haemodialysis patients (34,35). Smoking is a well-

established driver of systemic inflammation, oxidative stress, and atherosclerosis (10). In contrast to studies that have modelled T1 as a predictor of disease risk (12), our analysis considers T1 as the response variable to reflect structural effects associated with cardiometabolic risk in a population-based setting.

Limitations

Medical history in the NAKO cohort was obtained via self-reported questionnaires, which may introduce recall or reporting error. To mitigate this limitation, self-reports were complemented with measured elevations in relevant blood biomarkers. Absolute effect sizes were overall moderate; however, the observed differences can be aligned with early or subtle alterations in myocardial tissue composition. Given the high sensitivity of T1 mapping and the large, population-based sample, even small differences may indicate subclinical remodelling. The short interval between MR imaging and clinical phenotyping, together with extensive quality control and the confirmatory septal analysis, strengthens the internal validity of our findings.

Conclusions

Our analysis provides large-scale population-based evidence linking cardiometabolic risk factors to myocardial native T1 time. Diabetes, kidney disease and smoking were associated with prolonged T1 times, indicating potential early structural changes in myocardial tissue composition. Conversely, hyperlipidaemia was negatively associated with T1, requiring further investigation. We observed dynamic age-dependent differences in T1 time between women and men, suggesting a strong influence of sex on myocardial composition, especially before the age of 50. Our findings underscore the complex interplay between metabolic and myocardial health, emphasizing the importance of comprehensive cardiometabolic risk assessment.

Acknowledgements

This project was conducted with data (application numbers NAKO-744, NAKO-745) from the German National Cohort (NAKO). The NAKO is funded by the Federal Ministry of Education and Research (BMBF) [project funding reference numbers: 01ER1301A/B/C, 01ER1511D, 01ER1801A/B/C/D and 01ER2301A/B/C], federal states of Germany and the Helmholtz Association, the participating universities and the institutes of the Leibniz Association. We thank all participants who took part in the NAKO study and the staff of this research initiative. We thank Siemens Healthineers for providing the research segmentation algorithm.

Author Contributions

CA, CLS, CS, JG, and JS-M designed the study. CA drafted the initial manuscript. HS conducted manual quality control. SR and TP provided support with statistical analysis. HS, JG, JMN, JS-M, MBS, RH, SR, TH, TN, and TP substantially contributed to the preparation and revision of the manuscript. CLS, CS, FB, JAD, JS-M, KHG, KN, MBS, MD, RB, SR, SNW, TKe, TKü, TN, and TP contributed to data acquisition as part of the NAKO study. All authors reviewed and approved the final version.

Declarations

Disclosure of Interest

FB and JS-M hold research grants from Siemens Healthineers. FB and CLS have received honoraria from the speaker's bureau of Siemens Healthineers. All other authors declare no competing interests.

Data Availability

Data is available by application to the NAKO transfer office (https://transfer.nako.de).

Funding

JMN is supported by the Berta-Ottenstein-Programme for Clinician Scientists, Faculty of Medicine, University of Freiburg. RH receives funding from the German Research Foundation (project 402688427).

References

1. Messroghli DR, Moon JC, Ferreira VM, Grosse-Wortmann L, He T, Kellman P, et al. Clinical recommendations for cardiovascular magnetic resonance mapping of T1, T2, T2* and extracellular volume: A consensus statement by the Society for Cardiovascular Magnetic Resonance (SCMR) endorsed by the European Association for Cardiovascular Imaging (EACVI). J Cardiovasc Magn Reson. 2016 Dec;19(1):75.

2. Taylor AJ, Salerno M, Dharmakumar R, Jerosch-Herold M. T1 Mapping. JACC Cardiovasc Imaging. 2016 Jan;9(1):67–81.

3. Schelbert EB, Messroghli DR. State of the Art: Clinical Applications of Cardiac T1 Mapping. Radiology. 2016 Mar;278(3):658–76.

4. Puntmann VO, D'Cruz D, Smith Z, Pastor A, Choong P, Voigt T, et al. Native Myocardial T1 Mapping by Cardiovascular Magnetic Resonance Imaging in Subclinical Cardiomyopathy in Patients With Systemic Lupus Erythematosus. Circ Cardiovasc Imaging. 2013 Mar;6(2):295–301.

5. Peters A, German National Cohort (NAKO) Consortium, Peters A, Greiser KH, Göttlicher S, Ahrens W, et al. Framework and baseline examination of the German National Cohort (NAKO). Eur J Epidemiol. 2022 Oct;37(10):1107–24.

6. Bamberg F, Schlett CL, Caspers S, Ringhof S, Günther M, Hirsch JG, et al. Baseline MRI Examination in the NAKO Health Study. Dtsch Arztebl Int. 2024;121(18):587–93.

7. Salvador DB, Gamba MR, Gonzalez-Jaramillo N, Gonzalez-Jaramillo V, Raguindin PFN, Minder B, et al. Diabetes and Myocardial Fibrosis. JACC Cardiovasc Imaging. 2022 May;15(5):796–808.

8. Edwards NC, Moody WE, Yuan M, Hayer MK, Ferro CJ, Townend JN, et al. Diffuse Interstitial Fibrosis and Myocardial Dysfunction in Early Chronic Kidney Disease. Am J Cardiol. 2015 May;115(9):1311–7.

9. González A, Ravassa S, López B, Moreno MU, Beaumont J, San José G, et al. Myocardial Remodeling in Hypertension: Toward a New View of Hypertensive Heart Disease. Hypertension. 2018 Sept;72(3):549–58.

10. McEvoy JW, Blaha MJ, DeFilippis AP, Lima JAC, Bluemke DA, Hundley WG, et al. Cigarette Smoking and Cardiovascular Events: Role of Inflammation and Subclinical Atherosclerosis From the Multiethnic Study of Atherosclerosis. Arterioscler Thromb Vasc Biol. 2015 Mar;35(3):700–9.

11. Yao YS, Li TD, Zeng ZH. Mechanisms underlying direct actions of hyperlipidemia on myocardium: an updated review. Lipids Health Dis. 2020 Feb 8;19(1):23.

12. Raisi-Estabragh Z, McCracken C, Hann E, Condurache DG, Harvey NC, Munroe PB, et al. Incident Clinical and Mortality Associations of Myocardial Native T1 in the UK Biobank. JACC Cardiovasc Imaging. 2023 Apr;16(4):450–60.

13. Myhr KA, Andrés-Jensen L, Larsen BS, Kunkel JB, Kristensen CB, Vejlstrup N, et al. Sex- and age-related variations in myocardial tissue composition of the healthy heart: a native T1 mapping cohort study. Eur Heart J - Cardiovasc Imaging. 2024 July 31;25(8):1109–17.

14. Cavus E, Schneider JN, Bei der Kellen R, di Carluccio E, Ziegler A, Tahir E, et al. Impact of Sex and Cardiovascular Risk Factors on Myocardial T1, Extracellular Volume Fraction, and T2 at 3 Tesla: Results From the Population-Based, Hamburg City Health Study. Circ Cardiovasc Imaging. 2022 Sept 1;15(9):e014158.

15. Bamberg F, Kauczor HU, Weckbach S, Schlett CL, Forsting M, Ladd SC, et al. Whole-Body MR Imaging in the German National Cohort: Rationale, Design, and Technical Background. Radiology. 2015 Oct;277(1):206–20.

16. Popescu AB, Seitz A, Mahrholdt H, Wetzl J, Jacob A, Itu LM, et al. Deep learning-based segmentation of T1 and T2 cardiac MRI maps for automated disease detection [Internet]. arXiv; 2025 [cited 2025 July 3]. Available from: https://arxiv.org/abs/2507.00903

17. American Heart Association Writing Group on Myocardial Segmentation and Registration for Cardiac Imaging: Cerqueira MD, Weissman NJ, Dilsizian V, Jacobs AK, Kaul S, et al. Standardized Myocardial Segmentation and Nomenclature for Tomographic Imaging of the Heart: A Statement for Healthcare Professionals From the Cardiac Imaging Committee of the Council on Clinical Cardiology of the American Heart Association. Circulation. 2002 Jan 29;105(4):539–42.

18. Ammann C, Hadler T, Reisdorf P, Hickstein R, Noyan H, Gröschel J, et al. Cardiometry: Open-Source, Precise and Efficient Computation of Quantitative Parameters in Cardiovascular Magnetic Resonance. In: Proceedings of the ISMRM 33rd Annual Meeting and Exhibition. Honolulu, HI, USA: International Society for Magnetic Resonance in Medicine, Berkley, CA, USA; 2025.

19. Kuss O, Becher H, Wienke A, Ittermann T, Ostrzinski S, Schipf S, et al. Statistical Analysis in the German National Cohort (NAKO) – Specific Aspects and General Recommendations. Eur J Epidemiol. 2022 Apr;37(4):429–36.

20. Kellman P, Hansen MS. T1-mapping in the heart: accuracy and precision. J Cardiovasc Magn Reson. 2014 Dec;16(1):2.

21. Mach F, Baigent C, Catapano AL, Koskinas KC, Casula M, Badimon L, et al. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. Eur Heart J. 2020 Jan 1;41(1):111–88.

22. McEvoy JW, McCarthy CP, Bruno RM, Brouwers S, Canavan MD, Ceconi C, et al. 2024 ESC Guidelines for the management of elevated blood pressure and hypertension. Eur Heart J. 2024 Oct 7;45(38):3912–4018.

23. Rosmini S, Bulluck H, Captur G, Treibel TA, Abdel-Gadir A, Bhuva AN, et al. Myocardial native T1 and extracellular volume with healthy ageing and gender. Eur Heart J - Cardiovasc Imaging. 2018 June 1;19(6):615–21.

24. Kawel-Boehm N, Hetzel SJ, Ambale-Venkatesh B, Captur G, Francois CJ, Jerosch-Herold M, et al. Reference ranges ("normal values") for cardiovascular magnetic resonance (CMR) in adults and children: 2020 update. J Cardiovasc Magn Reson. 2020 Jan;22(1):87.

25. Ross R, Harker L. Hyperlipidemia and Atherosclerosis: Chronic hyperlipidemia initiates and maintains lesions by endothelial cell desquamation and lipid accumulation. Science. 1976 Sept 17;193(4258):1094–100.

26. Pathak RK, Mahajan R, Lau DH, Sanders P. The Implications of Obesity for Cardiac Arrhythmia Mechanisms and Management. Can J Cardiol. 2015 Feb;31(2):203–10.

27. Sado DM, White SK, Piechnik SK, Banypersad SM, Treibel T, Captur G, et al. Identification and Assessment of Anderson-Fabry Disease by Cardiovascular Magnetic Resonance Noncontrast Myocardial T1 Mapping. Circ Cardiovasc Imaging. 2013 May;6(3):392–8.

28. Juhasz V, Quinaglia T, Drobni ZD, Heemelaar JC, Neuberg DS, Han Y, et al. Atorvastatin and Myocardial Extracellular Volume Expansion During Anthracycline-Based Chemotherapy. JACC CardioOncology. 2025 Feb;7(2):125–37.

29. Colafella KMM, Denton KM. Sex-specific differences in hypertension and associated cardiovascular disease. Nat Rev Nephrol. 2018 Mar;14(3):185–201.

30. Tadic M, Cuspidi C, Celic V, Ivanovic B, Pencic B, Grassi G. The influence of sex on left ventricular strain in hypertensive population. J Hypertens. 2019 Jan;37(1):50–6.

31. Canciello G, Piccolo R, Izzo R, Bossone E, Pacella D, Lembo M, et al. Sex-Related Differences in Left Ventricular Geometry Patterns in Patients With Arterial Hypertension. JACC Adv. 2024 Oct;3(10):101256.

32. Lam B, Stromp TA, Hui Z, Vandsburger M. Myocardial native-T1 times are elevated as a function of hypertrophy, HbA1c, and heart rate in diabetic adults without diffuse fibrosis. Magn Reson Imaging. 2019 Sept;61:83–9.

33. Fishman SL, Sonmez H, Basman C, Singh V, Poretsky L. The role of advanced glycation end-products in the development of coronary artery disease in patients with and without diabetes mellitus: a review. Mol Med. 2018 Dec;24(1):59.

34. Rutherford E, Talle MA, Mangion K, Bell E, Rauhalammi SM, Roditi G, et al. Defining myocardial tissue abnormalities in end-stage renal failure with cardiac magnetic resonance imaging using native T1 mapping. Kidney Int. 2016 Oct;90(4):845–52.

35. Aoki J, Ikari Y, Nakajima H, Mori M, Sugimoto T, Hatori M, et al. Clinical and pathologic characteristics of dilated cardiomyopathy in hemodialysis patients. Kidney Int. 2005 Jan;67(1):333–40.