Validation of the Perimenopause Symptom Scale (Peri-SS) for Digital Self-Assessment: Psychometric Evaluation Among U.S. Women Aged 35 to 59 years

Yihan Xu PhD[1], Irina Llyich MD[1], Carley Prentice MSc[1], Yella Hewings-Martin PhD[1], Adam Cunningham PhD[1], Chrisandra Shufelt MD, MS[2-4], Stephanie Faubion MD, MBA[2-4], Liudmila Zhaunova PhD[1]

[1] Flo Health Inc., London
[2] Division of General Internal Medicine, Mayo Clinic, Jacksonville, Florida
[3] Mayo Clinic Center for Women's Health, Rochester, Minnesota
[4] Women's Health Research Center, Mayo Clinic, Rochester, Minnesota

Correspondence should be addressed to Yihan Xu (yihan_xu@flo.health)
27 Old Gloucester Street,
London,
WC1N 3AX,
United Kingdom

Abstract

Background
Perimenopause is a transitional life stage that ultimately affects all women with ovaries, yet it remains under-recognized in clinical care and poorly measured by existing tools. Most symptom scales were designed for postmenopausal populations, overlooking the unique and fluctuating symptom profiles that emerge in perimenopause. This gap hinders early recognition and timely intervention.

Objective
To address this unmet need, we developed and validated the Perimenopause Symptom Scale (Peri-SS): a concise, phase-specific, and psychometrically robust instrument intended for digital self-assessment and research use among U.S. women aged 35-59 who self-reported in perimenopause.

Methods
We conducted a longitudinal validation study with a final analytic sample of 1,255 U.S. women aged 35-59 (633 identified as perimenopausal) recruited from an online survey platform Prolific. Participants completed the Peri-SS along with the Menopause Rating Scale (MRS) and EuroQol 5-Dimension, 5-Level (EQ-5D-5L) at baseline and again 7-14 days later. We assessed structural validity, internal consistency, test-retest reliability, criterion and construct validity, and responsiveness to change, following COSMIN guidelines.

Results
The Peri-SS demonstrated strong psychometric performance. Confirmatory factor analysis confirmed a four-domain structure (vasomotor, psychological, sexual, physical), with excellent model fit (CFI = 0.932, RMSEA = 0.069). Internal consistency was excellent (α = 0.90 overall; domain range 0.73-0.86), and test-retest reliability was high (ICC = 0.82). Criterion validity was strong, with Peri-SS scores being highly correlated with MRS (r = 0.88). Construct validity was supported through moderate correlations with EQ-5D-5L utilities (ρ = –0.50) and effective discrimination between known groups by age, menopausal status, and hormone therapy use. The scale was also responsive to symptom changes over time (η^2 = 0.277).

Conclusion
The Peri-SS is a reliable, valid, and responsive tool for assessing symptom burden in perimenopause. Its concise format and strong psychometric performance make it well-suited for research, clinical care, and digital health applications.

Keywords. Perimenopause, perimenopause symptom scale (Peri-SS), psychometric evaluation, reliability, validity, symptoms, digital health

Introduction

Perimenopause is a transitional life stage that typically begins in a woman's 40s and often lasts
4 to 10 years before the onset of menopause, defined retrospectively after 12 consecutive
months of amenorrhea [1]. During this phase, women experience marked fluctuations of ovarian
hormones, particularly erratic and declining serum concentrations of 17β-estradiol as well as
changes in progesterone and follicle-stimulating hormone (FSH), which can trigger a wide range
of disruptive physical, cognitive, and psychological symptoms, such as hot flashes, night
sweats, disrupted sleep, mood changes, fatigue, and sexual health concerns [2–5]. Although
severity and duration vary, these symptoms can substantially impair daily functioning and quality
of life [6,7].

Importantly, emerging research suggests that perimenopause is not simply an early or less
intense version of menopause, but a distinct phase with its own symptom patterns, triggers, and
psychosocial impact [8]. For example, data from the Study of Women's Health Across the Nation
(SWAN) identified four distinct trajectories of vasomotor symptoms across the reproductive
stages, many of which begin in perimenopause and exhibit patterns not seen in
postmenopause [9]. A large-scale analysis of over 145,000 digital symptom logs also found that
perimenopausal women exhibit a unique pattern that is marked by the coexistence of persistent
menstrual cycle-related symptoms and emerging menopause symptoms, distinct from both pre-
and postmenopausal groups [10].

Despite being a universal and distinct experience that affects the quality of life for women,
perimenopause remains poorly recognized in clinical practice and under-represented in health
policy and research agendas [11,12]. And many women feel underprepared for this life stage and
report inadequate support from healthcare providers and workplaces [13,14].

A major barrier to timely care is lack of awareness: many women are unfamiliar with the full
range of symptoms that often begin in perimenopause and may mistakenly attribute them to
stress, unrelated health issues, or part of "normal aging" [13,15]. This challenge is compounded if
those changes occur in the absence of classic vasomotor symptoms or menstrual cycles are still
present [1,13], especially among women in their 30s or early 40s who may fall outside conventional
expectations of menopausal age [16].

Another key obstacle to early recognition and better support lies in the lack of perimenopause-
specific measurement tools. Whereas several instruments assess symptoms associated with
menopause, such as the Menopause Rating Scale (MRS) [17], the Menopause-specific Quality of
Life (MENQOL) Questionnaire [18], the Greene Climacteric Scale [19,20], and the Menopause
Transition Scale (MTS) [21] – these were developed for assessment of postmenopausal women,
rely on paper-based administration, and often omit emerging perimenopausal symptoms such
as digestive issues and sexual health concerns [10].

As more women turn to digital tools to track and manage their reproductive health, including
menstrual changes and perimenopausal symptoms [22,23], there is growing need for validated,
user-friendly instruments designed for scalable deployment in digital health settings.

The MenoScale represents a promising step toward modernisation, introducing a digital self-
assessment tool for menopausal symptoms measurement [24]. However, it was not specifically
developed for perimenopausal populations and may not fully capture their distinct symptom
experiences.

Taken together, existing tools fall short in capturing the multifaceted and fluctuating symptom
experience of women in perimenopause, which limits their usefulness for symptom tracking,
early intervention, or integration into digital care platforms.

In response to these gaps and aligned with the research priorities in menopause [12], we
developed and validated the Perimenopause Symptom Scale (Peri-SS): a concise,
perimenopause-specific, and psychometrically robust instrument designed for use in clinical
research and digital health contexts.

Methods
103

Scale development and validation study design
104

The development of the Peri-SS included initial item generation from literature review and
clinical input, followed by item reduction through exploratory factor analysis (EFA) in a separate
sample. This paper focuses primarily on validating the resulting 16-item scale using a
prospective, longitudinal cohort study.

The validation procedures followed the COnsensus-based Standards for the selection of health
Measurement INstruments (COSMIN) methodology for evaluating patient-reported outcome
measures (PROMs) [25] and adhered to key reporting elements from the STROBE guidelines [26].
The validation study was adequately powered based on the COSMIN-recommended sample
size of at least 10 participants per item [27], and the final analytic sample exceeded these
requirements. The study protocol was reviewed and approved by the WCB Institutional Review
Board (IRB No. 20250554).

Participants and recruitment
116

Participants were recruited via Prolific, an online research platform, between February 27 and
March 16, 2025. Eligible individuals were women aged 35 to 59 years, residing in the United
States, fluent in English. The surveys were administered via the survey platform SurveyMonkey.
Women were excluded if they were currently pregnant, had experienced pregnancy loss within
the past 90 days, had given birth in the past 12 months, had undergone a hysterectomy, or did
not provide informed electronic consent. These criteria were selected to ensure that participants
were representative of the population likely to experience perimenopause symptoms, while
minimizing potential confounding effects from recent pregnancy or potential surgical
menopause. Participants were remunerated at a rate of $12-15 USD per hour for their time
completing the surveys.
127

Participants completed a baseline survey (T1) and a follow-up survey (T2), 7 to 14 days later.
This interval was chosen to minimize recall bias and avoid substantial natural variation in
symptom severity while allowing for an assessment of temporal reliability. Participants
completed the Peri-SS, Menopause Rating Scale, and EuroQol five-dimensional questionnaire
(EQ-5D-5L) at both T1 and T2, whereas they answered existing medical conditions,
demographics, and other questions only at T1. The self-reported menopause stage was
assessed via a single-item question, where participants selected the stage that best described
them after reviewing standardized definitions and a visual timeline of the reproductive stages
(see **Supplementary material A**).

The development of the Perimenopause Symptom Scale (Peri-SS)

The Peri-SS was developed to address the lack of concise, perimenopause-specific instruments
suitable for digital self-assessment. Initial item generation drew from a targeted review of more
than 15 validated scales assessing menopause and perimenopause symptoms, including the
Menopause Rating Scale (MRS) [17,28], the Greene Climacteric Scale (GCS) [19,20], MENQOL [18],
and the Women's Health Questionnaire (WHQ) [29]—as well as input from clinical experts.

From an initial pool of 25 items, 16 were retained following exploratory factor analysis (EFA) in a
sample of 948 U.S.-based women who self-reported as perimenopausal as part of the Global
Perimenopause Survey among Flo users (data collected from December 4, 2024 to January 6,
2025, IRB# 20244604). Item selection was based on factor loadings, internal consistency, and
clinical interpretability, resulting in a four-domain structure: vasomotor symptoms (2 items),
psychological and emotional stress (5 items), sexual health (3 items), and physical/somatic
changes (6 items).

This scale structure was reviewed by clinicians to ensure clinical validity and relevancy.
Each item is rated on a 5-point Likert scale ranging from 0 (none) to 4 (very severe), reflecting
the impact of symptoms over the past 30 days. Domain scores are computed as the mean of
items within each domain, and the overall Peri-SS score is calculated as the average of domain
scores and scaled to a 0-100 range, with higher scores reflecting greater symptom burden (the
final item list are detailed in **Supplementary material B**).

To support interpretation, Peri-SS score is categorized into four severity bands, conceptually
aligned with the MRS: none to mild (0-8), mild to moderate (9-18), moderate to severe (19-35),
and severe (≥36). These thresholds were empirically derived using a data-driven approach and
calibrated based on their alignment with benchmark measures of symptom burden and quality
of life (see **Supplementary Material C** for derivation procedure details).

Validation procedures

We evaluated the following psychometric properties of the Peri-SS: structural validity, criterion
validity, construct validity (including hypothesis testing and known-group comparisons), internal
consistency, and test-retest reliability. Where relevant, thresholds for interpretation followed
COSMIN and related psychometric conventions [25], and we present the COSMIN checklist in

Supplementary material D. All statistical analyses were conducted using R (version 4.4.1),
using the "lavaan" package [30] for structural equation modeling and "psych" package [31] for
reliability analyses. All reliability and validity estimates were calculated using complete-case
data from T1 survey, except for test-retest reliability and responsiveness, which used data from
both T1 and T2.

Structural validity. Construct validity assesses whether the Peri-SS accurately measures the
underlying concept of perimenopause symptoms. We conducted confirmatory factor analysis
(CFA) to test whether the four-factor model identified during instrument development adequately
fit the observed data. The model was estimated using full-information maximum likelihood
(FIML) to account for missing data and applied standardized latent variable variances. Model fit
was assessed using the Comparative Fit Index (CFI) and the Root Mean Square Error of
Approximation (RMSEA). Fit was considered excellent if CFI > 0.95 or RMSEA < 0.06, and
acceptable if CFI > 0.90 or RMSEA < 0.08, in line with COSMIN criteria [25].

Internal consistency. Internal consistency refers to how well the items on a scale measure the
same underlying concept. We assessed it using Cronbach's alpha (α) for the total score and
each of the four domains. Values of $\alpha \geq 0.70$ were considered acceptable, ≥ 0.80 good, and $\geq$
0.90 excellent. Unidimensionality within domains was verified by reviewing inter-item
correlations and factor loadings from the CFA.

Test-retest reliability. To evaluate the temporal stability of the Peri-SS, we assessed test-
retest reliability over a 7-14 day interval using data from participants who completed both T1
and T2 surveys. We calculated the intraclass correlation coefficient (ICC) using a two-way
mixed-effects model with absolute agreement for the total Peri-SS score.

Responsiveness. Responsiveness refers to the ability of an instrument to detect meaningful
change over time in the construct it intends to measure. We assessed responsiveness of the
Peri-SS by examining whether changes in Peri-SS scores reflected changes in symptom
severity, using the MRS as an external anchor. Responsiveness of the Peri-SS was evaluated
using an anchor-based approach. Change in total MRS score between T1 and T2 surveys
served as the external anchor. Participants were categorized into three groups based on MRS
score change between baseline and follow-up: *improved* (≥4-point reduction), *no change*
(change between –4 and +4 points), or *worsened* (≥4-point increase). We then compared Peri-
SS score changes across these groups using the Kruskal-Wallis test. Effect size was quantified
using eta-squared based on the H-statistic ($\eta^2[H]$), with thresholds of 0.01, 0.06, and 0.14
representing small, medium, and large effects, respectively [32].

Criterion validity. Criterion validity examines whether Peri-SS scores align with an established
domain-specific measure. We assessed criterion validity by comparing the Peri-SS to the MRS,
a widely used instrument in clinical and research contexts to evaluate menopausal symptom
burden. While the MRS is not a gold standard for perimenopause-specific symptom burden, it

includes overlapping domains such as psychological, somatic, and urogenital symptoms,
making it a suitable reference for perimenopausal symptom burden [17,28].

We computed Pearson's correlation coefficient (r) to quantify the correlation between Peri-SS
and the two established scales, and we considered the correlation to be strong if r ≥ 0.70.
We also computed Cohen's kappa (κ) to assess agreement in categorical symptom severity
classifications (e.g., minimal, mild, moderate, severe). The correlation coefficient was
considered strong if r or κ is ≥ 0.70 [25].

Construct validity. Construct validity examines whether the Peri-SS measures the theoretical
construct it is intended to measure—namely, the perimenopause symptom burden. We
assessed construct validity using two complementary approaches: convergent validity and
known-group validity.

For convergent validity, we evaluated the relationship between Peri-SS total scores and the EQ-
5D-5L utility, a widely used measure of general health-related quality of life [33]. The EQ-5D-5L
instrument includes five dimensions (mobility, self-care, usual activities, pain/discomfort, and
anxiety/depression), each rated on five levels of severity. The responses were converted into
utility values using the U.S. population-specific value set [34]. We hypothesized that greater
symptom burden, as captured by the Peri-SS, would be associated with lower EQ-5D utility
values. We computed correlation using Spearman's ρ as the distribution of EQ-5D-5L scores
were highly skewed. Based on COSMIN thresholds, the convergent validity was considered
strong if |r| or |ρ| ≥ 0.50; weak if they are below 0.30; moderate if they're between 0.3 and 0.5 [25].

For known-group validity, we compared Peri-SS scores across subgroups expected to differ in
symptom severity, including age group, menopausal status, and hormone therapy (HT) use. As
Peri-SS scores were not normally distributed, we used Kruskal-Wallis tests to assess overall
group differences. Where significant main effects were detected, we conducted post hoc
pairwise comparisons using Wilcoxon rank-sum tests. All *p*-values from pairwise comparisons
were adjusted using the Bonferroni method to account for multiple testing. We hypothesized that
symptom scores would be higher among women in their mid-40s to early 50s, among those in
perimenopause or postmenopause compared to premenopausal women, and among those not
using HT.

Results

Participant characteristics

245 Figure 1 outlines the participant recruitment and eligibility flow. Of 1,537 individuals screened,
246 1,255 completed the baseline (T1) survey and met all eligibility criteria, including providing their

247   menopausal status. These participants formed the primary analytic sample for all baseline
248   psychometric analyses.
249

250
251   ***Figure 1.*** Recruitment flow
252
The majority self-identified as perimenopausal (50.4%), with the remainder identifying as
premenopausal (26.1%), postmenopausal (9.5%), or unsure (14.0%). The perimenopause
subgroup (n = 633) was used for confirmatory factor analysis and internal consistency
assessments. All menopausal status groups were included in construct validity analyses.
Table 1 summarizes baseline sociodemographic and clinical characteristics of the baseline
sample. Among the 1,255 eligible participants who completed the T1 survey, 1,107 completed
the T2 survey, of whom 1,099 remained eligible and were included in test-retest reliability
analyses.
261

262 **Table 1.** Characteristics of eligible participants at baseline

	Overall (N=1255)	By menopausal status			
		Peri menopause (N=633)	Post menopause (N=119)	Pre menopause (N=327)	Unsure (N=176)
Age (years)					
Mean (SD)	45.5 (5.38)	46.2 (4.30)	53.3 (4.42)	42.8 (4.68)	42.8 (4.70)
BMI (kg/m2)					
Mean (SD)	28.7 (7.59)	29.1 (7.65)	28.7 (7.53)	28.0 (7.42)	28.4 (7.66)
Median [Min, Max]	26.9 [10.7, 63.5]	27.4 [15.1, 59.1]	26.4 [16.1, 51.8]	25.9 [12.6, 63.5]	26.8 [10.7, 60.9]
Education level					
Bachelor's degree	443 (35.3%)	224 (35.4%)	38 (31.9%)	120 (36.7%)	61 (34.7%)
College/some college	210 (16.7%)	104 (16.4%)	24 (20.2%)	51 (15.6%)	31 (17.6%)
High school or below	336 (26.8%)	164 (25.9%)	33 (27.7%)	89 (27.2%)	50 (28.4%)
Postgraduate degree	261 (20.8%)	138 (21.8%)	24 (20.2%)	65 (19.9%)	34 (19.3%)
Prefer not to answer	5 (0.4%)	3 (0.5%)	0 (0%)	2 (0.6%)	0 (0%)
Race/Ethnicity					
Asian	51 (4.1%)	24 (3.8%)	1 (0.8%)	20 (6.1%)	6 (3.4%)
Black	159 (12.7%)	64 (10.1%)	17 (14.3%)	52 (15.9%)	26 (14.8%)
Hispanic	35 (2.8%)	15 (2.4%)	4 (3.4%)	11 (3.4%)	5 (2.8%)
Mixed/Other	98 (7.8%)	36 (5.7%)	9 (7.6%)	33 (10.1%)	20 (11.4%)
White	912 (72.7%)	494 (78.0%)	88 (73.9%)	211 (64.5%)	119 (67.6%)
Currently use of HT					
No	1190 (94.8%)	588 (92.9%)	104 (87.4%)	324 (99.1%)	174 (98.9%)
Yes	65 (5.2%)	45 (7.1%)	15 (12.6%)	3 (0.9%)	2 (1.1%)

263 * BMI = Body Mass Index

Structural validity

265 Structural validity was evaluated using confirmatory factor analysis (CFA) on the
266 perimenopausal subgroup (n = 633) to assess whether the hypothesized four-factor model
267 (vasomotor, psychological, sexual health, and physical / somatic domains) fit the observed data.

268
269   The CFA results supported the hypothesized four-factor structure of the Peri-SS. Model fit was
270   acceptable, with CFI = 0.932 and RMSEA = 0.069. All items loaded significantly on their
271   intended factors, with standardized factor loadings ranging from 0.536 to 0.872 (see Table 2).
272   These results suggest that the Peri-SS captures four distinct but correlated domains of
273   perimenopause symptoms.
274
For comparative purposes, we also conducted CFA on the MRS. It had a slightly lower model fit,
with CFI = 0.927 and RMSEA = 0.087. While all MRS items also loaded significantly on all three
domains, standardized loadings were lower in the somatic and urogenital domains (e.g., 0.470-
0.620 and 0.545-0.680, respectively). This comparison shows the structural coherence of the
Peri-SS and its potential for capturing a broader and more differentiated range of
perimenopause symptoms.

281
Figure 2. Path diagram of the final four-factor model of the Perimenopause Symptom Scale
(Peri-SS) based on confirmatory factor analysis. Circles represent latent constructs (domains),
and rectangles represent observed symptom items. Standardized factor loadings are shown on
each path and were all statistically significant at p < .001. Double-headed arrows represent
correlations between latent domains. This structure supports the theorized domains of
vasomotor, sexual, psychological, and physical symptoms in the Peri-SS.

Internal consistency

The Peri-SS demonstrated excellent overall reliability (standardized Cronbach's α = 0.90). Subscale reliability was also strong, with α values of 0.73 for the vasomotor domain, 0.78 for the sexual domain, 0.86 for the psychological domain, and 0.83 for the physical domain. These results indicate good to excellent internal consistency across all domains, suggesting that the items within each domain reliably measure a coherent underlying construct.

For comparison, the MRS showed an overall α of 0.86. Domain-specific values were 0.88 for psychological, 0.67 for somatic, and 0.62 for urogenital symptom domains. These results show that Peri-SS outperformed MRS in reliability among women who self-identified as perimenopausal.

Table 2. Internal consistency and standardized factor loadings for the four-factor structure of the Perimenopause Symptom Scale (Peri-SS)

	Vasomotor	Sexual	Psychological	Physical/Somatic
Cronbach's α	0.73	0.78	0.86	0.83
Hot flashes	0.729			
Night sweats	0.790			
Vaginal dryness		0.872		
Pain during sex		0.761		
Low sex drive		0.604		
Feeling down			0.825	
Irritability			0.756	
Anxiety			0.824	
Sleep difficulties			0.555	
Physical exhaustion			0.752	
Digestive issues				0.691
Hair changes				0.700
Skin changes				0.748
Joint discomfort				0.654
Bladder problems				0.536
Weight changes				0.691

Note: All factor loadings are standardized and statistically significant at $p < .001$.

Table 3 presents the descriptive summary of Peri-SS total score and domain scores, stratified by age group. The total score is scaled to range from 0 to 100, reflecting overall symptom burden, while domain scores range from 0 to 4, with higher scores indicating greater symptom

severity. The average total score across all participants was 32.6 (SD = 16.6), with higher
scores among those aged above 45 years compared with those aged 35-44 years. Across the
four symptom domains, psychological and emotional stress emerged as the most prominent
symptom domain across all age groups, while sexual health symptoms were rated lowest.
Vasomotor and sexual symptom severity generally increased with age, whereas psychological
symptoms peaked in midlife and declined slightly in the oldest age group. In contrast, physical
and somatic symptom scores remained relatively stable across age groups. A more detailed
breakdown of item-level scores is presented in Supplementary S-Table 2.
315
316 **Table 3.** Peri-SS total score and domain scores by age group among perimenopausal
317 participants (N = 633)
318

	35-44 (N=184)	45-54 (N=435)	55-64 (N=14)	Overall (N=633)
Total score	30.7 (16.6)	33.4 (16.6)	31.6 (14.7)	32.6 (16.6)
Vasomotor symptoms	1.1 (0.86)	1.2 (0.93)	1.3 (1.1)	1.2 (0.91)
Psychological and emotional stress	1.7 (0.83)	1.7 (0.83)	1.2 (0.72)	1.7 (0.83)
Sexual health	1.0 (0.94)	1.2 (0.94)	1.3 (1.1)	1.1 (0.94)
Physical and somatic changes	1.1 (0.81)	1.2 (0.76)	1.2 (0.84)	1.2 (0.78)

319 *Note:* Values are presented as Mean (SD). Total score ranges from 0 (minimal symptom) to 100 (severe
320 symptom), whereas domain scores range from 0 (minimal symptom) to 4 (severe symptom).

Test-retest reliability

We assessed test-retest reliability among 552 participants who remained eligible and completed
both T1 and T2 surveys within a 7-14 day interval. The intraclass correlation coefficient (ICC) for
the total Peri-SS score was 0.82 (95% CI: 0.70-0.88), indicating excellent reliability. Domain-
specific ICCs also demonstrated excellent temporal stability, ranging from 0.77 (physical), 0.80
(vasomotor), 0.80 (sexual), to 0.81 (psychological). In comparison, the total MRS score yielded
an ICC of 0.79 (95% CI: 0.66-0.85), with domain-specific ICCs ranging from 0.75 (urogenital),
0.77 (psychological), to 0.77 (somatic). These results support the Peri-SS as a stable and
reliable instrument for assessing perimenopause symptom burden over short intervals.

Responsiveness

The Peri-SS was responsive to changes in symptom burden over time. As shown in Figure 3,
Peri-SS score changes differed significantly across the MRS anchor categories (Kruskal-Wallis
χ^2 = 153.1, $p < 0.001$). Participants classified as *Improved* (n = 195) showed a mean reduction
of –9.5 points (SD = 8.9), while those who *Worsened* (n = 42) showed a mean increase of +8.1

points (SD = 8.7). Those with *No change* (n = 311) had a small mean decrease of –2.8 points
(SD = 6.4).

The effect size for the association between MRS change category and Peri-SS score change
was η^2 = 0.277, indicating a large magnitude of responsiveness. These results suggest that
although the Peri-SS captures symptom impact over the past 30 days, it was sensitive to
changes detected within a shorter interval, supporting its use for monitoring symptom
progression over time.

342
Figure 3. Peri-SS score change by MRS change category. Boxplots illustrate changes in Peri-
SS scores from baseline to follow-up, grouped by whether participants' symptom severity—
anchored to changes in the Menopause Rating Scale (MRS)—*improved*, showed *no change*, or
worsened. Negative values indicate symptom improvement. Peri-SS scores decreased
significantly in the "improved" group and increased in the "worsened" group, supporting the
scale's responsiveness to clinically meaningful change.

Criterion validity

The Peri-SS showed strong criterion validity. We found strong associations between the Peri-SS
and both domain-specific and generic health-related quality of life measures. There was a
strong positive correlation between Peri-SS and MRS total scores (Pearson's *r* = 0.88, *p* <
0.001), indicating substantial overlap in symptom severity ratings across the two instruments. A
supplementary analysis using Cohen's Kappa (**Supplementary material E** and S-Table 1) also
showed moderate agreement between MRS and Peri-SS severity categories (κ = .58).

Construct validity

The construct validity of the Peri-SS was supported by both convergent and known-group validity analyses. For convergent validity, the Peri-SS scores were moderately and significantly negatively correlated with EQ-5D utility values (Spearman's $\rho = -0.50$, $p < 0.001$), suggesting that greater perimenopause symptom burden was associated with poorer health-related quality of life. For comparison, the MRS demonstrated a slightly stronger correlation with EQ-5D utility values (Spearman's $\rho = -0.58$, $p < 0.001$).

The Peri-SS demonstrated strong known-group validity, with significant differences observed across subgroups expected to vary in symptom burden (Figure 4, Table 4). As hypothesized, symptom scores were significantly different by age group ($\chi^2 = 31.4$, $p < 0.001$), with women aged 45-54 years reporting higher symptom burden than those aged 35-44 (adjusted $p < 0.001$), consistent with the typical onset of perimenopause symptoms in midlife. Scores among women aged 55-64 did not significantly differ from either of the younger age groups after correction for multiple comparisons.

Peri-SS scores varied significantly by menopausal status ($\chi^2 = 71.2$, $p < 0.001$), with perimenopausal women reporting greater symptom burden than both premenopausal and "unsure" participants (adjusted $p < 0.001$ for both comparisons). Postmenopausal women also reported significantly higher scores than premenopausal and unsure groups (adjusted $p < 0.001$), but there was no significant difference between perimenopausal and postmenopausal participants. These findings are in line with expectations that women undergoing or having recently gone through perimenopause would experience more intense symptoms than those not yet in transition or uncertain of their menopausal status.

Finally, contrary to what we hypothesized, Peri-SS scores were significantly higher among women who reported current use of HT compared to non-users ($p = 0.002$). While this might initially appear counterintuitive, additional analysis supports a reverse causation explanation: in a multivariable logistic regression model adjusting for age, education, body mass index (BMI), and race/ethnicity, higher Peri-SS scores were significantly associated with greater likelihood of HT use – each 1-point increase in Peri-SS score was associated with a 2.4% increase in the odds of HT use (OR = 1.024, 95% CI: 1.009–1.039, $p = 0.002$). This suggests that women experiencing more severe symptoms were more likely to use HT. Together, these results strengthen the known-group validity of the Peri-SS by demonstrating its ability to discriminate between clinically relevant subgroups, in line with theoretical expectations and real-world treatment behavior.

For comparison, we also examined known-group differences for the MRS (see Supplementary S-Table 3 and 4). While overall patterns were broadly similar, Peri-SS detected additional group differences not identified by MRS, such as between ages 35-44 years and 55-59 years, and between postmenopause and unsure groups.

Table 4. Descriptive statistics and known-group differences in Peri-SS total scores by age group, menopausal status, and HT use (Kruskal-Wallis Test)

Group Variable	Category	N	Mean (SD)	Median [Min, Max]	Test statistic	p-value
Overall		1255	29.5 (16.8)	27.7 [0, 100]		
Age group	35-44	497	26.3 (16.2)	23.8 [0, 85.2]	$\chi^2 = 31.4$	<0.001
	45-54	691	31.6 (17.0)	30.0 [0, 100]		
	55-59	67	30.9 (16.3)	28.3 [2.1, 80.2]		
Menopausal status	Premenopause	327	24.5 (15.6)	21.9 [0, 84.8]	$\chi^2 = 71.2$	<0.001
	Perimenopause	633	32.6 (16.6)	30.6 [0, 89.6]		
	Postmenopause	119	33.5 (18.8)	30.8 [1.25, 100]		
	Unsure	176	24.8 (15.3)	22.6 [0, 85.2]		
HT use	No	1190	29.1 (16.8)	27.3 [0, 100]	$\chi^2 = 9.67$	0.018
	Yes	65	35.8 (17.1)	33.1 [3.33, 95.0]		

400   ***Figure 4.*** Perimenopause Symptom Scale (Peri-SS) total scores by a) age group (n = 1255), b) menopausal status (n = 1255), and

401   c) current hormone therapy (HT) use (n = 1255). Boxplots show the median, interquartile range, and individual data points (jittered).

402   Scores were significantly higher in the 45-54 year age group (p < .001), in the perimenopause and postmenopause group (p < .001),

403   and in women using HT (p = .002).

Discussion

Principal findings

Our findings provide robust evidence that the Peri-SS, a new digital self-assessment instrument designed to measure multidimensional symptom burden in perimenopause, meets key psychometric standards. Below, we summarize the key measurement properties demonstrated through this validation study.

Structurally, the Peri-SS was confirmed to have a four-factor model (vasomotor, psychological, sexual, and physical/somatic symptoms), supported by confirmatory factor analysis with good model fit (CFI = 0.932, RMSEA = 0.069). Internal consistency was excellent, with a total Cronbach's alpha of 0.90, and subscale alphas ranging from 0.73 to 0.86. On average, women in perimenopause reported moderate symptom burden as measured by Peri-SS, with some variation by age group. At the domain level, psychological symptoms were the most commonly reported, while sexual health symptoms were the least reported. Vasomotor and sexual symptoms showed a trend of increasing severity with age, whereas psychological symptoms appeared to peak during midlife and decline slightly in older age groups. Physical and somatic symptoms remained relatively stable across age groups.

The Peri-SS also demonstrated high test-retest reliability over a 7-14 day interval (ICC = 0.82), suggesting stable symptom reporting over short timeframes. In addition, the Peri-SS demonstrated large responsiveness to change (η^2 = 0.277), with symptom scores decreasing significantly among those who improved on an external anchor (MRS) and increasing among those whose symptoms worsened. In terms of criterion validity, the scale correlated strongly with the MRS (r = 0.88), supporting its ability to capture perimenopause symptom burden consistent with established instruments. Finally, construct validity was established through both convergent and known-group approaches: the Peri-SS was moderately negatively correlated with EQ-5D utility scores (ρ = –0.50), indicating that higher symptom burden was associated with lower quality of life, and was significantly associated with differences in age, menopausal status, and HT use.

Taken together, these findings provide strong initial support for the Peri-SS as a reliable, valid, and responsive instrument for assessing perimenopause symptoms among U.S. women aged 35-59 years in clinical, research and digital health settings.

Comparison with existing instruments

To better understand the contribution of the Peri-SS, it is important to consider how it compares to existing symptom measurement tools for menopause and midlife women's health.

A wide range of instruments exist to measure menopause symptoms, but most were
developed for postmenopausal populations or general midlife health rather than the
evolving needs of women in perimenopause [35]. There remains a significant knowledge
gap in effective strategies to help women prepare for and recognize the start of
perimenopause [12]. While existing instruments have proved useful, they share key
limitations ([17–20,24], validation was often done among postmenopausal women, early
perimenopause symptoms were underrepresented, reliance on paper-based
administration, and psychometric evaluation did not consistently adhere to best-practice
psychometric standards, such as the COSMIN framework [25].

Among these, the MRS is the most widely validated tool and serves as the primary comparator
for this study [17]. The Peri-SS shares structural similarities with MRS—both include
psychological, somatic, and urogenital symptoms—and demonstrated strong convergent validity
in our analysis. Like the MRS, the Peri-SS offers clearly defined domains supported by factor-
analytic results, allowing for interpretable and actionable domain-specific scores. However, the
Peri-SS appears to offer improved scope and relevance for perimenopausal populations. This is
because several commonly reported symptoms such as low sex drive, weight changes, and
digestive issues are underrepresented in MRS. Conversely, some of its somatic items (e.g.,
heart palpitations) may be more characteristic of later menopause stages and thus less reliable
in perimenopause populations [36].
Overall, Peri-SS may more accurately capture the earlier, fluctuating symptomatology typical of
perimenopause, where symptom clustering may differ from that observed postmenopause [10].

More recently, the MenoScale [24] represents an important step forward in menopause
symptom measurement in digital contexts. Developed using data from digital health users,
it captures a broad range of symptoms across 21 items, with particular emphasis on
somatic symptoms (10 items). Its factor structure—comprising vasomotor, sexual,
psychological, and somatic domains—is similar to that of the Peri-SS. However,
MenoScale was validated primarily among older women (average age over 50) and a
demographically homogeneous sample (95% White), which may limit its applicability to
younger or more diverse perimenopausal populations. In contrast, the Peri-SS was
developed and validated in a younger (mean age 46), more demographically diverse
cohort (73% White), and achieved robust psychometric performance across all domains
while balancing symptom comprehensiveness and survey burden.

Other legacy instruments further highlight the gap Peri-SS fills. The Greene Climacteric
Scale (GCS) [19,20], while widely used, was developed in the 1970s and lacks sexual
function nuances (only 1 item) and did not define explicit thresholds to categorize overall
symptom severity. The Women's Health Questionnaire (WHQ) [29] includes broader
symptoms but is lengthy with 36 items and was validated in a very small sample (n=48).
The MENQOL [18], though menopause-specific, did not define symptom domains thus was
not able to track women's domain-specific changes in symptoms. The Day-to-Day Impact
of Vaginal Aging (DIVA) [37] was developed for treatment-seeking postmenopausal women
and focuses narrowly on vaginal symptoms. Collectively, these tools reflect historical

strengths but are limited in their specificity, psychometric rigor, and inclusiveness of the
perimenopause experience.

In summary, the Peri-SS is a concise and psychometrically robust measure of
perimenopause symptom burden. By addressing the limitations of earlier instruments and
offering digital deployment, it provides a modern and actionable tool for both research and
clinical practice.

Strengths and implications

In addition to strong psychometric performance, this study offers several methodological and
practical strengths that support the real-world relevance and utility of the Peri-SS.

The validation process followed the COSMIN framework [25], ensuring rigorous evaluation across
reliability, construct validity, criterion validity, and responsiveness to change. The longitudinal
design enabled robust assessment of test-retest reliability and sensitivity to change—two
properties often underreported in prior instruments. The sample size was adequately powered
to detect differences across known groups, and the inclusion of women aged 35-59 years
includes women across the reproductive stages which is broader than typically captured. While
the sample was drawn from a single geographic and linguistic context (U.S.-based, English-
speaking), it was younger and more racially/ethnically diverse than many existing studies,
supporting the scale's relevance to earlier and more heterogeneous perimenopause
experiences.

Practically, the Peri-SS provides a digital-first, user-friendly tool that enables scalable and
nuanced symptom tracking across key domains of the perimenopause experience. Its four-
domain structure is clinically interpretable and supports a range of clinical and research
priorities in the care of midlife women [12]. These include evaluating and managing symptom
burden, assessing the effectiveness of non-hormonal and lifestyle-based interventions,
monitoring the progression of symptoms such as sleep disruption and fatigue, and identifying
symptom phenotypes to inform tailored support and treatment pathways [1]).

For digital health tools, the Peri-SS enables dynamic segmentation of users based on symptom
burden and supports personalization of educational content, thereby enhancing user
engagement and perceived relevance [38]. The brevity of the 16-item instrument ensures it is
practical for repeated use without overwhelming users, making it especially suitable for mobile
platforms and longitudinal tracking.

Furthermore, the Peri-SS has potential value beyond individual symptom monitoring. It could be
deployed in workplace health initiatives or employee wellbeing platforms to assess the impact of
perimenopause symptoms on productivity, absenteeism, or quality of life [39]. This opens avenues
for evaluating support interventions (e.g., policy changes, access to symptom management
tools), and normalizing perimenopause and menopause-related health as a part of occupational
health discourse [40].

By offering a validated, perimenopause-specific, and digitally deployable scale, the Peri-SS
addresses a critical gap in measurement and enables more responsive, personalized, and
evidence-based approaches to supporting women through perimenopause.

Limitations and future directions

Despite its strengths, we acknowledge several limitations which would inform future
research and application. First, our sample was recruited online. Although this mode of
recruitment aligns with the intended context of use (digital health platforms), it
nevertheless limits generalizability to individuals with lower digital literacy or limited
internet access. To enhance the broader applicability of the Peri-SS, future validation
studies should include populations with more diverse geographic, educational, and digital
literacy backgrounds.

Second, in this initial validation study, we relied on self-reported rather than clinically
confirmed menopause stages. However, we observed clear known-group differences in
symptom burden across menopause stages, supported by both the Peri-SS and MRS
scores, which adds confidence to the validity of the self-reported categories (see
Supplementary S-Tables 3 and S-Table 4). Furthermore, we conducted additional
analysis on participants' medical-seeking behaviors and found that 45.0% of participants
who self-identified as perimenopausal had consulted a medical professional in the past 12
months about their symptoms. Among these, 56.8% received a formal confirmation. This
finding further supports that the self-reported menopause stage in our sample has clinical
grounding. To mitigate this limitation, further research is needed to validate Peri-SS in
populations with clinically confirmed menopausal status.

Third, while the Peri-SS includes a broad range of symptoms reported by women in
perimenopause, it does not capture every possible experience. Some common symptoms
during perimenopause, such as headache, fatigue, and brain fog [10] were excluded due to poor
psychometric performance (high correlation with other items or high cross domain factor
loadings) during development. Additionally, menstrual cycle changes were not included as a
scored symptom domain in the Peri-SS, despite being a hallmark of perimenopause [5]. During
scale development, we treated menstrual cycle irregularity as a contextual indicator for
classifying women as likely perimenopausal, rather than as a symptom in its own right.
However, we acknowledge that the extent and pattern of menstrual changes vary widely and
may themselves be perceived by women as a meaningful part of the symptom experience. We
plan to consider incorporating bleeding pattern changes as either a standalone domain or a
moderating factor in future iterations, especially as more nuanced cycle-related data become
available through digital tracking tools.

Finally, symptom experiences may vary across cultural and linguistic contexts. The current
version of the Peri-SS was developed and validated in English among U.S.-based participants,
and certain symptom concepts may not translate seamlessly. To address this, we are
conducting qualitative work to adapt the scale for use in multiple languages, starting with

German, French, Spanish, Brazilian Portuguese, and Italian, to ensure cultural relevance and
cross-contextual validity.

557 Conclusion

This study provides strong initial validation of the Peri-SS, a concise and digitally deployable
self-assessment tool designed to measure the multidimensional symptom burden experienced
during perimenopause. The scale demonstrated excellent psychometric properties across
structural validity, internal consistency, test-retest reliability, construct validity, and
responsiveness to change. Compared to legacy instruments, the Peri-SS is more inclusive of
perimenopause-specific symptoms and better aligned with modern psychometric standards. Its
digital-first format and clinical interpretability position it as a promising tool for research, clinical
care, and digital health applications.

References

References

1. Delamater L, Santoro N. Management of the Perimenopause. Clin Obstet Gynecol. 2018
Sep;61(3):419–32.

2. Burger HG, Dudley EC, Robertson DM, Dennerstein L. Hormonal changes in the
menopause transition. Recent Prog Horm Res. 2002;57:257–75.

3. Fang Y, Liu F, Zhang X, Chen L, Liu Y, Yang L, et al. Mapping global prevalence of
menopausal symptoms among middle-aged women: a systematic review and meta-
analysis. BMC Public Health. 2024 Jul 2;24(1):1767.

4. Li R xia, Ma M, Xiao X rong, Xu Y, Chen X ying, Li B. Perimenopausal syndrome and
mood disorders in perimenopause: prevalence, severity, relationships, and risk factors.
Medicine (Baltimore). 2016 Aug;95(32):e4466.

5. Santoro N. Perimenopause: From Research to Practice. J Womens Health 2002. 2016
Apr;25(4):332–9.

6. Coronado PJ, Fasero M, Mendoza N, González SP, Sánchez-Méndez S, Presa J, et al.
Quality of life at peri- and postmenopause: analysis of the results from app 'Mi
Menopausia'. Maturitas. 2025 Jun;197:108267.

7. Li S, Holm K, Gulanick M, Lanuza D. Perimenopause and the Quality of Life. Clin Nurs
Res. 2000 Feb 1;9(1):6–23.

8. Santoro N, Roeca C, Peters BA, Neal-Perry G. The Menopause Transition: Signs,
Symptoms, and Management Options. J Clin Endocrinol Metab. 2021 Jan 1;106(1):1–15.

9. Tepper PG, Brooks MM, Randolph JF, Crawford SL, El Khoudary SR, Gold EB, et al.
Characterizing the Trajectories of Vasomotor Symptoms Across the Menopausal
Transition. Menopause N Y N. 2016 Oct;23(10):1067–74.

10. Aras SG, Grant AD, Konhilas JP. Clustering of > 145,000 symptom logs reveals distinct
pre, peri, and menopausal phenotypes. Sci Rep. 2025 Jan 3;15(1):640.

11. Allen JT, Laks S, Zahler-Miller C, Rungruang BJ, Braun K, Goldstein SR, et al. Needs
assessment of menopause education in United States obstetrics and gynecology
residency training programs. Menopause N Y N. 2023 Oct 1;30(10):1002–5.

12. Nash Z, Christmas M, Gronlund T, Hickey M, Burgin J, Carpenter J, et al. Top ten
menopause research priorities. The Lancet. 2024 Dec 21;404(10471):2535–6.

13. Harper JC, Phillips S, Biswakarma R, Yasmin E, Saridogan E, Radhakrishnan S, et al. An
online survey of perimenopausal women to determine their attitudes and knowledge of the
menopause. Womens Health Lond Engl. 2022;18:17455057221106890.

14. O'Reilly K, McDermid F, McInnes S, Peters K. An exploration of women's knowledge and
experience of perimenopause and menopause: An integrative literature review. J Clin
Nurs. 2023 Aug;32(15–16):4528–40.

15.   Barber K, Charles A. Barriers to Accessing Effective Treatment and Support for
603   Menopausal Symptoms: A Qualitative Study Capturing the Behaviours, Beliefs and
604   Experiences of Key Stakeholders. Patient Prefer Adherence. 2023;17:2971–80.

605   16.   Cunningham AC, Hewings-Martin Y, Wickham AP, Prentice C, Payne JL, Zhaunova L.
606   Perimenopause symptoms, severity, and healthcare seeking in women in the US. Npj
607   Womens Health. 2025 Feb 25;3(1):12.

608   17.   Heinemann K, Ruebig A, Potthoff P, Schneider HP, Strelow F, Heinemann LA, et al. The
609   Menopause Rating Scale (MRS) scale: A methodological review. Health Qual Life
610   Outcomes. 2004 Sep 2;2(1):45.

611   18.   Hilditch JR, Lewis J, Peter A, van Maris B, Ross A, Franssen E, et al. A menopause-
612   specific quality of life questionnaire: development and psychometric properties. Maturitas.
613   1996 Jul;24(3):161–75.

614   19.   Greene JG. A factor analytic study of climacteric symptoms. J Psychosom Res.
615   1976;20(5):425–30.

616   20.   Greene JG. Constructing a standard climacteric scale. Maturitas. 1998 May 20;29(1):25–
617   31.

618   21.   Andersen NJ, Parker JL, Pettigrew S, Bitner D. Validation of the Menopause Transition
619   Scale (MTS). Menopause N Y N. 2022 Jul 1;29(7):868–76.

620   22.   Levy J, Romo-Avilés N. "A good little tool to get to know yourself a bit better": a qualitative
621   study on users' experiences of app-supported menstrual tracking in Europe. BMC Public
622   Health. 2019 Sep 3;19(1):1213.

623   23.   Nascimento IJB do, Abdulazeem HM, Weerasekara I, Marquez J, Vasanthan LT, Deeken
624   G, et al. Transforming women's health, empowerment, and gender equality with digital
625   health: evidence-based policy and practice. Lancet Digit Health [Internet]. 2025 Jun 1
626   [cited 2025 Jul 16];7(6). Available from:
627   https://www.thelancet.com/journals/landig/article/PIIS2589-7500(25)00022-6/fulltext

628   24.   Kim C, Marples L, Platts A, Bermingham KM, Amati F, Hamoda H, et al. MenoScale: A
629   novel digital tool to measure menopause symptoms and subjective quality of life -
630   validation, preliminary insights on the menopausal experience and association with diet
631   quality. Post Reprod Health. 2025 Apr 22;20533691251335807.

632   25.   Gagnier JJ, de Arruda GT, Terwee CB, Mokkink LB, Consensus group. COSMIN reporting
633   guideline for studies on measurement properties of patient-reported outcome measures:
634   version 2.0. Qual Life Res Int J Qual Life Asp Treat Care Rehabil. 2025 Mar 28;

635   26.   von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP.
636   Strengthening the reporting of observational studies in epidemiology (STROBE) statement:
637   guidelines for reporting observational studies. BMJ. 2007 Oct 20;335(7624):806–8.

638   27.   Morgado FFR, Meireles JFF, Neves CM, Amaral ACS, Ferreira MEC. Scale development:
639   ten main limitations and recommendations to improve future research practices. Psicol
640   Reflex E Crítica Rev Semest Dep Psicol UFRGS. 2017 Jan 25;30:3.

28.   Heinemann LA, Potthoff P, Schneider HP. International versions of the Menopause Rating
Scale (MRS).

29.   Hunter MS. The Women's Health Questionnaire (WHQ): Frequently Asked Questions
(FAQ). Health Qual Life Outcomes. 2003 Sep 10;1:41.

30.   Rosseel Y, Jorgensen TD, Wilde LD, Oberski D, Byrnes J, Vanbrabant L, et al. lavaan:
Latent Variable Analysis [Internet]. 2024 [cited 2025 Jun 27]. Available from: https://cran.r-
project.org/web/packages/lavaan/index.html

31.   Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research
[Internet]. 2025 [cited 2025 Jun 27]. Available from: https://cran.r-
project.org/web/packages/psych/index.html

32.   Jacob Cohen. Statistical Power Analysis for the Behavioral Sciences. 2nd Edition. New
York: Routledge; 1988.

33.   Feng YS, Kohlmann T, Janssen MF, Buchholz I. Psychometric properties of the EQ-5D-5L:
a systematic review of the literature. Qual Life Res. 2021 Mar 1;30(3):647–73.

34.   Pickard AS, Law EH, Jiang R, Pullenayegum E, Shaw JW, Xie F, et al. United States
Valuation of EQ-5D-5L Health States Using an International Protocol. Value Health J Int
Soc Pharmacoeconomics Outcomes Res. 2019 Aug;22(8):931–41.

35.   Jenabi E, Shobeiri F, Hazavehei SMM, Roshanaei G. Assessment of Questionnaire
Measuring Quality of Life in Menopausal Women: A Systematic Review. Oman Med J.
2015 May;30(3):151–6.

36.   Freeman EW, Sammel MD, Lin H, Gracia CR, Pien GW, Nelson DB, et al. Symptoms
associated with menopausal transition and reproductive hormones in midlife women.
Obstet Gynecol. 2007 Aug;110(2 Pt 1):230–40.

37.   Huang AJ, Gregorich SE, Kuppermann M, Nakagawa S, Van Den Eeden SK, Brown JS, et
al. The Day-to-Day Impact of Vaginal Aging Questionnaire: A Multidimensional Measure of
the Impact of Vaginal Symptoms on Functioning and Well-being in Postmenopausal
Women. Menopause N Y N. 2015 Feb;22(2):144–54.

38.   Murray E, Hekler EB, Andersson G, Collins LM, Doherty A, Hollis C, et al. Evaluating
digital health interventions: key questions and approaches. Am J Prev Med. 2016
Nov;51(5):843–51.

39.   Schei TS, Abernethy K. The impact of a specialist-led digital health application on
menopause symptoms in the workplace: A single-arm, longitudinal evaluation. Maturitas.
2024 Sep;187:108005.

40.   DePree B, Houghton K, Shiozawa A, Esterberg E, King DD, Kim J, et al. Treatment and
resource utilization for menopausal symptoms in the United States: a retrospective review
of real-world evidence from US electronic health records. Menopause N Y N. 2023 Jan
1;30(1):70–9.

List of abbreviations

BMI – Body Mass Index

CFI – Comparative Fit Index

CFA – Confirmatory Factor Analysis

COSMIN – COnsensus-based Standards for the selection of health Measurement INstruments

DIVA – Day-to-Day Impact of Vaginal Aging Questionnaire

EFA – Exploratory Factor Analysis

EQ-5D-5L – EuroQol five-dimensional questionnaire (five-level version)

FIML – Full-Information Maximum Likelihood

HT – Hormone Therapy

ICC – Intraclass Correlation Coefficient

IRB – Institutional Review Board

MenoScale – Menopause Scale

MENQOL – Menopause-specific Quality of Life Questionnaire

MRS – Menopause Rating Scale

MTS – Menopause Transition Scale

Peri-SS – Perimenopause Symptom Scale

PROMs – Patient-Reported Outcome Measures

RMSEA – Root Mean Square Error of Approximation

STROBE – Strengthening the Reporting of Observational studies in Epidemiology

WHQ – Women's Health Questionnaire

Acknowledgements

We are grateful for Flo users and other women that participated in our studies.
We would also like to thank Dzmitry Kuzmiankou (backend engineer) and Sasha Pautaran
(project manager) for their valuable contributions to the technical implementation of the Peri-SS.

Author contributions

Y.X. led the conceptualization, study design, and data analysis. She also drafted the study
protocol and manuscript. I.I. contributed to the literature review, critically appraised existing
menopause measurement scales, and supported the development of the Peri-SS. C.P. co-
developed the study protocol, assisted with the ethics application, and contributed to the
literature review. Y.H-M supported refinement of the study protocol and contributed to the
development of the Peri-SS instrument. A.C. supported the design and implementation of the
survey. C.S. and S.F. reviewed the Peri-SS scale and provided critical revisions to the
manuscript. L.Z. supervised the overall study design and implementation.

All authors reviewed and approved the final manuscript.

Statements and declarations

Ethical considerations

This study was conducted in two phases, both of which received ethical approval from an
independent institutional review board. The initial scale development phase, including item
generation and exploratory factor analysis, was approved by the WCB Institutional Review
Board (IRB No. 20244604). The subsequent validation study, involving confirmatory factor
analysis and psychometric testing, was approved under a separate protocol (IRB No.
20250554). All participants provided informed consent electronically prior to participation and
were informed of their right to withdraw at any time without penalty. We compensated
participants at a fair hourly rate ($12-15 USD/hour) in line with ethical guidelines for online
research platforms.

Consent to participate

All participants provided electronic informed consent prior to participation, in accordance with
the approved IRB protocols. The consent form, administered via SurveyMonkey, described the
purpose of the study, eligibility criteria, voluntary nature of participation, data privacy
safeguards, and estimated time commitment. Participants indicated their agreement by
selecting a consent checkbox before proceeding to the survey. No participants that did not
consent was included in the study.

737 ## Consent for publication
738 Not applicable. All data were collected anonymously or de-identified prior to analysis, and no
739 individual-level or identifiable information is included in this manuscript.

740 ## Declaration of conflicting interest

743
744 ## Funding statement

745 The study was funded by Flo Health Inc.

746
Data availability
Aggregated data analyses and deidentified individual-level data may be shared upon
reasonable request for the purpose of replication, secondary analysis, or further methodological
development, pending permission from Flo Health UK Inc.

Licensing information
The Perimenopause Symptom Scale is licensed under a Creative Commons Attribution-
NonCommercial 4.0 International License (CC BY-NC 4.0). This means it is freely available for
non-commercial research and academic use, provided appropriate credit is given.
Written permission is required from Flo Health UK Limited for any commercial use, including
incorporation into digital products, health apps, or services offered by commercial entities.