Post-discharge Oral Home Antibiotics Versus No Oral Home Antibiotics in Complicated

Pediatric Appendicitis: A Systematic Review and Meta-analysis

Running head: Oral Discharge Antibiotics for Complicated PAA

Authors: **Javier Arredondo Montero** (MD, PhD) [1] and **María Rico-Jiménez** (MD) [2]

Affiliations:

(1) Pediatric Surgery Department, Complejo Asistencial Universitario de León, León, Spain.

(2) Pediatric Surgery Department, Hospital Universitario Niño Jesús, Madrid, Spain.

Corresponding author:

Javier Arredondo Montero, MD, PhD

      Address: Department of Pediatric Surgery, Complejo Asistencial Universitario de León.

      c/Altos de Nava s/n, 24008 León, Castilla y León, Spain

      Email: jarredondo@saludcastillayleon.es/ javier.montero.arredondo@gmail.com

      Phone: +34 987 23 74 00

ORCID ID:

      Javier Arredondo Montero: 0000-0002-0943-6367

CRediT authorship contribution statement:

JAM: Conceptualization and study design; literature search and selection; data curation and extraction; formal analysis; investigation; methodology; project administration; resources; validation; visualization; writing – original draft; writing – review and editing.

MRJ: Literature search and selection; data curation and extraction; writing, review, and editing.

CONFLICTS OF INTEREST:

The authors declare that they have no conflict of interest.

FINANCIAL STATEMENT/FUNDING:

This review did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors, and none of the authors has external funding to declare.

ETHICAL APPROVAL:

This study did not involve the participation of human or animal subjects, and therefore, IRB approval was not sought.

STATEMENT OF AVAILABILITY OF THE DATA USED DURING THE SYSTEMATIC REVIEW:

All data used for the meta-analytical models are available in the accompanying supplementary dataset file.

Post-discharge Oral Antibiotics Versus No Oral Home Antibiotics in Complicated Pediatric Appendicitis: A Systematic Review and Meta-analysis

ABSTRACT

Background: The use of oral home antibiotics (OHA) after discharge in children undergoing surgery for complicated acute appendicitis (CAA) remains controversial. This systematic review and meta-analysis aimed to evaluate whether OHA reduces the risk of infectious complications or readmissions compared to patients discharged without antibiotics (NHA).

Methods: This systematic review was prospectively registered in PROSPERO (CRD420251049919). We searched PubMed, Web of Science, Scopus, Ovid, and Cochrane CENTRAL from inception to March 2025. Two independent reviewers screened the studies, extracted the data, and assessed the methodological quality using the ROBINS-I tool. Four random-effects meta-analyses and four leave-one-out analyses were conducted for intra-abdominal abscesses (IAA), surgical site infections (SSI), organ/space infections (OSI), and hospital readmissions (RA). Three exploratory random-effects meta-regression models were performed for RA.

Results: Fourteen studies comprising 26,174 pediatric patients with CAA were included. Meta-analyses showed no significant differences between intervention (IG) and comparator (CG) groups for IAA (RR 0.97; 95% CI: 0.38–2.47), SSI (RR 0.89; 95% CI: 0.58–1.38), OSI (RR 1.19; 95% CI: 0.73–1.93). or RA (RR 1.02; 95% CI: 0.73–1.41), with no evidence of publication bias. A sensitivity analysis restricted to studies reporting crude data for OHA vs. NHA revealed a significantly lower RA risk in NHA patients (RR 0.75; 95% CI: 0.62–0.90, P < 0.01). Meta-

regression exploring differences in predischarge total leukocyte counts showed a non-significant trend toward increased RA in patients receiving OHA.

Conclusions: OHA after discharge do not appear to reduce the risk of postoperative complications in children treated surgically for CAA. Given the lack of consistent benefit and potential for unnecessary harm, routine use of post-discharge OHA is not supported. A risk-adapted, stewardship-oriented approach is warranted.

Funding: None.

Registration: PROSPERO (CRD420251049919).

Keywords: *complicated acute appendicitis;pediatric;oral antibiotics;discharge; surgical site infection;readmission;stewardship*

Post-discharge Oral Home Antibiotics Versus No Oral Home Antibiotics in Complicated Pediatric Appendicitis: A Systematic Review and Meta-analysis

Introduction

Acute appendicitis (AA) is the most common surgical emergency in the pediatric population, with an estimated annual incidence of 9.68 per 10,000 children [1]. While most cases present as non-complicated appendicitis (NCAA), approximately 15–25% present as complicated acute appendicitis (CAA) [2,3], including gangrenous or perforated forms, which are associated with higher postoperative morbidity. Patients with CAA are more susceptible to postoperative complications and typically require prolonged antibiotic therapy and extended hospitalization, contributing to a substantial burden on healthcare systems and resource utilization [4].

Despite consensus on the need for intravenous antibiotic therapy during hospitalization for CAA, and a growing trend toward institutional protocols promoting shorter intravenous courses or combined intravenous–oral regimens—with outcomes shown to be non-inferior to traditional approaches [5,6]—substantial variation persists in post-discharge management, particularly regarding the use of oral antibiotics. Current guidelines provide limited or no specific recommendations on the continuation of antibiotics after discharge, resulting in heterogeneous institutional practices and variability among surgeons. Several studies have reported marked inter-professional variability in this aspect of care, underscoring both the lack of standardization and the ongoing uncertainty regarding its clinical benefit [7].

The rationale for prescribing oral antibiotics after hospital discharge in cases of CAA is primarily based on the assumption that continued antimicrobial coverage may reduce the risk of delayed infectious complications, such as surgical site infections (SSI), intra-abdominal abscesses (IAA),

organ space infections (OSI), or unplanned readmissions (RA). Concerns over early hospital discharge and residual contamination in perforated or gangrenous appendicitis have traditionally justified this strategy. Oral home antibiotics (OHA) are often perceived as a low-cost, accessible intervention that may provide a safety buffer during the vulnerable post-discharge period.

However, prolonged antibiotic exposure is not without risks. The unnecessary use of antimicrobials contributes to adverse drug reactions, disruption of the gut microbiota, an increased risk of *Clostridioides difficile* infection, and a broader public health threat from antimicrobial resistance [8]. Given their developing immune and microbial systems, these concerns are particularly relevant in pediatric populations. Notably, emerging evidence suggests that post-discharge oral antibiotic therapy may offer limited or no benefit in reducing complications—and may even be associated with overtreatment and an increased risk of adverse events.

Given the widespread but inconsistent use of post-discharge OHA in children with surgically treated CAA—and the growing body of evidence questioning their effectiveness—there is a pressing need to synthesize the available data to inform clinical decision-making. The absence of clear guideline recommendations and the potential for both overuse and harm further underscore this gap. Therefore, the objective of this systematic review is to evaluate the impact of oral antibiotic therapy after hospital discharge on outcomes in pediatric patients with surgically treated CAA, with a specific focus on infectious complications (including SSI, OSI, and IAA) and readmissions.

Methods

Literature Search and Selection

We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [10] and the Cochrane Handbook for Systematic Reviews of Interventions (version 6.5) [11]. The PRISMA checklist is provided in Supplementary File 1. This review was prospectively registered in the International Prospective Register of Systematic Reviews (PROSPERO ID: CRD420251049919).

Eligible studies were identified by searching the primary existing medical bibliography databases (PubMed, Web of Science, Scopus, Ovid, and Cochrane Central). Supplementary File 2 shows the detailed search strategy for each bibliographic database. The search was last executed on 11.05.2025.

JAM and MRJ selected articles using the COVIDENCE ® tool. The search results were imported into the platform, and both authors independently screened the articles. Disagreements were resolved by consensus. Supplementary File 3 shows the inclusion and exclusion criteria.

Quality Assessment

The Cochrane Risk of Bias 2 (RoB 2) tool was selected to evaluate the methodological quality and risk of bias of randomized controlled trials (RCTs) [12]. In contrast, the Risk of Bias in Non-randomized Studies - of Interventions (ROBINS-I) tool was chosen to assess the methodological quality and risk of bias of non-randomized studies of interventions (NRSIs) [13].

Data Extraction and Synthesis

Two reviewers (JAM, MRJ) independently extracted data from the included studies using a standardized and piloted form. Extracted variables included: first author, year of publication, country of study, study design, sample size, age, sex distribution, definitions of the intervention

and comparator groups, discharge criteria, total leukocyte count (TLC) admission values, TLC predischarge values, antibiotics employed (type, duration, and route), events by group including SSI, IAA, OSI, RA, reoperations (RO), and visits to the Emergency Department (EDV), and p-values for between-group comparisons. Disagreements during extraction were resolved by discussion or consultation with a third reviewer. Study authors were contacted for clarification when critical data were missing or unclear. In studies where raw data were available disaggregated by treatment group — OHA versus no home antibiotics (NHA) — these values were directly used; when not available, they were inferentially estimated, and if this was not possible, events were reported according to treatment protocol groups (e.g., "old protocol" versus "new protocol"), explicitly stating the lack of specific data for OHA and NHA. Unit standardization was not required.

Meta-analysis

Four primary random-effects meta-analyses (MAs) were conducted using the restricted maximum likelihood (REML) estimator for both the pooled effect size and the between-study variance (τ^2), following the Cochrane Handbook recommendations (version 6.5) [11, 14–16]. For each outcome, the choice of confidence interval around the pooled effect was based on the number of included studies and the τ^2. When τ^2 was close to zero or only two studies were available, conventional Wald-type intervals were deemed acceptable. When more than two studies were included and τ^2 was greater than zero, confidence intervals were computed using both the Hartung–Knapp–Sidik–Jonkman (HKSJ) [15] and the truncated modified Knapp–Hartung (mKH) methods [16]. The approach providing the most robust and interpretable inference was selected. Exploratory leave-one-out sensitivity analyses were conducted for each one of the primary meta-analyses to assess the influence of individual studies on overall

estimates. Where sufficient studies (≥10) were available, univariable random-effects meta-regression analyses were performed to explore potential sources of heterogeneity.

Publication Bias and Small Study Effects

In meta-analyses including at least ten studies, publication bias and small-study effects were explored through visual inspection of contour-enhanced funnel plots, which included significance contours at conventional thresholds ($p < 0.01$, $p < 0.05$, $p < 0.10$, and $p > 0.10$), as well as through the application of Egger's and Begg's statistical tests [17,18]. To further investigate asymmetry and between-study heterogeneity, Galbraith radial plots and L'Abbé plots were also generated. In cases where funnel plot asymmetry was observed, the trim-and-fill method was applied to estimate the potential impact on the pooled effect size [19].

GRADE Assessment

The certainty of evidence for each meta-analytic outcome was evaluated using the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) framework. The assessment considered five core domains: risk of bias, inconsistency, indirectness, imprecision, and publication bias. For each outcome, certainty was rated as high, moderate, low, or very low based on the quality of included studies, the magnitude of heterogeneity (I^2 and τ^2), the width of confidence intervals, and the presence of small-study effects.

Results

Summary of the Included Studies

The search yielded 331 articles (Scopus, n = 93; PubMed, n = 57; Web of Science, n = 106; Ovid, n = 12; Cochrane CENTRAL, n = 63). After removing 150 duplicates, 181 records remained. Of these, 167 were excluded based on the predefined inclusion and exclusion criteria. Ultimately, this systematic review included 14 studies encompassing data from 26,174 patients (15,488 males, 10,686 females) [20-33]. Minimal discrepancies (<0.5%) were identified between the number of patients included per group (IC and CG), the number of patients per gender (male/female), and the total number of patients reported in the review, attributable to the lack of explicit reporting in some studies. Concerning study design, no randomized controlled trials were identified; all included studies were observational. The flowchart of the search and selection process is shown in Figure 1.

Sociodemographic and Clinical Characteristics

All studies were conducted between 2015 and 2025 in pediatric populations [20-33]. Thirteen were from the United States [20-29,31-33], and one was from Japan [30]. One prospective cohort study was identified [20], along with eight retrospective cohort studies [22–27, 29, 31], and five studies based on national databases, including the NSQIP Pediatric Database [21, 28, 32, 33] and the Diagnosis Procedure Combination Database [30].

All included studies focused on patients with complicated acute appendicitis (CAA), although moderate variability was observed in the definitions applied. Several authors defined CAA as perforated acute appendicitis (PAA) identified intraoperatively as a "visible hole in the appendix" [20-26,28,29,31-33], or as the presence of free fecalith in the abdominal cavity [20,21,23,24,28,31-33]. Others considered CAA to include the presence of intra-abdominal abscess [21,23,28,31-33], diffuse fibrinopurulent exudate within the peritoneal cavity [21,33],

fibrinopurulent contamination extending beyond the right lower quadrant and pelvis [28], or involving more than two quadrants, including the pelvis [32]. Additional criteria used were the presence of diffuse peritonitis [23], intra-abdominal stool [24], and endoluminal content leaking outside the bowel during surgery [27]. Some authors also included gangrenous appendicitis (GA) under the CAA definition [22], or based the diagnosis on histopathological confirmation of PAA [29]. One study did not specify the criteria used to define CAA [30]. It is worth noting that many studies applied mixed criteria for defining CAA, and only a few explicitly referred to histopathological findings, despite histopathology being considered the diagnostic gold standard for acute appendicitis.

Regarding surgical technique, the information reported across studies is highly limited and heterogeneous. Some authors mentioned at some point in the manuscript that a laparoscopic approach was used [20,23], but without specifying whether any conversions occurred or if 100% of patients underwent laparoscopy. Other studies included only laparoscopic procedures and explicitly excluded open appendectomies or conversions [24,31,33]. Authors using national databases adjusted odds ratios for the "laparoscopic technique" variable and reported the proportion of patients undergoing laparoscopy, with significant differences between groups (p < 0.01) [21]. In one study, propensity score matching was used to balance the number of open and laparoscopic procedures between groups, while retaining both techniques in the analysis [30]. In contrast, some authors provided no details on surgical technique [22,28,32], while others included both open and laparoscopic appendectomies indiscriminately [25,27]. One study explicitly stated that the intraoperative technique was not standardized [26]. Finally, another study specified that all procedures were laparoscopic, although one conversion occurred in the NHA group and was not excluded [29].

Post-Discharge Oral Home Antibiotics vs. No-Home Antibiotics

The intervention group (IG) was defined as the NHA group or as those managed under a protocol specifically implemented to reduce or eliminate post-discharge antibiotic use. In most studies, this corresponded to a "new protocol" cohort explicitly designed to minimize or avoid OHA. In some cases, raw data were reported specifically for NHA patients; in others, outcomes reflected the entire cohort managed under the revised protocol, even if not all patients were strictly NHA. The comparator group (CG) included patients who either received OHA at discharge or were managed according to standard protocols that did not restrict post-discharge antibiotic use. However, these protocols did not always guarantee that all patients in the CG actually received OHA. We clearly specified in the manuscript whether raw data were available or not for each study, and analyses were conducted accordingly.

A total of six studies directly compared patients discharged with OHA versus those discharged without them (NHA) [21,27,28–30,33], while eight studies compared outcomes across two distinct clinical protocols—typically a pre-implementation (baseline) phase versus a post-implementation phase [20,22–26,31,32]. In most cases, the pre-protocol group reflected routine or institutionally guided administration of OHA, whereas the post-protocol group incorporated a clinical modification aimed at reducing OHA use. Although these constituted targeted interventions, they were primarily implemented as part of institutional practice changes rather than formal research initiatives, and most studies retained an observational retrospective design.

The specific characteristics of these institutional protocols, including their rationale and implications for antibiotic stewardship and discharge practices, are detailed in Table 1. Notably, one study (Ferguson [24]) inverted this structure: the authors standardized the routine

prescription of OHA and designated that cohort as the intervention group. For the purposes of this review, we reclassified this group as the comparator (CG) to maintain consistency in operational definitions across studies. These criteria and their application are detailed in Table 1.

Table 1 summarizes the data extracted from the fourteen studies that evaluated the role of post-discharge OHA in CAA.

Risk of Bias Assessment

In the domain of bias due to confounding, seven studies were rated as having a serious risk of bias [20,21,24,25,27,29,33], while the remaining seven were rated as moderate [22,23,26,28,30,31,32]. None were considered at low risk. One of the main reasons for assigning a serious risk rating was the presence of significant differences in baseline patient characteristics, with systematically sicker patients more likely to receive OHA. For instance, in the study by Anderson et al. [21], patients in the OHA group had significantly higher preoperative leukocyte counts, a greater prevalence of systemic inflammatory response syndrome, and higher rates of predischarge SSI. This phenomenon—known as *confounding by indication*—precludes a clear distinction between the effects of the treatment and the underlying severity of illness. Another key reason was the presence of concurrent interventions; for example, the study by Ferguson et al. [24] coincided with a quality improvement initiative on the use of intraoperative abdominal drains and a separate pilot randomized trial of intra-abdominal irrigation with povidone-iodine, making it difficult to isolate the effect of the antibiotic protocol alone. Other reasons included the use of historical controls, which introduces temporal confounding, and massive, unadjusted baseline differences in disease severity between hospital cohorts being compared (such as in the case of Desai et al.) [20].

For bias in the selection of participants into the study, no studies were identified as having a serious risk, 5 were rated as moderate [20,24,28,30,31], and 9 were rated as low risk [21-23,25-27,29,32,33].

In the domain of bias in the classification of interventions, no studies were found to pose a serious risk, 7 were rated as moderate [21,23,25,27,28,30,33], and 7 were rated as low risk [20,22,24,26,29,31,32].

Regarding bias due to deviations from intended interventions, one study was rated as having a serious risk [23], nine were rated as moderate [21,24,25,27,28-32], and four were rated as low risk [20,22,26,33]. The serious risk rating was assigned because the study explicitly reported that "overall compliance with protocol guidelines was not ideal," with participation being at the discretion of the surgeon. This led to substantial crossover, where patients in the new protocol group still received older treatments, thus contaminating the comparison groups and biasing the results toward finding no difference.

In terms of bias due to missing data, no studies had a serious risk, 8 were rated as moderate [20,22,23,25-27,29,33], and 6 were rated as low risk [21,24,28,30-32].

Regarding bias in the measurement of outcomes, none of the studies posed a serious risk, one study posed a moderate risk [20], and 14 were rated as low risk [21-33].

Finally, in the domain of bias in the selection of the reported result, no study was judged to be at serious or moderate risk; all 14 studies were rated as having a low risk of bias [20-33].

Figure 2 visually displays the ROBINS-I risk of bias assessments across individual domains for each included study.

Intraabdominal Abscess (Oral Home Antibiotics vs. No Home Antibiotics).

Six studies reported IAA rates stratified by treatment group [20,24,25,26,27,28]. Bonasso et al. [22] presented two separate cohorts. Four studies found no significant differences in IAA rates between groups [20,25,26,28]. One study reported a significantly higher IAA rate in the IG compared to the CG [24], whereas another study reported significantly higher IAA rates in the CG than in the IG [27].

Notably, the only study reporting increased IAA rates in the IG was Ferguson et al. [24], which uniquely implemented a protocol to standardize the routine use of OHA, rather than to reduce them. This contrasts with the rest of the literature, where interventions consistently aimed to minimize post-discharge antibiotic use.

Meta-Analysis for Intraabdominal Abscess (Oral Home Antibiotics vs. No Home Antibiotics).

For the meta-analysis of IAA, a random-effects model was fitted using restricted maximum likelihood (REML) with truncated modified Knapp–Hartung (mKH) adjustments. Seven studies were included, encompassing 788 patients in the IG and 1,011 patients in the CG. The pooled risk ratio (RR) comparing IG versus CG was 0.97 (95% CI: 0.38-2.47; model p = 0.94), indicating no significant difference in IAA rates between the groups. Between-study heterogeneity was moderate-to-substantial (I^2 = 66.98%; Cochran's Q p = 0.01), and the estimated between-study variance was $\tau^2 = 0.47$. Leave-one-out sensitivity analysis revealed that excluding any single study did not materially alter the overall result or achieve statistical significance in any iteration, supporting the robustness of the null finding (Figure 3). This meta-analysis included a combination of studies reporting raw group-level data (OHA vs NHA) and others presenting outcomes for cohorts managed under pre- or post-intervention protocols.

Supplementary File 4 and Table 1 provide detailed information on the data used for each study, including whether raw exposure data were available. Given that fewer than ten studies were included, between-study heterogeneity was not further explored using funnel plots or formal statistical tests for small-study effects (e.g., Egger's or Begg's tests). Similarly, no meta-regression analyses were conducted due to the low number of studies and limited statistical power.

Surgical Site Infection (Oral Home Antibiotics vs. No Home Antibiotics).

Seven studies reported SSI rates stratified by treatment group [21,23,24,29,31-33]. Five studies found no significant differences in SSI rates between groups [23,24,29,32,33]. One study reported a significantly higher SSI rate in the IG compared to the CG [31], whereas another study reported significantly higher SSI rates in the CG than in the IG [21].

Meta-Analysis for Surgical Site Infection (Oral Home Antibiotics vs. No Home Antibiotics).

For the meta-analysis of SSI, a random-effects model was fitted using REML with mKH adjustments. Seven studies were included, encompassing 3,858 patients in the IG and 4,610 in the CG. The pooled RR comparing IG versus CG was 0.89 (95% CI: 0.58-1.38; model p = 0.54), indicating no significant difference in SSI rates. Between-study heterogeneity was low (I^2 = 21.91%; Cochran's Q p = 0.33), and the estimated between-study variance was $\tau^2 = 0.05$. Leave-one-out sensitivity analysis revealed no iteration reached statistical significance. The exclusion of Russell et al. produced the most pronounced change, yielding a borderline non-significant effect (RR = 0.76, 95% CI: 0.56-1.02; p = 0.06), suggesting a potential trend toward higher SSI rates in the IG, although still inconclusive (Figure 4). This meta-analysis included a combination of studies reporting raw group-level data (OHA vs NHA) and others presenting outcomes for

cohorts managed under pre- or post-intervention protocols. Given that fewer than ten studies were included, between-study heterogeneity was not further explored using funnel plots or formal statistical tests for small-study effects (e.g., Egger's or Begg's tests). Similarly, no meta-regression analyses were conducted due to the low number of studies and limited statistical power.

<u>Organ Space Infection (Oral Home Antibiotics vs. No Home Antibiotics).</u>

Three studies reported OSI rates stratified by treatment group [28,29,32]. None of them found significant differences in OSI rates between groups [28,29,32].

<u>Meta-Analysis for Organ Space Infection (Oral Home Antibiotics vs. No Home Antibiotics).</u>

For the meta-analysis of OSI, a random-effects model was fitted using REML. Three studies were included, encompassing 685 patients in the IG and 558 in the CG. In the case of Anandalwar et al., only data from the propensity-matched cohort were included to avoid duplicating data from the same population and to ensure study independence. The pooled RR comparing IG versus CG was 1.19 (95% CI: 0.73-1.93; model $p = 0.48$), indicating no significant difference in OSI rates. Between-study heterogeneity was negligible ($I^2 = 0\%$; Cochran's Q $p = 0.47$), and the estimated between-study variance was $\tau^2 = 0$. Accordingly, confidence intervals were calculated using conventional Wald-type methods, as recommended when no heterogeneity is detected. Leave-one-out sensitivity analysis revealed that no single study significantly altered the overall result or achieved statistical significance in any iteration (Figure 5). This meta-analysis included a combination of studies reporting raw group-level data (OHA vs NHA) and others presenting outcomes for cohorts managed under pre- or post-intervention protocols. Given that fewer than ten studies were included, between-study

heterogeneity was not further explored using funnel plots or formal statistical tests for small-study effects (e.g., Egger's or Begg's tests). Similarly, no meta-regression analyses were conducted due to the low number of studies and limited statistical power.

Readmissions (Oral Home Antibiotics vs. No Home Antibiotics).

Thirteen studies reported RA rates stratified by treatment group [21–33]. Ten studies found no significant differences in RA rates between groups [22–29, 31, 32]. One study reported a significantly higher RA rate in the IG compared to the CG [24], whereas two studies reported significantly higher RA rates in the CG than in the IG [21, 30]. In the case of Patwardhan et al. [33], although the difference did not reach statistical significance, a marginal p-value (0.06) was observed, suggesting a higher rate of events in the IG. Although most studies defined RA as occurring within 30 days of initial discharge, some—such as Morita et al. [30]—used a 60-day timeframe, which may influence effect estimates due to the extended follow-up window.

Meta-Analysis for Readmissions (Oral Home Antibiotics vs. No Home Antibiotics).

For the meta-analysis of RA, a random-effects model was fitted using REML with mKH adjustments. Twelve studies were included, encompassing 7,259 patients in the IG and 8,053 in the CG. In the cases of Anandalwar et al. (2021) and Morita et al. (2022) [28,30], only data from the propensity-matched cohorts were included to ensure independence of observations. The pooled RR comparing IG versus CG was 1.02 (95% CI: 0.73-1.41; model p = 0.92), indicating no significant difference in readmission rates. Between-study heterogeneity was substantial ($I^2 =$ 68.68%; Cochran's Q p = 0.00), with a between-study variance of $\tau^2 = 0.13$. Leave-one-out sensitivity analysis revealed that no single study materially affected the overall result or achieved statistical significance. The exclusion of Ferguson et al. (2020) produced the most pronounced

shift, but the result remained non-significant (Figure 6). Publication bias and small-study effects were assessed for the readmissions meta-analysis using Egger's and Begg's tests, which yielded non-significant results (Egger $p = 0.18$; Begg $p = 0.19$), indicating no statistical evidence of asymmetry. Visual inspection of funnel, Galbraith, and contour-enhanced plots also did not reveal patterns suggestive of publication bias (Figure 7). This meta-analysis included a combination of studies reporting raw group-level data (OHA vs NHA) and others presenting outcomes for cohorts managed under pre- or post-intervention protocols.

A univariable meta-regression was conducted to explore whether the availability of crude group-level data comparing OHA versus NHA influenced the pooled effect estimates. This analysis showed a non-significant trend toward lower risk ratios in studies reporting crude OHA/NHA data compared to those using indirect comparisons ($\beta = -0.56$; $p = 0.081$). Based on this finding, all meta-analyses (IAA, SSI, OSI, and RA) were repeated, including only studies with clearly defined OHA and NHA groups and events. All restricted models remained non-significant except for RA, where the analysis yielded a pooled risk ratio of 0.75 (95% CI: 0.62-0.90; $p < 0.01$) under a random-effects model using REML with mKH adjustments. Heterogeneity was negligible ($I^2 = 0\%$; Cochran's Q $p = 0.55$), suggesting that studies directly comparing OHA and NHA consistently found a higher rate of readmissions among patients who received oral antibiotics at discharge (Figure 8). This result supports the hypothesis that OHA may not reduce—and could even increase—the risk of hospital readmission.

Another univariable meta-regression was conducted using predischarge total leukocyte count (TLC) differences as a categorical moderator to explore whether inflammatory status at discharge influenced the association between OHA and the risk of RA. The analysis compared studies reporting no differences in TLC between groups (reference category) with those reporting

significant differences—most often reflecting higher TLC levels in the comparator group (CG, i.e., OHA)—and studies that did not specify this information. Studies with significant intergroup TLC differences showed a non-significant trend toward lower pooled risk ratios ($\beta = –1.15$; $p = 0.14$), suggesting fewer readmissions in the intervention group (IG, i.e., NHA). Importantly, these differences likely reflect clinical decision-making, whereby patients with signs of residual inflammation were more prone to receive antibiotics. Therefore, the lower RR observed in these studies in favor of NHA does not indicate a protective effect of OHA but rather the opposite: that despite being clinically selected for antibiotic treatment, OHA patients experienced more readmissions. This supports the interpretation that OHA does not reduce—and may even increase—the risk of hospital readmission, independent of predischarge inflammatory status. However, as the category with significant TLC differences included only two studies, this trend should be interpreted with caution, given the potential influence of individual study effects and limited statistical power.

Lastly, a univariable meta-regression comparing studies at high versus moderate risk of bias revealed a non-significant trend toward more favorable outcomes in the NHA group among high-risk studies (logRR difference: –0.34; 95% CI: –1.03 to 0.36; $p = 0.30$). However, this association did not reach statistical significance, explained none of the observed heterogeneity ($R^2 = 0\%$), and should therefore be interpreted as clinically and methodologically uninformative.

Secondary Outcomes (Oral Home Antibiotics vs. No Home Antibiotics).

Some studies also examined alternative outcomes such as composite post-discharge morbidity [21], reoperations (RO) [21,23,24], and Emergency Department visits (EDV) [21,23,24,29]. Detailed outcome data are available in Table 1. Given the low number of studies reporting these

outcomes and the heterogeneity and dispersion of available data, no meta-analytic models were fitted for these endpoints.

GRADE Assessment

Finally, we evaluated the certainty of evidence using the GRADE approach across primary outcomes. Overall, the certainty was low for IAA, SSI, and RA, and very low for OSI. The main reasons for downgrading were the observational nature of all studies, the presence of heterogeneity in some analyses, and imprecision due to wide confidence intervals. No outcome reached a moderate or high level of certainty, reinforcing the need for future randomized trials to address this gap.

Discussion

This systematic review and meta-analysis provides the most comprehensive synthesis to date of the impact of OHA following surgical management of CAA in children. Across multiple outcomes—including IAA, SSI, OSI, and RA—no consistent benefit was observed in favor of OHA use. Both individual study estimates and pooled risk ratios failed to support routine administration of post-discharge OHA. Despite their widespread use, this is the first synthesis to assess their effectiveness across these core outcomes. Notably, several analyses even trended in favor of NHA discharge strategies, despite these patients often presenting with lower predischarge inflammatory markers.

Our findings challenge the traditional rationale for extended antibiotic coverage after discharge in pediatric CAA, particularly in cases with adequate source control. Previous guidelines and institutional practices often endorsed OHA as a safeguard against latent infections or undetected

abscess formation. However, the current evidence, including recent protocol-based studies aimed at restricting OHA, suggests that this approach may offer no additional benefit—and may contribute to overtreatment, antimicrobial resistance, and unnecessary drug exposure. This underscores the importance of aligning postoperative management with principles of antimicrobial stewardship, especially in pediatric patients.

Interestingly, the only study reporting a significantly higher RA and IAA rate in the NHA group [24] was also the only one to implement a protocol aimed at standardizing rather than limiting post-discharge OHA. This divergence contrasts with the broader trend in the literature, where interventions were designed to reduce or eliminate OHA use. Thus, this deviation highlights how institutional culture and protocol framing may shape not only practices but also outcomes. It also raises the possibility of confounding by protocol context—underscoring the need for randomized comparisons within standardized care frameworks.

From a clinical standpoint, these results reinforce the safety and feasibility of discharging selected pediatric patients without OHA following appendectomy for CAA, provided that intraoperative findings, clinical stability, and objective inflammatory markers indicate resolution. The evidence suggests that individualized clinical judgment and predischarge inflammatory markers may suffice to guide safe discharge decisions, rather than adhering to routine antibiotic protocols. This supports the development of risk-adapted discharge pathways integrating objective markers of resolution. These findings align with broader principles of antibiotic stewardship in pediatric surgery.

Another potential source of bias stems from clinical decision-making criteria for prescribing antibiotics at discharge. Although some studies reported predischarge TLC and their

comparability between groups, this information was frequently absent or insufficiently detailed. This limits the ability to determine whether the inflammatory status at discharge systematically influenced treatment allocation. In our exploratory meta-regression, studies reporting statistically significant differences in TLC between groups—typically indicating higher values in the antibiotic group—showed a trend toward lower pooled RR in the NHA. While this finding should be interpreted with caution due to the limited power and potential confounding, it suggests that patients perceived as being at higher risk (e.g., those with elevated TLC) may have been selectively prescribed antibiotics, yet did not show a corresponding reduction in readmission rates. This paradoxical trend may reflect residual confounding and highlight the complexity of interpreting observational data. This hypothesis-generating observation highlights the need for future studies to document clinical parameters that inform discharge decisions rigorously and to control for these factors in comparative analyses. Structured discharge algorithms incorporating inflammatory markers may help reduce selection bias and improve comparability in future research.

Another important consideration is the limited reporting and control of variables that may substantially influence outcomes but were inconsistently addressed across studies. For example, none of the included articles provided objective measures of patient compliance with prescribed oral antibiotics—a critical determinant of treatment efficacy. Similarly, the relationship between discharge antibiotic strategy and surgical technique (e.g., laparoscopic versus open approach), intraoperative findings, or postoperative management protocols was rarely explored in depth. These factors could interact with the effect of OHA and confound outcome estimates. Additionally, key elements such as discharge timing, the presence of drains, or concomitant antibiotic regimens (e.g., IV continuation) were often insufficiently detailed. Potential

differences in microbiological etiology or resistance patterns were also largely unaddressed, limiting biological interpretability and external applicability. These gaps highlight the need for future prospective studies to standardize and report such variables systematically, enabling more precise estimation of the clinical utility—and limitations—of routine OHA in CAA.

From a biological standpoint, the rationale for OHA in this setting remains biologically questionable. The peritoneal cavity, particularly in the context of localized postoperative inflammation and fibrinous exudate, may offer limited penetration for orally administered antibiotics—especially in the absence of systemic signs of infection. Furthermore, the oral route introduces variability in absorption, particularly in children with reduced appetite, postoperative ileus, or gastrointestinal dysmotility. These pharmacokinetic limitations, combined with incomplete data on compliance and dosing, raise concerns about the real-world efficacy of oral agents in preventing deep or loculated infections. Additionally, unnecessary antibiotic exposure may select for resistant microbial strains, potentially altering the child's microbiome and increasing future vulnerability to infection. Altogether, these pharmacological considerations further support a risk-adapted rather than uniform approach to OHA use.

Moreover, none of the included studies systematically reported adverse events associated with oral antibiotic use, such as diarrhea, candidiasis, or intestinal dysbiosis, which represents an important limitation. This omission prevents quantification of the potential collateral harms of unnecessary antibiotic administration, but it indirectly reinforces one of the core principles of antibiotic stewardship: avoiding treatments that provide no clinical benefit and may instead cause preventable adverse effects. In this context, our findings are particularly applicable to patients with favorable postoperative evolution—namely, those who are afebrile, tolerating oral intake, in good general condition, and with normalized inflammatory markers at discharge—for whom

continued antibiotic treatment beyond discharge may offer no additional benefit. It should also be emphasized that the majority of included studies administered intravenous antibiotics during hospitalization, and thus, these results should not be extrapolated to settings in which early discharge with oral antibiotics alone is considered—an approach not evaluated in this review. Avoiding unnecessary OHA in these patients may also reduce the risk of avoidable harm in an already recovering pediatric population.

Lastly, an important methodological concern highlighted by this review is the substantial variability in the reporting of surgical technique. We observed marked differences both in terms of inclusion/exclusion criteria based on surgical approach (open vs. laparoscopic) and in the overall quality of reporting. While some studies explicitly excluded open appendectomies or documented conversions, others merely stated that laparoscopy was used without specifying whether this applied to all patients or whether any conversions occurred. Several studies failed to report surgical approach altogether. This lack of standardization is concerning, given well-established evidence that open appendectomies carry a higher risk of surgical site infection compared to laparoscopic procedures, potentially introducing uncontrolled confounding in the interpretation of results. Similarly, relevant operative variables such as surgical duration—which may reflect case complexity and is associated with postoperative outcomes—were scarcely reported across the included studies. Given the well-established association between surgical approach and postoperative infection risk, this lack of standardization represents a critical gap in current evidence. Future research should systematically collect and transparently report key surgical variables to allow for more accurate risk adjustment and more robust interpretation of findings.

This review has several strengths. It adheres to PRISMA and Cochrane methodological standards. It applies modern, rigorous statistical techniques, such as random-effects REML meta-analysis with robust confidence intervals (HKSJ and modified Knapp–Hartung), leave-one-out sensitivity analyses, and assessment of publication bias when feasible. Univariable meta-regression was used to explore sources of heterogeneity, including differences in predischarge TLC and the type of comparator group (raw OHA/NHA vs protocol-defined). Particular care was taken to distinguish studies reporting individual-level data from those reporting outcomes based on pre- and post-protocol cohorts, and all analyses were conducted accordingly. Importantly, statistical results were interpreted in light of the clinical context, considering how discharge practices and patient selection may have influenced both antibiotic use and outcomes. However, relevant limitations must be acknowledged. There was variability in the definition of CAA across studies, and individual-level OHA and NHA raw data were often unavailable. Many studies lacked control for key variables such as antibiotic type, dosage, or precise discharge criteria. Additionally, the complications—particularly IAA and OSI—were not extensively characterized across articles, e.g., omitting severity grading or volumetric analysis of abscesses. Most studies were retrospective and based in the USA, and those using large databases, while offering substantial sample sizes, often lacked granular clinical information. Notably, only one study [27] incorporated solid microbiological data, which limits the biological interpretability of the outcomes. Accordingly, the certainty of evidence was rated as low or very low across all primary outcomes using the GRADE framework [34], primarily due to the observational nature of the studies, moderate to high heterogeneity in some analyses, and imprecision of effect estimates. These conclusions should be interpreted with caution given the overall low certainty of evidence.

In light of these findings, the routine prescription of OHA after discharge in pediatric patients with surgically treated CAA appears unwarranted. Across over 26,000 patients, no consistent benefit was observed in terms of IAA, SSI, OSI or RA. Importantly, subgroup crude analyses even suggest that OHA may increase the risk of readmission, particularly in patients selected for treatment based on residual inflammation (TLC). However, the current body of evidence is based exclusively on observational studies, with heterogeneous definitions, limited adjustment for confounders, and no data on patient adherence. There is an urgent need for adequately powered, multicenter randomized controlled trials that rigorously control for clinical, surgical, and microbiological variables to definitively establish the role—or lack thereof—of oral home antibiotics in this setting. Future studies should clearly specify the surgical technique employed (e.g., open vs laparoscopic, use of peritoneal lavage), characterize microbiological findings and antibiotic susceptibility profiles, and distinguish individual-level antibiotic exposure (OHA vs NHA) rather than relying solely on protocol-based groupings. Additionally, consistent definitions for outcomes such as IAA, SSI, and readmissions are essential to ensure comparability across studies. Until such trials are available, a cautious, individualized approach to OHA prescription—grounded in objective resolution criteria—may best balance safety with antimicrobial stewardship in pediatric CAA.

Figure legends

Figure 1. Flowchart illustrating the search strategy and study selection process.

Figure 2. Graphical summary of the risk of bias assessment using the ROBINS-I tool for the included non-randomized studies. Green circles with a plus sign indicate low risk of bias, yellow circles with a question mark represent moderate risk, and red circles with a minus sign denote high risk of bias.

Figure 3. <u>Top</u>: Forest plot of the random-effects meta-analysis for intra-abdominal abscesses (IAA) using REML with Knapp–Hartung adjustment for confidence intervals (IG vs. CG). <u>Bottom</u>: Leave-one-out forest plot for the same analysis.

Figure 4. <u>Top</u>: Forest plot of the random-effects meta-analysis for surgical site infections (SSI) using REML with Knapp–Hartung adjustment (IG vs. CG). <u>Bottom</u>: Leave-one-out forest plot for the same analysis.

Figure 5. <u>Top</u>: Forest plot of the random-effects meta-analysis for organ/space infections (OSI) using REML (IG vs. CG). <u>Bottom</u>: Leave-one-out forest plot for the same analysis.

Figure 6. <u>Top</u>: Forest plot of the random-effects meta-analysis for hospital readmissions (RA) using REML with Knapp–Hartung adjustment (IG vs. CG). <u>Bottom</u>: Leave-one-out forest plot for the same analysis.

Figure 7. <u>Top</u>: Contour-enhanced funnel plot for the RA meta-analysis, showing significance contours at $p < 0.01$, $p < 0.05$, $p < 0.10$, and $p > 0.10$. <u>Center</u>: Galbraith (radial) plot for the RA meta-analysis. <u>Bottom</u>: L'Abbé plot for the RA meta-analysis.

Figure 8. Forest plot of the sensitivity meta-analysis for hospital readmissions (RA) based on crude IG vs. CG data, using REML with Knapp–Hartung adjustment.

Supplementary Materials

Supplementary File 1. PRISMA checklist

Supplementary File 2. Full search strategy.

Supplementary File 3. Inclusion and exclusion criteria.

Supplementary File 4. Full dataset

References

[1].   Chionye R. Ossai, Lu Pu, David Kaelber; Using Aggregated Data from 1.4 Million Pediatric Patients to Describe the Epidemiology and Demographic Characteristics of Appendicitis. *Pediatrics* July 2020; 146 (1_MeetingAbstract): 232–234. 10.1542/peds.146.1MA3.232b

[2].   Omling E, Salö M, Saluja S, Bergbrant S, Olsson L, Persson A, Björk J, Hagander L. Nationwide study of appendicitis in children. Br J Surg. 2019 Nov;106(12):1623-1631. doi: 10.1002/bjs.11298. Epub 2019 Aug 6. PMID: 31386195; PMCID: PMC6852580.

[3].   Georgeades C, Bodnar C, Bergner C, Van Arendonk KJ. Association of complicated appendicitis with geographic and socioeconomic measures in children. Surgery. 2024 Nov;176(5):1475-1484. doi: 10.1016/j.surg.2024.07.044. Epub 2024 Sep 3. PMID: 39232975.

[4].   Michelson KA, Bucher BT, Neuman MI. Cost and Late Hospital Care of Publicly Insured Children After Appendectomy. J Surg Res. 2024 May;297:41-46. doi: 10.1016/j.jss.2024.02.003. Epub 2024 Mar 1. PMID: 38430861; PMCID: PMC11023751.

[5].   Wang C, Li Y, Ji Y. Intravenous versus intravenous/oral antibiotics for perforated appendicitis in pediatric patients: a systematic review and meta-analysis. BMC Pediatr. 2019 Nov 4;19(1):407. doi: 10.1186/s12887-019-1799-6. PMID: 31684906; PMCID: PMC6827245.

[6].   Lam JY, Beaudry P, Simms BA, Brindle ME. Impact of implementing a fast-track protocol and standardized guideline for the management of pediatric appendicitis. Can J Surg. 2021 Jul 5;64(4):E364-E370. doi: 10.1503/cjs.005420. PMID: 34223740; PMCID: PMC8410463.

[7].   Muehlstedt SG, Pham TQ, Schmeling DJ. The management of pediatric appendicitis: a survey of North American Pediatric Surgeons. J Pediatr Surg. 2004 Jun;39(6):875-9; discussion 875-9. doi: 10.1016/j.jpedsurg.2004.02.035. PMID: 15185217.

[8].   Huttner B, Harbarth S, Carlet J, et al. Antibiotics are not automatic anymore—the French national campaign to cut antibiotic overuse. PLoS Med. 2010;7(7):e1000353. doi:10.1371/journal.pmed.1000353

[9].   Huttner B, Harbarth S, Carlet J, et al. Antibiotics are not automatic anymore—the French national campaign to cut antibiotic overuse. PLoS Med. 2010;7(7):e1000353. doi:10.1371/journal.pmed.1000353

[10].   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021 Mar 29;372:n71. doi: 10.1136/bmj.n71. PMID: 33782057; PMCID: PMC8005924.

[11].   Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Cochrane Handbook for Systematic Reviews of Interventions version 6.5 (updated August 2024). Cochrane, 2024. Available from www.cochrane.org/handbook.

[12].   Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ. 2019 Aug 28;366:l4898. doi: 10.1136/bmj.l4898. PMID: 31462531.

[13].   Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. BMJ. 2016 Oct 12;355:i4919. doi: 10.1136/bmj.i4919. PMID: 27733354; PMCID: PMC5062054.

[14].   Veroniki AA, Jackson D, Viechtbauer W, Bender R, Bowden J, Knapp G, et al. Methods to estimate the between-study variance and its uncertainty in meta-analysis. Res Synth Methods. 2016;7(1):55–79. doi:10.1002/jrsm.1164

[15].   IntHout J, Ioannidis JP, Borm GF. The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. BMC Med Res Methodol. 2014 Feb 18;14:25. doi: 10.1186/1471

[16].   Röver C, Knapp G, Friede T. Hartung-Knapp-Sidik-Jonkman approach and its modification for random-effects meta-analysis with few studies. BMC Med Res Methodol. 2015 Nov 14;15:99. doi: 10.1186/s12874-015-0091-1. PMID: 26573817; PMCID: PMC4647507

[17]. 2288-14-25. PMID: 24548571; PMCID: PMC4015721Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. Biometrics. 1994;50(4):1088-1101.

[18]. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. BMJ. 1997;315(7109):629-634.

[19]. Shi L, Lin L. The trim-and-fill method for publication bias: practical guidelines and recommendations based on a large database of meta-analyses. Medicine (Baltimore). 2019 Jun;98(23):e15987. doi: 10.1097/MD.0000000000015987. PMID: 31169736; PMCID: PMC6571372.

[20]. Desai AA, Alemayehu H, Holcomb GW 3rd, St Peter SD. Safety of a new protocol decreasing antibiotic utilization after laparoscopic appendectomy for perforated appendicitis in children: A prospective observational study. J Pediatr Surg. 2015 Jun;50(6):912-4. doi: 10.1016/j.jpedsurg.2015.03.006. Epub 2015 Mar 14. PMID: 25812441.

[21]. Anderson KT, Bartz-Kurycki MA, Kawaguchi AL, Austin MT, Holzmann-Pazgal G, Kao LS, Lally KP, Tsao K. Home Antibiotics at Discharge for Pediatric Complicated Appendicitis: Friend or Foe? J Am Coll Surg. 2018 Aug;227(2):247-254. doi: 10.1016/j.jamcollsurg.2018.04.004. Epub 2018 Apr 20. PMID: 29680415.

[22]. Bonasso PC, Dassinger MS, Wyrick DL, Smith SD, Burford JM. Evaluation of white blood cell count at time of discharge is associated with limited oral antibiotic therapy in children with complicated appendicitis. Am J Surg. 2019 Jun;217(6):1099-1101. doi: 10.1016/j.amjsurg.2018.12.071. Epub 2019 Jan 3. PMID: 30639131.

[23]. Rossidis AC, Brown EG, Payton KJ, Mattei P. Implementation of an evidence-based protocol after appendectomy reduces unnecessary antibiotics. J Pediatr Surg. 2020 Nov;55(11):2379-2386. doi: 10.1016/j.jpedsurg.2020.07.001. Epub 2020 Jul 9. PMID: 32753275.

[24]. Ferguson DM, Parker TD, Arshad SA, Garcia EI, Hebballi NB, Tsao K. Standardized Discharge Antibiotics May Reduce Readmissions in Pediatric Perforated Appendicitis. J Surg Res. 2020 Nov;255:388-395. doi: 10.1016/j.jss.2020.05.086. Epub 2020 Jun 29. PMID: 32615311.

[25]. Gordon AJ, Choi JH, Ginsburg H, Kuenzler K, Fisher J, Tomita S. Oral Antibiotics and Abscess Formation After Appendectomy for Perforated Appendicitis in Children. J Surg Res. 2020 Dec;256:56-60. doi: 10.1016/j.jss.2020.05.082. Epub 2020 Jul 16. PMID: 32683057.

[26]. Ketha B, Stephenson KJ, Dassinger MS 3rd, Smith SD, Burford JM. Eliminating Use of Home Oral Antibiotics in Pediatric Complicated Appendicitis. J Surg Res. 2021 Jul;263:151-154. doi: 10.1016/j.jss.2020.12.059. Epub 2021 Feb 27. PMID: 33652177.

[27]. Plattner AS, Newland JG, Wallendorf MJ, Shakhsheer BA. Management and Microbiology of Perforated Appendicitis in Pediatric Patients: A 5-Year Retrospective Study. Infect Dis Ther. 2021 Dec;10(4):2247-2257. doi: 10.1007/s40121-021-00502-x. Epub 2021 Jul 21. PMID: 34287780; PMCID: PMC8572942.

[28]. Anandalwar SP, Graham DA, Kashtan MA, Hills-Dunlap JL, Rangel SJ. Influence of Oral Antibiotics Following Discharge on Organ Space Infections in Children With

Complicated Appendicitis. Ann Surg. 2021 Apr 1;273(4):821-825. doi: 10.1097/SLA.0000000000003441. PMID: 31274648.

[29]. Jen J, Hwang R, Mattei P. Post-discharge antibiotics do not prevent intra-abdominal abscesses after appendectomy in children. J Pediatr Surg. 2023 Feb;58(2):258-262. doi: 10.1016/j.jpedsurg.2022.10.024. Epub 2022 Oct 29. PMID: 36428182.

[30]. Morita K, Fujiogi M, Michihata N, Matsui H, Fushimi K, Yasunaga H, Fujishiro J. Oral Antibiotics and Organ Space Infection after Appendectomy and Intravenous Antibiotics Therapy for Complicated Appendicitis in Children. Eur J Pediatr Surg. 2023 Feb;33(1):74-80. doi: 10.1055/a-1958-7915. Epub 2022 Oct 11. PMID: 36220134.

[31]. Theodorou CM, Lee SY, Lawrence Y, Saadai P, Hirose S, Brown EG. The Utility of Discharge Antibiotics in Pediatric Perforated Appendicitis Without Leukocytosis. J Surg Res. 2022 Jul;275:48-55. doi: 10.1016/j.jss.2022.01.024. Epub 2022 Feb 23. PMID: 35219251; PMCID: PMC10032146.

[32]. Russell KW, Skarda DE, Jones TW, Barnhart DC, Short SS. Cessation of Antibiotics for Complicated Appendicitis at Discharge Does Not Increase Risk of Post-operative Infection. J Pediatr Surg. 2024 Jan;59(1):91-95. doi: 10.1016/j.jpedsurg.2023.09.023. Epub 2023 Sep 26. PMID: 37858398.

[33]. Patwardhan UM, Kahan A, Eldredge RS, Russell KW, Lee J, Short SS, Padilla B, Cairo SB, Acker SN, Jensen AR, Kelley-Quon LI, Rothstein DH, Fialkowski EA, Chao SD, Gillory L, Pandya S, Diaz-Miron J, Ignacio RC Jr. Comparison of Postoperative Antibiotic Protocols for Pediatric Complicated Appendicitis: A Western Pediatric Surgery Research Consortium

Study. J Pediatr Surg. 2025 Apr;60(4):162165. doi: 10.1016/j.jpedsurg.2025.162165. Epub 2025 Jan 9. PMID: 39827485.

[34]. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, Schünemann HJ; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. BMJ. 2008 Apr 26;336(7650):924-6. doi: 10.1136/bmj.39489.470347.AD. PMID: 18436948; PMCID: PMC2335261.

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases and registers only

Study	IG Yes	IG No	CG Yes	CG No		Risk ratio with 95% CI	Weight (%)
Desai et al. (2015)	11	124	6	130		1.85 [0.70, 4.85]	15.49
Bonasso et al. (2019) old protocol	1	6	7	68		1.53 [0.22, 10.72]	7.57
Bonasso et al. (2019) new protocol	1	68	7	21		0.06 [0.01, 0.45]	7.05
Ferguson et al. (2020)	19	194	18	386		2.00 [1.07, 3.73]	19.36
Gordon et al. (2020)	7	184	14	148		0.42 [0.18, 1.03]	16.42
Ketha et al. (2021)	7	51	9	62		0.95 [0.38, 2.40]	15.95
Plattner et al. (2021)	14	101	12	123		1.37 [0.66, 2.84]	18.16
Overall						0.97 [0.38, 2.47]	

Heterogeneity: $t^2 = 0.47$, $I^2 = 66.98\%$, $H^2 = 3.03$

Test of $q_i = q_j$: Q(6) = 17.50, p = 0.01

Test of q = 0: t(6) = −0.08, p = 0.94

1/128   1/16   1/2   4

Random−effects REML model
Truncated Knapp Hartung standard errors

Omitted study	Risk ratio with 95% CI	p−value
Desai et al. (2015)	0.84 [0.26, 2.65]	0.708
Bonasso et al. (2019) old protocol	0.91 [0.29, 2.85]	0.836
Bonasso et al. (2019) new protocol	1.21 [0.62, 2.36]	0.487
Ferguson et al. (2020)	0.81 [0.27, 2.48]	0.650
Gordon et al. (2020)	1.28 [0.53, 3.13]	0.502
Ketha et al. (2021)	0.93 [0.27, 3.18]	0.880
Plattner et al. (2021)	0.86 [0.25, 2.92]	0.764

1/2   1   2

Random−effects REML model
Truncated Knapp Hartung standard errors

		IG		CG			Risk ratio	Weight
Study		Yes	No	Yes	No		with 95% CI	(%)
Anderson et al. (2018)		92	2,706	163	3,451		0.73 [0.57, 0.94]	48.28
Rossidis et al. (2020)		8	167	11	152		0.68 [0.28, 1.64]	12.52
Ferguson et al. (2020)		2	211	3	401		1.26 [0.21, 7.51]	3.62
Jen et al. (2022)		5	272	0	86		3.44 [0.19, 61.63]	1.43
Theodorou et al. (2022)		8	108	3	91		2.16 [0.59, 7.92]	6.50
Russell et al. (2024)		13	108	14	171		1.42 [0.69, 2.91]	17.24
Patwardhan et al. (2025)		9	149	6	58		0.61 [0.23, 1.64]	10.40
Overall							0.89 [0.58, 1.38]	

Heterogeneity: $t^2 = 0.05$, $I^2 = 21.91\%$, $H^2 = 1.28$

Test of $q_i = q_j$: $Q(6) = 6.89$, $p = 0.33$

Test of $q = 0$: $t(6) = -0.65$, $p = 0.54$

1/4   1   4   16

Random−effects REML model
Truncated Knapp Hartung standard errors

	Risk ratio	
Omitted study	with 95% CI	p−value
Anderson et al. (2018)	1.07 [0.58, 1.97]	0.786
Rossidis et al. (2020)	0.98 [0.55, 1.75]	0.921
Ferguson et al. (2020)	0.88 [0.55, 1.41]	0.521
Jen et al. (2022)	0.87 [0.56, 1.35]	0.444
Theodorou et al. (2022)	0.81 [0.55, 1.19]	0.215
Russell et al. (2024)	0.76 [0.56, 1.02]	0.062
Patwardhan et al. (2025)	0.98 [0.56, 1.71]	0.917

0.55   1.97

Random−effects REML model
Truncated Knapp Hartung standard errors

Study	IG Yes	IG No	CG Yes	CG No		Risk ratio with 95% CI	Weight (%)
Anandalwar et al. (2021) after propensity matching	19	268	12	275		1.58 [0.78, 3.20]	47.33
Jen et al. (2022)	9	268	4	82		0.70 [0.22, 2.21]	17.65
Russell et al. (2023)	9	112	13	172		1.06 [0.47, 2.40]	35.02
Overall				

 | 1.19 [0.73, 1.93] | |

Heterogeneity: $t^2 = 0.00$, $I^2 = 0.00\%$, $H^2 = 1.00$

Test of $q_i = q_j$: $Q(2) = 1.53$, $p = 0.47$

Test of $q = 0$: $z = 0.70$, $p = 0.48$

Random−effects REML model

Omitted study		Risk ratio with 95% CI	p−value
Anandalwar et al. (2021) after propensity matching		0.92 [0.47, 1.79]	0.809
Jen et al. (2022)		1.33 [0.78, 2.28]	0.290
Russell et al. (2023)		1.20 [0.56, 2.56]	0.637

Random−effects REML model

Study	IG Yes	IG No	CG Yes	CG No		Risk ratio with 95% CI	Weight (%)
Anderson et al. (2018)	153	2,645	286	3,328		0.69 [0.57, 0.84]	15.41
Ferguson et al. (2020)	28	185	26	378		2.04 [1.23, 3.39]	10.98
Gordon et al. (2020)	10	181	17	145		0.50 [0.24, 1.06]	7.83
Rossidis et al. (2020)	9	166	5	158		1.68 [0.57, 4.90]	5.08
Ketha et al. (2021)	10	48	9	62		1.36 [0.59, 3.12]	7.02
Plattner et al. (2021)	9	106	11	124		0.96 [0.41, 2.24]	6.89
Anandalwar et al. (2021) after propensity matching	17	270	16	271		1.06 [0.55, 2.06]	8.88
Jen et al. (2022)	10	267	6	80		0.52 [0.19, 1.38]	5.72
Morita et al. (2022) after propensity matching	153	2,616	205	2,564		0.75 [0.61, 0.91]	15.27
Theodorou et al. (2022)	11	86	6	107		2.14 [0.82, 5.56]	5.92
Russell et al. (2023)	9	112	11	174		1.25 [0.53, 2.93]	6.84
Patwardhan et al. (2025)	12	146	3	61		1.62 [0.47, 5.55]	4.16
Overall						1.02 [0.73, 1.41]	

Heterogeneity: $t^2 = 0.13$, $I^2 = 68.78\%$, $H^2 = 3.20$

Test of $q_i = q_j$: $Q(11) = 28.70$, $p = 0.00$

Test of $q = 0$: $t(11) = 0.10$, $p = 0.92$

Random−effects REML model
Truncated Knapp Hartung standard errors

Omitted study		Risk ratio with 95% CI	p−value
Anderson et al. (2018)		1.09 [0.76, 1.56]	0.607
Ferguson et al. (2020)		0.87 [0.66, 1.13]	0.267
Gordon et al. (2020)		1.08 [0.77, 1.51]	0.641
Rossidis et al. (2020)		0.99 [0.70, 1.39]	0.944
Ketha et al. (2021)		1.00 [0.70, 1.41]	0.982
Plattner et al. (2021)		1.03 [0.72, 1.47]	0.876
Anandalwar et al. (2021) after propensity matching		1.02 [0.71, 1.46]	0.914
Jen et al. (2022)		1.06 [0.75, 1.49]	0.715
Morita et al. (2022) after propensity matching		1.08 [0.74, 1.56]	0.665
Theodorou et al. (2022)		0.97 [0.69, 1.34]	0.816
Russell et al. (2023)		1.00 [0.71, 1.43]	0.980
Patwardhan et al. (2025)		1.00 [0.71, 1.40]	0.980

Random−effects REML model
Truncated Knapp Hartung standard errors

Study	IG		CG			Risk ratio with 95% CI	Weight (%)
	Yes	No	Yes	No			
Gordon et al. (2020)	10	181	17	145		0.50 [0.24, 1.06]	5.73
Plattner et al. (2021)	9	106	11	124		0.96 [0.41, 2.24]	4.54
Anandalwar et al. (2021) after propensity matching	17	270	16	271		1.06 [0.55, 2.06]	7.39
Jen et al. (2022)	10	267	6	80		0.52 [0.19, 1.38]	3.36
Morita et al. (2022) after propensity matching	153	2,616	205	2,564		0.75 [0.61, 0.91]	78.98
Overall						0.75 [0.62, 0.90]	

Heterogeneity: $t^2 = 0.00$, $I^2 = 0.00\%$, $H^2 = 1.00$

Test of $q_i = q_j$: Q(4) = 3.07, p = 0.55

Test of q = 0: z = −3.16, p = 0.00

Random−effects REML model