Quantifying H5N1 outbreak potential and control effectiveness in high-risk

agricultural populations

Izel Avkan[1*], Suzanne Gokool[2], Louise E. Smith[3], Genevieve Clapp[1], Rachel Cox[4], Amy C.

Thomas[1], Ellen Brooks-Pollock[1]

[1] Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

[2] UKHSA, UK Health Security Agency, London, UK

[3] Behavioural Science and Insights Unit, UK Health Security Agency, London, UK

[4] School of Physiology, Pharmacology and Neuroscience, University of Bristol, Bristol, UK

[*] Corresponding author

E-mail: df23614@bristol.ac.uk

Abstract

Avian influenza is a global public health threat. Since 2021, the ongoing H5N1 panzootic has

brought a major shift in H5Nx epidemiology, including unprecedented spread, wide host

range and lack of seasonality. Infections in marine mammals, wildlife and livestock have

heightened concern for human-to-human transmission and pandemic potential. Contact

tracing and self-isolation are used as public health measures in the UK to manage contacts of

confirmed human cases of avian influenza. In this study, we aimed to estimate potential

outbreak sizes and evaluate the effectiveness of contact tracing and self-isolation in

managing community outbreaks of H5N1 following spillover from birds to people.

We characterised contact patterns from an underrepresented agricultural population at high

risk of avian influenza exposure through contact with birds (Avian Contact Study). Informed

by these realistic social contact data, we modelled outbreak sizes using a stochastic

branching process model.

Most simulations resulted in small-scale outbreaks, ranging from 0 to 10 cases. When the

basic reproduction number was 1.1, contact tracing and self-isolation reduced the average

outbreak size from 41 cases (95% Confidence Interval (CI): 37-46 cases) to 7 cases (95% CI: 6-

8 cases), preventing, on average, 8 out of every 10 infections. However, controls became less

effective in reducing the outbreak size when a higher proportion of cases were

asymptomatic. Overall, our findings suggest that contact tracing and self-isolation can be

effective at preventing zoonotic infections. Increasing awareness, encouraging self-isolation,

and detecting asymptomatic cases through routine surveillance are important components

of zoonotic infection containment strategies.

Introduction

Highly Pathogenic Avian Influenza (HPAI) continues to pose a threat to both animal and human public health. It is a highly contagious viral disease that affects both domestic and wild birds, with occasional spillover to mammals, including humans (1,2). The H5N1 virus was first detected in southern China in 1996, causing large poultry outbreaks in Hong Kong in 1997. The virus re-emerged in 2003 spreading among wild birds across Asia, and later in Africa, Europe, and the Middle East (3). In October 2020, the reassortment of poultry and wild bird viruses led to the emergence of a novel H5N1 strain of clade 2.3.4.4b in the Netherlands and spread across Europe (4). Between 2022 and 2024, this H5N1 variant spread to mammals—including foxes, seals, sea lions, and mink (5). In March 2024, the virus was detected for the first time in dairy cows in the USA with widespread cattle-to-cattle transmission and spillover to cats and humans (5). In March 2025, it was detected for the first time in a sheep in the UK (6). Given the continued evolution of the virus and its increasing ability to infect mammals, there is a growing concern about its potential for human-to-human transmission.

Avian influenza is a viral infection that can cause mild to severe illness, including death in humans. Since 1997, a total of 998 human cases has been reported in more than 23 countries, with approximately half resulting in death (3,7). Common symptoms in humans include conjunctivitis (eye redness), fever, and respiratory symptoms such as cough, sore throat, and shortness of breath (8). However, there is a big variability in the proportion of asymptomatic infections and the factors influencing symptomatology are not well understood due to limited human cases (8,9).

The current absence of sustained human-to-human transmission categorises the situation as

Phase 3 of the WHO pandemic phases (8,10,11). However, cases without zoonotic exposure

and limited human-to-human transmission have been documented (12,13). Data from family

clusters found evidence of human-to-human transmission with an estimated reproduction

number of 1.14 (13). Although the current global public health risk is low, reassortment

between avian and human seasonal influenza viruses could result in antigenic changes that

facilitate human-to-human transmission, potentially transitioning the situation to the

next phase (14). This emphasises the need for continuous surveillance and the

development of effective control strategies.

Contact tracing is a key public health strategy used in early-stage pandemic response to

control the spread of infectious diseases (15,16). Its success depends on factors such as a

small proportion of asymptomatic cases, long incubation periods, prompt identification of

contacts, and public willingness to share information (16). Currently, contact tracing and self-

isolation are used by UK public health teams to manage contacts of confirmed human cases

of avian influenza (17). Depending on the type of exposure, contacts are placed under either

active or passive follow-up for 10 days after their last exposure. Active follow-up involves

daily phone calls to monitor for symptoms, while passive follow-up involves providing

guidance on what to do if symptoms develop. If a contact develops symptoms, they are

assessed as a possible case and are advised to self-isolate while arrangements are made for

clinical assessment and investigation.

The accuracy of models investigating the effectiveness of contact tracing relies on

understanding contact networks, such as the number of contacts people have and with

whom. Several factors influence contact networks, such as occupation, residential setting

(urban or rural), and age (18,19). However, rural communities are often underrepresented in

social contact surveys (20,21). Given the global epizootic of H5N1 avian influenza over the

past 5 years, individuals working closely with birds are at increased risk of potential

exposure, yet their contact patterns remain poorly understood. Understanding their contact

patterns is crucial for assessing transmission dynamics.

In this paper, we used a novel dataset on contact patterns from an underrepresented UK

agricultural population at risk of exposure to avian influenza through contact with birds. We

applied a branching process model to simulate potential outbreak sizes for different

scenarios and assessed the effectiveness of contact tracing and self-isolation to manage

outbreaks of avian influenza among humans given known parameters for human-to-human

transmission.

Methods

Data description

We used data from the Avian Contact Study—a questionnaire open to individuals aged 18 or

over in the UK who have contact with domestic and/or wild birds to investigate the potential

transmission dynamics of HPAI and inform public health policies (22). Although the

questionnaire was open to people aged 18 and above, no individuals aged 18 responded, so

our dataset includes only those aged 19 and over. Respondents generally had high levels of

bird exposure: 20% reported flock sizes over 10000 birds, and more than 60% reported daily

contact with birds (22). The questionnaire was launched in-person at the British Pig and

Poultry Fair (15-16 May 2024) and subsequently circulated online via various poultry forums

and networks by word-of-mouth (15 May to 31 October 2024). The questionnaire was

divided into four sections:

Section A: "About you"

Section B: "Your contact with any type of domestic or wildlife birds"

Section C: "Your contact with other people"

Section D: "Your awareness of avian influenza (bird flu)"

Details of the eligibility criteria, questionnaire format, patient and public involvement, and

key results are described in Thomas AC, Gokool S, Clapp G et al. (2024) (22).

We extracted self-reported contact data with other people to inform branching process

models. Direct contact was defined as either physical contact (such as a handshake, hug,

kiss, or contact sport), conversational contact (exchanging at least a few words), or contact

by distance (being within 2 metres of another person). Respondents were asked how many

direct contacts they had in the last 24 hours, with the response options "1-5", "6-10", "11-

15", "16-20", and "More than 20". Respondents then provided further details about each

contact, such as relationship to the contact, age group of the contact, location, duration, and

type of contact (physical, conversational, or by distance). Although the categorical response

included an option of "More than 20 contacts", respondents could only provide further

information for up to twenty contacts in the following questions.

Additionally, we used contact data from CoMix, a large-scale contact survey started during

the COVID-19 pandemic and conducted in several European countries, including the UK, as a

comparator dataset from the general population (23). We took data relating to the final

survey round collected post-pandemic in November 2022. The data processing for CoMix is

described in the Supplementary Material.

Data analysis

Since the number of reported contacts was a categorical variable, we counted the number of

contacts listed by each participant to obtain an integer variable. This allowed calculation of

summary statistics, degree distributions, and informed the branching process model. We

used the 'Relationship to the contact' column to count the listed contacts, due to its lower

proportion of missing data. We then compared the number of listed contacts to the

reported contact category. No exclusions were made if the number of listed contacts did not

align with the reported contact category. The number of listed contacts was used in our

model.

We calculated the average number of contacts per person per day overall and stratified by

age group and compared these values to CoMix (23). Distributions of contacts were

calculated for three age groups (19-44, 45-59, 60+) representing young adults, adults, and

older adults, as well as for all age groups combined. We fitted negative binomial and Poisson

distributions to the data using the Maximum Likelihood Estimate method from the

'fitdistrplus' package version 1.2-1 in R and compared both fits using the likelihood ratio test.

Social networks commonly exhibit structural heterogeneity, and contact matrices are a

widely used method for representing contact heterogeneities across age groups (24,25). We

constructed age-stratified contact matrices to illustrate the average number of contacts

made between different age groups, both overall and stratified for family contacts. We

grouped ages into 10-year intervals to balance the level of detail without compromising

sample sizes in each age group.

Model framework

We used a stochastic branching process model to estimate the potential outbreak size and

effectiveness of contact tracing and self-isolation in preventing the spread of avian influenza

(Fig 1) (26–28).

We defined outbreak size as the number of individuals in a cluster including the index case

and analysed how it changed with the basic reproduction number, R_0, the average number

of secondary infections generated by an infected individual in a wholly susceptible

population (29). We varied R_0 from 0 to 1.1 in increments of 0.1, consistent with estimates

from previous studies on avian and pandemic influenza (13,30). To capture contact

heterogeneities with age, we divided the population into four age groups (0-18, 19-44, 45-

59, 60+). Since our dataset did not include individuals under 19 years old, we did not explore

scenarios where index cases were under 19, but they are included in transmission chains

using contact data from CoMix.

Fig 1. Contact Tracing Schematic. Example of a simulated transmission chain starting with person A, the index case, who was infected through contact with an infected bird. The number of secondary infections generated by each person is drawn from a negative binomial distribution. The time of infection for new cases is drawn from a gamma distribution, relative to the symptom onset of the infector. In this example, Person A infects 2 individuals, B and C, before isolating. If a contact is traced, they isolate on the day they develop symptoms and do not cause further infections. They may still be able to infect other people before showing symptoms. If a traced contact does not develop symptoms, or a contact is not traced, they would not isolate.

Number of secondary cases

The number of contacts made by an infected individual was drawn from an age-specific negative binomial distribution. The number of secondary infections (R) generated by that case was calculated as the product of the number of contacts (n), a fixed infectious period (d), and the probability of transmission ($\frac{R_0}{\lambda_c d}$, where R_0 is the basic reproduction number, λ_c is the dominant eigenvalue of the contact matrix). Mathematically, the number of secondary

infections generated by an individual in age group a (for age groups 19-44 years, 45-59 years

and 60+ years) is given by:

$$R_a = n_a \frac{R_0}{\lambda_c d} d, \text{ where } a \in \{19\text{-}44, 45\text{-}59, 60+\}$$

The age group of each secondary case was determined by sampling from a multinomial

distribution, using probabilities from the corresponding row of the contact matrix,

normalised to one. Time of infection for each secondary case relative to the symptom onset

of the primary case was drawn from a shifted Gamma distribution taken to reproduce

estimates from the literature (shape parameter 13, rate parameter 2, resulting in a mean of

3.5 days (SD: 1.80 days) between the symptom onset and infection) (8,13,29–32). We

adjusted the distribution to allow infections to happen up to one day prior to the onset of

symptoms.

Asymptomatic infections

The probability of being asymptomatic may depend on factors such as age and the strain of

the virus. However, due to limited data on human cases of H5N1 avian influenza, the

proportion of asymptomatic cases and the impact of age on this probability are not well

understood. For the baseline scenario, we assumed 40% of individuals developed

asymptomatic infections (8,9). However, as the proportion of asymptomatic infections has a

major impact on the effectiveness of contact tracing and self-isolation, we explored a full

range of scenarios from 0% of infections being asymptomatic to 100%

Contact tracing and self-isolation

We investigated the impact of contact tracing and self-isolation on the outbreak size. We

assumed that 40% of contacts were traced in the baseline scenario, based on an estimate

from reported figures during the COVID-19 pandemic (33). We determined whether new

cases would be traced based on their symptom status. By default, the index case and the

contacts of asymptomatic cases were not traced. Contacts of symptomatic cases could be

traced with probabilities varying from 0 (no contact traced) to 1 (all contacts traced). For the

purposes of this model, we defined self-isolation as the separation from others of

symptomatic individuals who had been traced by public health teams, beginning on the day

of symptom onset, in accordance with current guidelines. Traced contacts who do not

develop symptoms, and untraced contacts, would not self-isolate.

We ran the model 9000 times per parameter set with the maximum time for each simulation

of 100 days.

We used R version 4.4.3 in our analyses.

Ethics Statement

211 Ethical approval for the study was obtained from the University of Bristol, Faculty of Health

212 Sciences Research Ethics Committee, approval number 17048 on 16 January 2024.

213 Informed written consent (using e-consent hosted on REDCap) for the use of data collected

214 via the questionnaire was obtained from respondents. A copy of the consent form and

215 participant information sheet is given in the accompanying underlying data.

Results

A total of 268 respondents completed the questionnaire between May and October 2024.

We excluded 31 respondents who did not report a categorical number of contacts, and 40

who selected a category but did not provide further details, as we could not obtain an

integer value from their responses. Among the 197 respondents who provided data on the

categorical question about number of contacts and who listed individual contacts, 82% were

consistent.

Contact patterns

The distribution of numbers of contacts is shown in Fig 2A. The average number of contacts

was 4.64 (standard deviation (SD): 3.40). A Negative Binomial (NB) model provided a better

fit than a Poisson model based on Akaike Information Criterion (NB: 968, Poisson: 1052),

Bayesian Information Criterion (NB: 974, Poisson: 1056), and likelihood ratio test (p-value <

0.001). The final NB model had a mean of 4.64 and a dispersion parameter of 4.02.

Additionally, we fitted NB distributions to the number of contacts stratified by age group to

inform the branching process model, shown in S2 Fig.

Contact matrices suggest intergenerational living in agricultural workers

We constructed age-structured contact matrices to show the average number of contacts

made between different age groups, both for all contacts and family members (Fig 2B, C).

We observe broadly similar patterns to previous contact surveys, with respondents

interacting more frequently with individuals of a similar age. Respondents aged 19-29

reported an average of 6.8 contacts per day, with nearly 90% of their contacts occurring with

individuals aged 20-59. Respondents aged 30 and above had an average of 4.4 contacts per

day.

Fig 2C illustrates contact patterns among family members. Respondents reported an average

of 1.8 family contacts per day, with the 19-29 age group having the lowest (mean: 1.35

contacts, SD: 0.18 contacts) and the 40-49 age group the highest (mean: 1.97 contacts, SD:

0.14 contacts). We observed three distinct diagonal patterns: younger individuals frequently

interacting with their parents, older individuals with their children and grandchildren, and

individuals interacting with those of a similar age, reflecting intergenerational living

arrangements.

Fig 2. Contact patterns. Contact patterns reported in the Avian Contact Study. **(A)** the distribution of the

number of contacts per person. The red line indicates the fitted negative binomial distribution, and the pink

region represents the 95% confidence interval. The fitted negative binomial distribution had a mean of 4.64

and a dispersion parameter of 4.02. **(B)** Average number of contacts between age groups, while **(C)** Average

number of contacts between family members only. In both **(B)** and **(C)**, darker colours indicate higher average

contacts.

A comparison of the Avian Contact Study and CoMix is shown in Fig 3. Respondents in the

Avian Contact Study had higher average contact levels than respondents of CoMix across all

age groups. The smaller sample size in the Avian Contact Study compared to CoMix may

explain the wider confidence intervals around the mean. Statistical comparisons of the

contact patterns are provided in the Supplementary Materials.

Fig 3. Average number of contacts from the Avian Contact Study (AvCos) and CoMix for all age groups. Black

lines indicate the 95% confidence interval, and grey error bars indicate the interquartile range (IQR).

Influence of age on outbreak size

As expected, outbreak size was highly dependent on the reproduction number. Infections

with reproduction numbers less than 0.2 did not lead to onward transmission. Over 75% of

simulations resulted in small-scale outbreaks (0-10 cases). The average outbreak size

increased sharply from 4 cases at R_0=1 to 41 cases at R_0=1.1 (Fig S3).

We also examined how the age group of the index case influenced outbreak dynamics. For

R_0 between 0.7 and 1, half of simulations with index cases aged 60+ resulted in no

secondary cases, compared to 40% of simulations for those aged 19-59. When $R_0 = 1.1$, 9%

of simulations with index cases in the 19-59 age group led to outbreaks larger than 50 cases,

compared to 6% for those with index cases aged 60+.

For index cases in the 19-44 and 45-59 age groups, R_0 did not significantly affect the age

distribution of the cases. In contrast, for index cases aged 60+, we observed an increase in

the percentage of cases in the 0-18 age group from 0 to 13% as R_0 increased. This is due to

limited direct contact between the 60+ and 0-18 age groups, meaning individuals aged 0-18

were infected further down the transmission chain through interactions with other age

groups (Figs S3 and S4).

Reduced effectiveness of contact tracing and self-isolation with

higher proportion of asymptomatic infections

We compared how outbreak sizes changed when 40% of infections were asymptomatic,

under both no control and control (contact tracing and self-isolation) scenarios, depending

on R_0 (Fig 4A). Under no control, no secondary cases were observed when R_0 was below 0.2

- the only case was the index case. Over 75% of simulations resulted in small-scale outbreaks

(less than 10 cases). Contact tracing and self-isolation reduced the maximum outbreak size

from 5705 cases to below 1000, and the average outbreak size from 41 cases to below 10,

when R_0=1.1.

We also evaluated how the average outbreak size changed under different levels of control

measures and varying proportions of asymptomatic infections (Fig 4B). When the proportion

of asymptomatic infections was low, even low to moderate levels of contact tracing and self-

isolation (0 to 40%) were sufficient to reduce the average outbreak size from 41 cases to

below 10. However, the reduction in the average outbreak size became less pronounced as

the proportion of asymptomatic infections increased.

Fig 4. The impact of contact tracing and self-isolation on final outbreak size. (A) The distribution of outbreak

sizes for varying levels of R_0 under no control and control (contact tracing and self-isolation). The orange

line indicates the average outbreak size, and the shaded region represents the 95% prediction interval. We

assumed that 40% of cases were asymptomatic and that 40% of contacts were traced. Among those traced, all

symptomatic cases were assumed to self-isolate. **(B)** The average outbreak size across 9000 simulations for

varying levels of contact tracing and proportions of asymptomatic infections when R_0 = 1.1. Shaded regions

represent the 95% confidence interval.

Discussion

Principal findings

Our model provides insights into the potential outbreak size of avian influenza among

humans under different levels of transmission. Most simulations resulted in small-scale

outbreaks, with outbreaks exceeding 100 cases when the basic reproduction number was

above 0.8, consistent with previous estimates (34). Contact tracing and self-isolation

reduced outbreak size, but their potential effectiveness in preventing cases decreased as the

proportion of asymptomatic infections increased, since they rely on symptomatic cases.

Previous modelling studies similarly concluded that contact tracing and self-isolation alone

was highly unlikely to prevent large outbreaks, particularly when subclinical infections were

present (28,35). Our findings support and extend these conclusions to a different pathogen.

Translating findings into action

Asymptomatic infections limit the effectiveness of contact tracing and self-isolation, and

they are optimally identified and controlled through asymptomatic testing programmes,

which have previously shown to be successful at detecting cases (36). Reintroducing such

programmes could enhance current interventions by improving detection of asymptomatic

infections.

Moreover, we observed that the index cases aged 19-59 play an important role in

transmission dynamics, leading to larger outbreaks based solely on contact data.

Additionally, analysis of previous H5N1 infections suggests that the infection is more

prevalent among younger adults, and older age may not be a risk factor for severe outcomes

(37). Taken together, these findings suggest that younger adults may benefit more from

targeted interventions.

Strengths and limitations

We used a novel dataset from the Avian Contact Study, which focused on individuals at

higher risk of avian influenza and was developed with the help of public contributors to

ensure relevance and accessibility (22). To our knowledge, it is the only contact survey

conducted in the UK that has focused on farming communities since the COVID-19

pandemic. This allowed us to examine post-pandemic contact patterns in a rural, high-risk

groups, often underrepresented in contact surveys, who may have different mixing patterns

compared to the general population. We used age-stratified mixing patterns in our model to

estimate outbreak sizes, extending previous models that did not incorporate age structure.

This allowed us to capture variation in outbreak sizes across age groups and identify which

groups were more likely to drive larger outbreaks.

In addition to common limitations of other contact surveys (21,38), our data did not include

individuals under the age of 19, so we could not compare their contact patterns to those of

the general population. Although we included 0-18 age group from CoMix in the

transmission chain, this may not fully reflect the contact patterns due to differences in the

cohorts and study periods. Additionally, our analysis was based on data from individuals

living in the UK, most of whom were poultry farmers, hence our findings may not be

generalised to other countries or to farming communities with different livestock practices.

Although we included age in our model, we could not explore its relationship with all the

parameters. For instance, the influence of age on symptomatology is not well understood. As

a result, we kept the proportion of asymptomatic infections constant across all age groups.

Our model assumed that people with symptoms would isolate on the same day, although

this may not be the case in practice with people adopting a 'wait and see' approach (39).

Areas for future work

356 Future research is required to further characterise contact patterns within agricultural

357 demographics and other related groups. In addition to farmers who make up 60% of the

358 agricultural workforce in the UK, 12% is covered by casual workers (40). Despite their high

359 exposure risk, it can be challenging to engage with this group due to language and trust

360 barriers.

Funding statement

Funding for the Avian Contact Study was awarded by PolicyBristol from the Research

England QR Policy Support Fund (QR PSF) 2022-24 for investigating 'Zoonotic spillover of

avian influenza'. AT is funded by the Wellcome Trust, Early Career

Award [227041/Z/23/Z]. EBP and IA acknowledges support from the National Institute for

Health Research Health Protection Research Unit (NIHR HPRU) in Evaluation and Behavioural

Science at the University of Bristol (NIHR207385).

Acknowledgements

We are grateful to all respondents who completed the questionnaire and to the public

contributors for their input into study design and questionnaire development. We thank all

members of the wider Avian Contact Study team for input into study design, data collection

and interpretation. We thank Dr Anastasia Chatzilena (University of Bristol) for her guidance

regarding the statistical analysis of contact data from the Avian Contact Study and CoMix.

Data and code availability

Underlying data

Repository data.bris: The Avian Contact Study: questionnaire data 15 May – 31 July
2024. Data are openly available at the University of Bristol Research Data
Repository data.bris, at https://doi.org/10.5523/bris.3nmqsrbv5ruom2abn0ql6e8yh2
(41).

This project contains the following underlying data:

- Data file 1. (Raw underlying questionnaire data – csv file)
- Data file 2. (Raw underlying questionnaire data - .RDS file)
- Data file 3. (Associated data dictionary – csv file)
- Data file 4. (Code for importing underlying data in csv format into R for setting
up labelled data - .r file)
- Data file 5. (Blank consent form and participant information sheet – pdf file)

Data are available under the terms of National Archives' Non-Commercial

Government Licence for public sector information.

Extended data

Repository Zenodo: The Avian Contact Study Questionnaire and Data Dictionary
[10.5281/zenodo.13617061]
This project contains the following extended data:

- AvianInfluenzaSocialContactSu.pdf (The final questionnaire REDCap – PDF)
- AvianInfluenzaSocialContactSur_DataDictionary_080824v1.csv (Associated
data dictionary -csv file)

Data are available under the terms of the <u>Creative Commons Attribution 4.0</u>

<u>International license</u> (CC-BY 4.0).

Software availability

Source code available from: <u>https://github.com/amythomas/aviancontactstudy</u>

License: <u>Creative Commons Attribution 4.0 International license</u> (CC-BY 4.0).

References:

1. WOAH. Avian Influenza [Internet]. WOAH - World Organisation for Animal Health. Available from: https://www.woah.org/en/disease/avian-influenza/

2. World Health Organisation. https://www.who.int/emergencies/disease-outbreak-news/item/2025-DON575. 2025. Avian Influenza A(H5N1) - Cambodia.

3. CDC. Past Reported Global Human Cases with Highly Pathogenic Avian Influenza A(H5N1) (HPAI H5N1) by Country, 1997-2025 [Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-flu/php/avian-flu-summary/chart-epi-curve-ah5n1.html

4. Graziosi G, Lupini C, Catelli E, Carnaccini S. Highly Pathogenic Avian Influenza (HPAI) H5 Clade 2.3.4.4b Virus Infection in Birds and Mammals. Animals [Internet]. 2024 Jan;14(9):1372. Available from: https://www.mdpi.com/2076-2615/14/9/1372

5. CDC. 2020-2024 Highlights in the History of Avian Influenza (Bird Flu) Timeline [Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-flu/avian-timeline/2020s.html

6. Mahase E. H5N1: UK reports world's first case in a sheep. BMJ. 2025 Mar;388:r591.

7. CDC. Highly Pathogenic Avian Influenza A(H5N1) Virus: Interim Recommendations for Prevention, Monitoring, and Public Health Investigations [Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-flu/prevention/hpai-interim-recommendations.html

8. Garg S, Reinhart K, Couture A, Kniss K, Davis CT, Kirby MK, et al. Highly Pathogenic Avian Influenza A(H5N1) Virus Infections in Humans. New England Journal of

Medicine [Internet]. 2025 Feb;392(9):843–54. Available from:

https://www.nejm.org/doi/full/10.1056/NEJMoa2414610

9.   CDC. Technical Report: June 2024 Highly Pathogenic Avian Influenza A(H5N1) Viruses

[Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-

flu/php/technical-report/h5n1-06052024.html

10.   NHS. https://www.nhs.uk/conditions/bird-flu/. 2022. Bird flu.

11.   Geneva: World Health Organization. Pandemic Influenza Preparedness and Response:

A WHO Guidance Document. [Internet]. 2009 [cited 2025 Jul 24]. Available from:

https://www.ncbi.nlm.nih.gov/books/NBK143061/

12.   CDC. H5 Bird Flu: Current Situation [Internet]. Avian Influenza (Bird Flu). 2025.

Available from: https://www.cdc.gov/bird-flu/situation-summary/index.html

13.   Yang Y, Halloran ME, Sugimoto JD, Longini IM. Detecting Human-to-Human

Transmission of Avian Influenza A (H5N1). Emerg Infect Dis [Internet]. 2007

Sep;13(9):1348–53. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2857285/

14.   World Health Organisation. https://www.who.int/publications/m/item/updated-joint-

fao-who-woah-assessment-of-recent-influenza-a(h5n1)-virus-events-in-animals-and-

people_dec2024. 2020. Updated joint FAO/WHO/WOAH public health assessment of

recent influenza A(H5) virus events in animals and people.

15.   Brandt AM. The History of Contact Tracing and the Future of Public Health. Am J

Public Health [Internet]. 2022 Aug;112(8):1097–9. Available from:

https://ajph.aphapublications.org/doi/full/10.2105/AJPH.2022.306949

16.   El-Sadr WM, Platt J, Bernitz M, Reyes M. Contact Tracing: Barriers and Facilitators. Am

J Public Health [Internet]. 2022 Jul;112(7):1025–33. Available from:

https://ajph.aphapublications.org/doi/full/10.2105/AJPH.2022.306842

17.   UKHSA. Investigation and initial clinical management of possible human cases of avian

influenza with potential to cause severe human disease [Internet]. GOV.UK. Available

from: https://www.gov.uk/government/publications/avian-influenza-guidance-and-

algorithms-for-managing-human-cases/investigation-and-initial-clinical-management-

of-possible-human-cases-of-avian-influenza-with-potential-to-cause-severe-human-

disease

18.   Thomas A, Danon L, Christensen H, Northstone K, Smith D, Nixon E, et al. Limits of

lockdown: characterising essential contacts during strict physical distancing. 2021

Mar; Available from:

https://www.medrxiv.org/content/10.1101/2021.03.12.21253484v1

19.   McCreesh N, Morrow C, Middelkoop K, Wood R, White RG. Estimating age-mixing

patterns relevant for the transmission of airborne infections. Epidemics [Internet].

2019 Sep;28:100339. Available from:

https://www.sciencedirect.com/science/article/pii/S1755436518301646

20.   Edelman LS, Yang R, Guymon M, Olson LM. Survey Methods and Response Rates

Among Rural Community Dwelling Older Adults. Nurs Res [Internet]. 2013

Aug;62(4):286. Available from:

https://journals.lww.com/nursingresearchonline/FullText/2013/07000/Survey_Metho

ds_and_Response_Rates_Among_Rural.10.aspx

21.   Hoang T, Coletti P, Melegaro A, Wallinga J, Grijalva CG, Edmunds JW, et al. A

Systematic Review of Social Contact Surveys to Inform Transmission Models of Close-

contact Infections. Vol. 30, Epidemiology. Lippincott Williams and Wilkins; 2019. p.

723–36.

22.   Thomas AC, Gokool S, Clapp G, Whitlow H, McGrath C, Moore P, et al. Data note on

the Avian Contact Study: a questionnaire resource for avian influenza public health

planning. Wellcome Open Res. 2025;9.

23.   Jarvis CI, Van Zandvoort K, Gimma A, Prem K, Auzenbergs M, O'Reilly K, et al.

Quantifying the impact of physical distance measures on the transmission of COVID-

19 in the UK. BMC Med. 2020 May;18.

24.   Koltai J, Vásárhelyi O, Röst G, Karsai M. Reconstructing social mixing patterns via

weighted contact matrices from online and representative surveys. Sci Rep. 2022

Dec;12.

25.   Dall'Amico L, Kleynhans J, Gauvin L, Tizzoni M, Ozella L, Makhasi M, et al. Estimating

household contact matrices structure from easily collectable metadata. PLoS One.

2024 Mar;19.

26.   Farrington CP, Grant AD. The distribution of time to extinction in subcritical branching

processes: Applications to outbreaks of infectious disease. J Appl Probab.

1999;36:771–9.

27.   Feng J, Zhu W, Ye X, Liu Z, Zhu Y, Wu Q, et al. Effects of contact tracing and nucleic acid

testing on the COVID-19 outbreak in Zunyi, China: data-driven study using a branching

process model. BMC Infect Dis. 2023 Jan;23:10.

28.   Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of

controlling COVID-19 outbreaks by isolation of cases and contacts. Lancet Glob Health

[Internet]. 2020 Apr;8(4):e488–e496. Available from:

https://www.thelancet.com/article/S2214-109X(20)30074-7/fulltext

29.   Ungchusak K, Auewarakul P, Dowell SF, Kitphati R, Auwanit W, Puthavathana P, et al.

505   Probable Person-to-Person Transmission of Avian Influenza A (H5N1). New England

506   Journal of Medicine. 2005 Jan;352:333–40.

507   30.   Wang H, Feng Z, Shu Y, Yu H, Zhou L, Zu R, et al. Probable limited person-to-person

508   transmission of highly pathogenic avian influenza A (H5N1) virus in China. The Lancet.

509   2008;371:1427–34.

510   31.   Cowling BJ, Jin L, Lau EHY, Liao Q, Wu P, Jiang H, et al. Comparative epidemiology of

511   human infections with avian influenza A H7N9 and H5N1 viruses in China: A

512   population-based study of laboratory-confirmed cases. The Lancet. 2013;382:129–37.

513   32.   Aditama TY, Samaan G, Kusriastuti R, Sampurno OD, Purba W, Misriyah, et al. Avian

514   influenza H5N1 transmission in households, Indonesia. PLoS One. 2012 Jan;7.

515   33.   Department of Health and Social Care. Chapter 7: contact tracing and isolation. 2023

516   Jan.

517   34.   Ward J, Lambert JW, Russell TW, Azam JM, Kucharski AJ, Funk S, et al. Estimates of

518   epidemiological parameters for H5N1 influenza in humans: a rapid review. 2024.

519   35.   Davis EL, Lucas TCD, Borlase A, Pollington TM, Abbott S, Ayabina D, et al. Contact

520   tracing is an imperfect tool for controlling COVID-19 transmission and relies on

521   population adherence. Nat Commun [Internet]. 2021 Sep;12(1):5412. Available from:

522   https://www.nature.com/articles/s41467-021-25531-5

523   36.   Capelastegui F, Smith J, Kumbang J, Humphreys C, Padfield S, Turner J, et al. Pilot of

524   asymptomatic swabbing of humans following exposures to confirmed avian influenza

525   A(H5) in avian species in England, 2021/2022. Influenza Other Respir Viruses. 2023

526   Aug;17.

527   37.   Garretson TA, Liu J, Li SH, Scher G, Santos JJS, Hogan G, et al. Immune history shapes

528   human antibody responses to H5N1 influenza viruses. Nat Med. 2025;

529   38.   Nixon E, Silvonen T, Barreaux A, Kwiatkowska R, Trickey A, Thomas A, et al. A mixed

530   methods analysis of participation in a social contact survey. Epidemics. 2022 Dec;41.

531   39.   Mowbray F, Woodland L, Smith LE, Amlôt R, Rubin GJ. Is My Cough a Cold or Covid? A

532   Qualitative Study of COVID-19 Symptom Recognition and Attitudes Toward Testing in

533   the UK. Front Public Health. 2021 Aug;9.

534   40.   https://www.gov.uk/government/statistics/agricultural-workforce-in-england-at-1-

535   june/agricultural-workforce-in-england-at-1-june-2023 [Internet]. Agricultural

536   workforce in England at 1 June 2024.

537   41.   Thomas A, Brooks-Pollock E. Data collected from the Avian Contact Study

538   (https://bristol.ac.uk/avian-contact) questionnaire between 15 May 2024 and 31 July

539   2024. First data release. [Internet]. data.bris. 2024 [cited 2024 Sep 16]. Available

540   from: https://data.bris.ac.uk/data/dataset/3nmqsrbv5ruom2abn0ql6e8yh2

545