Controlling avian influenza spillover events: a modelling study

Izel Avkan[1*], Suzanne Gokool[2], Louise E. Smith[3], Genevieve Clapp[1], Rachel Cox[4], Amy C.

Thomas[1], Ellen Brooks-Pollock[1]

[1] Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

[2] UKHSA, UK Health Security Agency, London, UK

[3] Behavioural Science and Insights Unit, UK Health Security Agency, London, UK

[4] School of Physiology, Pharmacology and Neuroscience, University of Bristol, Bristol,

UK

[*]Corresponding author

E-mail: df23614@bristol.ac.uk

Abstract

Avian influenza is a highly contagious viral disease that affects both domestic and wild

birds, with occasional spillover to mammals, including humans. As of June 2025, 117

H5N1 infections in humans have been reported worldwide since 2020. Given the ability

of the virus to infect mammals, there is a growing concern about its potential for

human-to-human transmission. Currently, contact tracing and self-isolation are used in

the UK to manage contacts of confirmed human cases of avian influenza. In this study,

we aimed to estimate potential outbreak sizes and evaluate the effectiveness of contact

tracing and self-isolation in managing avian influenza spillover events.

We used a novel dataset from the Avian Contact Study to analyse contact patterns

within an underrepresented agricultural population at high risk of avian influenza

exposure through contact with birds. We modelled outbreak sizes using a stochastic

branching process model with measured contact data.

Most simulations resulted in small-scale outbreaks, ranging from 0 to 10 cases. When

the basic reproduction number was 1.1, contact tracing and self-isolation reduced the

average outbreak size from 41 cases (95% Confidence Interval (CI): 37-46 cases) to 7

cases (95% CI: 6-8 cases), preventing, on average, 8 out of every 10 infections. However,

they became less effective in reducing the outbreak size when a higher proportion of

cases were asymptomatic. Overall, our findings suggest that contact tracing and self-

isolation can be effective at preventing zoonotic infections. Increasing awareness,

encouraging self-isolation, and detecting asymptomatic cases through routine

surveillance are important components of zoonotic infection containment strategies.

Introduction

Highly Pathogenic Avian Influenza (HPAI) continues to pose a threat to both animal and human public health. It is a highly contagious viral disease that affects both domestic and wild birds, with occasional spillover to mammals, including humans (1,2). The H5N1 virus was first detected in southern China in 1996, causing large poultry outbreaks in Hong Kong in 1997. The virus re-emerged in 2003 spreading among wild birds across Asia, and later in Africa, Europe, and the Middle East (3). In October 2020, the reassortment of poultry and wild bird viruses led to the emergence of a novel H5N1 strain of clade 2.3.4.4b in the Netherlands and spread across Europe (4). Between 2022 and 2024, this H5N1 variant spread to mammals—including foxes, seals, sea lions, and mink (5). In March 2024, the virus was detected for the first time in dairy cows in the USA with widespread cattle-to-cattle transmission and spillover to cats and humans (5). In March 2025, it was detected for the first time in a sheep in the UK (6). Given the continued evolution of the virus and its increasing ability to infect mammals, there is a growing concern about its potential for human-to-human transmission.

Avian influenza is a viral infection that can cause mild to severe illness, including death in humans. Since 1997, a total of 998 human cases have been reported in more than 23 countries, with approximately half resulting in death (3,7). Common symptoms in humans include conjunctivitis (eye redness), fever, and respiratory symptoms such as cough, sore throat, and shortness of breath (8). However, there is a big variability in the proportion of asymptomatic infections and the factors influencing symptomatology are not well understood due to limited human cases (8,9).

Currently, avian influenza transmission is not sustained in human populations (8,10).

However, cases without zoonotic exposure and limited human-to-human transmission

have been documented (11,12). Data from family clusters found evidence of human-to-

human transmission with an estimated reproduction number of 1.14 (12). Although the

current global public health risk is low, reassortment between avian and human

influenza viruses could result in antigenic changes that facilitate human-to-human

transmission, with epidemic or pandemic potential (13). This emphasises the need for

continuous surveillance and the development of effective control strategies.

Contact tracing is a key public health strategy used in early-stage pandemic response to

control the spread of infectious diseases (14,15). Its success depends on factors such

as a small proportion of asymptomatic cases, long incubation periods, prompt

identification of contacts, and public willingness to share information (15). Currently,

contact tracing and self-isolation are used by UK public health teams to manage

contacts of confirmed human cases of avian influenza (16). Depending on the type of

exposure, contacts are placed under either active or passive follow-up for 10 days after

their last exposure. Active follow-up involves daily phone calls to monitor for symptoms,

while passive follow-up involves providing guidance on what to do if symptoms develop.

If a contact develops symptoms, they are assessed as a possible case and are advised

to self-isolate while arrangements are made for clinical assessment and investigation.

The accuracy of models investigating the effectiveness of contact tracing relies on

understanding contact networks, such as the number of contacts people have and with

whom. Several factors influence contact networks, such as occupation, residential

setting (urban or rural), and age (17,18). However, rural communities are often

underrepresented in social contact surveys (19,20). Given the global epizootic of H5N1

avian influenza over the past 5 years, individuals working closely with birds are at

increased risk of potential exposure, yet their contact patterns remain poorly

understood. Understanding their contact patterns is crucial for assessing transmission

dynamics.

In this paper, we used a novel dataset on contact patterns from an underrepresented UK

agricultural population at risk of exposure to avian influenza through contact with birds.

We applied a branching process model to simulate potential outbreak sizes for different

scenarios and assessed the effectiveness of contact tracing and self-isolation to

manage outbreaks of avian influenza among humans given known parameters for

human-to-human transmission.

Methods

Data description

We used data from the Avian Contact Study—a questionnaire open to individuals aged

18 or over in the UK who have contact with domestic and/or wild birds to investigate the

potential transmission dynamics of HPAI and inform public health policies (21).

Although the questionnaire was open to people aged 18 and above, no individuals aged

18 responded, so our dataset includes only those aged 19 and over. The questionnaire

was launched in-person at the British Pig and Poultry Fair (15-16 May 2024) and

subsequently circulated online via various poultry forums and networks by word-of-

mouth (May to October 2024). The questionnaire was divided into four sections:

Section A: "About you"

Section B: "Your contact with any type of domestic or wildlife birds"

Section C: "Your contact with other people"

Section D: "Your awareness of avian influenza (bird flu)"

Details of the eligibility criteria, questionnaire format, patient and public involvement,

and key results are described in Thomas AC, Gokool S, Clapp G et al. (2024) (21).

We extracted self-reported contact data with other people to inform branching process

models. Direct contact was defined as either physical contact (such as a handshake,

hug, kiss, or contact sport), conversational contact (exchanging at least a few words), or

contact by distance (being within 2 metres of another person). Respondents were asked

how many direct contacts they had in the last 24 hours, with the response options "1-5",

"6-10", "11-15", "16-20", and "More than 20". Respondents then provided further details

about each contact, such as relationship to the contact, age group of the contact,

location, duration, and type of contact (physical, conversational, or by distance).

Although the categorical response included an option of "More than 20 contacts",

respondents could only provide further information for up to twenty contacts in the

following questions.

Additionally, we used contact data from CoMix, a large-scale contact survey started

during the COVID-19 pandemic and conducted in several European countries, including

the UK, as a comparator dataset from the general population (22). We took data relating

to the final survey round collected post-pandemic in November 2022. The data

processing for CoMix is described in the Supplementary Material.

Data analysis

Since the number of reported contacts was a categorical variable, we counted the

number of contacts listed by each participant to obtain an integer variable. This allowed

calculation of summary statistics, degree distributions, and informed the branching

process model. We used the 'Relationship to the contact' column to count the listed

contacts, due to its lower proportion of missing data. We then compared the number of

listed contacts to the reported contact category. No exclusions were made if the

number of listed contacts did not align with the reported contact category. The number

of listed contacts was used in our model.

We calculated the average number of contacts per person per day overall and stratified

by age group and compared these values to CoMix (22). Distributions of contacts were

calculated for three age groups (19-44, 45-59, 60+) representing young adults, adults,

and older adults, as well as for all age groups combined. We fitted negative binomial

and Poisson distributions to the data using the Maximum Likelihood Estimate method

from the 'fitdistrplus' package version 1.2-1 in R and compared both fits using the

likelihood ratio test.

Social networks commonly exhibit structural heterogeneity, and contact matrices are a

widely used method for representing contact heterogeneities across age groups (23,24).

We constructed age-stratified contact matrices to illustrate the average number of

contacts made between different age groups, both overall and stratified for family

contacts. We grouped ages into 10-year intervals to balance the level of detail without

compromising sample sizes in each age group.

Model framework

We used a stochastic branching process model to estimate the potential outbreak size

and effectiveness of contact tracing and self-isolation in preventing the spread of avian

influenza (Fig 1) (25–27).

We defined outbreak size as the number of individuals in a cluster including the index

case and analysed how it changed with the basic reproduction number, R_0, the average

number of secondary infections generated by an infected individual in a wholly

susceptible population (28). We varied R_0 from 0 to 1.1 in increments of 0.1, consistent

with estimates from previous studies on avian and pandemic influenza (12,29). To

capture contact heterogeneities with age, we divided the population into four age

groups (0-18, 19-44, 45-59, 60+). Since our dataset did not include individuals under 19

years old, we did not explore scenarios where index cases were under 19, but they are

included in transmission chains using contact data from CoMix.

Fig 1. Contact Tracing Schematic. Example of a simulated transmission chain starting with person A, the

index case, who was infected through contact with an infected bird. The number of secondary infections

generated by each person is drawn from a negative binomial distribution. The time of infection for new

cases is drawn from a gamma distribution, relative to the symptom onset of the infector. In this example,

Person A infects 2 individuals, B and C, before isolating. If a contact is traced, they isolate on the day they

develop symptoms and do not cause further infections. They may still be able to infect other people

before showing symptoms. If a traced contact does not develop symptoms, or a contact is not traced,

they would not isolate.

171   Number of secondary cases

The number of contacts made by an infected individual was drawn from an age-specific

negative binomial distribution. The number of secondary infections (R) generated by

that case was calculated as the product of the number of contacts (n), a fixed infectious

period (d), and the probability of transmission ($\frac{R_0}{\lambda_c d}$, where R_0 is the basic reproduction

number, λ_c is the dominant eigenvalue of the contact matrix). Mathematically, the

number of secondary infections generated by an individual in age group a (for age

groups 19-44 years, 45-59 years and 60+ years) is given by:

179
$$R_a = n_a \frac{R_0}{\lambda_c d} d, \text{ where } a \in \{19\text{-}44, 45\text{-}59, 60+\}$$

The age group of each secondary case was determined by sampling from a multinomial

distribution, using probabilities from the corresponding row of the contact matrix,

normalised to one. Time of infection for each secondary case relative to the symptom

onset of the primary case was drawn from a shifted Gamma distribution taken to

reproduce estimates from the literature (shape parameter 13, rate parameter 2,

resulting in a mean of 3.5 days (SD: 1.80 days) between the symptom onset and

infection) (8,12,30–33). We adjusted the distribution to allow infections to happen up to

one day prior to the onset of symptoms.

188   Asymptomatic infections

The probability of being asymptomatic may depend on factors such as age and the

strain of the virus. However, due to limited data on human cases of H5N1 avian

influenza, the proportion of asymptomatic cases and the impact of age on this

probability are not well understood. For the baseline scenario, we assumed 40% of

individuals developed asymptomatic infections (8,9). However, as the proportion of

asymptomatic infections has a major impact on the effectiveness of contact tracing and

self-isolation, we explored a full range of scenarios from 0% of infections being

asymptomatic to 100%

Contact tracing and self-isolation

We investigated the impact of contact tracing and self-isolation on the outbreak size.

We assumed that 40% of contacts were traced in the baseline scenario, based on an

estimate from reported figures during the COVID-19 pandemic (34). We determined

whether new cases would be traced based on their symptom status. By default, the

index case and the contacts of asymptomatic cases were not traced. Contacts of

symptomatic cases could be traced with probabilities varying from 0 (no contact traced)

to 1 (all contacts traced). For the purposes of this model, we defined self-isolation as

the separation from others of symptomatic individuals who had been traced by public

health teams, beginning on the day of symptom onset, in accordance with current

guidelines. Traced contacts who do not develop symptoms, and untraced contacts,

would not self-isolate.

We ran the model 9000 times per parameter set with the maximum time for each

simulation of 100 days.

We used R version 4.4.3 in our analyses.

Results

A total of 268 respondents completed the questionnaire between May and October

2024. We excluded 31 respondents who did not report a categorical number of

contacts, and 40 who selected a category but did not provide further details, as we

could not obtain an integer value from their responses. Among the 197 respondents

who provided data on the categorical question about number of contacts and who

listed individual contacts, 82% were consistent.

Contact patterns

The distribution of numbers of contacts is shown in Fig 2A. The average number of

contacts was 4.64 (standard deviation (SD): 3.40). A Negative Binomial (NB) model

provided a better fit than a Poisson model based on Akaike Information Criterion (NB:

968, Poisson: 1052), Bayesian Information Criterion (NB: 974, Poisson: 1056), and

likelihood ratio test (p-value < 0.001). The final NB model had a mean of 4.64 and a

dispersion parameter of 4.02. Additionally, we fitted NB distributions to the number of

contacts stratified by age group to inform the branching process model, shown in S2

Fig.

Contact matrices suggest intergenerational living in agricultural workers

We constructed age-structured contact matrices to show the average number of

contacts made between different age groups, both for all contacts and family members

(Fig 2B, C). We observe broadly similar patterns to previous contact surveys, with

respondents interacting more frequently with individuals of a similar age. Respondents

aged 19-29 reported an average of 6.8 contacts per day, with nearly 90% of their

contacts occurring with individuals aged 20-59. Respondents aged 30 and above had an

average of 4.4 contacts per day.

Fig 2C illustrates contact patterns among family members. Respondents reported an

average of 1.8 family contacts per day, with the 19-29 age group having the lowest

(mean: 1.35 contacts, SD: 0.18 contacts) and the 40-49 age group the highest (mean:

1.97 contacts, SD: 0.14 contacts). We observed three distinct diagonal patterns:

younger individuals frequently interacting with their parents, older individuals with their

children and grandchildren, and individuals interacting with those of a similar age,

reflecting intergenerational living arrangements.

Fig 2. Contact patterns. Contact patterns reported in the Avian Contact Study. **(A)** the distribution of the

number of contacts per person. The red line indicates the fitted negative binomial distribution, and the

pink region represents the 95% confidence interval. The fitted negative binomial distribution had a mean

of 4.64 and a dispersion parameter of 4.02. **(B)** Average number of contacts between age groups, while **(C)**

Average number of contacts between family members only. In both **(B)** and **(C)**, darker colours indicate

higher average contacts.

A comparison of the Avian Contact Study and CoMix is shown in Fig 3. Respondents in

the Avian Contact Study had higher average contact levels than respondents of CoMix

across all age groups. The smaller sample size in the Avian Contact Study compared to

CoMix may explain the wider confidence intervals around the mean. Statistical

comparisons of the contact patterns are provided in the Supplementary Materials.

Influence of age on outbreak size

As expected, outbreak size was highly dependent on the reproduction number.

Infections with reproduction numbers less than 0.2 did not lead to onward

transmission. Over 75% of simulations resulted in small-scale outbreaks (0-10 cases).

The average outbreak size increased sharply from 4 cases at R_0=1 to 41 cases at R_0=1.1

(Fig S3).

We also examined how the age group of the index case influenced outbreak dynamics.

For R_0 between 0.7 and 1, half of simulations with index cases aged 60+ resulted in no

secondary cases, compared to 40% of simulations for those aged 19-59. When $R_0 =$

1.1, 9% of simulations with index cases in the 19-59 age group led to outbreaks larger

than 50 cases, compared to 6% for those with index cases aged 60+.

For index cases in the 19-44 and 45-59 age groups, R_0 did not significantly affect the age

distribution of the cases. In contrast, for index cases aged 60+, we observed an increase

in the percentage of cases in the 0-18 age group from 0 to 13% as R_0 increased. This is

due to limited direct contact between the 60+ and 0-18 age groups, meaning individuals

aged 0-18 were infected further down the transmission chain through interactions with

other age groups (Figs S3 and S4).

Reduced effectiveness of contact tracing and self-isolation

with higher proportion of asymptomatic infections

We compared how outbreak sizes changed when 40% of infections were asymptomatic,

under both no control and control (contact tracing and self-isolation) scenarios,

depending on R_0 (Fig 4A). Under no control, no secondary cases were observed when R_0

was below 0.2 - the only case was the index case. Over 75% of simulations resulted in

small-scale outbreaks (less than 10 cases). Contact tracing and self-isolation reduced

the maximum outbreak size from 5705 cases to below 1000, and the average outbreak

size from 41 cases to below 10, when R_0=1.1.

We also evaluated how the average outbreak size changed under different levels of

control measures and varying proportions of asymptomatic infections (Fig 4B). When

the proportion of asymptomatic infections was low, even low to moderate levels of

contact tracing and self-isolation (0 to 40%) were sufficient to reduce the average

outbreak size from 41 cases to below 10. However, the reduction in the average

outbreak size became less pronounced as the proportion of asymptomatic infections

increased.

Fig 4. The impact of contact tracing and self-isolation on final outbreak size. (A) The distribution of

outbreak sizes for varying levels of R_0 under no control and control (contact tracing and self-isolation).

The orange line indicates the average outbreak size, and the shaded region represents the 95% prediction

interval. We assumed that 40% of cases were asymptomatic and that 40% of contacts were traced.

Among those traced, all symptomatic cases were assumed to self-isolate. **(B)** The average outbreak size

across 9000 simulations for varying levels of contact tracing and proportions of asymptomatic infections

when R_0 = 1.1. Shaded regions represent the 95% confidence interval.

Discussion

Principal findings

Our model provides insights into the potential outbreak size of avian influenza among humans under different levels of transmission. Most simulations resulted in small-scale outbreaks, with outbreaks exceeding 100 cases when the basic reproduction number was above 0.8, consistent with previous estimates (35). Contact tracing and self-isolation reduced outbreak size, but their potential effectiveness in preventing cases decreased as the proportion of asymptomatic infections increased, since they rely on symptomatic cases. Previous modelling studies similarly concluded that contact tracing and self-isolation alone was highly unlikely to prevent large outbreaks, particularly when subclinical infections were present (27,36). Our findings support and extend these conclusions to a different pathogen.

Translating findings into action

Asymptomatic infections limit the effectiveness of contact tracing and self-isolation, and they are optimally identified and controlled through asymptomatic testing programmes, which have previously shown to be successful at detecting cases (37). Reintroducing such programmes could enhance current interventions by improving detection of asymptomatic infections.

Moreover, we observed that the index cases aged 19-59 play an important role in transmission dynamics, leading to larger outbreaks based solely on contact data. Additionally, analysis of previous H5N1 infections suggests that the infection is more

prevalent among younger adults, and older age may not be a risk factor for severe

outcomes (38). Taken together, these findings suggest that younger adults may benefit

more from targeted interventions.

Strengths and limitations

We used a novel dataset from the Avian Contact Study, which focused on individuals at

higher risk of avian influenza and was developed with the help of public contributors to

ensure relevance and accessibility (21). To our knowledge, it is the only contact survey

conducted in the UK that has focused on farming communities since the COVID-19

pandemic. This allowed us to examine post-pandemic contact patterns in a rural, high-

risk groups, often underrepresented in contact surveys, who may have different mixing

patterns compared to the general population. We used age-stratified mixing patterns in

our model to estimate outbreak sizes, extending previous models that did not

incorporate age structure. This allowed us to capture variation in outbreak sizes across

age groups and identify which groups were more likely to drive larger outbreaks.

In addition to common limitations of other contact surveys (20,39), our data did not

include individuals under the age of 19, so we could not compare their contact patterns

to those of the general population. Although we included 0-18 age group from CoMix in

the transmission chain, this may not fully reflect the contact patterns due to differences

in the cohorts and study periods. Additionally, our analysis was based on data from

individuals living in the UK, most of whom were poultry farmers, hence our findings may

not be generalised to other countries or to farming communities with different livestock

practices. Although we included age in our model, we could not explore its relationship

with all the parameters. For instance, the influence of age on symptomatology is not

well understood. As a result, we kept the proportion of asymptomatic infections

constant across all age groups. Our model assumed that people with symptoms would

isolate on the same day, although this may not be the case in practice with people

adopting a 'wait and see' approach (40).

Areas for future work

358 Future research is required to further characterise contact patterns within agricultural

359 demographics and other related groups. In addition to farmers who make up 60% of the

360 agricultural workforce in the UK, 12% is covered by casual workers (41). Despite their

361 high exposure risk, it can be challenging to engage with this group due to language and

362 trust barriers.

Funding statement

Funding for the Avian Contact Study was awarded by PolicyBristol from the Research

England QR Policy Support Fund (QR PSF) 2022-24 for investigating 'Zoonotic spillover

of avian influenza'. AT is funded by the Wellcome Trust, Early Career

Award [227041/Z/23/Z]. EBP and IA acknowledges support from the National Institute

for Health Research Health Protection Research Unit (NIHR HPRU) in Evaluation and

Behavioural Science at the University of Bristol (NIHR207385).

Acknowledgements

We are grateful to all respondents who completed the questionnaire and to the public

contributors for their input into study design and questionnaire development. We thank

all members of the wider Avian Contact Study team for input into study design, data

collection and interpretation. We thank Dr Anastasia Chatzilena (University of Bristol)

for her guidance regarding the statistical analysis of contact data from the Avian

Contact Study and CoMix.

Data and code availability

Underlying data

Repository data.bris: The Avian Contact Study: questionnaire data 15 May – 31 July
2024 Due to the sensitivity of the data involved, these data are published as a
restricted dataset at the University of Bristol Research Data Repository data.bris,
at https://doi.org/10.5523/bris.3nmqsrbv5ruom2abn0ql6e8yh2 (42). The metadata
record published openly by the repository at this location clearly states how data can
be accessed by *bona fide* researchers. Requests for access will be considered by
the University of Bristol Research Data Service, who will assess the motives of
potential data re-users before deciding to grant access to the data. No authentic
request for access will be refused and re-users will not be charged for any part of
this process.

This project contains the following underlying data:

- Data file 1. (Raw underlying questionnaire data – csv file)
- Data file 2. (Raw underlying questionnaire data - .RDS file)
- Data file 2. (Associated data dictionary – csv file)
- Data file 3. (Code for importing underlying data in csv format into R for setting
up labelled data - .r file)
- Data file 4. (Blank consent form and participant information sheet – pdf file)

Data are available under the terms of National Archives' Non-Commercial

Government Licence for public sector information.

Extended data

Repository Zenodo: The Avian Contact Study Questionnaire and Data Dictionary
[10.5281/zenodo.13617061]
This project contains the following extended data:

- AvianInfluenzaSocialContactSu.pdf (The final questionnaire REDCap – PDF)
- AvianInfluenzaSocialContactSur_DataDictionary_080824v1.csv (Associated
data dictionary -csv file)

Data are available under the terms of the Creative Commons Attribution 4.0

International license (CC-BY 4.0).

Software availability

Source code available from: https://github.com/amythomas/aviancontactstudy

License: Creative Commons Attribution 4.0 International license (CC-BY 4.0).

References:

1. WOAH. Avian Influenza [Internet]. WOAH - World Organisation for Animal Health. Available from: https://www.woah.org/en/disease/avian-influenza/

2. World Health Organisation. https://www.who.int/emergencies/disease-outbreak-news/item/2025-DON575. 2025. Avian Influenza A(H5N1) - Cambodia.

3. CDC. Past Reported Global Human Cases with Highly Pathogenic Avian Influenza A(H5N1) (HPAI H5N1) by Country, 1997-2025 [Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-flu/php/avian-flu-summary/chart-epi-curve-ah5n1.html

4. Graziosi G, Lupini C, Catelli E, Carnaccini S. Highly Pathogenic Avian Influenza (HPAI) H5 Clade 2.3.4.4b Virus Infection in Birds and Mammals. Animals [Internet]. 2024 Jan;14(9):1372. Available from: https://www.mdpi.com/2076-2615/14/9/1372

5. CDC. 2020-2024 Highlights in the History of Avian Influenza (Bird Flu) Timeline [Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-flu/avian-timeline/2020s.html

6. Mahase E. H5N1: UK reports world's first case in a sheep. BMJ. 2025 Mar;388:r591.

7. CDC. Highly Pathogenic Avian Influenza A(H5N1) Virus: Interim Recommendations for Prevention, Monitoring, and Public Health Investigations [Internet]. Avian Influenza (Bird Flu). 2025. Available from: https://www.cdc.gov/bird-flu/prevention/hpai-interim-recommendations.html

8.   Garg S, Reinhart K, Couture A, Kniss K, Davis CT, Kirby MK, et al. Highly

Pathogenic Avian Influenza A(H5N1) Virus Infections in Humans. New England

Journal of Medicine [Internet]. 2025 Feb;392(9):843–54. Available from:

https://www.nejm.org/doi/full/10.1056/NEJMoa2414610

9.   CDC. Technical Report: June 2024 Highly Pathogenic Avian Influenza A(H5N1)

Viruses [Internet]. Avian Influenza (Bird Flu). 2025. Available from:

https://www.cdc.gov/bird-flu/php/technical-report/h5n1-06052024.html

10.   NHS. https://www.nhs.uk/conditions/bird-flu/. 2022. Bird flu.

11.   CDC. H5 Bird Flu: Current Situation [Internet]. Avian Influenza (Bird Flu). 2025.

Available from: https://www.cdc.gov/bird-flu/situation-summary/index.html

12.   Yang Y, Halloran ME, Sugimoto JD, Longini IM. Detecting Human-to-Human

Transmission of Avian Influenza A (H5N1). Emerg Infect Dis [Internet]. 2007

Sep;13(9):1348–53. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2857285/

13.   World Health Organisation. https://www.who.int/publications/m/item/updated-

joint-fao-who-woah-assessment-of-recent-influenza-a(h5n1)-virus-events-in-

animals-and-people_dec2024. 2020. Updated joint FAO/WHO/WOAH public

health assessment of recent influenza A(H5) virus events in animals and people.

14.   Brandt AM. The History of Contact Tracing and the Future of Public Health. Am J

Public Health [Internet]. 2022 Aug;112(8):1097–9. Available from:

https://ajph.aphapublications.org/doi/full/10.2105/AJPH.2022.306949

15.   El-Sadr WM, Platt J, Bernitz M, Reyes M. Contact Tracing: Barriers and Facilitators.

467   Am J Public Health [Internet]. 2022 Jul;112(7):1025–33. Available from:

468   https://ajph.aphapublications.org/doi/full/10.2105/AJPH.2022.306842

469   16.   UKHSA. Investigation and initial clinical management of possible human cases of

470   avian influenza with potential to cause severe human disease [Internet]. GOV.UK.

471   Available from: https://www.gov.uk/government/publications/avian-influenza-

472   guidance-and-algorithms-for-managing-human-cases/investigation-and-initial-

473   clinical-management-of-possible-human-cases-of-avian-influenza-with-

474   potential-to-cause-severe-human-disease

475   17.   Thomas A, Danon L, Christensen H, Northstone K, Smith D, Nixon E, et al. Limits

476   of lockdown: characterising essential contacts during strict physical distancing.

477   2021 Mar; Available from:

478   https://www.medrxiv.org/content/10.1101/2021.03.12.21253484v1

479   18.   McCreesh N, Morrow C, Middelkoop K, Wood R, White RG. Estimating age-mixing

480   patterns relevant for the transmission of airborne infections. Epidemics [Internet].

481   2019 Sep;28:100339. Available from:

482   https://www.sciencedirect.com/science/article/pii/S1755436518301646

483   19.   Edelman LS, Yang R, Guymon M, Olson LM. Survey Methods and Response Rates

484   Among Rural Community Dwelling Older Adults. Nurs Res [Internet]. 2013

485   Aug;62(4):286. Available from:

486   https://journals.lww.com/nursingresearchonline/FullText/2013/07000/Survey_Me

487   thods_and_Response_Rates_Among_Rural.10.aspx

20.   Hoang T, Coletti P, Melegaro A, Wallinga J, Grijalva CG, Edmunds JW, et al. A

Systematic Review of Social Contact Surveys to Inform Transmission Models of

Close-contact Infections. Vol. 30, Epidemiology. Lippincott Williams and Wilkins;

2019. p. 723–36.

21.   Thomas AC, Gokool S, Clapp G, Whitlow H, McGrath C, Moore P, et al. Data note

on the Avian Contact Study: a questionnaire resource for avian influenza public

health planning. Wellcome Open Res. 2025;9.

22.   Jarvis CI, Van Zandvoort K, Gimma A, Prem K, Auzenbergs M, O'Reilly K, et al.

Quantifying the impact of physical distance measures on the transmission of

COVID-19 in the UK. BMC Med. 2020 May;18.

23.   Koltai J, Vásárhelyi O, Röst G, Karsai M. Reconstructing social mixing patterns via

weighted contact matrices from online and representative surveys. Sci Rep. 2022

Dec;12.

24.   Dall'Amico L, Kleynhans J, Gauvin L, Tizzoni M, Ozella L, Makhasi M, et al.

Estimating household contact matrices structure from easily collectable

metadata. PLoS One. 2024 Mar;19.

25.   Farrington CP, Grant AD. The distribution of time to extinction in subcritical

branching processes: Applications to outbreaks of infectious disease. J Appl

Probab. 1999;36:771–9.

26.   Feng J, Zhu W, Ye X, Liu Z, Zhu Y, Wu Q, et al. Effects of contact tracing and nucleic

acid testing on the COVID-19 outbreak in Zunyi, China: data-driven study using a

branching process model. BMC Infect Dis. 2023 Jan;23:10.

27. Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of

controlling COVID-19 outbreaks by isolation of cases and contacts. Lancet Glob

Health [Internet]. 2020 Apr;8(4):e488–e496. Available from:

https://www.thelancet.com/article/S2214-109X(20)30074-7/fulltext

28. Dietz K. The estimation of the basic reproduction number for infectious diseases.

Stat Methods Med Res. 1993;2:23–41.

29. Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the

reproduction number for seasonal, pandemic, and zoonotic influenza: A

systematic review of the literature. BMC Infect Dis. 2014 Sep;14.

30. Ungchusak K, Auewarakul P, Dowell SF, Kitphati R, Auwanit W, Puthavathana P, et

al. Probable Person-to-Person Transmission of Avian Influenza A (H5N1). New

England Journal of Medicine. 2005 Jan;352:333–40.

31. Wang H, Feng Z, Shu Y, Yu H, Zhou L, Zu R, et al. Probable limited person-to-

person transmission of highly pathogenic avian influenza A (H5N1) virus in China.

The Lancet. 2008;371:1427–34.

32. Cowling BJ, Jin L, Lau EHY, Liao Q, Wu P, Jiang H, et al. Comparative epidemiology

of human infections with avian influenza A H7N9 and H5N1 viruses in China: A

population-based study of laboratory-confirmed cases. The Lancet.

2013;382:129–37.

33. Aditama TY, Samaan G, Kusriastuti R, Sampurno OD, Purba W, Misriyah, et al.

Avian influenza H5N1 transmission in households, Indonesia. PLoS One. 2012

Jan;7.

34.   Department of Health and Social Care. Chapter 7: contact tracing and isolation.

533   2023 Jan.

534   35.   Ward J, Lambert JW, Russell TW, Azam JM, Kucharski AJ, Funk S, et al. Estimates

535   of epidemiological parameters for H5N1 influenza in humans: a rapid review.

536   2024.

537   36.   Davis EL, Lucas TCD, Borlase A, Pollington TM, Abbott S, Ayabina D, et al. Contact

538   tracing is an imperfect tool for controlling COVID-19 transmission and relies on

539   population adherence. Nat Commun [Internet]. 2021 Sep;12(1):5412. Available

540   from: https://www.nature.com/articles/s41467-021-25531-5

541   37.   Capelastegui F, Smith J, Kumbang J, Humphreys C, Padfield S, Turner J, et al. Pilot

542   of asymptomatic swabbing of humans following exposures to confirmed avian

543   influenza A(H5) in avian species in England, 2021/2022. Influenza Other Respir

544   Viruses. 2023 Aug;17.

545   38.   Garretson TA, Liu J, Li SH, Scher G, Santos JJS, Hogan G, et al. Immune history

546   shapes human antibody responses to H5N1 influenza viruses. Nat Med. 2025;

547   39.   Nixon E, Silvonen T, Barreaux A, Kwiatkowska R, Trickey A, Thomas A, et al. A

548   mixed methods analysis of participation in a social contact survey. Epidemics.

549   2022 Dec;41.

550   40.   Mowbray F, Woodland L, Smith LE, Amlôt R, Rubin GJ. Is My Cough a Cold or

551   Covid? A Qualitative Study of COVID-19 Symptom Recognition and Attitudes

552   Toward Testing in the UK. Front Public Health. 2021 Aug;9.

553   41.   https://www.gov.uk/government/statistics/agricultural-workforce-in-england-at-

554   1-june/agricultural-workforce-in-england-at-1-june-2023 [Internet]. Agricultural

555   workforce in England at 1 June 2024.

556   42.   Thomas A, Brooks-Pollock E. Data collected from the Avian Contact Study

557   (https://bristol.ac.uk/avian-contact) questionnaire between 15 May 2024 and 31

558   July 2024. First data release. [Internet]. data.bris. 2024 [cited 2024 Sep 16].

559   Available from:

560   https://data.bris.ac.uk/data/dataset/3nmqsrbv5ruom2abn0ql6e8yh2