Prevalence of Risk Factors and Established Cardiovascular Disease Among All of Us Participants: Benchmarking Against National Estimates

Mengjing Yang MPH[1], Tinsae Admassu MD MPH[1], Harlan M Krumholz MD SM[1,2,3], *Chenxi Huang PhD[2,3], *Karthik Murugiah MBBS MHS [2,3]

[1]Yale School of Public Health, New Haven, CT
[2]Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of Medicine, New Haven, CT
[3]Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, New Haven, CT

*Co-senior authors

Correspondence: Karthik Murugiah MD, 195 Church St, 6th Floor, New Haven, CT 06510; 203-487-1239 (tel). 203-764-5653 (fax).

Funding: Dr Murugiah received support from the National Heart, Lung, and Blood Institute of the National Institutes of Health (under award K08HL157727).
Disclosures: Dr Krumholz has received options for Element Science and Identifeye and payments from F-Prime for advisory roles; he is a co-founder of and holds equity in Hugo Health, Refactor Health, and ENSIGHT-AI; and he is associated with research contracts through Yale University from Janssen, Kenvue, Novartis, and Pfizer.

Abstract Word Count: 249

Manuscript Word Count: 2198

ABSTRACT

Introduction:
Knowing how All of Us (AoU) Research Program participants differ in demographics and prevalence of cardiovascular risk factors and established cardiovascular disease (CVD) from the U.S population is important for cardiovascular researchers using these data, and to provide context to their findings.

Methods:
We included AoU participants enrolled between May 2017-June 2022. 'AoU Survey' cohort included participants who completed the Personal and Family Health History survey (N=185,232). 'AoU EHR' included participants with linked EHR data (N = 287,012). We identified cardiovascular risk factors and established CVD through survey questions in 'AoU Survey' and National Health and Nutrition Examination Survey (NHANES), and SNOMED codes for 'AoU EHR'. Prevalence was compared to the weighted NHANES 2017-March 2020 cycle (N=9,683 representing 248 million), overall and by demographic sub-groups.

Results:
'AoU Survey' and 'AoU EHR' were both older than NHANES (56.1±17 years and 57.6±17 years versus 47.5±18 years) with more female participants. Black participants were underrepresented in 'AoU Survey'. Prevalence of cardiovascular comorbidities in 'AoU Survey' and NHANES were similar overall, but lower when stratified by age and among female participants. Black participants had higher rates of many comorbidities like hypertension, diabetes, coronary artery disease and congestive heart failure than the U.S Black population, but lower chronic kidney disease. 'AoU EHR' with its code-based identification had distinctly higher prevalence of most cardiovascular comorbidities.

Conclusion:
AoU includes more historically underrepresented groups, as intended. However, researchers should understand that demographic composition and disease prevalence change based on data availability and how conditions are ascertained.

INTRODUCTION

The All of Us (AoU) Research Program, launched by the National Institutes of Health, is a large and diverse U.S. cohort designed to accelerate precision medicine research.(1) AoU integrates multiple data modalities, including electronic health record (EHR), detailed participant surveys, biospecimens and genomics, physical measurements, and digital health data, while prioritizing enrollment of groups historically underrepresented in biomedical research.(2) For cardiovascular research, these resources can enable new insights into risk factors, disease progression, and treatment responses across diverse communities.

At the same time, AoU enrollment relies on convenience sampling and broad outreach rather than probability-based methods. Consequently, its participants may systematically differ from the general U.S. adult population. Further, AoU analyses may leverage survey responses or linked EHR data, depending on the research question and on data availability. These sources define distinct AoU sub-cohorts whose demographic composition and measured cardiovascular disease (CVD) burden can differ, both from each other and from the U.S. population, potentially affecting the interpretation of study findings. Recognizing these differences can help investigators select the appropriate AoU sub-cohort for a given aim and calibrate expectations about external validity.

Accordingly, this study benchmarks AoU against a national reference by comparing the demographic composition and prevalence of major cardiovascular risk factors and established CVD in AoU sub-cohorts defined by available survey data and by available EHR data with contemporaneous estimates from the National Health and Nutrition Examination Survey (NHANES).(3) We conduct the analyses overall and by key demographic subgroups. Our comprehensive analyses provide critical context for interpreting future cardiovascular research leveraging this landmark cohort.

METHODS

Data Sources and Study population

AoU Research Program

We used the Controlled Tier Dataset v7 and included participants aged ≥18 years enrolled between May 31, 2017, and June 30, 2022. Cardiovascular comorbidities were identified either from self-reported diagnoses in the Personal and Family Health History (PFHH) survey or from diagnosis codes in linked electronic health records (EHR). Based on data availability, we defined three analytic cohorts: (1) AoU Overall: all eligible

participants regardless of survey/EHR availability; (2) AoU Survey: participants who completed PFHH; and (3) AoU EHR: participants with linked EHR data. Each participant provided written informed consent, and oversight was provided by the AoU Institutional Review Board.

NHANES

NHANES is a continuous program conducted by the National Center for Health Statistics using a complex, multistage, stratified, clustered design to produce nationally representative estimates for the U.S. population. All NHANES analyses incorporate provided sample weights and design variables to account for unequal selection probabilities and nonresponse. We analyzed adults (≥18 years) from the 2017–March 2020 cycle who completed the following questionnaires: demographics (DEMO), smoking (SMQ), medical conditions (MCQ), blood pressure (BPQ), kidney conditions (KIQ), occupation (OCQ), health insurance (HIQ), diabetes (DIQ), and health status (HUQ). Use of publicly available, de‐identified NHANES data was deemed exempt by the Yale University Institutional Review Board.

Patient characteristics

AoU definitions

Self‐reported demographics (e.g., sex, race/ethnicity, employment, income, insurance) were obtained from the "Basics" survey. We examined eight cardiovascular conditions: hypertension (HTN), hyperlipidemia (HLD), diabetes mellitus (DM), coronary artery disease (CAD), acute myocardial infarction (AMI), congestive heart failure (CHF), cerebrovascular accident/transient ischemic attack (CVA/TIA), and chronic kidney disease (CKD). In AoU Survey, a condition was considered 'Present' if the participant selected Self to the relevant PFHH item; 'Absent' if only family members were selected; 'Did not answer' if Skip was selected; and 'Missing' if no response was provided. In AoU EHR, a condition was 'Present' if at least one corresponding Systematized Nomenclature of Medicine Clinical Terms (SNOMED CT)(4) code appeared any time prior to the participant's PFHH survey date; otherwise, it was classified 'Absent'. This approach aligns with the "ever told by a health professional" wording used in survey items and avoids imposing an arbitrary look‐back window given heterogeneous EHR observation period. The full SNOMED CT code lists are provided in Supplementary Table 1.

NHANES definitions

Sociodemographic variables were collected via structured interviews. Cardiovascular comorbidities were defined by self‐report using targeted questions: CHF (MCQ.160b), coronary heart disease (MCQ.160c), angina (MCQ.160d), AMI (MCQ.160e), and stroke (MCQ.160f); CKD and dialysis were assessed with KIQ.022 and KIQ.025, respectively; diabetes, hypertension, and hyperlipidemia were identified with DIQ.010, BPQ.020, and BPQ.080. Responses were categorized as 'Present' for 'Yes', 'Absent' for 'No', 'Did not answer' for 'Refused'/'Don't know', and 'Missing' when no response was recorded.

Race/ethnicity harmonization

Race and ethnicity were self‐reported in both datasets. For cross‐dataset comparability, we harmonized categories to Non‐Hispanic White, Non‐Hispanic Black, Hispanic, Asian, and Other. In NHANES, "Other race including multiracial" aggregates several less common groups without further breakdown. AoU allows more granular self‐identification (e.g., American Indian or Alaska Native, Native Hawaiian or other Pacific Islander, Middle Eastern or North African, multiracial). To maintain consistency, these detailed AoU categories were grouped under Other; because the underlying composition differs between datasets, comparisons involving "Other" are not interpreted.

Statistical analysis

Analyses were conducted in R 4.4.0. For NHANES, we used the survey package to account for the complex design, incorporating weights, strata, and primary sampling units to derive nationally representative estimates and variance. For AoU, which is non‐probabilistic, we reported unweighted means, proportions, and 95% confidence intervals.

We first compared key sociodemographic characteristics (age, sex, race/ethnicity, income, insurance) across NHANES and the three AoU cohorts (AoU Overall, AoU Survey, AoU EHR) using two‐sample t‐tests for continuous variables and chi‐square tests for categorical variables. For comorbidity prevalence, comparisons were restricted to NHANES vs AoU Survey and NHANES vs AoU EHR. We further examined group differences within strata defined by age (18-40, 40-65, >65 years), sex, and race/ethnicity using the same framework. Two‐sided P<0.05 was considered statistically significant; given the large sample size and multiple comparisons, significance testing was used to highlight patterns rather than to draw definitive causal inferences.

RESULTS

The AoU Overall cohort included 413,457 participants (Figure 1). Of these, 185,232 (44.8%) completed the PFHH survey (AoU Survey), and 287,012 (69.4%) had linked electronic health records (AoU EHR). The NHANES cohort included 9,683 adults, representing an estimated 248 million U.S. adults after weighting.

Compared with NHANES, AoU participants were older (mean±SD 56.1±17.1 vs 47.5±17.9 years) and more often female (60.4% vs 51.8%), with a higher proportion of Black participants (18.6% vs 11.5%; all P<0.001; Table 1). Both AoU sub-cohorts were also older (AoU Survey 58.1±17.1; AoU EHR 57.6±17.0 years) and had higher percentage of females (AoU Survey 64.0%; AoU EHR 60.1%) than NHANES. Black representation differed by sub-cohort with AoU Survey having 9.1%, lower than NHANES (11.5%) and AoU EHR having 19.9%, higher than NHANES. The higher mean age of AoU Survey versus NHANES was observed across demographic subgroups, with men 62.6 vs 46.7 years in NHANES; women 56.9 vs 48.3; White 61.4 vs 49.8; Black 56.8 vs 45.3; all P<0.001 (Supplementary Table 2).

Relative to AoU EHR, AoU Survey participants were slightly older and more often female and had higher proportions of White participants (69.7% vs 52.4%), greater educational attainment (>12 years: 83.5% vs 67.3%), and higher employment (53.3% vs 45.8%).

The overall prevalence of CV risk factors and established CVD is shown in Figure 2 and Supplementary Table 3. Overall, self-reported prevalence in the AoU Survey cohort were similar to or marginally lower than NHANES estimates (e.g. hypertension 29.6% in AoU Survey and 31.7% in NHANES, DM 11.3% in both, CAD 4.1% in both). In contrast, the AoU EHR cohort with its code-based disease identification had a significantly higher prevalence of all comorbidities except for CVA/TIA, which was lower than in NHANES (e.g. hypertension 41.6%, DM 19.6%, CAD 11.7% etc).

The prevalence of CV risk factors and established CVD stratified by age, sex and race/ethnicity is shown in Figures 3 and 4 and Supplementary Table 4. Across age groups, the prevalence of CV risk factors and established CVD was lower in the AoU Survey cohort compared to NHANES, and this pattern was consistent across most comorbidities. A similar pattern was also observed among women. Stratified by race, the prevalence of CV risk factors and established CVD was consistently lower among White participants in the AoU Survey cohort when compared to the White population in NHANES, despite a higher mean age of White participants in AoU. However, Black participants deviated from this pattern, with higher rates of hypertension, diabetes, CAD, CHF, but lower prevalence of CKD.

DISCUSSION

In this benchmarking study, All of Us (AoU) participants differed from the U.S. adult population in composition, with older age, a higher proportion of women and Black adults, and these differences varied based on participant availability of health survey data and EHR data. CVD burden also differed by ascertainment methods. Survey‑based AoU estimates were broadly similar to national figures overall, but within age strata was consistently lower across most conditions. In contrast, EHR‑based prevalence in AoU exceeded both AoU survey and NHANES estimates for most conditions. Subgroup patterns were heterogeneous: women in AoU Survey reported fewer conditions than women in NHANES, whereas Black participants in AoU Survey had higher hypertension, diabetes, and heart failure than Black adults in NHANES.

These findings should be interpreted considering how AoU is assembled and how conditions are captured. AoU is a convenience sample with intentional outreach to groups historically underrepresented in research, which is reflected in a higher overall proportion of Black participants than NHANES; however, representation differs across sub-cohorts, with the AoU Survey cohort in fact including a lower proportion of Black participants than both AoU Overall and NHANES. Differences between the survey‑defined and EHR‑defined cohorts extended beyond race/ethnicity to education and employment, indicating that choosing a data source effectively selects into different AoU populations. These differences also underscore the persistent socioeconomic and logistic barriers to completing surveys which continue to limit our ability to obtain self-reported health behaviors and health experiences of underrepresented populations.(5) At the same time, disease ascertainment by self‑report and clinical coding have their different sets of biases.(6) Self‑report can under‑ascertain conditions because of recall or health‑literacy barriers(7), whereas EHR capture depends on care access, encounter frequency, and coding practices.(8) Our EHR definitions used any prior SNOMED CT code recorded before the participant's baseline survey date to align conceptually with the "ever diagnosed" wording in both AoU and NHANES surveys, avoiding arbitrary look‑back windows in the face of heterogeneous EHR observation periods. Together, these compositional and measurement factors could explain the lower survey‑based prevalence within age strata and the higher EHR‑based prevalence.

This study adds clarity to the literature(9) by providing a methods‑aligned benchmark for AoU CVD burden against a contemporaneous, probability‑sampled national reference and by presenting AoU survey‑based and EHR‑based estimates separately rather than combining sources. Prior AoU comparisons that contrasted EHR‑only AoU definitions with NHANES survey estimates(10), or that used non‑survey references

such as Global Burden of Diseases (GBD)(11), are difficult to interpret because of non‑parallel case definitions, broader multimorbidity scopes, or misaligned time windows. By aligning "ever diagnosed" definitions, matching time frames, and stratifying by age, sex, and race/ethnicity, our analysis isolates where AoU diverges most from the national benchmark and shows how those divergences depend on the data source a study elects to use. Relative to legacy cardiovascular cohorts such as MESA(12) and ARIC(13), AoU offers unprecedented scale and multimodal depth; our results clarify how its convenience sampling and multi‑source ascertainment should be considered when generalizing AoU‑based findings to the U.S. population.

These benchmarks have practical implications. Investigators using AoU for cardiovascular research should prespecify whether analyses will rely on survey or EHR data and recognize that this choice entails different demographic compositions and measured burdens of CVD. Because AoU estimates are unweighted and derived from a convenience sample, they should not be interpreted as national prevalence; instead, contrasts with NHANES can guide expectations about direction and magnitude of differences when translating AoU findings. Stratification by age, sex, and race/ethnicity is essential for interpretation and for assessing whether observed associations are likely to hold in the broader U.S. population.

This work has limitations. It is cross‑sectional, using a single survey timepoint and an EHR snapshot, so we do not assess disease onset or progression. Misclassification remains possible in both sources: survey measures may be affected by recall or comprehension, and EHR measures by incomplete capture or coding variability. AoU's non‑probability sampling and unweighted analysis limit direct comparability to NHANES despite the benchmarking approach, and the harmonized "Other" race/ethnicity category differs in composition across datasets and is not interpreted. EHR data are available only for participants who consented and who could be linked through participating healthcare provider organizations; expanding linkage may change the composition of the EHR‑defined cohort in future AoU releases. Further, AoU Survey in our study refers to completion of the Personal/Family Health History which was used to identify self-reported comorbidities. Response rates and demographic distributions will vary for other surveys. Finally, we did not examine within‑person concordance between survey and EHR because our objective was to compare population burden across data‑defined AoU cohorts, not to reconcile sources.

CONCLUSION

AoU's breadth and diversity make it a powerful platform for cardiovascular research, but who is included and how conditions are ascertained shape measured CVD burden. By

benchmarking AoU survey-defined and EHR-defined cohorts against NHANES using aligned definitions and time frames, we provide actionable context to help researchers choose the appropriate AoU data source for their aims and to judge the generalizability of AoU-based cardiovascular findings.

ACKNOWLEDGEMENTS

We gratefully acknowledge All of Us participants for their contributions, without whom this research would not have been possible. We also thank the National Institutes of Health's All of Us Research Program for making available the participant data examined in this study.

TABLES AND FIGURES

Table 1. Comparison of Socio-Demographic Factors between NHANES and AoU cohorts

Socio-demographic Status	NHANES Weighted		AoU General (N = 413,457)		AoU Survey (N = 185,232)		AoU EHR (N = 287,012)	
Age (mean, SD, in years)	47.49 (17.87)		56.06(17.11)		58.14(17.14)		57.63(16.96)	
	N	%	N	%	N	%	N	%
Age Group								
18-40	94125616	37.96	91731	22.19	37045	20.00	57725	20.11
40-65	103189321	41.62	174886	42.30	71991	38.87	120291	41.91
>=65	50632123	20.42	146840	35.52	76196	41.14	108996	37.98
Sex								
Male	119592918	48.23	155169	37.53	62560	33.77	108739	37.89
Female	128354142	51.77	249565	60.36	118491	63.97	172401	60.07
Did not answer	0	0.00	4439	1.07	1237	0.67	3169	1.10
Missing	0	0.00	4284	1.04	2944	1.59	2703	0.67
Race/ Ethnicity								
White	154162965	62.18	222646	53.85	129090	69.69	150402	52.40
Black	28420787	11.46	77069	18.64	16918	9.13	57177	19.92
Asian	14781102	5.96	13838	3.35	6086	3.29	8038	2.80
Hispanic/Latino/a/x	40485465	16.33	64672	15.64	17236	9.31	47699	16.62
Other/ >1	10096742	4.07	23443	5.67	10266	5.54	15690	5.47
Did not answer	0	0.00	11692	2.83	5636	3.04	8005	2.79
Missing	0	0.00	97	0.02	0	0.00	1	0.00
Marital Status								
Married	148072184	59.72	204011	49.34	107790	58.19	139534	48.62
Widowed, Separated or divorced	45281859	18.26	92150	22.29	35758	19.30	67086	23.37
Never married	47000622	18.96	102480	24.79	36484	19.70	70297	24.49

Did not answer	94828	0.04	14719	3.56	5200	2.81	10094	3.52
Missing	7497567	3.02	97	0.02	0	0.00	1	0.00
Education (years)								
<12 y	26538169	10.70	36432	8.81	5976	3.23	27867	9.71
12 y	64776211	26.13	77014	18.63	19654	10.61	56778	19.78
>12 y	148884269	60.05	286481	69.29	154659	83.49	193184	67.31
Did not answer	2508442	0.10	13433	3.25	4943	2.67	9182	3.20
Missing	7497567	3.02	97	0.00	0	0.00	1	0.00
Self-rated Health								
Good	201938577	81.44	293368	70.95	144907	78.23	205987	71.77
Fair	39302718	15.85	81000	19.59	31524	17.02	60044	20.92
Poor	6419188	2.59	20555	4.97	7352	3.97	15358	5.35
Did not answer	286576	0.12	17288	4.18	1449	0.78	5467	1.90
Missing	0	0.00	1246	0.30	0	0.00	156	0.05
Health Insurance								
Covered	214599200	86.55	371633	89.88	173855	93.86	260359	90.71
Not covered	32892008	13.27	26779	6.48	6058	3.27	16733	5.83
Did not answer	455852	0.18	13158	3.18	3690	1.99	8685	3.03
Missing	0	0.00	1887	0.46	1629	0.88	1235	0.43
Smoking								
Yes	102779026	41.45	155901	37.71	65107	22.68	114595	39.93
No	145104923	58.52	232630	56.26	116753	40.68	164021	57.15
Did not answer	63111	0.03	23680	5.73	3372	1.17	8240	2.87
Missing	0	0.00	1246	0.3	0	0.00	156	0.05
Employment Status								
Employed	158614770	63.97	199392	48.23	98660	53.26	131320	45.75
Unemployed	89276381	36.01	197741	47.83	81166	43.82	144650	50.40
Did not answer	55909	0.02	16227	3.92	5406	2.92	11041	3.85
Missing	0	0.00	97	0.02	0	0.00	1	0.00

Figure 1: Venn diagram representing the overlap of participants with Person and Family Survey data and EHR data

Figure 2: Prevalence of CV risk factors and established CVD

This figure compares the estimated prevalence of major cardiometabolic conditions including acute myocardial infarction (AMI), coronary artery disease (CAD), congestive heart failure (CHF), chronic kidney disease (CKD), cerebrovascular accident/transient ischemic attack (CVA/TIA), diabetes mellitus (DM), hyperlipidemia (HLD), and hypertension (HTN) across three data sources: NHANES (self-report and exam-based), All of Us (AoU) EHR, and AoU survey responses. EHR-based prevalence estimates were consistently higher than survey-based or NHANES estimates for most conditions, particularly for AMI, CAD, and CKD.

Figure 3: Prevalence of CV risk factors and established CVD by age and sex

This figure presents the prevalence of eight cardiometabolic conditions by demographic subgroups (age groups: 18–40, 40–65, ≥65; and sex: Male, Female) across three datasets. EHR-based data from All of Us generally show higher prevalence rates than self-reported survey responses and NHANES, with the largest age-associated

increases observed for hypertension, diabetes, and CAD. The sex-specific trends vary by condition and data source.

Figure 4: Prevalence of CV risk factors and established CVD by race/ethnicity

This figure shows race/ethnicity-stratified prevalence for key cardiovascular and metabolic conditions, comparing NHANES and both EHR and survey data from All of Us. Disparities in disease burden are evident across racial/ethnic groups, with Black participants generally exhibiting higher prevalence of hypertension and diabetes across all datasets, while Asian participants show lower rates for most conditions.

Supplementary Table 1: SNOMED CT codes used to identify each disease condition

Condition	SNOMED ID
Hypertension	"194791005", "10725009", "10757401000119104", "118781000119108", "1201005",
	"123799005", "132721000119104", "140121000119100", "16229371000119106", "194783001",
	"194785008", "194788005", "198941007", "198942000", "198944004", "198945003",
	"198946002", "198947006", "198949009", "198965005", "198966006", "198967002",
	"198983002", "198984008", "198985009", "198986005", "198997005", "198999008",
	"199000005", "199002002", "199005000", "199007008", "199008003", "206596003",
	"23130000", "237279007", "28119000", "288250001", "307632004", "31992008",
	"367390009", "371125006", "37618003", "38341003", "397748008", "398254007",
	"40521000119100", "41114007", "429457004", "443482000", "461301000124109",
	"46764007", "48146000", "48194001", "52698002", "541000119105", "5501000119106",
	"56218007", "57684003", "59621000", "63287004", "65518004", "67359005",
	"697929007", "697930002", "70272006", "706882009", "712832005", "71421000119105",
	"72022006", "73410007", "765182005", "78808002", "78975002", "8218002",
	"82771000119102", "86041002", "8762007", "89242004", "95605009"
Hyperlipidemia	"55822004", "129589009", "299465007", "13644009","238076009",
	"398036000","238079002","238080004","267432004","302870006",
	"238083002","34528009","267435002","34349009","238085009",
	"402473001","267433009","129591001","267434003","238088006",
	"238040008","238089003","1571000119104","701000119103","402727002",
	"426161002","137941000119106","137931000119102"
Diabetes	"5368009", "5969009", "8801005", "11687002", "23045005",

	"28032008", "44054006", "46635009", "46894009", "49817004",
	"51002006", "59079001", "73211009", "75022004", "76751001",
	"81531005", "91352004", "111552007", "190368000", "190389009",
	"190447002", "199223000", "199225007", "199226008", "199227004",
	"199228009", "199229001", "199230006", "237599002", "237601000",
	"237627000", "313435000", "313436004", "359642000", "408540003",
	"413183008", "426705001", "426875007", "427089005", "445260006",
	"609561005", "609562003", "609563008", "609564002", "609566000",
	"609567009", "609568004", "609570008", "703136005", "703138006",
	"737212004", "1481000119100", "31321000119102", "40791000119105",
	"40801000119106", "105401000119101", "106281000119103",
	"10753491000119101", "10754881000119104"
CAD	"53741008", "371804009", "429673002", "11018701000119109", "139011000119104",
	"15960061000119102", "15960141000119102", "16891151000119103", "194842008",
	"194843003", "233817007", "285141000119106", "285151000119108", "371803003",
	"371805005", "429245005", "442224005", "442240008", "442421004", "443502000",
	"444855007", "451041000124103", "451361000124102", "719678003", "724431008",
	"791000119109", "92517006"
AMI	"22298006", "57054005", "401303003", "703164000", "15713081000119108",
	"15713121000119105", "16837681000119104", "17531000119105", "1755008",
	"194802003", "194856005", "194857001", "194858006", "23311000119105",
	"233835003", "233837006", "233840006", "285981000119103", "307140009",
	"314207007", "380001000004106", "401314000", "418044006", "54329005",
	"58612006", "59063002", "65547006", "70211005", "703164000",
	"703211006", "703360004", "70422006", "73795002", "76593002",
	"79009004", "836294006"
CHF	"42343007", "10633002", "194767001", "92506005", "88805009",
	"426263006", "426611007", "67441000119101", "23341000119109",
	"82523003", "96311000119109", "194779001", "194781004",
	"5148006", "698594003", "698296002", "66989003",

	"15629541000119106", "15781000119107", "83105008",
	"7401000175100", "16838951000119100"
CVA/TIA	"10349009", "14070001", "25772007", "56267009", "57981008",
	"70936005", "195213000", "195217004", "230713003", "230715005",
	"371040005", "371121002", "16371781000119100", "16000351000119109",
	"16000471000119107", "16000511000119103", "16002031000119102",
	"16002071000119104", "16002111000119106", "16002151000119100",
	"16002231000119106", "16002271000119109", "16002351000119105",
	"16002391000119100", "16002431000119105", "16002471000119108",
	"16002511000119104", "16023911000119108", "16024111000119109",
	"16026951000119102", "195212005", "195216008", "230690007",
	"230691006", "281240008", "292671000119104", "292681000119101",
	"292691000119103", "329401000119103", "329621000119105",
	"329641000119104", "329651000119102", "329671000119106",
	"330791000119108", "371041009", "373606000", "39925003",
	"413758000", "422504002", "434141000124103", "457551000124104",
	"5571000124103", "716051003", "78569004", "9901000119100",
	"99451000119105", "12237911000119109", "12237951000119105",
	"195200006", "195209007", "195210002", "195211003",
	"230716006", "230717002", "266257000", "64009001",
	"710575003", "88032003"
CKD	"709044004", "46177005", "49708008", "90688005", "236433006",
	"236436003", "431855005", "431856006", "431857002", "433144002",
	"433146000", "700378005", "700379002", "713313000", "714152005",
	"714153000", "723190009", "711000119100", "721000119107",
	"731000119105", "741000119101", "771000119108", "90741000119107",
	"90761000119106", "90791000119104", "96441000119101",
	"129151000119102", "129161000119100", "129181000119109",
	"691421000119108", "104931000119100"

Supplementary Table 2: Mean age of demographic subgroups

Group	Subgroup	NHANES		AoU Survey		AoU EHR	
		Mean	SD	Mean	SD	Mean	SD
Gender	Female	48.28	18.02	56.9	16.86	56.91	16.93
	Male	46.65	17.67	62.6	17.04	60.63	16.78
Race/Ethnicity	White	49.79	18.12	61.41	16.82	62.13	17.04
	Black	45.30	17.23	56.84	14.78	56.91	14.51
	Asian	45.05	16.75	48.23	16.51	50.4	16.92
	Hispanic	41.56	16.21	50.34	15.67	51.94	15.91
	Other	46.04	16.92	49.59	16.73	50.58	16.95

Supplementary Table 3. Prevalence of CV risk factors and established CVD

	NHANES Weighted		AoU Survey (N=185232)		AoU EHR (N=287012)	
	N	%	N	%	N	%
Hypertension						
Yes	78613893	31.71	54917	29.65	119554	41.65
No	169132190	68.21	65520	35.37	167458	58.35
Do not answer	200977	0.08	49763	26.87		
Missing	0	0.00	15032	8.12		
Hyperlipidemia						
Yes	82319763	33.20	56749	30.64	109435	38.13
No	164499677	66.34	46095	24.89	177577	61.87
Do not answer	1127620	0.45	62836	33.92		
Missing	0	0.00	19552	10.56		
Diabetes						
Yes	27931576	11.27	20988	11.33	56223	19.59
No	219839616	88.66	56709	30.62	230789	80.41
Do not answer	175869	0.07	84751	45.75		

Missing	0	0.00	22784	12.30		
CAD						
Yes	10166914	4.10	7661	4.14	33521	11.68
No	229863340	92.71	36314	19.60	253491	88.32
Do not answer	419240	0.17	107980	58.29		
Missing	7497567	3.02	33277	17.97		
AMI						
Yes	8977335	3.62	4997	2.70	14875	5.18
No	231146169	93.22	65960	35.61	272137	94.82
Do not answer	325989	0.13	86627	46.77		
Missing	7497567	3.02	27648	14.93		
CHF						
Yes	6287028	2.54	3890	2.10	15994	5.57
No	233775925	94.28	40928	22.10	271018	94.43
Do not answer	386549	0.16	107241	57.90		
Missing	7497567	3.02	33173	17.91		
CVA/TIA						
Yes	9087945	3.67	6156	3.32	7969	2.78
No	231020064	93.17	46124	24.90	279043	97.22
Do not answer	341484	0.14	102600	55.39		
Missing	7497567	3.02	30352	16.39		
CKD						
Yes	7687625	3.10	3629	1.96	26221	9.14
No	232461842	93.75	1219	0.66	260791	90.86
Do not answer	300026	0.12	139414	75.26		
Missing	7497567	3.02	40970	22.12		

by demographic subgroups

	Category	NHANES Weighted	AOU Survey (N=185232)	AOU EHR (N=287012)
		%	%	%
Subgroup: Age Group				
Hypertension	18-40y	12.29	8.16	12.96
	40-65y	35.61	26.84	38.31
	>=65y	59.85	42.74	60.55
Hyperlipidemia	18-40y	11.79	8.27	6.85
	40-65y	40.50	27.50	32.26
	>=65y	58.14	44.47	61.17
Diabetes	18-40y	2.21	3.38	6.82
	40-65y	13.18	11.97	19.53
	>=65y	24.20	14.59	26.42

CAD	18-40y	0.42	0.11	0.48
	40-65y	3.71	1.98	6.95
	>=65y	11.73	8.14	22.83
AMI	18-40y	0.51	0.17	0.61
	40-65y	3.76	1.81	4.06
	>=65y	9.12	4.77	8.84
CHF	18-40y	0.32	0.27	0.70
	40-65y	2.34	1.69	4.03
	>=65y	7.05	3.37	9.86
CVA/TIA	18-40y	0.53	0.64	0.29
	40-65y	3.21	2.78	1.70
	>=65y	10.41	5.14	5.28
CKD	18-40y	1.74	0.59	1.53
	40-65y	2.62	1.99	6.64
	>=65y	6.60	3.57	15.91
Subgroup: Gender				
Hypertension	Male	32.70	36.18	45.29
	Female	30.78	26.08	39.22
Hyperlipidemia	Male	34.92	36.97	42.01
	Female	31.60	27.18	35.59
Diabetes	Male	12.58	13.56	20.79
	Female	10.04	10.11	18.79
CAD	Male	5.84	7.84	16.81
	Female	2.48	2.14	8.39
AMI	Male	5.06	4.90	7.61
	Female	2.28	1.51	3.64
CHF	Male	2.95	2.95	7.25
	Female	2.15	1.64	4.50
CVA/TIA	Male	3.23	3.97	3.04
	Female	4.07	2.97	2.61
CKD	Male	3.33	2.51	11.94
	Female	2.88	1.66	7.33
Subgroup: Race				
Hypertension	White	32.98	30.26	42.45
	Black	39.04	42.23	47.89
	Asian	24.98	15.72	23.05
	Hispanic/Latino/a/x	27.73	21.12	37.22
	Other/ >1	24.41	22.04	32.38
Hyperlipidemia	White	36.16	33.08	44.73
	Black	26.96	26.35	30.76
	Asian	28.66	22.00	26.96

	Hispanic/Latino/a/x	31.90	22.83	30.87
	Other/ >1	27.23	23.52	28.02
Diabetes	White	10.91	10.20	16.64
	Black	12.49	19.20	24.60
	Asian	10.86	7.03	13.13
	Hispanic/Latino/a/x	9.68	13.92	24.86
	Other/ >1	13.08	10.09	16.32
CAD	White	5.40	4.88	14.09
	Black	1.72	2.31	9.87
	Asian	2.12	1.74	6.07
	Hispanic/Latino/a/x	2.17	1.55	7.90
	Other/ >1	1.98	2.64	8.56
AMI	White	4.28	2.96	5.45
	Black	3.04	2.70	5.90
	Asian	1.45	1.02	2.38
	Hispanic/Latino/a/x	2.54	1.64	4.21
	Other/ >1	2.61	1.81	3.95
CHF	White	2.81	2.10	5.51
	Black	3.56	3.75	7.69
	Asian	0.83	0.59	2.08
	Hispanic/Latino/a/x	1.78	1.20	4.21
	Other/ >1	1.54	1.44	4.07
CVA/TIA	White	3.97	3.56	3.31
	Black	4.83	4.33	2.40
	Asian	1.50	1.15	1.34
	Hispanic/Latino/a/x	2.61	1.68	2.00
	Other/ >1	2.79	2.46	1.94
CKD	White	3.13	2.08	9.25
	Black	3.93	2.59	11.12
	Asian	1.85	0.77	4.50
	Hispanic/Latino/a/x	2.87	1.01	7.72
	Other/ >1	2.91	1.40	6.77

REFERENCES

1. Investigators TAoURP. The "All of Us" Research Program. New England Journal of Medicine 2019;381:668–676.
2. Mapes B, Foster C, Kusnoor S et al. Diversity and inclusion for the All of Us research program: A scoping review. PLOS ONE 2020;15:e0234962.
3. Prevention UCfDCa. US National Health and Nutrition Examination Survey.
4. SNOMED International. (2024). SNOMED CT. Retrieved from https://www.nlm.nih.gov/healthit/snomedct/snomed_overview.html.
5. Jang M, Vorderstrasse A. Socioeconomic Status and Racial or Ethnic Differences in Participation: Web-Based Survey. JMIR Res Protoc 2019;8:e11865.
6. Sulieman L, Cronin RM, Carroll RJ et al. Comparing medical history data derived from electronic health records and survey answers in the All of Us Research Program. J Am Med Inform Assoc 2022;29:1131–1141.
7. Newell SA, Girgis A, Sanson-Fisher RW, Savolainen NJ. The accuracy of self-reported health behaviors and risk factors relating to cancer and cardiovascular disease in the general population11The full text of this article is available via AJPM Online at http://www.elsevier.com/locate/ajpmonline.: A critical review. American Journal of Preventive Medicine 1999;17:211–229.
8. Bower JK, Patel S, Rudy JE, Felix AS. Addressing Bias in Electronic Health Record-based Surveillance of Cardiovascular Disease Risk: Finding the Signal Through the Noise. Current Epidemiology Reports 2017;4:346–352.
9. Acosta JN, Leasure AC, Both CP et al. Cardiovascular Health Disparities in Racial and Other Underrepresented Groups: Initial Results From the All of Us Research Program. Journal of the American Heart Association 2021;10:e021724.
10. Wang J, Ferguson EL, Buto P et al. Sociodemographic, health-related, and clinical characteristics and their associations with mortality among all of us participants compared with the United States general population. Am J Epidemiol 2025.
11. Zeng C, Schlueter DJ, Tran TC et al. Comparison of phenomic profiles in the All of Us Research Program against the US general population and the UK Biobank. J Am Med Inform Assoc 2024;31:846–854.
12. Blaha Michael J, DeFilippis Andrew P. Multi-Ethnic Study of Atherosclerosis (MESA). JACC 2021;77:3195–3216.
13. Wright Jacqueline D, Folsom Aaron R, Coresh J et al. The ARIC (Atherosclerosis Risk In Communities) Study. JACC 2021;77:2939–2959.