Association of device-based measured sleep duration, sleep efficiency and sleep regularity with cardiometabolic health: the Prospective Physical Activity, Sitting, and Sleep (ProPASS) consortium

Annemarie Koster[1*], Raaj Kishore Biswas[2,3*], Matthew N Ahmadi[2,3], Joanna M Blodgett[4,5], Nicholas A Koemel[2,3], Wenxin Bian[2,3], Andrew J Atkin[6], Richard M Pulsford[7], Borja del Pozo Cruz[8], Carlos Celis-Morales[9], John Mitchell[4], Pasan Hettiarachchi[10], Peter J Johansson[10], Magnus Svartengren[10], Hsiu-Wen Chan[11], Kristin Suorsa[12,13], Esmée A Bakker[14], Sari Stenholm[12,13], Thijs M.H. Eijsvogels[15], Hans HCM Savelberg[16], Vegar Rangul[17], Alun D Hughes[18,19,5], I-Min Lee[20,21], Peter A. Cistulli[22,23,24], Andreas Holtermann[25], Mark Hamer[,4,5**], Emmanuel Stamatakis[1,2**], ProPASS Collaboration[***]

*Joint first authors; **Joint senior authors;

***ProPASS collaboration:
- Carla JH van der Kallen: Department of Internal Medicine, CARIM School for Cardiovascular Diseases Maastricht University, Maastricht, the Netherlands.
- Dick HJ Thijssen: Department of Medical BioSciences, Exercise Physiology research group, Radboud university medical center, Nijmegen, the Netherlands
- Nidhi Gupta: National Research Centre for the Working Environment, Copenhagen, Denmark
- Lauren B Sherar: School of Sport, Exercise and Health Sciences, Loughborough University, UK

1. Department of Social Medicine, CAPHRI Care and Public Health Research Institute, Maastricht University, Maastricht, the Netherlands.
2. Mackenzie Wearables Research Hub, Charles Perkins Centre, University of Sydney, Sydney, NSW, Australia
3. School of Health Sciences, Faculty of Medicine and Health, University of Sydney, Sydney, NSW, Australia
4. Institute of Sport Exercise and Health, Division of Surgery and Interventional Sciences, UCL, United Kingdom.
5. University College London Hospitals NIHR Biomedical Research Centre, London, UK
6. School of Health Sciences, University of East Anglia, Norwich, UK
7. Faculty of Health and Life Sciences, University of Exeter, Exeter, UK.
8. Department of Sport Sciences, Faculty of Medicine, Health, and Sports, Universidad Europea de Madrid, Madrid, Spain.
9. School of Cardiovascular and Metabolic Health, University of Glasgow, Glasgow, United Kingdom
10. Occupational and Environmental Medicine, Department of Medical Sciences, Uppsala University, Uppsala, Sweden
11. Australian Women and Girls' Health Research Centre, School of Public Health, The University of Queensland, Brisbane, Queensland, Australia
12. Department of Public Health, University of Turku and Turku University Hospital, Turku, Finland
13. Centre for Population Health Research, University of Turku and Turku University Hospital; Turku, Finland
14. Department of Primary and community care, Radboud university medical center, Nijmegen, the Netherlands
15. Department of Medical BioSciences, Exercise Physiology Research Group, Radboud University Medical Center, Nijmegen, the Netherlands
16. Department of Nutrition and Movement Science, NUTRIM School for Nutrition and Translational Research in Metabolism, Maastricht University, Maastricht, the Netherlands
17. HUNT Research Centre, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU), Levanger, Norway
18. MRC Unit for Lifelong Health and Ageing, UCL Institute of Cardiovascular Science, UCL, United Kingdom.
19. UCL BHF Research Accelerator, University College London, London, UK
20. Division of Preventive Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA
21. Department of Epidemiology, Harvard TH Chan School of Public Health, Boston, Massachusetts, USA
22. Sleep Research Group, Charles Perkins Centre, University of Sydney, Camperdown, New South Wales, Australia
23. Northern Clinical School, Faculty of Medicine and Health, University of Sydney, Camperdown, New South Wales, Australia
24. Department of Respiratory and Sleep Medicine, Royal North Shore Hospital, St Leonards, New South Wales, Australia
25. National Research Centre for the Working Environment, Copenhagen, Denmark

Abstract

Background: Sleep is critical for cardiometabolic health, but evidence on the independent and joint associations of device-measured sleep parameters is lacking. The aim of this study is to examine the independent and joint associations of sleep duration, regularity, and efficiency, with cardiometabolic health outcomes.

Methods: Cross-sectional data from six cohorts (n = 14,085 participants; five countries, ≥4-day wear time) from the Prospective Physical Activity, Sitting and Sleep consortium (ProPASS) were used. Sleep duration (short: <7 h/day; adequate: 7-8 h/day; long: >8 h/day), sleep regularity index as a measure of day to day variability in sleep-wake pattern (regular: 87.3%; slightly irregular: 75.9-87.3%; irregular: <75.9%) , and sleep efficiency as the ratio of total sleep time to total time in bed (high, medium, low; based on tertiles) were derived from thigh-worn accelerometer data. Cardiometabolic health markers included body mass index, waist circumference, HDL and LDL cholesterol, glycated haemoglobin, systolic and diastolic blood pressure, and a composite cardiometabolic risk z-score was computed. Generalized linear regression was used to examine individual and joint associations of sleep parameters with cardiometabolic health, adjusted for age, sex, cohort, smoking, alcohol consumption, medication, prevalent cardiovascular disease and moderate to vigorous intensity physical activity.

Results: Short sleep duration (β: 0.05, 95%CI: 0.03-0.07), an irregular sleep pattern (0.14, 0.11-0.18), and low sleep efficiency (0.10, 0.07-0.13) were associated with a higher cardiometabolic risk score compared to adequate sleep duration, regular sleep patterns, and high sleep efficiency respectively. A long sleep duration was not associated with cardiometabolic risk score (0.01, -0.03-0.03). The joint analysis of all three sleep parameters shows that individuals with an irregular and low sleep efficiency, regardless of sleep duration, had worse cardiometabolic health (in long sleepers: 0.21, 95%CI: 0.15-0.28; in adequate sleepers: 0.23, 95%CI: 0.16-0.30; in short sleepers: 0.28, 95%CI: 0.22-0.35) compared to adequate, regular and highly efficient sleepers

Conclusion: Our study suggests that sleep regularity and efficiency seem important for cardiometabolic health beyond sleep duration. Multidimensional indices of sleep incorporating sleep duration, regularity, and efficiency should be evaluated across diverse populations in future longitudinal studies and trials.

Introduction

Sleep is widely recognized as critical for maintaining overall health. [1] There is established evidence for associations between sleep duration and health outcomes including cardiometabolic conditions such as type 2 diabetes, excess adiposity and high blood pressure.[1] Typically, studies report a U-shaped association between sleep duration and health, indicating that both shorter and longer sleep duration are associated with increased risk of mortality and disease risk, including obesity, diabetes and cardiovascular disease.[2-4] While most research has focused on sleep duration, more qualitative aspects of sleep have also shown to be independently associated with health. For example, a meta-analysis of 17 prospective cohort studies showed that lower self-reported sleep quality (e.g. defined as difficulty maintaining sleep or disturbed sleep) has been associated with coronary heart disease risk but not with other cardiovascular outcomes and mortality.[5] Poor sleep efficiency (i.e. spending more time awake while lying in bed), quantified by in-home polysomnography, has been associated with increased cardiovascular disease risk.[6]

In addition to sleep duration and efficiency, emerging evidence suggests that the stability of sleep-wake cycles also is important for health. [7] The variability of sleep patterns, particularly the duration and timing of sleep, has the potential to disrupt the circadian rhythm of biological processes. Such disruption is known to, for example, increase cardiovascular risk.[8] A systematic review examining the associations of sleep timing and sleep regularity with health outcomes in adults reported that higher sleep variability was associated with adverse health outcomes.[7] However, the evidence was judged to be of very low to moderate quality, mainly due to reliance on self-reporting of sleep behaviours which are limited by reporting bias and which prevent translation into public health guidance for sleep.[7] A systematic review of sleep variability and cardiometabolic health shows that variability is likely associated with obesity, weight gain, cardiovascular disease, and metabolic syndrome but not with insulin resistance.[9] It is important to note that many studies that have examined associations between sleep variability and health have not adjusted for sleep duration. More recently, findings from the Multi-Ethnic Study of Atherosclerosis have shown that a greater device-measured sleep irregularity was cross-sectionally associated with obesity, hypertension, fasting glucose, and haemoglobin A1C[10], and prospectivity with metabolic syndrome[11], and incident cardiovascular disease.[12] A prospective study in the Hispanic Community Health Study/Study of Latinos, did not observe associations between device-based sleep regularity with glucose biomarkers and incident diabetes[13] while in a recent study using data from the UK Biobank a high sleep irregularity was associated with higher risk of type 2 diabetes.[14]

Most epidemiological studies to date have relied on self-reported sleep characteristics, which are subject to numerous limitations and device-based assessment of sleep characteristics is preferable in terms of validity and data resolution.[15] Further, the use of 24-hour assessment protocols is especially appealing for the assessment of sleep regularity over multiple nights.[16] Wrist-worn accelerometers are commonly used for sleep assessment, and validation studies comparing them to polysomnography, the gold standard for sleep measurement, have shown that while they are highly sensitive in detecting sleep, their specificity in identifying wakefulness is relatively low.[17] Recently, Johansson et al. have shown that sleep duration can be measured with comparable accuracy using a thigh-worn accelerometer.[18] Using data from the Prospective Physical Activity, Sitting and Sleep (ProPASS) consortium, we examined the independent and joint associations of sleep duration, regularity and efficiency with cardiometabolic health markers. This ProPASS resource consist of harmonized individual participant data of six cohorts from the Netherlands, UK, Australia, Denmark, and Finland with thigh-worn accelerometer data.

Methods

<u>Consortium sample</u>

ProPASS is an international research collaboration platform consisting of observational cohort studies with thigh-worn accelerometry data.[19] For this project, we included data from six participating studies that were pooled and harmonized: The Maastricht Study (TMS; The Netherlands, n=7273)[20], the 1970 British Birth Cohort Study (BCS70; UK, n=4786)[21], the Australian Longitudinal Study on Women's Health (ALSWH; Australia, n=918)[22], the Danish PHysical ACTivity cohort with Objective measurements cohort (DPhacto; Denmark, n=426)[23], the Nijmegen Exercise Study (NES; The Netherlands, n=535)[24], and the Finnish Retirement and Aging Study (FIREA; Finland, n=147)[26]. **Supplementary table 1** presents detailed information for each cohort. Data were physically pooled at the University of Sydney adhering to cohort-specific requirements and necessary data transfer agreements; this included harmonization of covariates and outcomes, as well as cleaning and processing of raw accelerometer data.[27,28] Ethical approval and informed consent were obtained at the cohort level.

<u>Sleep</u>

All cohorts collected movement behaviour data using a 7-day, 24 h/day thigh-worn accelerometer protocol; four studies used ActivPAL devices (BCS70, TMS, ALSWH, NES), one used Axivity devices (FIREA), and one used ActiGraph devices (DPhacto). Raw accelerometer data were centrally

processed using previously validated software, ActiPASS v 1.32[29]. ActiPASS identifies behaviours in 2 s windows with a 50% overlap, resulting in a resolution of 1 s epochs, and implements algorithms for non-wear, sleep detection, posture, and activity intensity.[18,30,31] Validation against polysomnography has shown good sensitivity (0.84) and accuracy (0.80) to detect sleep.[18,31] Participants with a minimum of four valid wear days (≥20 hours per day), including at least one weekend day were included in the analyses. We estimated sleep duration using a validated algorithm that identified thresholds for wakefulness and sleep.[18] Sleep duration (h/day) was categorized into short (<7 h/day), adequate (7-8 h/day), and long (>8 h/day).[1] Sleep efficiency (%) defined as the ratio of total sleep time to total time in bed, was subsequently categorized into tertiles (high: efficiency >91.83; medium, 85.3≤ efficiency ≤ 91.83; low: efficiency<85.3).[32]The Sleep Regularity Index (SRI, %) represents the percentage probability of an individual maintaining the same sleep/wake state on adjacent days, with a scale from 0 (totally irregular) to 100 (perfectly regular).[33] Based on previous research using device data from the UK Biobank, we classified participants into regular (>87.3), slightly irregular (75.9-87.3), and irregular (<75.9) sleep regularity groups[8, 34].

Cardiometabolic health markers

Harmonized cardiometabolic health markers included body mass index, waist circumference, cardiometabolic blood markers, and blood pressure.[27] Body mass index (BMI, kg/m^2; calculated from height and weight) and waist circumference (cm) were assessed by trained nurses or researchers during home or clinic-based visits. Cardiometabolic blood biomarkers were measured in five studies (not available in DPhacto) and included: total and HDL cholesterol (mmol/L), triglycerides (mmol/L), and glycated haemoglobin (HbA1c, mmol/mol; measured in ALSWH, BCS70, and TMS only). LDL cholesterol was calculated based on the Friedewald equation.[35] Measurement and assay methodology were similar across cohorts, with consistently low coefficients of variation (< 6%). Blood pressure measurement protocols were consistent across the six cohorts, each of which used automated blood pressure monitors administered by a research nurse or study staff member to assess systolic and diastolic blood pressure (mmHg). Details on the assessment of blood biomarkers and the measurements across cohort are presented in **supplementary table 2 and 3**. A composite cardiometabolic risk score was computed by averaging the standardized values (z-scores) of eight biomarkers: BMI, waist circumference, HDL cholesterol, LDL cholesterol, triglycerides, HbA1c, diastolic blood pressure, and systolic blood pressure. The score was computed when data were available for at least six of the eight biomarkers, allowing for up to two missing values per participant. To ensure consistency in directionality— where higher scores indicate poorer cardiometabolic health—, the standardized HDL cholesterol values were inverted prior to averaging.

Covariates

Covariates were selected as confounders in the analyses a priori based on data availability and known associations with sleep characteristics and cardiometabolic health markers. The following covariates were collected in all cohorts: age (years), sex (male, female), device-measured moderate to vigorous physical activity (including running, cycling, inclined stepping, and walking with cadence ≥100 steps/min, min/day), smoking status (non-smoker, current smoker), alcohol consumption (tertiles based on self-reported weekly consumption), lipid-modifying, antihypertensive or glucose-lowering medications (yes, no), and history of cardiovascular disease (yes, no). Additionally, a subset of cohorts had data on education (four cohorts; ranging from none or lower than high school to university degree or higher), self-rated health (five-point Likert scale), self-reported diet quality (low, medium, or high vegetable and fruit consumption) and physical function score; which were adjusted for as part of sensitivity analysis. Full details of the measurement and subsequent harmonization of covariates in each cohort are provided in **Supplementary table 3**.

Statistical analyses

The range of all cardiometabolic markers and three sleep indices were Winsorized at the 2.5^{th} and 97.5^{th} percentile to alleviate the impact of sparse data or outliers. We conducted a pooled individual participant data analysis using adjusted generalized linear regression to examine the associations between three categorical sleep exposures and composite cardiometabolic health, as well as individual markers including BMI, waist circumference, HDL cholesterol, LDL cholesterol, triglycerides, HbA1c, and both diastolic and systolic blood pressure. Results are presented as visualizations of beta coefficients (β) with 95% confidence intervals. Additionally, we performed joint association analyses to assess all combinations of sleep duration, regularity, and efficiency consisting of 9 mutually exclusive categories for each combination. Further, we examined a three-way interaction analysis of all sleep parameter (27 combinations) with cardiometabolic health markers, utilizing adjusted generalized linear regression models. Regression assumptions were evaluated through residual diagnostics, including residual plots and leverage versus residual squared plots. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication, previous cardiovascular incidence and moderate to vigorous physical activity; and mutually adjusted for sleep duration or regularity or efficiency, as appropriate.

To assess the associations of sleep exposures as continuous variables, we employed restricted cubic spline adjusted generalized linear models with knots placed at the 10^{th}, 50^{th}, and 90^{th} percentiles.

Non-linearity was examined using Wald tests. Sensitivity analyses were conducted to assess the robustness of findings by further adjusting for diet, education, self-reported health, and physical function in the three-way joint models from available participants. To reduce the influence of reverse causation analyses were repeated after excluding participants with self-reported health as poor or fair (excluded n = 1,789), excluding participants with prevalent or history of cardiovascular disease (excluded n = 1,341), and excluding people with a BMI over 35kg/m^2 (included n = 11,591). We added sensitivity analysis with an alternative defitnion of composite cardiometabolic risk score uses the same set of eight biomarkers on complete cases, that it applies a stricter criterion by excluding individuals with missing data in any biomarkers; (included n= 9,927). We further sensitivity including only those with at least 7 days of valid wear days (≥20 hours per day); (included n= 9,331). Interaction effects were tested between each sleep exposure, age, and sex. All statistical analyses were performed using R (version 4.4.0), with modelling conducted using the rms package (version 6.8.1).

Results

The analytical sample comprised 14,085 participants, with a mean age of 54.3 years and 54.9% were female. Average sleep duration in the pooled cohort was 7.2 (SD: 1.0) hours, with sleep regularity 78.4% and sleep efficiency at 86.8%. **Table 1** shows the baseline characteristics of participants stratified by sleep duration. Compared to those with adequate sleep duration (7–8 hours), short sleepers (<7 hours) were older, had a lower proportion of females, higher smoking rates, poorer self-rated health, greater medication use, a higher prevalence of cardiovascular disease, and higher levels of MVPA. Short sleepers also had a lower level of education and more often had a lower occupational class. Long sleepers (>8 hours), had a higher proportion of females, reported poorer self-rated health, and had lower physical activity levels. The baseline characteristics stratified by SRI and sleep efficiency are shown in **supplemental table 4 and 5**. The correlation between sleep duration and both sleep efficiency and sleep regularity was low (r = 0.28 and r = −0.16, respectively). Sleep efficiency showed a moderate positive correlation with sleep regularity (r = 0.45)(not tabulated)

Sleep duration and cardiometabolic health
Compared to those with adequate sleep duration, participants with shorter sleep duration had significantly higher composite cardiometabolic risk z-score (β: 0.05, 95%CI: 0.03- 0.07) in the fully adjusted model including adjustment for sleep regularity and efficiency (**Figure 1A**). Long sleep

duration was also associated with higher composite cardiometabolic risk score (0.01, -0.03-0.03), albeit not statistically significant. **Supplemental Figure 1** shows the association of sleep duration with the individual cardiometabolic health markers. Compared to an adequate sleep duration, short sleep duration but not long sleep duration was associated with a significantly higher BMI (0.56, 0.39-0.74); waist circumference (1.65, 1.18-2.12); HbA1c (0.58, 0.31- 0.85); and diastolic blood pressure (0.43, 0.02-0.84). There was no association between sleep duration and HDL, LDL cholesterol and triglyceride levels.

Sleep regularity and cardiometabolic health

Participants who had a slightly irregular and irregular sleeping pattern had a higher composite cardiometabolic risk z-score compared to those with regular SRI in the fully adjusted model which included adjustment for sleep duration and efficiency (slightly irregular: 0.07, 0.04-0.09; irregular: 0.14, 0.11-0.18 **Figure 1B**). As for the individual cardiometabolic health markers (**Supplemental Figure 2**), compared to a regular sleep pattern, both slightly irregular and irregular sleep patterns were associated with a higher BMI, a larger waist circumference, lower HDL levels; higher LDL; higher levels of triglycerides; higher HbA1c; and higher diastolic blood pressure. There was no association with systolic blood pressure.

Sleep efficiency and cardiometabolic health

A lower sleep efficiency was also associated with a higher composite cardiometabolic risk z-score in the fully adjusted model which included adjustment for sleep duration and regularity (**Figure 1C**). Compared to participants with high sleep efficiency, those with a medium and low sleep efficiency had a higher composite cardiometabolic risk score (medium: 0.05, 0.03- 0.07, low: 0.10, 0.07-0.13). As for the individual cardiometabolic health markers (**Supplemental Figure 3**), a medium and low efficiency was associated with a higher BMI, a larger waist circumference, lower HDL levels (only statistically significant for low sleep efficiency); higher levels of triglycerides; higher HbA1c (only statistically significant for low sleep efficiency); and higher systolic and diastolic blood pressure.

Continuous sleep characteristics and cardiometabolic health

A reverse J-shaped association was observed for sleep duration (**Figure 2A**) and a more linear association for sleep regularity (**Figure 2B**) and efficiency (**Figure 2C**). Consistent with the categorization above, short sleep duration was associated with a higher composite cardiometabolic risk score with the most favourable cardiometabolic health in those sleeping 7-8 hours. Greater sleep regularity and efficiency were linked to better cardiometabolic profiles.

Joint associations of sleep duration, regularity and efficiency with cardiometabolic health

The joint associations of sleep duration and sleep regularity are shown in **Figure 3**. Compared to adequate sleepers with a regular SRI, individuals with more irregular sleep patterns tended to show higher cardiometabolic risk scores in long, adequate, and short sleepers; the highest cardiometabolic risk score was in short sleepers with an irregular sleep pattern (0.19, 0.14-0.25). **Figure 4** shows the joint association between sleep duration and efficiency indicating that compared to individuals with an adequate sleep duration and a high sleep efficiency, those with a medium and low sleep efficiency had a significantly higher cardiometabolic risk score in all sleep duration groups. Finally **Figure 5** shows joint associations of sleep regularity and sleep efficiency. Compared to those with regular and highly efficient sleep patterns, worse cardiometabolic health was observed in those with a more irregular sleep pattern in those with a medium and low sleep efficiency. The highest cardiometabolic risk score was observed in people with an irregular sleep pattern and a low sleep efficiency (0.24, 0.20-0.28).

The three-way joint associations of sleep characteristics are presented in **Figure 6**. Compared to the group with an adequate sleep duration, regular sleep pattern, and high sleep efficiency, cardiometabolic risk score was highest in groups with medium or low sleep efficiency, combined with an irregular sleep pattern in people with an adequate and long sleep duration. In the short sleepers, all combinations of sleep regularity and efficiency presented worse cardiometabolic health except for the group with a regular SRI and a high sleep efficiency.

Sensitivity analyses

Adjustment for diet and levels of education or physical function did not materially change the results of the joint associations of sleep duration, regularity and efficiency with cardiometabolic risk score (**Supplemental Figure 4** and **Supplemental Figure 6**). Additional adjustment for body fat percentage, which was only available in two of the six cohorts (n=3,373), reduced the regression coefficients; the combination of a long sleep duration with a slightly/irregular patterns and a medium/low sleep efficiency retained significantly higher composite cardiometabolic risk z-scores compared to those with an adequate sleep duration, a regular SRI and a high sleep efficiency (**Supplemental Figure 5**). Results remained similar where participants with poor or fair self-reported health were excluded (**Supplemental Figure 7**); in those with a history of cardiovascular disease were excluded (**Supplemental Figure 8**); in participant with a BMI>35 kg/m^2 excluded (**Supplemental Figure 9**); with the application of an alternative composite cardiometabolic risk score (**Supplemental Figure 10**); or in participants with at least 7 days of valid wear days (**Supplemental Figure 11**). Multiplicative

interactions between each sleep parameter and sex and age were tested; however, no consistent evidence of effect modification by sex or age was observed (**Supplemental Table 6** and **Supplemental Table 7**)

Discussion

In this individual participant data pooled analysis of six cohort studies within the ProPASS consortium, we investigated the independent and joint associations of device-measured sleep duration, regularity, and efficiency with cardiometabolic health outcomes. In these cross-sectional analyses, shorter sleep duration, greater irregularity in sleep patterns, and lower sleep efficiency were independently associated with poorer cardiometabolic health outcomes compared to adequate sleep duration, regular sleep patterns, and high sleep efficiency respectively. Both the 2-way and 3-way joint associations of sleep duration with sleep regularity and sleep efficiency indicated that, regardless of sleep duration, both greater sleep irregularity and lower sleep efficiency were linked to adverse cardiometabolic health profiles. These associations were particularly pronounced in the 3-way joint analysis, suggesting possible additive effects of sleep efficiency and regularity i.e. individuals with adequate, long, or short sleep duration had worse cardiometabolic health when exhibiting either a more irregular sleep pattern or lower sleep efficiency. To our knowledge, our study is the first to investigate the 3-way joint associations of any sleep parameters with cardiometabolic health. Our finding that irrespective of sleep duration, both sleep regularity and sleep efficiency are associated with cardiometabolic health is novel, emphasising the importance of sleep parameters other than duration for cardiometabolic health.

Our study suggests that, compared to an adequate sleep duration, short sleep duration—but not long sleep duration—was associated with poorer cardiometabolic health. This association was consistently observed across individual cardiometabolic health markers, except for elevated triglyceride levels in long sleepers, as well as the composite cardiometabolic risk score. This is consistent with findings from a systematic review and meta-analyses showing that short sleep duration was associated with increased risk of mortality, diabetes, hypertension, cardiovascular disease, and obesity.[4] Our findings that long sleep duration was not associated with cardiometabolic health contrast with previous evidence from a review, which reported that long sleep duration is associated with increased incidence of mortality and several conditions, including stroke, diabetes, and cardiovascular disease, although no significant association was found with hypertension.[3] An umbrella review synthesizing evidence from 11 systematic reviews - 96% of the underlying studies

relied on self-reported sleep data - found a U-shaped association between sleep duration and health outcomes.[1] The dose–response curves from that review align with the findings of the current study indicating that the most favourable sleep duration was approximately 7–8 hours per day which we choose as the reference group in our analyses.

Our results provide novel empirical evidence suggesting that an irregular sleep pattern and low sleep efficiency are both associated with poorer cardiometabolic health, independent of each other and sleep duration. Meta-analyses and systematic reviews have shown that poor sleep quality - as assessed by self-reported measures in the majority of studies - was associated with coronary heart disease,[5] hypertension[36], arterial stiffness[37, 38] and endothelial dysfunction[38] but not with all-cause or cardiovascular disease mortality[5] and atherosclerosis.[38] Systematic reviews on sleep variability - based on self-reported data or accelerometry - and cardiometabolic health have shown associations with obesity, [9, 39] cardiovascular disease, and metabolic syndrome[9,40] but not with insulin resistance.[9] It is important to note that sleep variability in the mostly observational studies underlying these reviews has been assessed with different indicators such as 'social jetlag' as the variation between working days and free days, and the standard deviation of sleep duration of sleep across multiple days. We chose to operationalize the SRI as this parameter is increasingly prioritized as an objective metric for day-to-day variability in sleep wake cycle. A recent study using data from the UK Biobank investigating joint associations between sleep duration and sleep regularity shows that meeting sleep duration recommendations did not offset the adverse effect of an irregular sleep patterns for the risk type 2 diabetes and major adverse cardiovascular events.[8]

In a 2023 consensus statement by the US National Sleep Foundation, the expert panel acknowledged the importance of sleep regularity for health and performance.[40] Further, a recent 2025 scientific statement from the American Heart Association outlines the concept of multidimensional sleep health and highlights the multiple dimensions of sleep that are relevant for cardiometabolic health.[41] It emphasizes that while sleep duration has been widely studied and thereby informed the development of guidelines,[42] more research is needed to examine multiple sleep dimensions. This broader focus is necessary to support the creation of comprehensive public health guidelines and clinical interventions.

While there is substantial evidence to support the biological causal link between short sleep duration and adverse health outcomes, the biological pathway of long sleep and health remains less clear. Long sleep may be related to poorer sleep quality and sleep fragmentation imposed by other

health conditions; therefore there is a possibility that it reflects reverse causation or residual confounding more than causal effects on cardiometabolic health.[43] In the current study, our finding indicate that individuals with long sleep duration combined with either an irregular sleep pattern or low sleep efficiency exhibit poorer cardiometabolic health. This finding suggests that it is not long sleep duration, but its interaction with other adverse sleep characteristics that is associated with negative health outcomes. Sleep irregularity may result in circadian misalignment, a disconnect between the central and peripheral clocks. Short-term circadian misalignment has been associated with acute metabolic disturbances including an increase in postprandial glucose, insulin and mean arterial pressure, decreases in leptin and sleep efficiency,[44] increases in 24-hour blood pressure, and inflammation.[10] These mechanisms may explain how circadian misalignment may lead to increased cardiovascular disease risk.[9]

Strengths of our study include the large sample size (n= 14,085) drawn from six cohorts across five countries and the device-based assessment of multiple sleep parameters. By examining the joint associations of three sleep characteristics with cardiometabolic health, our study contributes to the growing body of evidence that sleep is a multidimensional construct, which should be evaluated beyond sleep duration alone. Our study also has some limitations that should be acknowledged. First, our study was cross-sectional, and reverse causation is possible - that is, poorer health may have led to less optimal sleep parameters. A limitation of relying on accelerometry alone to estimate sleep parameters is that it is less accurate in distinguishing between quiet reclined wakefulness and actual sleep, potentially leading to exposure misclassification and underestimation of any true associations that may exist. Further, we did not have data on clinical sleeping disorders such as sleep apnoea and insomnia. Finally, adjustments were limited to selected confounders that were available in all cohorts and we cannot exclude residual confounding. Some of the additional adjustments, e.g. for shift work was not possible because of data availability and additional adjustment for fat mass was only possible in a smaller sample as it was not measured in all cohorts.

Conclusion

These findings suggest the importance of adopting a multidimensional approach to sleep health, demonstrating that sleep duration alone may not fully capture its relationship with cardiometabolic risk. By demonstrating the joint associations of sleep duration, regularity, and efficiency, this study contributes to the growing evidence that multidimensional sleep profiles are more informative than single sleep parameter. Future longitudinal studies are needed to further clarify these associations

across diverse populations and to evaluate the effectiveness of interventions aimed at promoting consistent sleep patterns and enhancing sleep efficiency as strategies to reduce cardiometabolic risk.

Acknowledgements, Sources of Funding, Disclosures

Acknowledgements

The data on which this research is based were drawn from six observational studies in the Netherlands, UK, Australia, Denmark, and Finland. We are grateful to all participants who provided the survey data.

Sources of Funding

This project was funded by a British Heart Foundation Special Grant (SP/F/20/150002) and National Health and Medical Research Council (Australia) Investigator (APP1194510) and Ideas (APP1180812) Grants. The establishment of the ProPASS consortium was supported by an unrestricted 2018-20 grant by PAL Technologies (Glasgow, UK). ActiPASS development was partly funded by FORTE, Swedish Research Council for Health, Working Life and Welfare (2021-01561). M.N.A is supported by The National Heart Foundation (APP 107158). E.S. is funded by a National Health and Medical Research Council Investigator Grant (APP1194510). G.M. is supported by a National Health and Medical Research Council Principal Research Fellowship (APP1121844). A.D.H. receives support from the British Heart Foundation, the Horizon 2020 Framework Programme of the European Union, the National Institute for Health Research University College London Hospitals Biomedical Research Centre, the UK Medical Research Council, the National Institute for Health Research, and the Wellcome Trust and works in a unit that receives support from the UK Medical Research Council. E.A.B. and T.M.H.E are supported by the ActiveLIFE consortium grant (01-001-2024-0623) of the Dutch Heart Foundation

Disclosures

All authors declare no disclosure of interest for this contribution.

Details of individual cohort funding

The Maastricht Study was supported by the European Regional Development Fund via OP-Zuid, the Province of Limburg, the Dutch Ministry of Economic Affairs (grant 31O.041), Stichting De Weijerhorst (Maastricht, the Netherlands), the Pearl String Initiative Diabetes (Amsterdam, the Netherlands), the Cardiovascular Center (CVC, Maastricht, the Netherlands), CARIM School for Cardiovascular Diseases (Maastricht, the Netherlands), CAPHRI Care and Public Health Research Institute (Maastricht, the Netherlands), NUTRIM School for Nutrition and Translational Research in

Metabolism (Maastricht, the Netherlands), Stichting Annadal (Maastricht, the Netherlands), Health Foundation Limburg (Maastricht, the Netherlands), and by unrestricted grants from Janssen-Cilag BV (Tilburg, the Netherlands), Novo Nordisk Farma BV (Alphen aan den Rijn, the Netherlands), and Sanofi-Aventis Netherlands BV (Gouda, the Netherlands).

The 1970 British Cohort Study is funded by the Economic and Social Research Council (ES/M001660/1). The age 46 sweep was also funded by the British Heart Foundation (grant SP/15/6/31397) and a joint award from the Economic and Social Research Council and the Medical Research Council (grant RES-579-47-0001).

The ALSWH is funded by the Australian Government Department of Health, Disability and Ageing and its substudy (Menarche-to-PreMenopause), from which accelerometry and clinical data were obtained, was funded by the National Health and Medical Research Council Project Grant (APP1129592); and partly supported by NHMRC Investigator (APP1194510) and Ideas (APP1180812) Grants.

DPhacto is funded by The Danish Working Environment Research Fund.

NES is supported by the Department of Physiology of the Radboud University Medical Center, the Dutch Heart Foundation (2020T063), and Siemens Healthcare Diagnostics (the Hague, Netherlands).

FIREA is supported by the Academy of Finland (286294, 294154, 319246, 332030, 361780), Ministry of Education and Culture, Juho Vainio Foundation and Finnish State Grants for Clinical Research.

References

1. Chaput, J.-P. *et al.* Sleep duration and health in adults: an overview of systematic reviews. *Appl. Physiol. Nutr. Metab.* **45**, S218–S231 (2020).

2. Krittanawong, C. *et al.* Association between short and long sleep durations and cardiovascular outcomes: a systematic review and meta-analysis. *Eur. Heart J. Acute Cardiovasc. Care* **8**, 762–770 (2019).

3. Jike, M., Itani, O., Watanabe, N., Buysse, D. J. & Kaneita, Y. Long sleep duration and health outcomes: A systematic review, meta-analysis and meta-regression. *Sleep Med. Rev.* **39**, 25–36 (2018).

4. Itani, O., Jike, M., Watanabe, N. & Kaneita, Y. Short sleep duration and health outcomes: a systematic review, meta-analysis, and meta-regression. *Sleep Med* **32**, 246–256 (2017).

5. Kwok, C. S. *et al.* Self-Reported Sleep Duration and Quality and Cardiovascular Disease and Mortality: A Dose-Response Meta-Analysis. *J. Am. Heart Assoc.* **7**, e008552 (2018).

6. Yan, B. *et al.* Objective Sleep Efficiency Predicts Cardiovascular Disease in a Community Population: The Sleep Heart Health Study. *J. Am. Heart Assoc.* **10**, e016201 (2021).

7. Chaput, J.-P. *et al.* Sleep timing, sleep consistency, and health in adults: a systematic review. *Appl. Physiol. Nutr. Metab.* **45**, S232–S247 (2020).

8. Chaput, J.-P. *et al.* Sleep regularity and major adverse cardiovascular events: a device-based prospective study in 72 269 UK adults. *J. Epidemiol. Community Health* **79**, 257–264 (2025).

9. Zhu, B. *et al.* Associations between sleep variability and cardiometabolic health: A systematic review. *Sleep Med. Rev.* **66**, 101688 (2022).

10. Lunsford-Avery, J. R., Engelhard, M. M., Navar, A. M. & Kollins, S. H. Validation of the Sleep Regularity Index in Older Adults and Associations with Cardiometabolic Risk. *Sci. Rep.* **8**, 14158 (2018).

11. Huang, T. & Redline, S. Cross-sectional and Prospective Associations of Actigraphy-Assessed Sleep Regularity With Metabolic Abnormalities: The Multi-Ethnic Study of Atherosclerosis.

Diabetes Care **42**, 1422–1429 (2019).

12. Huang, T., Mariani, S. & Redline, S. Sleep Irregularity and Risk of Cardiovascular Events. *J. Am. Coll. Cardiol.* **75**, 991–999 (2020).

13. Fritz, J. *et al.* Cross-sectional and prospective associations between sleep regularity and metabolic health in the Hispanic Community Health Study/Study of Latinos. *Sleep* **44**, zsaa218 (2021).

14. Chaput, J.-P. *et al.* Sleep Irregularity and the Incidence of Type 2 Diabetes: A Device-Based Prospective Study in Adults. *Diabetes Care* **47**, 2139–2145 (2024).

15. Lauderdale, D. S., Knutson, K. L., Yan, L. L., Liu, K. & Rathouz, P. J. Self-Reported and Measured Sleep Duration: How Similar Are They? *Epidemiology* **19**, 838–845 (2008).

16. Smith, M. T. *et al.* Use of Actigraphy for the Evaluation of Sleep Disorders and Circadian Rhythm Sleep-Wake Disorders: An American Academy of Sleep Medicine Clinical Practice Guideline. *J. Clin. Sleep Med.* **14**, 1231–1237 (2018).

17. Conley, S. *et al.* Agreement between actigraphic and polysomnographic measures of sleep in adults with and without chronic conditions: A systematic review and meta-analysis. *Sleep Med. Rev.* **46**, 151–160 (2019).

18. Johansson, P. J. *et al.* Development and performance of a sleep estimation algorithm using a single accelerometer placed on the thigh: an evaluation against polysomnography. *J. Sleep Res.* **32**, e13725 (2023).

19. Stamatakis, E. *et al.* Emerging collaborative research platforms for the next generation of physical activity, sleep and exercise medicine guidelines: the Prospective Physical Activity, Sitting, and Sleep consortium (ProPASS). *Br. J. Sports Med.* **54**, 435–437 (2020).

20. Schram, M. T. *et al.* The Maastricht Study: an extensive phenotyping study on determinants of type 2 diabetes, its complications and its comorbidities. *Eur. J. Epidemiol.* **29**, 439–451 (2014).

21. Elliott, J. & Shepherd, P. Cohort Profile: 1970 British Birth Cohort (BCS70). *Int. J. Epidemiol.* **35**, 836–843 (2006).

22. Chan, H.-W. *et al.* Cohort profile: a prospective Australian cohort study of women's reproductive characteristics and risk of chronic disease from menarche to premenopause (M-PreM). *BMJ Open* **12**, e064333 (2022).

23. Jørgensen, M. B. *et al.* The DPhacto cohort: An overview of technically measured physical activity at work and leisure in blue-collar sectors for practitioners and researchers. *Appl. Ergon.* **77**, 29–39 (2019).

24. Van Der Sluijs, K. M. *et al.* Cohort Profile: The Nijmegen Exercise Study (NES). *Int. J. Epidemiol.* **54**, (2025).

25. Maessen, M. F. H. *et al.* Lifelong Exercise Patterns and Cardiovascular Health. *Mayo Clin. Proc.* **91**, 745–754 (2016).

26. Leskinen, T. *et al.* Changes in non-occupational sedentary behaviours across the retirement transition: the Finnish Retirement and Aging (FIREA) study. *J. Epidemiol. Community Health* **72**, 695–701 (2018).

27. Blodgett, J. M. *et al.* Device-measured physical activity and cardiometabolic health: the Prospective Physical Activity, Sitting, and Sleep (ProPASS) consortium. *Eur. Heart J.* **45**, 458–471 (2024).

28. Ahmadi, M. N. *et al.* Relationship of device measured physical activity type and posture with cardiometabolic health markers: pooled dose–response associations from the Prospective Physical Activity, Sitting and Sleep Consortium. *Diabetologia* **67**, 1051–1065 (2024).

29. Hettiarachchi, P. & Johansson, P. ActiPASS. Zenodo https://doi.org/10.5281/zenodo.15181460 (2025).

30. Skotte, J., Korshøj, M., Kristiansen, J., Hanisch, C. & Holtermann, A. Detection of physical activity types using triaxial accelerometers. *J. Phys. Act. Health* **11**, 76–84 (2014).

31. Hettiarachchi, P. *et al.* Validity of a Non-Proprietary Algorithm for Identifying Lying Down Using Raw Data from Thigh-Worn Triaxial Accelerometers. *Sensors* **21**, 904 (2021).

32. Ibáñez, V., Silva, J. & Cauli, O. A survey on sleep assessment methods. *PeerJ* **6**, e4849 (2018).

33. Phillips, A. J. K. *et al.* Irregular sleep/wake patterns are associated with poorer academic performance and delayed circadian and sleep/wake timing. *Sci. Rep.* **7**, 3216 (2017).

34. Windred, D. P. *et al.* Objective assessment of sleep regularity in 60 000 UK Biobank participants using an open-source package. *Sleep* **44**, zsab254 (2021).

35. Samuel, C. *et al.* Accuracy of 23 Equations for Estimating LDL Cholesterol in a Clinical Laboratory Database of 5,051,467 Patients. *Glob. Heart* **18**, 36 (2023).

36. Lo, K., Woo, B., Wong, M. & Tam, W. Subjective sleep quality, blood pressure, and hypertension: a meta-analysis. *J. Clin. Hypertens.* **20**, 592–605 (2018).

37. Saz-Lara, A. Association between sleep duration and sleep quality with arterial stiffness: A systematic review and meta-analysis. *Sleep Health* (2022).

38. Aziz, M. *et al.* Association of Subjective and Objective Sleep Duration as well as Sleep Quality with Non-Invasive Markers of Sub-Clinical Cardiovascular Disease (CVD): A Systematic Review.

39. Arab, A., Karimi, E. & Garaulet, M. Social jetlag and obesity: A systematic review and meta-analysis.

40. Sletten, T. L. The importance of sleep regularity_ a consensus statement of the National Sleep Foundation sleep timing and variability panel. *Sleep Health* (2023).

41. St-Onge, M.-P. *et al.* Multidimensional Sleep Health: Definitions and Implications for Cardiometabolic Health: A Scientific Statement From the American Heart Association. *Circ. Cardiovasc. Qual. Outcomes* e000139 (2025) doi:10.1161/HCQ.0000000000000139.

42. Lloyd-Jones, D. M. *et al.* Life's Essential 8: Updating and Enhancing the American Heart Association's Construct of Cardiovascular Health: A Presidential Advisory From the American Heart Association.

43. Chaput, J., Dutil, C. & Sampasa-Kanyinga, H. Sleeping hours: what is the ideal number and how does age impact this. *Nat Sci Sleep* **10**, 421–430 (2018).

44. Scheer, F. A. J. L., Hilton, M. F., Mantzoros, C. S. & Shea, S. A. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc. Natl. Acad. Sci.* **106**, 4453–4458

(2009).

45. Morris, C. J., Purvis, T. E., Hu, K. & Scheer, F. A. J. L. Circadian misalignment increases cardiovascular disease risk factors in humans. *Proc. Natl. Acad. Sci.* **113**, (2016).

Association of device-based measured sleep duration, sleep efficiency and sleep regularity with cardiometabolic health: the Prospective Physical Activity, Sitting, and Sleep (ProPASS) consortium

Table 1: Baseline characteristics of participants stratified by sleep duration.

Baseline characteristic	Short (< 7 hours)	Adequate (7-8 hours)	Long (>9 hours)	Full sample
Sample size	6,008	5,179	2,898	14,085
Age (mean (SD))	55.3 (9.5)	53.7 (9.4)	53.2 (9.4)	54.3 (9.5)
Sex = Female, n (%)	2,714 (45.2)	3,064 (59.2)	1,954 (67.4)	7,732 (54.9)
Cohort, n (%)				
ALSWH	250 (4.2)	384 (7.4)	284 (9.8)	918 (6.5)
BCS70	1,769 (29.4)	1,822 (35.2)	1,195 (41.2)	4,786 (34.0)
DPhacto	231 (3.8)	158 (3.1)	37 (1.3)	426 (3.0)
FIREA	91 (1.5)	45 (0.9)	11 (0.4)	147 (1.0)
NES	269 (4.5)	201 (3.9)	65 (2.2)	535 (3.8)
TMS	3,398 (56.6)	2,569 (49.6)	1,306 (45.1)	7,273 (51.6)
Alcohol consumption[7], n (%)				
Tertile 1 (least)	1,667 (31.6)	1,529 (34.0)	951 (39.2)	4,147 (34.0)
Tertile 2 (moderate)	1,778 (33.7)	1,559 (34.7)	808 (33.3)	4,145 (34.0)
Tertile 3 (highest)	1,829 (34.7)	1,407 (31.3)	665 (27.4)	3,901 (32.0)
Smoking = Current smoker, n (%)	932 (15.6)	651 (12.6)	392 (13.6)	1,975 (14.1)
Education, n (%)				
None or less than high school	595 (10.6)	518 (10.5)	412 (14.6)	1,525 (11.4)
High school (⬜16 years)	1,500 (26.7)	1,313 (26.7)	870 (30.9)	3,683 (27.6)
Further education (⬜16–18 years)	2,338 (41.6)	1,929 (39.2)	937 (33.3)	5,204 (39.0)
University degree or higher	1,182 (21.1)	1,157 (23.5)	595 (21.1)	2,934 (22.0)
Self-rated health, n (%)				
Excellent	666 (11.3)	665 (13.0)	335 (11.8)	1,666 (12.0)
Very good	1,826 (31.0)	1,733 (34.0)	870 (30.5)	4,429 (32.0)
Good	2,648 (44.9)	2,127 (41.7)	1,181 (41.5)	5,956 (43.0)

Fair	661 (11.2)	507 (9.9)	365 (12.8)	1,533 (11.1)
Poor	93 (1.6)	65 (1.3)	98 (3.4)	256 (1.8)
Cardiometabolic medication use[1], n (%)	1,973 (35.0)	1,319 (26.9)	805 (28.5)	4,097 (30.7)
Prevalent CVD, n (%)	660 (11.1)	452 (8.8)	292 (10.1)	1,404 (10.0)
MVPA mins per day (mean (SD)) [5]	78.1 (29.5)	74.4 (27.6)	63.3 (26.5)	73.7 (28.7)
Diet[6], n (%)				
Low quality	1,393 (38.5)	1,135 (38.8)	691 (45.3)	3,219 (39.9)
Medium quality	1,370 (37.9)	1,134 (38.7)	569 (37.3)	3,073 (38.1)
High quality	852 (23.6)	659 (22.5)	264 (17.3)	1,775 (22.0)
Body Mass Index, kg/m^2 (mean (SD))	27.2 (4.5)	26.4 (4.3)	27.0 (4.6)	26.8 (4.5)
Waist circumference, cm (mean (SD))	95.7 (13.2)	92.0 (12.8)	93.1 (13.3)	93.8 (13.2)
HDL[2], mmol/L (mean (SD))	1.5 (0.4)	1.6 (0.4)	1.6 (0.4)	1.6 (0.4)
LDL[3], mmol/L (mean (SD))	3.4 (0.9)	3.4 (0.9)	3.4 (0.9)	3.4 (0.9)
Triglycerides, mmol/L (mean (SD))	1.5 (0.8)	1.4 (0.7)	1.5 (0.8)	1.4 (0.8)
HbA1C[4], mmol/mol (mean (SD))	38.4 (7.4)	37.0 (6.5)	37.5 (6.9)	37.7 (7.0)
Systolic blood pressure, mmHg (mean (SD))	131.7 (16.6)	128.7 (16.4)	127.6 (16.6)	129.8 (16.6)
Diastolic blood pressure, mmHg (mean (SD))	77.1 (9.9)	76.0 (9.8)	75.9 (9.8)	76.5 (9.8)
Sleep regularity index (mean (SD))	79.4 (10.1)	79.6 (9.6)	73.9 (11.9)	78.4 (10.6)
Sleep duration, h/day (mean (SD))	6.2 (0.6)	7.5 (0.3)	8.6 (0.5)	7.2 (1.0)
Sleep efficiency (mean (SD))	84.4 (9.7)	89.1 (7.1)	87.7 (8.0)	86.8 (8.7)

SRI: Sleep Regularity Index; SD: standard deviation.

[1]Lipid-modifying, hypertensive, and glucose-lowering medications; [2]High-density lipoprotein cholesterol; [3]Low-density lipoprotein cholesterol, [4]Glycated Hemoglobin, [5]Accelerometer-derived moderate-to-vigorous physical activity, [6] Self-reported daily intake of fruits and vegetables served as a proxy for dietary, [7]Self-reported alcohol consumption.

Figures

Figure 1: Association of sleep duration (A), regularity (B), and efficiency (C) with composite cardiometabolic risk score.

Legend: Composite cardiometabolic risk score, z-score, was derived from the composite sample distribution of BMI, waist circumference, HDL cholesterol, LDL cholesterol, triglycerides, glycated haemoglobin (HbA1c), systolic and diastolic blood pressure. Here, a higher score represents poorer cardiometabolic health. Sleep duration was categorized into short (<7h/day), adequate (7-8h/day) and long (>8h/day), sleep regularity (irregular, SRI<75.9; slightly irregular, 75.9≤SRI≤ 87.3; regular, SRI> 87.3) and sleep efficiency (low, efficiency<85.3; medium, 85.3≤ efficiency ≤ 91.83; high, efficiency >91.83) were categorized into tertiles. Reference group was set to adequate, regular and

high, for sleep duration, SRI and efficiency. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication, previous cardiovascular incidence and moderate to vigorous physical activity. All sleep characteristics were mutually adjusted. GLM coefficients represent the mean differences between the reference group and each of the other groups. N= 12,307. SRI: Sleep regularity index, BMI: body mass index.

Figure 2: Association of (A) sleep duration, (B) regularity, and (C) efficiency with composite cardiometabolic risk score.

Legend: Composite cardiometabolic risk score, z-score, was derived from the composite sample distribution of BMI, waist circumference, HDL cholesterol, LDL cholesterol, triglycerides, glycated haemoglobin (HbA1c), systolic and diastolic blood pressure. Here, a higher score represents poorer cardiometabolic health. Sleep duration, sleep regularity and sleep efficiency were continuous. Reference for each group was the lowest value in the distribution. Knots for each exposure were placed at 10th, 50th and 90th percentiles. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication, previous cardiovascular incidence and moderate to vigorous physical activity. All sleep characteristics were mutually adjusted. N= 12,307. SRI: Sleep regularity index, BMI: body mass index.

Figure 3: Joint association of sleep duration and regularity with composite cardiometabolic risk score.

Legend: Sleep duration was categorized into short (<7h/day), adequate (7-8h/day) and long (>8h/day) and sleep regularity (irregular, SRI<75.9; slightly irregular, 75.9≤SRI≤ 87.3; regular, SRI> 87.3) were categorized into tertiles. Reference group was set to adequate duration with regular sleep. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication, previous

cardiovascular incidence, moderate to vigorous physical activity and sleep efficiency. N= 12,307. GLM coefficients represent the mean differences between the reference group and each of the other groups. SRI: sleep regularity index.

Figure 4: Joint association of sleep duration and efficiency with composite cardiometabolic risk score.

Legend: Sleep duration was categorized into short (<7h/day), adequate (7-8h/day) and long (>8h/day) and sleep efficiency (low, efficiency<85.3; medium, 85.3≤ efficiency ≤ 91.83; high, efficiency >91.83) were categorized into tertiles. Reference group was set to adequate duration with, high efficiency of sleep. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication,

previous cardiovascular incidence, moderate to vigorous physical activity and sleep regularity. N= 12,307. GLM coefficients represent the mean differences between the reference group and each of the other groups. SRI: sleep regularity index.

Figure 5: Joint association of sleep regularity and efficiency with composite cardiometabolic risk score.

Legend: Sleep regularity (irregular, SRI<75.9; slightly irregular, 75.9≤SRI≤ 87.3; regular, SRI> 87.3) and sleep efficiency (low, efficiency<85.3; medium, 85.3≤ efficiency ≤ 91.83; high, efficiency >91.83) were categorized into tertiles. Reference group was set to regular with high efficiency of sleep. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication, previous

cardiovascular incidence, moderate to vigorous physical activity and sleep duration. N= 12,307. GLM coefficients represent the mean differences between the reference group and each of the other groups. SRI: sleep regularity index.

Figure 6: Three-way Joint association of sleep duration, regularity and efficiency with composite cardiometabolic risk score.

Legend: Sleep duration was categorized into short (<7h/day), adequate (7-8h/day) and long (>8h/day), sleep regularity (irregular, SRI<75.9; slightly irregular, 75.9≤SRI≤ 87.3; regular, SRI>87.3) and sleep efficiency (low, efficiency<85.3; medium, 85.3≤ efficiency ≤ 91.83; high, efficiency >91.83) were categorized into tertiles. Reference group was set to adequate duration with regular and high

efficiency of sleep. All models were adjusted for age, sex, cohort, smoking, alcohol consumption, medication, previous cardiovascular incidence and moderate to vigorous physical activity. N= 12,307. GLM coefficients represent the mean differences between the reference group and each of the other groups. SRI: sleep regularity index.