TITLE PAGE

Title of the article:

CDS-PD: A Novel Clinical Decision Support Platform for Parkinson's Disease

Authors:

Deepak K. Gupta, MD[1,2,3*], Pedram Golnari, MD[4], Katrina Prantzalos MS[4], Ian Zurlo[1], Vivikta Iyer, MBBS[1,2], Brenna M. Lobb, MS, MPH, CCRP[5,6], Cole Zweber, MS[1,2], Manu Bulusu, MS[4], Dakota Clarke, BS MPA[5,6], James T. Boyd, MD[1,2], Curtis Tatsuoka, PhD[7], Amie L. Hiller, MD[5,6], Satya S. Sahoo, PhD[4*]

[1]Department of Neurology, University of Vermont Medical Center, Burlington, USA

[2]Department of Neurological Sciences, College of Medicine, University of Vermont, Burlington, USA

[3]Clinical Informatics, University of Vermont Health Network, Burlington, USA

[4]Department of Population & Quantitative Health Sciences, School of Medicine, Case Western Reserve University, Cleveland, USA

[5]Department of Neurology, Oregon Health & Sciences University, Portland, USA

[6]Department of Neurology, Parkinson's Disease Research Education and Clinical Center, VA Portland Health Care System, Portland, USA

[7]Director, University of Maryland Marlene & Stewart Greenebaum Comprehensive Cancer Center Biostatistics Shared Resource Center

* Corresponding Authors: drdkgupta@gmail.com (802-489-6886), satya.sahoo@case.edu (216-368-0532)

Keywords:

Informatics, Diagnosis, Parkinson Disease

Word count (excluding title page, abstract, references, figures and tables): 2146

ABSTRACT

Objectives: The Movement Disorders Society clinical diagnostic criteria for Parkinson's disease (MDS-PD) allow highly sensitive and specific diagnosis of Parkinson's disease. However, their adoption has been limited due to lack of a clinical decision support (CDS) tool to support clinicians and researchers in systematically and accurately applying the MDS-PD criteria

Materials and Methods: We have developed and evaluated a CDS platform for PD (CDS-PD) as a modular and extensible informatics platform with comprehensive functionalities for recording relevant patient information. We have performed real time application of diagnostic algorithm of the MDS-PD criteria.

Results: The CDS-PD platform shows high concordance with application of the MDS-PD criteria by experienced movement disorders neurologists.

Discussion: The CDS-PD platform is a transformative step towards realizing the standardized electronic implementation of the MDS-PD criteria for PD patient care and clinical trials at point-of-care.

Conclusion: The CDS-PD platform can be accessed after registration at weblink https://www.cdspd.org

LAY SUMMARY

Diagnostic criteria for Parkinson's disease (PD) have been developed to improve accuracy of clinical diagnosis of PD. However, these criteria are inherently complex to apply using a manual approach with pen-and-paper and are not supported in electronic health record systems. There is a need for a clinical decision support (CDS) system to support clinicians and researchers alike in the implementation of these criteria. We developed a novel technology platform to address this unmet need.

MAIN TEXT

BACKGROUND & SIGNIFICANCE

Parkinson's disease (PD) is the second most common neurodegenerative disorder after Alzheimer's disease[1] and it is the fastest growing neurological disorder globally in terms of prevalence and disability[2]. There is an increasing concern in the healthcare community that these worldwide prevalence trends point to a potential PD epidemic[3]. Therefore, an early and accurate diagnosis of PD is crucial for both clinical care and research using systematic application of validated formal diagnostic criteria. Specifically, the accuracy of clinical diagnosis without the application of diagnostic criteria can be low in primary care and even in general neurology clinics where movement disorders specialty expertise may not be readily available[4]. Achieving a high accuracy of diagnosis is particularly important as neuroprotective clinical trials[5] and preventative strategies[6] focus on early-stage PD[5-8], which is especially challenging to differentiate from related atypical parkinsonian disorders (APD), namely dementia with Lewy bodies[9], multiple system atrophy[10], progressive supranuclear palsy[11], and corticobasal degeneration[12].

The development of the International Parkinson and Movement Disorder Society (MDS) Clinical Diagnostic Criteria for Parkinson's disease (MDS-PD Criteria)[13] represents a critical step forward towards aiding accurate early diagnosis of PD in clinical care and trials. Since their publication over a decade ago in 2010, the MDS-PD criteria have been validated by expert neurologists with 10 years or more of experience in PD[14] and against post-mortem neuropathological diagnosis[15], and thus considered global standard for PD diagnosis both for clinical and research use[16]. However, applying the MDS-PD criteria using the existing paper-based format requires the neurologists to address two key challenges: (1) capturing a diverse set of patient information, including time dependent variables, with consistent quality; and (2) manually applying a complex, multi-step algorithm to generate an accurate diagnosis. In context of universal adoption of the electronic health record (EHR) systems in the United States[17], the current lack of support for the MDS-PD criteria in leading EHR systems has severely limited their adoption in real-world clinical practice and research. Thus, there is a clear and urgent unmet need to develop clinical decision support (CDS) to support the neurologist in applying the MDS-PD criteria at point-of-care.

We address this above gap through the first CDS platform for PD (CDS-PD) that implements the MDS-PD criteria using an advanced informatics software stack and a distributed deployment architecture for supporting multicenter research study. The unique CDS-PD platform comprehensively addresses the two integral components of CDS: (1) accurate capture of patient's information through a neurologist designed & structured user interface intended to represent existing clinical workflows; and (2) real-time application of algorithm on the recorded information to perform diagnostic classification on the MDS-PD criteria.

OBJECTIVES

The CDS-PD platform realizes the transformative potential of the MDS-PD criteria in enhancing early and accurate diagnosis of PD patients at the point of care. It is built for multi-institution, distributed deployment with modular design to evolve with changing diagnosis guidelines and clinical needs of PD patients.

METHODS

The MDS-PD criteria[13] consist of four categories of different criteria and an algorithm: essential criteria (*which is a perquisite*), absolute exclusion criteria and red flags criteria (*which collectively constitute negative features*) and supportive criteria (*which constitute positive features*); the algorithm involves establishing essential criteria, ruling out absolute exclusion criteria, and with weighing/counterbalancing of red flags criteria with supportive criteria to allow classification of the patient into two distinct levels of diagnostic certainty for PD, namely, "*Clinically Established PD*", "*Clinically Probable PD"*, or "*Not PD*" (figure 1). Of note, the MDS-PD criteria can only be operationalized on the patient information (history, neurological examination, and relevant diagnostic tests) captured by a clinician with sufficient training in neurology[16].

The CDS-PD platform incorporates an extensible design framework with integrated modules for data recording and analysis features that are implemented using the Django web development software stack[18]. The Django software uses a Model-View-Controller (MVC) architecture with the Model storing the data in an object relational model, the View encoding the user interface, and the Controller encoding the functionalities of the application, including user interactions. The CDS-PD platform uses PostgreSQL relational database management system to manage data and incorporates a role-based access control (RBAC) module to support user accounts and access. The CDS-PD platform is accessible as a browser-based web application with support for multi-institution clinical research studies.

We systematically evaluated the CDS-PD platform as part of a prospective clinical research study investigating diagnostic and prognostic differences between PD patients with or without history of exposure to agent orange. This multicenter study is approved by the single institutional review board (IRB), WCG IRB, and is being conducted at the University of Vermont Medical Center (UVMMC), the Oregon Health & Sciences University (OHSU), and the Veteran Affairs Portland Healthcare System (VAPORHCS), with grant funding from the Congressionally Directed Medical Research Programs (CDMRP) through the Department of Defense (DoD)[19].

RESULTS

The CDS-PD platform currently consists of two modules: (1) patient information recording module; and (2) patient information analysis module. The former module allows systematic and detailed entry of patient information while ensuring data quality rules, while the latter allows subsequent automated application of algorithm of the MDS-PD criteria (as detailed below and depicted in a video walkthrough of this module is provided in supplementary material 1).

Patient Information Recording Module: This module closely models clinical workflow typically used by movement disorders specialists for evaluation of a patient for PD diagnosis using the MDS-PD criteria. It is implemented using a tab-structured interface consisting of 11 sections/tabs: *demographics, motor symptoms, non-motor symptoms, family history, social & environmental history, occupational & military history, health profile, medications, exam, scales, domain-specific information (DSI), and diagnostics.* These sections also incorporate the National Institute of Health (NIH) Common Data Elements (CDE), for example, family history CDE for Parkinson's disease[20].

Each information category/tab is further organized into appropriate subsections/subtabs corresponding to the required level of granularity. For example, the Exam section/tab is organized into five subsections: *General Exam, Neurological Exam, Basic Movement Disorders Exam, Additional Movement Disorders Exam, and Advanced Movement Disorders Exam* (figure 2). The Basic Movement Disorders exam subsection/subtab captures the patient information corresponding to the most fundamental motor examination features of PD, specifically, parkinsonism (bradykinesia, rigidity, rest tremor), action tremor (postural, kinetic tremor), and dyskinesia.

The user interface (UI) consists of intuitive selection buttons, drop-down menus, calendar dates, and in a few cases free text fields. The UI incorporates jQuery validation libraries and rule-based validation checks for: (1) correct data type for a data entry field, (2) range value: to verify the validity of values within predefined limits for a given data element, and (3) value presence: to ensure that mandatory fields are completed by the user. These built-in data quality checks reduce the occurrence of missing information as well as incorrect data entries with minimal additional burden on users.

Patient Information Analysis Module: This module implements the four categories of criteria (essential, absolute exclusion, supportive, red flags) and the algorithm of the MDS-PD criteria as series of queries to the PostgreSQL database management system (DBMS) and leverages the DBMS query optimizer for fast execution of the classification algorithm. Specifically, each of the criteria under the four categories are subdivided into components and subcomponents (as applicable) and mapped to corresponding patient information variables stored in the database, followed by analysis of the variable values according to the condition/logic of the criteria. For example, parkinsonism is the only essential criteria category that consists of three components (*bradykinesia, rigidity, and rest tremor*). These components are first assessed in terms of body part (*upper limb, lower limb, neck*) and laterality (*right or left; for limb only*) as part of the Movement Disorder Society-Unified Parkinson's Disease Rating Scale (MDS-UPDRS)[21] and then evaluated to be true or false on the condition/logic of "*bradykinesia must be present with or without rigidity or rest tremor*" (figure 3, top panel). The rest of the 23 criteria under the other three categories have been similarly implemented.

Once all 24 criteria have been evaluated to a Boolean value (true or false) using the recorded patient information, corresponding values are then evaluated on the algorithm of the MDS-PD criteria to classify the two distinct levels of diagnostic certainty for PD, namely, "*Clinically Established PD*", "*Clinically Probable PD*", or "*Not PD*". For example, the "Clinically Established PD" level of diagnostic certainty requires the following: presence of requisite essential criteria, absence of negative features (all 9 absolute exclusion criteria and 10 red flags), and presence of 2 positive features (supportive criteria) (figure 3, bottom panel).

Variable	UVMMC Site (n=23)	OHSU/VAPORHCS Site (n=20)	All Patients (n=43)
Age (mean, years)	70.57	75.90	73.05
Sex (female/male)	7 / 16	4 / 16	11 / 32
Ethnicity (Not Hispanic or Latino)	22	19	41
MDS-UPDRS (mean) - Part I - Part II - Part III - Part IV	 10.52 16.00 38.09 1.17	 14.75 20.25 43.65 5.70	 12.49 17.98 40.67 3.28
MOCA (mean)	25.91	23.90	24.98
UPSIT (mean)	19.13	16.20	17.77

Table 1: Baseline demographics and clinical characteristics of the Parkinson's disease patients at the two sites. (MOCA – Montreal Cognitive Assessment, UPSIT – University of Pennsylvania Smell Identification Test).

Evaluation:

The deployment of the CDS-PD platform in the clinical research study involved the creation of standalone instances of the platform that were deployed on a touchscreen laptop (Microsoft Surface) at the UVMMC site and OHSU/VAPORHCS site. The study personnel (movement disorders neurologists and research coordinators) at both sites were trained in the use of CDS-PD platform with a user guide document and as part of regular virtual team meetings for eliciting their feedback during its development.

The CDS-PD platform was evaluated using a subset of study cohort (PD patients with disease duration >=5 years; table 1) against the standard of clinical diagnosis made by fellowship trained movement disorders neurologists (with over 4 years of independent practice experience at a minimum) using paper-based format of the MDS-PD criteria. Based on this evaluation, the concordance rate of the CDS-PD platform was 91.30% (21 out of 23 matched cases) for the UVMMC site, and 95.00% (19 out of 20 cases) for the OHSU/VAPORHCS site, for an overall 93.02% (40 out of 43 cases) for the two sites combined (two-tailed p value 0.248, McNemar's test).

Of the 3 mismatched cases, the CDS-PD platform classified the patients as "Clinically Probable PD" due to the presence of one red flag (*#4: Inspiratory respiratory dysfunction: either diurnal or nocturnal inspiratory stridor or frequent inspiratory sigh*) while the movement disorders neurologist classified the patients as "Clinically Established PD". Subsequently, the movement disorders neurologists acknowledged to have not recalled the symptom of inspiratory stridor (captured during the symptoms review) while applying the MDS-PD criteria using the paper sheet. One potential reason for this recall bias might be that inspiratory stridor is a rare motor symptom encountered in PD and anticipated to be present infrequently even in APD.

DISCUSSION

Despite publication of the MDS-PD criteria a decade ago, their usage has been limited at point of care in clinical practice despite becoming standard of care in PD clinical trials even in their current paper-based format. The lack of CDS approach for the MDS-PD criteria in context of EHR systems is likely a major factor contributing to this gap. Additionally, standardized and electronic application of the MDS-PD criteria through the CDS-PD platform can provide consistent and efficient application of objective diagnostic criteria and thus contribute to increasing the accuracy of PD diagnosis in both clinical care and research. For example, absolute exclusion criteria #1 (*Unequivocal cerebellar abnormalities*), involves various clinical elements of cerebellar gait, limb ataxia, and specific oculomotor abnormalities—e.g., sustained gaze-evoked nystagmus, macro square wave jerks, and hypermetric saccades. Notably, hypermetric saccades, can be observed in advanced PD[22], and thus complicate the differentiation of cerebellar involvement from basal ganglia dysfunction in terms of consistent application of absolute exclusion criteria in a clinically relevant context. These nuances underscore the need for a CDS approach in applying such clinically complex diagnostic criteria.

In contrast to manual paper-based application of the MDS-PD criteria, the CDS-PD platform has several inherent advantages: systematic and less error-prone collection of the input data needed on all four categories of criteria included in MDS-PD criteria; computerized modelling of the diagnostic algorithm of the MDS-PD criteria; and computer-assisted processing of the collected input data on the diagnostic algorithm for classifying the diagnostic level of the patient's diagnosis. Also, the CDS-PD platform automatically applies the Movement Disorder Society Criteria for Clinically Established Early Parkinson's Disease[23] for patients with disease duration < 5 years (supplementary material 2). Furthermore, the CDS-PD platform also allows for automatic characterization of the PD into clinically relevant subtypes, for example, tremor-dominant (TD) and postural instability and gait dysfunction (PIGD), and indeterminate (IT) subtypes based on the patient data from the MDS-UPDRS[21] (supplementary material 3).

A key aspect of the CDS-PD platform is it's modular and scalable architecture that enables it to be expanded to support additional use cases, for example, currently supporting a multicenter project developing novel paradigm of studying cognitive dysfunction against conventional schemes of mild cognitive impairment and dementia in PD[24]. In the short term, we plan to develop the CDS-PD platform for computer-assisted differential diagnosis of PD from APD for general neurologists, using practical algorithms[7, 8, 25], and for movement disorders neurologists, using specific criteria[9-12]. In the long term, we envision that the CDS-PD platform will be interoperable with EHR using SMART (Substitutable Medical Applications, Reusable Technologies) on the FHIR (Fast Healthcare Interoperability Resources) technology framework[26], as well as advance precision medicine for individual PD and APD patients[27, 28] through AI-based data mining of relevant data repositories[29, 30] at point-of-care.

AUTHOR CONTRIBUTIONS

Deepak Gupta: Conceptualization, Validation, Visualization, Investigation, Data Curation, Software, Formal Analysis, Project Administration, Writing – Original Draft, Writing – Review and Editing

Pedram Golnari: Data Curation, Methodology, Formal Analysis, Software, Writing—Review and Editing

Katrina Prantzalos: Methodology, Software

Ian Zurlo: Conceptualization, Methodology

Vivikta Iyer: Investigation, Data Curation, Formal Analysis, Writing – Review and Editing

Brenna M Lobb: Investigation, Data Curation, Formal Analysis, Writing – Review and Editing

Cole Zweber: Formal Analysis, Visualization, Writing—Review and Editing

Manu Bulusu: Software

Dakota Clarke: Investigation, Data Curation

James T. Boyd: Writing—Review and Editing

Curtis Tatsuoka: Formal Analysis

Amie L. Hiller: Conceptualization, Methodology, Writing—Review and Editing

Satya S. Sahoo: Conceptualization, Validation, Visualization, Investigation, Data Curation, Formal Analysis, Project Administration, Writing – Original Draft, Writing – Review and Editing

FUNDING

This study was funded by the project titled "Ontology-based, Real-time, Machine learning Informatics System for Parkinson's Disease (ORMIS-PD)", funded by the Early Investigator Research Award (EIRA) of the Parkinson Research Program (PRP) of the Congressionally Directed Medical Research Program (CDMRP) of the United States (US) Department of Defense (DOD). Award # W81XWH2110859. This work also benefited from the projects funded with grants from the Dravet Syndrome Foundation (DSF), US National Institutes of Health (NIH): U24EB029005, R01DA053028

CONFLICTS OF INTEREST

None

DATA AVAILABILITY

The dataset used in this publication will be made available be made available by request from any qualified investigator.

REFERENCES

1. Willis, A.W., et al., *Incidence of Parkinson disease in North America.* npj Parkinson's Disease, 2022. **8**(1): p. 170.
2. Feigin, V.L., et al., *Global, regional, and national burden of neurological disorders during 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015.* The Lancet Neurology, 2017. **16**(11): p. 877-897.
3. Dorsey, E.R., et al., *The Emerging Evidence of the Parkinson Pandemic.* J Parkinsons Dis, 2018. **8**(s1): p. S3-S8.
4. Raty, V., et al., *Stability and Accuracy of a Diagnosis of Parkinson Disease Over 10 Years.* Neurology, 2025. **104**(9): p. e213499.
5. Lang, A.E., et al., *Trial designs used to study neuroprotective therapy in Parkinson's disease.* Mov Disord, 2013. **28**(1): p. 86-95.
6. Crotty, G.F., et al., *Planning for Prevention of Parkinson Disease: Now Is the Time.* Neurology, 2022. **99**(7 Suppl 1): p. 1-9.
7. McFarland, N.R., *Diagnostic Approach to Atypical Parkinsonian Syndromes.* Continuum (Minneap Minn), 2016. **22**(4 Movement Disorders): p. 1117-42.
8. Chou, K. *Diagnosis and differential diagnosis of Parkinson disease.* 2024. **UpToDate**.
9. McKeith, I.G., et al., *Diagnosis and management of dementia with Lewy bodies: Fourth consensus report of the DLB Consortium.* Neurology, 2017. **89**(1): p. 88-100.
10. Wenning, G.K., et al., *The movement disorder society criteria for the diagnosis of multiple system atrophy.* Movement Disorders, 2022. **37**(6): p. 1131-1148.
11. Höglinger, G.U., et al., *Clinical diagnosis of progressive supranuclear palsy: the movement disorder society criteria.* Movement disorders, 2017. **32**(6): p. 853-864.
12. Armstrong, M.J., et al., *Criteria for the diagnosis of corticobasal degeneration.* Neurology, 2013. **80**(5): p. 496-503.
13. Postuma, R.B., et al., *MDS clinical diagnostic criteria for Parkinson's disease.* Mov Disord, 2015. **30**(12): p. 1591-601.
14. Postuma, R.B., et al., *Validation of the MDS clinical diagnostic criteria for Parkinson's disease.* Mov Disord, 2018. **33**(10): p. 1601-1608.
15. Virameteekul, S., et al., *Clinical Diagnostic Accuracy of Parkinson's Disease: Where Do We Stand?* Mov Disord, 2023. **38**(4): p. 558-566.
16. *MDS Position Paper: Diagnosis of Parkinson's Disease.* 2023; Available from: https://www.movementdisorders.org/MDS/News/Newsroom/Position-Papers/MDS-Position-Diagnosis-of-PD.htm.
17. IT, A.S.f.T.P.O.o.t.N.C.f.H. *National Trends in Hospital and Physician Adoption of Electronic Health Records.* Available from: https://www.healthit.gov/data/quickstats/national-trends-hospital-and-physician-adoption-electronic-health-records.
18. *Django.* Available from: https://www.djangoproject.com/.
19. *Ontology-based, Real-time, Machine learning Informatics System for PD (ORMIS-PD).* 2021; Available from: https://dtic.dimensions.ai/details/grant/grant.9843089?search_mode=content&search_text=ormis-pd&search_type=kws&search_field=full_search.

20.   Elements, N.C.D. *CDE Detailed Report - Family History*. Available from: https://cde-fe.ninds.nih.gov/ninds/cde-details/CRF/pd/F2948/Participant%20History%20and%20Family%20History/General%20Health%20History/Family%20History.

21.   Stebbins, G.T., et al., *How to identify tremor dominant and postural instability/gait difficulty groups with the movement disorder society unified Parkinson's disease rating scale: comparison with the unified Parkinson's disease rating scale.* Mov Disord, 2013. **28**(5): p. 668-70.

22.   Lal, V. and D. Truong, *Eye movement abnormalities in movement disorders.* Clin Park Relat Disord, 2019. **1**: p. 54-63.

23.   Berg, D., et al., *Movement disorder society criteria for clinically established early Parkinson's disease.* Mov Disord, 2018. **33**(10): p. 1643-1646.

24.   *Cognitive Functions Impairment as a Novel Paradigm for Delineating Cognitive Dysfunction in Parkinson's Disease (PD-CFI)*. 2023; Available from: https://dtic.dimensions.ai/details/grant/grant.13858982?search_mode=content&search_text=Cognitive%20Functions%20Impairment%20as%20a%20Novel%20Paradigm%20for%20Delineating%20Cognitive%20Dysfunction%20in%20Parkinson%27s%20Disease%20(PD-CFI)%0A&search_type=kws&search_field=full_search.

25.   Bruno, M.K., et al., *A general neurologist's practical diagnostic algorithm for atypical Parkinsonian disorders: a consensus statement.* Neurology: Clinical Practice, 2024. **14**(6): p. e200345.

26.   Mandel, J.C., et al., *SMART on FHIR: a standards-based, interoperable apps platform for electronic health records.* Journal of the American Medical Informatics Association, 2016. **23**(5): p. 899-908.

27.   Fereshtehnejad, S.M., et al., *Clinical criteria for subtyping Parkinson's disease: biomarkers and longitudinal progression.* Brain, 2017. **140**(7): p. 1959-1976.

28.   Dadu, A., et al., *Identification and prediction of Parkinson's disease subtypes and progression using machine learning in two cohorts.* NPJ Parkinsons Dis, 2022. **8**(1): p. 172.

29.   Marek, K., et al., *The Parkinson's progression markers initiative (PPMI) - establishing a PD biomarker cohort.* Ann Clin Transl Neurol, 2018. **5**(12): p. 1460-1477.

30.   (FNIH), F.f.t.N. *Accelerating Medicines Partnership Program for Parkinson's Disease and Related Disorders (AMP PDRD)*. Available from: https://amp-pd.org/data.

Essential Criteria

1. Parkinsonism, defined as bradykinesia, in combination with at least 1 of rest tremor or rigidity, in limbs.

Red Flags Criteria

1. Rapid progression of gait impairment requiring regular use of wheelchair.
2. A complete absence of progression of motor symptoms or signs over 5 or more y unless stability is related to treatment.
3. Early bulbar dysfunction: severe dysphonia or dysarthria or severe dysphagia within first 5 y.
4. Inspiratory respiratory dysfunction: either diurnal or nocturnal inspiratory stridor or frequent inspiratory sigh.
5. Severe autonomic failure in the first 5 y of disease, including orthostatic hypotension and severe urinary retention or urinary incontinence.
6. Recurrent (>1/y) falls because of impaired balance within 3 y of onset.
7. Disproportionate antecollis (dystonic) or contractures of hand or feet within the first 10 y.
8. Absence of any of the common nonmotor features of disease despite 5 y disease duration.
9. Otherwise-unexplained pyramidal tract signs, defined as pyramidal weakness or clear pathologic hyperreflexia, on examination.
10. Bilateral symmetric parkinsonism, reported by history and confirmed on examination.

Supportive Criteria

1. Clear and dramatic beneficial response to dopaminergic therapy.
2. Presence of levodopa-induced dyskinesia.
3. Rest tremor of a limb, documented on clinical examination (in past, or on current examination).
4. Presence of either olfactory loss or cardiac sympathetic denervation on MIBG scintigraphy.

Absolute Exclusion Criteria

1. Unequivocal cerebellar abnormalities, such as cerebellar gait, limb ataxia, or cerebellar oculomotor abnormalities.
2. Downward vertical supranuclear gaze palsy, or selective slowing of downward vertical saccades.
3. Diagnosis of probable behavioral variant frontotemporal dementia or primary progressive aphasia within the first 5 y of disease.
4. Parkinsonian features restricted to the lower limbs for more than 3 y.
5. Drug-induced parkinsonism.
6. Absence of observable response to high-dose levodopa despite at least moderate severity of disease.
7. Unequivocal cortical sensory loss, clear limb ideomotor apraxia, or progressive aphasia.
8. Normal functional neuroimaging of the presynaptic dopaminergic system.
9. Expert physician concludes that an alternative syndrome is more likely than PD.

Essential Criteria Present → Yes → No PD

#Absolute Exclusion Criteria >=1 → Yes → No PD

#Red Flags Criteria >2 → Yes → No PD

#Supportive Criteria = 0 → Yes → No PD

#Red Flags Criteria = 0 & #Supportive Criteria = 2 → Yes → Clinically Established PD

#Red Flags Criteria <= #Supportive Criteria → Yes → Probable PD

No → No PD

Reset | Demographics | Motor Symptoms | Non-Motor Symptoms | Family History | S & E History | O & M History | Health Profile | Medications | Exam | Scales | DSI | Diagnostics | Submit

General Exam | Neurological Exam | Movement Disorder Exam | Additional Movement Disorder Exam | Advanced Movement Exam

Bradykinesia

	0	1	2	3	4
Face	0	1	2	3	4
Neck	0	1	2	3	4
Trunk	0	1	2	3	4
RUL	0	1	2	3	4
RLL	0	1	2	3	4
LUL	0	1	2	3	4
LLL	0	1	2	3	4

Rest Tremor

	0	1	2	3	4
Lip	0	1	2	3	4
Chin	0	1	2	3	4
H & N	0	1	2	3	4
RUL	0	1	2	3	4
RLL	0	1	2	3	4
LUL	0	1	2	3	4
LLL	0	1	2	3	4

Rigidity

	0	1	2	3	4
Neck	0	1	2	3	4
RUL	0	1	2	3	4
RLL	0	1	2	3	4
LUL	0	1	2	3	4
LLL	0	1	2	3	4

Postural Action Tremor

	0	1	2	3	4
Neck	0	1	2	3	4
RUL	0	1	2	3	4
RLL	0	1	2	3	4
LUL	0	1	2	3	4
LLL	0	1	2	3	4

Kinetic Action Tremor

	0	1	2	3	4
Neck	0	1	2	3	4
RUL	0	1	2	3	4
RLL	0	1	2	3	4
LUL	0	1	2	3	4
LLL	0	1	2	3	4

Dyskinesia

	0	1	2	3	4
H & N	0	1	2	3	4
RUL	0	1	2	3	4
RLL	0	1	2	3	4
Trunk	0	1	2	3	4
LUL	0	1	2	3	4
LLL	0	1	2	3	4

Movement Exam Notes

```
parkinsonism_rigidity = {
    rigidity_neck >= 1 or
    rigidity_rue >= 1 or rigidity_lue >= 1 or
    rigidity_rle >= 1 or rigidity_lle >= 1
}

parkinsonism_bradykinesia = {
    bradykinesia_ft_rh >= 1 or bradykinesia_ft_lh >= 1 or
    bradykinesia_hm_rh >= 1 or bradykinesia_hm_lh >= 1 or
    bradykinesia_psm_rh >= 1 or bradykinesia_psm_lh >= 1 or
    bradykinesia_tt_rf >= 1 or bradykinesia_tt_lf >= 1 or
    bradykinesia_la_rl >= 1 or bradykinesia_la_ll >= 1
}

parkinsonism_rtremor = {
    rtremor_rue >= 1 or rtremor_lue >= 1 or
    rtremor_rle >= 1 or rtremor_lle >= 1
}

has_parkinsonism = (parkinsonism_bradykinesia and
                    (parkinsonism_rtremor or parkinsonism_rigidity)
                   )

if (essentialCount === "1" && absoluteExclusionCount === "0" &&
    redFlagsCount === "0" && supportiveCriteriaCount >= 2) {
    document.getElementById("diagnosis-item").textContent = "Clinically Established PD";
```

Figure 1: Four categories of criteria and algorithm of the MDS-PD criteria.

Figure 2: A screenshot of the Patient Information Recording Module depicting 11 data sections/tabs and "Movement Disorders Exam" subsection of the "Exam" data section/tab.

Figure 3: Pseudocode showing implementation of the essential criteria (top panel) and algorithm for "Clinically Established PD" diagnostic level (bottom panel) of the MDS-PD criteria.

Note: Subject Test0507 shown in figure 2 and supplementary material is a dummy/test subject.