Mindfulness Meditation Training Reshapes Somatomotor Edge-Centric Connectivity Underlying Somatic Awareness: Insights for Healthy and Depressed Individuals

Yuwen He[a,b,c], Seng Hang Lei[d], Weng Si Che[e], Matthew D. Sacchet[f], Amy Jing-Wen Yin[a,g,h], Wenyue Gao[a], Anqi Gu[i,j], Natalia Chan[a], Xinyun Xiao[a], Jiaze Li[a], Jieting Chen[k], Mek Wong[d,h], Elvo Kuai Long Sou[d,h], Yizhong Shen[l], Jingju Quan[l], Weng Ieong Tou[m], Lin Ian Lio[m], Chih Chiang Chou[m], Peng Zhang[c], Bin Liu[n], Zhen Yuan[a,b]*

[a] Centre for Cognitive and Brain Sciences, University of Macau, Macau

[b] Faculty of Health Sciences, University of Macau, Macau

[c] State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of Sciences, Beijing, China

[d] Student Counselling Section, University of Macau, Macau

[e] Tranquil Mindfulness Centre, Macau

[f] Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

[g] Department of Psychiatry, Affiliated Hospital of Guangdong Medical University, Zhanjiang, China

[h] Department of Psychology, Faculty of Social Sciences, University of Macau, Macau

[i] Centre for Data Science, Institute of Collaborative Innovation, University of Macau, Macau

[j] Department of Computer and Information Science, Faculty of Science and Technology, University of Macau, Macau

[k] Department of Radiology, State Key Laboratory of Oncology in South China, Guangdong Provincial Clinical Research Center for Cancer, Sun Yat-Sen University, Guangzhou, Guangdong Province, China

[l] Zhuhai Third People's Hospital, Zhuhai, China

[m] Macau Psychiatric Association, Macau

[n] Emergency Department, Zhujiang Hospital, Southern Medical University, Guangzhou, China

*Corresponding author: Prof. Zhen Yuan; Email: zhenyuan@um.edu.mo

Abstract

Background: Somatic awareness is a fundamental practice for novice mindfulness practitioners. It has been proposed that mindfulness meditation training (MMT) enhances brain function through somatic awareness in both non-clinical and clinical populations. However, the neural signatures underlying this mechanism remain poorly understood. **Methods**: In our study, we conducted mindfulness interventions in both healthy (51 subjects) and depressed (35 subjects) populations and examined whether somatomotor network (SMN)-related edge-centric functional connectivity (eFC) is modulated by MMT, uncovering the population common and population-specific functioning pathways that may underlie somatic awareness. In addition, we tested whether such SMN-related eFCs can predict the insomnia status across populations. **Results**: Our findings revealed population-common eFC changes involving the SMN and attentional networks, indicating MMT may modulate the attentional system across populations via somatic awareness. Additionally, we identified population-specific eFC profiles engaging SMN-subcortical networks in the healthy population, suggesting MMT may target more pervasively the automatic processes in the healthy population. The specific eFC profile engaged SMN-default mode network interactions in the depressed population, which may be the functioning mechanism explaining MMT is efficient for depressed individuals. These findings demonstrate the differential neural signatures underlying different focuses of mindfulness via somatic awareness in different populations. Furthermore, support vector regression analysis demonstrated that these altered eFCs significantly predicted improvements in insomnia after MMT in both populations. **Conclusions**: In summary, our results highlight both shared and

distinct neural functioning signatures underlying mindfulness-related mental health improvements via somatic awareness in both the healthy and depressed populations.

Key words: Mindfulness meditation training; Edge-centric functional connectivity; Somatic awareness; Major depressive disorder; Population-common and Population-specific Neutral Signatures.

1   Introduction

The effectiveness of mindfulness in improving both mental and physical health across

clinical and non-clinical populations has been well documented [1, 2]. As this evidence

base has grown, researchers have increasingly turned their attention to an equally

important question: How do mindfulness-based interventions (MBIs) yield such diverse

and far-reaching benefits [3, 4]? Neuroimaging studies have proposed that several

large-scale functional networks including the default mode network (DMN), salience

network (SN), and frontoparietal network (FPN) are the key neural substrates

underlying mindfulness [5, 6]. In addition, emerging evidence highlights the

involvement of fronto-limbic circuits[7], along with the insula's role [8] and the

cingulate cortex [9]. While these findings underscore the importance of higher-order

associative and limbic regions, the potential contributions of somatosensory cortices to

mindfulness remain underexplored [10]. More importantly, this gap is particularly

striking given that somatic awareness practices, such as body scans and breath-focused

attention, are foundational to many mindfulness training programs, particularly for

beginners [11, 12]. These practices cultivate present-moment awareness of bodily

sensations and serve as entry points for developing the attentional focus, self-awareness,

emotion regulation, and other key capacities linked to the therapeutic effects of

mindfulness meditation [4]. Therefore, somatosensory processing might play a crucial

mediating role in the early stages and effectiveness of mindfulness meditation.

However, the specific neural mechanism by which somatosensory regions contribute to

mindfulness remains poorly understood. The present study aims to address this gap by

examining the role of somatomotor network (SMN), including somatosensory cortices,

in MBIs.

SMN consists of the postcentral regions (somatosensory regions), superior temporal

regions, premotor and paracentral areas, as well as parts of the insula and superior

parietal lobule [13]. These regions maintain extensive connections with both frontal and

subcortical brain structures [14] positioning them as potential key mediators in

mindfulness-related processes. For example, mindfulness practices have been shown to

enhance somatosensory attention [10, 15] and improve perceptual decision-making [16].

Additional magnetoencephalography studies indicate that mindfulness practice

enhances local alpha modulation [17], with somatosensory regions playing a central

role in these neural changes. Meanwhile, advanced meditation has been related to

altered brain network activity and connectivity of somatomotor networks including

during advanced concentrative absorption meditation which includes experiences of

complete cessation of bodily awareness [18, 19]. Further, [10] emphasized that

mindfulness practice often begins with focused attention on bodily sensations,

highlighting the importance of somatic awareness. To investigate how mindfulness

meditation training (MMT) modulates connectivity between somatosensory regions

and other brain networks, SMN-related edge-centric functional connectivity (eFC) were

constructed in this study to uncover the neural substrates involved in the functioning

mechanisms of mindfulness.

The eFC is a recently developed neuroimaging method that captures intricate neural

encoding and processing patterns by examining connectivity between pairs of edges

(i.e., connections between brain nodes) [20]. This approach reveals rich information

about neural communication and has been shown to reliably involve sensorimotor and

attentional networks [20]. Recent work also illustrated eFC's utility in studying

cognitive functions [21], neurological disorders [22], and even its heritability in relation

to clinical symptoms [23]. Consequently, eFC is an ideal tool for exploring how MMT

modulates SMN-related functional connectivity, offering deeper insights into the neural

signatures underlying how MMT functions via somatic awareness.

Meanwhile, MBIs have demonstrated benefits for both clinical and non-clinical

populations, though their underlying neural mechanisms might vary depending on the

functioning pathways and differential psychopathological sources. In the healthy

population, mindfulness training has been associated with reduced stress, improved

sleep quality, and enhanced emotion regulation [24-26]. Among clinical groups, such

as individuals with major depressive disorder (MDD), mindfulness practices have been

shown to alleviate symptoms, including depressive symptoms, duration of relapse, and

improve cognitive function [27, 28]. Among them, MBIs could improve sleep quality

and attentional control in both the healthy population and MDD. On the other hand,

MBIs in general population may bring more emphasis on stress reduction, while MBIs

in MDD may focus more on the improvement of depressive symptoms. Such

discrepancies of MBIs in different populations may signify their different functioning

neural pathways. Uncovering the population-common and population-specific

functioning pathways can provide insights into the functioning mechanism of MBIs and

offer guidance to develop personalized MBIs for different populations. What's more, it

can also facilitate using neuromodulation approaches to enhance mindfulness practice

effects, which is an emerging field in mindfulness research [29].

Insomnia is common among various psychiatric conditions, including MDD,

schizophrenia, and so on [30]. It was demonstrated MBIs can improve sleep quality in

both the healthy population and MDD. Meanwhile, somatic awareness is a foundation

of mindfulness practice, it would be reasonable to infer that somatic awareness may

contribute significantly to improving sleep quality. In our study, we would like to

examine whether the SMN-related eFCs can predict the sleep status in both the healthy

population and MDD. Such a test can help us to examine the importance of SMN-

related eFCs in the improvement of symptoms across different populations, which may

demonstrate the importance of somatic awareness in the functioning mechanism of

MBIs.

In this study, we hypothesized that MMT would modulate SMN-related eFC networks.

The resulting changes in SMN-related eFCs can serve as neural signatures underlying

how mindfulness achieves its therapeutic effects through somatic awareness in both

healthy and MDD individuals. Accordingly, two MMT programs were conducted in

healthy and MDD populations separately to examine our hypotheses. Specifically, it is

expected from the present study that: 1) Mindfulness training regulates SMN-related

eFCs, reflecting on the functioning mechanism of mindfulness meditation via somatic

awareness, 2) both population-common and population-specific eFC profiles are

modulated by mindfulness meditation, explaining its common and differential

functioning effects across populations, and 3) these SMN-related eFCs can predict

improvements in mental health outcomes, such as insomnia, in the healthy and MDD

populations. By investigating these dynamics, we aim to elucidate how MMT reshapes

brain connectivity to improve mental health, offering insights into its neural

mechanisms in both healthy and clinical populations.

Methods

Participants and Study Protocol

MMT was conducted for the healthy population and the MDD population separately.

In the healthy population, forty participants were randomly assigned to a mindfulness

(MD) group, while twenty-five participants were assigned to a non-treatment control

(NC) group (Figure 1. A). After taking first set of tests (Figure 1. B, T1), Healthy

participants in the MD group took a 4-week brief mindfulness workshop for college

students to enhance stress management [31], while participants in the NC group

received no intervention during this period. At the two-month follow-up, all healthy

participants were invited to take the second set of tests (Figure 1. B, T2).

In the MDD population, twenty-two participants with MDD were randomly assigned to

an MD group, taking an 8-week MBCT workshop in addition to their usual treatment,

while another twenty-two participants with MDD were assigned to a waitlist (WL)

group, receiving no additional intervention apart from their usual treatment during the

8 weeks. All participants with MDD took a set of tests before and after the 8-week

period (Figure 1. D).

Finally, 30 healthy participants were left in the MD group, and 21 healthy participants

were left in the NC group; 16 participants with MDD were left in the MD group, and

19 participants with MDD were left in the WL group, after removing participants who

dropped out early or had incomplete data.

Clinical Measurement

The Generalized Anxiety Disorder Scale (GAD-7) [32] was used to assess the anxiety

level in the healthy population. The Beck depression inventory (BDI) [33] was used to

assess the depressive symptoms in the patients with MDD. The insomnia severity index

was used to evaluate the insomnia severity in both populations [34].

Regions of Interest

The Brainnetome atlas was developed based on the structural and functional

connectivity profiles of each brain voxel [35]. We took 246 subregions in this atlas for

our initial regions of interest (ROIs). In later analysis, the ROIs were assigned to

specific Yeo's resting networks [13], including the subcortical network (Sub), visual

network (Vis), SMN, dorsal attention network (DAN), ventral attention network (VAN),

limbic network (Lim), frontoparietal network (FPN), and default mode network (DMN).

Calculation of the Edge-centric Functional Connectivity

Image acquisition and preprocessing procedures were detailed in the supplementary

methods. We calculated the edge-centric functional connectivity (eFC) according to the

procedures proposed by [20]. The time series in each ROI was extracted and z-scored,

and then the dot product of each pair of ROIs was used to index the edge time series of

the pair of ROIs. For each pair of edges, the eFC was calculated with Pearson's

correlation with the time series of these two edges.

Statistical Analysis

In our current study, we focused on the SMN-related eFCs. We extracted these eFCs

that at least involved a brain region in the SMN. To reveal the eFCs that were modulated

by MMT, we performed repeated-measure ANOVA analysis in each population sample

for each eFC. It was followed by counting the eFCs, showing a significant group × time

interaction effect ($p < 0.001$ or $p < 0.01$). We then grouped these significant eFCs into

categories according to Yeo's network. For example, if an edge included a region in the

SMN and another region in the DMN, then this edge would be classified as category

SMN-DMN; then if an eFC included two edges, one edge was classified in SMN-DMN,

another was in SMN-DAN, this eFC would be grouped in the eFC category SMN-DMN-SMN-DAN. In addition, we also counted the frequency of a brain region involved in the eFCs showing significant interaction effects in the two data samples.

Permutation test

We used a random permutation algorithm built in MATLAB to shuffle the significant statistical p-values of all eFCs in the abovementioned repeated-measurement ANOVA. In each permutation, the abovementioned frequency of counted significant eFCs belonging to each category or involved in each ROI was calculated and recorded. These permutation procedures were repeated 10,000 times to generate a null hypothesis probability distribution for each eFC category. We then regarded the eFC category with frequency value higher than 95% of simulated frequences as a reliably significant result, showing reliable modulation effects from MMT

Support vector regression analysis

In the healthy and MDD populations, we included the eFCs with significant interaction ($p < 0.001$ or $p < 0.01$) to a support vector regression model following procedures used by previous studies [36, 37] to predict clinical measurements significantly improved by mindfulness meditation training. We conducted this multivariate prediction analysis with the Spider toolbox with a linear kernel function (C=1), epsilon was set to 0.1, and 10-fold cross-validation was used to estimate the prediction power. Meanwhile, we also adopted a bootstrapping method to determine the contribution of each eFC to the predictive results. We categorized the eFCs as significant contributors when their corresponding Bootstrapping $p < 0.005$.

Results

Demographic characteristics and behavioral results

Healthy individuals in the MD and NC groups were comparable in age, gender, and education (Table 1). Individuals with MDD in the MD and WL groups were comparable in age, gender, education, baseline severity of depressive symptoms, first-episode age, duration of disease, and medication status (Table 1).

MMT improved anxiety and insomnia in the healthy individuals in the MD group (Figure S1), while these effects were not observed in the NC group. MMT improved insomnia but did not improve depressive symptoms in individuals with MDD in the MD group (Figure S2).

The eFCs modulated by mindfulness meditation training in the healthy population

As shown in Figure 2. A & D, widespread eFCs were modulated by MMT in the healthy population ($p < 0.001$ or $p < 0.01$). Most of these SMN-related eFCs, showing group × time interaction, engaged brain regions in the inferior frontal gyrus (IFG), cuneus, medial, superior frontal gyrus (SFG), and posterior temporal cortex (Figure 2. B & E, Table S4). Figure 2. C & F show the network-related eFCs modulated by MMT corresponding to two threshold sets ($p < 0.001$ or $p < 0.01$; Tables S5 & S6). After projecting these eFCs into Yeo's large-scale brain networks and performing a permutation test in both threshold sets, it was evident that MMT mainly modulated eFCs in SMN-subcortical and SMN-attentional networks (DAN and VAN) related categories across different thresholds (highlighted in orange/red squares in Figure 2. C & F), indicating that MMT may significantly influence the bottom-up automatic process and the attentional process via somatic awareness in the healthy population.

The eFCs modulated by mindfulness meditation training in the MDD Population

As shown in Figure 3. A & D, diverse eFCs were modulated by MMT in patients with

MDD ($p < 0.001$ or $p < 0.01$). Most of these SMN-related eFCs, showing group × time

interaction, engaged brain regions in the bilateral middle frontal gyrus (MFG), interior

parietal lobule, temporoparietal junction, IFG, cuneus, right posterior cingulate cortex

(PCC), and anterior cingulate cortex (ACC), right precuneus, and medial frontal cortex

(mPFC) (Figure 3. B & E; Table S4). Figure 3. C & F show the network-related eFCs

modulated by MMT corresponding to two threshold sets ($p < 0.001$ or $p < 0.01$; Tables

S9 & S10). After the permutation test at the network category level in both threshold

sets, it was observed that MMT mainly modulated eFCs in SMN-DMN, and SMN-

attentional networks related categories. Similar to healthy population, MMT modulated

widespread eFCs engaging SMN and attentional networks, which may be population

common neural signatures underlying MMT effects. Of note, widespread eFCs

engaging edges in SMN-DMN categories, including eFCs in the SMN-DMN-SMN-

Sub, SMN-DMN-SMN-Vis, SMN-DMN-SMN-VAN, SMN-DMN-SMN-FPN, and

SMN-DMN-SMN-DMN categories, were reliably modulated by MMT in both

threshold sets ($p < 0.001$ or $p < 0.01$), indicating that MMT may influence widespread

DMN-related functions via somatic awareness in the MDD population.

The prediction results of insomnia with eFCs in the healthy population

As shown in Figure 4. A, the support vector regression prediction model revealed that

eFCs (in the set of p < 0.001) can significantly predict the insomnia severity in the

healthy population (r=0.28, $p = 0.004$), indicating the eFCs modulated by MMT may

be involved in the improvement of insomnia in the healthy population. The other set of

eFCs ($p < 0.01$) cannot significantly predict any clinical measurements. The edges of

eFCs in the SMN-Vis, SMN-VAN categories showed evident contributions in

predicting insomnia (Figure 4. B, Bootstrapping p < 0.005; Table S8).

The prediction results of insomnia with eFCs in the MDD Population

The support vector regression prediction model revealed that eFCs (in the set of $p <$

0.01) can significantly predict the insomnia severity in patients with MDD (Figure 4.

C, r = 0.36, p = 0.003). The edges of eFCs in widespread categories, including SMN-

DAN, SMN-VAN, SMN-Vis, and SMN-DMN, contribute significantly to predicting

insomnia (Figure 4. D, Bootstrapping $p < 0.005$; Table S11), indicating the eFCs

modulated by MMT may be involved in the improvement of insomnia in patients with

MDD.

Discussion

Our study identified both shared and population-specific profiles of SMN-related eFCs

underlying MMT effects in the healthy and MDD populations. These findings provide

novel insights into the neural signatures underlying the functional mechanisms of

mindfulness—particularly those involving somatic awareness—across different

populations. Further analyses revealed that these eFCs significantly predicted insomnia

severity in the healthy and MDD populations. These results suggest that the eFCs

modulated by MMT may play a critical role in the clinical improvements observed in

both populations.

Population-common eFCs modulated by MMT—particularly those involving SMN-DAN and SMN-VAN—suggest a potentially shared neural mechanism across populations. These findings support the hypothesis that enhancement of attentional control via somatic awareness is a core function of diverse mindfulness-based interventions [3, 4, 10]. We found that MMT modulated eFCs engaging SMN and attentional networks in both the healthy and MDD population. A foundational review by Kerr et al. [10] first proposed that somatic awareness in mindfulness practices, such as the Koru Mindfulness curriculum and Mindfulness-Based Cognitive Therapy (MBCT), facilitates the filtering of sensory inputs to the primary sensory cortex and helps organize the flow of sensory information in the brain, thereby enhancing attentional regulation. Extending this hypothesis, our findings reveal specific eFC patterns modulated by MMT that may serve as neural signatures of its mechanisms in attentional functions through somatic awareness across different populations. The dorsal attention network (DAN) plays a central role in attentional control and executive functioning [38], while the ventral attention network (VAN) is involved in salience detection and attentional reorientation [39]. Our observation that mindfulness meditation robustly modulated eFCs in SMN-DAN and SMN-VAN across populations underscores complex interactions among the SMN, DAN, and VAN during mindfulness practice. These interactions may underlie the enhanced attentional control associated with somatic awareness, providing further neural evidence in support of Kerr et al.'s (2013) hypothesis.

The specific eFC profile, regulated pervasively by MMT in the healthy population,

involved SMN and subcortical network (Sub). The subcortical network, as defined in

our study, includes the amygdala, hippocampus, basal ganglia, and thalamus. These

brain regions are fundamental in many automatic processes, including emotion,

memory, reward, and perception [40-42]. Uncovering numerous SMN-Sub eFCs

modulated by MMT in the healthy population suggests that these connections may

underlie the influence of mindfulness on automatic behavioral patterns, basic

perception, and salience processing. Thus, these altered eFCs may underlie MMT

effects on the 'autopilot mode' in healthy individuals [12], encouraging individuals to

pay attention to and be aware of what they would perform. Although prior studies have

reported changes in functional connectivity involving subcortical regions—such as the

amygdala, putamen, and thalamus [43-45]—few have specifically examined the

connectivity between the SMN and subcortical networks in the context of mindfulness

meditation. Our study provided convincing evidence that MMT would regulate the

interactions engaging SMN and attention networks across different populations, which

may be the neural signatures underlying MMT effects on attentional control and

reducing autopilot behaviors.

Notably, MMT uniquely modulated widespread eFCs in the categories involving edges

between SMN and DMN. These changes may underlie improvements in self-referential

processes among patients with MDD and provide insight into the functioning

mechanism underlying the efficiency of MBCT for patients with MDD. Rumination is

characterized in patients with MDD and it is about self-referential processes [46]. Our

findings provide neural underpinnings to support the proposal of Kerr et al. (2013) that

somatic attention might compete with internally focused rumination and thus efficiently

help patients with MDD to improve symptoms. The widespread modulation of SMN-

DMN eFCs indicates that self-referential processes are prone to MMT in the MDD

population. While many studies have documented DMN alterations associated with

MMT [5, 48, 49], relatively few have examined DMN changes following MMT

specifically in patients with MDD. Our study provides strong evidence suggesting

DMN-related networks are widely modulated by MMT, which may be one of the most

important functioning mechanism underlying MMT effects on patients with MDD.

Our further analysis revealed that SMN-related eFCs modulated by MMT in both the

healthy and MDD populations significantly predicted changes in insomnia severity.

These findings suggest that the modulated SMN-related eFCs may serve as neural

signatures underlying the mental health improvements associated with mindfulness

meditation across populations. Specifically, in the healthy population, eFCs involving

the SMN-VAN, SMN-DMN, SMN-caudate, and SMN-fusiform contributed

significantly to predictive performance. This implies that interactions between the SMN

and these brain regions may be critical for supporting the mental health benefits of

mindfulness in non-clinical individuals. In the MDD sample, eFCs involving the SMN,

VAN, SMN-caudate, and SMN-visual cortex (SMN-Vis) were most predictive,

highlighting the importance of these interactions in mediating symptom reduction

through MBCT in clinical populations. Collectively, these findings indicate that the

eFCs identified in our study may represent essential neural mechanisms through which

mindfulness meditation, facilitated by somatic awareness, contributes to clinical

improvement in both healthy and MDD populations.

Our study has several limitations. First, the mindfulness workshops were tailored for

different populations and did not follow identical intervention durations. This

difference may have introduced confounding effects in identifying population-specific

eFC profiles. However, we posit that intervention duration likely affects the magnitude

of connectivity changes more than the pattern of eFC alterations. Second, the sample

size for the MDD group was relatively small. This limitation stemmed from a high

dropout rate. However, several intervention studies with similar sample size also

observed robust interventional effects [57-59], indicating our findings also provide

relatively reliable evidence. Future studies with larger clinical samples will be

necessary to validate our findings and assess their generalizability.

In conclusion, our study identified population-common changes in eFCs involving

SMN and attentional networks after MMT, highlighting the attention regulation in

MMT across different populations— Meanwhile, we uncovered population-specific

alterations: the healthy population showed pervasive modulations in SMN-Sub-related

eFCs, indicating the importance of MMT effects on automatic processes via somatic

awareness in the healthy population; whereas the MDD population exhibited

widespread changes in SMN-DMN-related eFCs, demonstrating DMN in patients with

MDD is prone to MMT. Importantly, these eFCs were associated with improvements

in insomnia in both populations. These results offer novel insights into the distinct and

shared neural mechanisms underlying mindfulness-based interventions in non-clinical

and clinical populations.

Acknowledgements

We thank Professor Yuan Zhou's lab for their valuable suggestions; thank Ms Joanna

Kong for her assistance in recruiting participants.

References

[1] Creswell JD, Way BM, Eisenberger NI, Lieberman MD. Neural correlates of dispositional mindfulness during affect labeling. Psychosom Med. 2007;69(6):560-5. https://doi.org/10.1097/PSY.0b013e3180f6171f.

[2] Piet J, Hougaard E. The effect of mindfulness-based cognitive therapy for prevention of relapse in recurrent major depressive disorder: a systematic review and meta-analysis. Clin Psychol Rev. 2011;31(6):1032-40. https://doi.org/10.1016/j.cpr.2011.05.002.

[3] Shapiro SL, Carlson LE, Astin JA, Freedman B. Mechanisms of mindfulness. Journal of clinical psychology. 2006;62(3):373-86.

[4] Tang YY, Holzel BK, Posner MI. The neuroscience of mindfulness meditation. Nat Rev Neurosci. 2015;16(4):213-25. https://doi.org/10.1038/nrn3916.

[5] Rahrig H, Vago DR, Passarelli MA, Auten A, Lynn NA, Brown KW. Meta-analytic evidence that mindfulness training alters resting state default mode network connectivity. Sci Rep. 2022;12(1):12260. https://doi.org/10.1038/s41598-022-15195-6.

[6] Sezer I, Pizzagalli DA, Sacchet MD. Resting-state fMRI functional connectivity and mindfulness in clinical and non-clinical contexts: A review and synthesis. Neurosci Biobehav Rev. 2022;135:104583. https://doi.org/10.1016/j.neubiorev.2022.104583.

[7] Melis M, Schroyen G, Pollefeyt J, Raes F, Smeets A, Sunaert S, et al. The Impact of Mindfulness-Based Interventions on Brain Functional Connectivity: a Systematic Review. Mindfulness. 2022;13(8):1857-75. https://doi.org/10.1007/s12671-022-01919-2.

[8] Young KS, van der Velden AM, Craske MG, Pallesen KJ, Fjorback L, Roepstorff A, et al. The impact of mindfulness-based interventions on brain activity: A systematic review of functional magnetic resonance imaging studies. Neurosci Biobehav Rev. 2018;84:424-33. https://doi.org/10.1016/j.neubiorev.2017.08.003.

[9] Zsadanyi SE, Kurth F, Luders E. The Effects of Mindfulness and Meditation on the Cingulate Cortex in the Healthy Human Brain: A Review. Mindfulness. 2021;12(10):2371-87. https://doi.org/10.1007/s12671-021-01712-7.

[10] Kerr CE, Sacchet MD, Lazar SW, Moore CI, Jones SR. Mindfulness starts with the body: somatosensory attention and top-down modulation of cortical alpha rhythms in mindfulness meditation. Frontiers in human neuroscience. 2013;7:12.

[11] Segal Z, Williams M, Teasdale J. Mindfulness-based cognitive therapy for depression. Guilford press; 2012.

[12] Williams M, Penman D. Mindfulness: A practical guide to finding peace in a frantic world. Hachette UK; 2011.

[13] Yeo BT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. J Neurophysiol. 2011;106(3):1125-65. https://doi.org/10.1152/jn.00338.2011.

[14] Kropf E, Syan SK, Minuzzi L, Frey BN. From anatomy to function: the role of the somatosensory cortex in emotional regulation. Brazilian Journal of Psychiatry. 2018;41(03):261-9.

[15] Bokk O, Forster B. The effect of a short mindfulness meditation on somatosensory attention. Mindfulness. 2022;13(8):2022-30.

[16] Mirams L, Poliakoff E, Brown RJ, Lloyd DM. Brief body-scan meditation practice
improves somatosensory perceptual decision making. Consciousness and
Cognition. 2013;22(1):348-59.
[17] Kerr CE, Jones SR, Wan Q, Pritchett DL, Wasserman RH, Wexler A, et al. Effects
of mindfulness meditation training on anticipatory alpha modulation in primary
somatosensory cortex. Brain research bulletin. 2011;85(3-4):96-103.
[18] Treves IN, Yang WFZ, Sparby T, Sacchet MD. Dynamic brain states underlying
advanced concentrative absorption meditation: A 7-T fMRI-intensive case study.
Netw Neurosci. 2025;9(1):125-45. https://doi.org/10.1162/netn_a_00432.
[19] Yang WFZ, Chowdhury A, Bianciardi M, van Lutterveld R, Sparby T, Sacchet MD.
Intensive whole-brain 7T MRI case study of volitional control of brain activity in
deep absorptive meditation states. Cereb Cortex. 2024;34(1).
https://doi.org/10.1093/cercor/bhad408.
[20] Faskowitz J, Esfahlani FZ, Jo Y, Sporns O, Betzel RF. Edge-centric functional
network representations of human cerebral cortex reveal overlapping system-
level architecture. Nat Neurosci. 2020;23(12):1644-54.
https://doi.org/10.1038/s41593-020-00719-y.
[21] Jiang L, Yang Q, He R, Wang G, Yi C, Si Y, et al. Edge-centric functional network
predicts risk propensity in economic decision-making: evidence from a resting-
state fMRI study. Cerebral Cortex. 2023;33(14):8904-12.
[22] Yang B, Wang M, Zhou W, Wang X, Chen S, Yuan L-X, et al. Edge-centric functional
network analyses reveal disrupted network configuration in autism spectrum
disorder. Journal of Affective Disorders. 2023;336:74-80.
[23] Chen Z, Tang Y, Liu X, Li W, Hu Y, Hu B, et al. Edge-centric connectome-genetic
markers of bridging factor to comorbidity between depression and anxiety.
Nature Communications. 2024;15(1). https://doi.org/10.1038/s41467-024-
55008-0.
[24] Brown KW, Ryan RM. The benefits of being present: Mindfulness and its role in
psychological well-being. Journal of Personality and Social Psychology.
2003;84(4):822-48. https://doi.org/10.1037/0022-3514.84.4.822.
[25] Winbush NY, Gross CR, Kreitzer MJ. The effects of mindfulness-based stress
reduction on sleep disturbance: a systematic review. Explore (NY). 2007;3(6):585-
91. https://doi.org/10.1016/j.explore.2007.08.003.
[26] Li S, Liu ZY, Che LP, Gong Y, Liu SJ, Guo XY. The neural mechanisms underlying
the modulation of attentional deployment on emotional stability. Exp Brain Res.
2022;240(6):1823-32. https://doi.org/10.1007/s00221-022-06361-3.
[27] Kuyken W, Byford S, Taylor RS, Watkins E, Holden E, White K, et al. Mindfulness-
based cognitive therapy to prevent relapse in recurrent depression. J Consult Clin
Psychol. 2008;76(6):966-78. https://doi.org/10.1037/a0013786.
[28] Cernasov P, Greene R, Kinard J, Halverson T, Lowery S, McLamb M, et al.
Attenuated Default Mode Network Functional Connectivity is Associated With
Improvement in Depressive Symptoms Following Mindfulness-Based Cognitive
Therapy in a Transdiagnostic Anhedonic Sample. Biol Psychiat.
2019;85(10):S158-S9. https://doi.org/DOI 10.1016/j.biopsych.2019.03.401.
[29] Abellaneda-Perez K, Potash RM, Pascual-Leone A, Sacchet MD.
Neuromodulation and meditation: A review and synthesis toward promoting well-
being and understanding consciousness and brain. Neurosci Biobehav Rev.
2024;166:105862. https://doi.org/10.1016/j.neubiorev.2024.105862.
[30] Harvey AG. Insomnia, psychiatric disorders, and the transdiagnostic perspective.
Current Directions in Psychological Science. 2008;17(5):299-303.

[31] Greeson JM, Juberg MK, Maytan M, James K, Rogers H. A randomized controlled
trial of Koru: a mindfulness program for college students and other emerging
adults. J Am Coll Health. 2014;62(4):222-33.
https://doi.org/10.1080/07448481.2014.887571.
[32] Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing
generalized anxiety disorder -: The GAD-7. Arch Intern Med. 2006;166(10):1092-7.
https://doi.org/DOI 10.1001/archinte.166.10.1092.
[33] Beck AT, Ward C, Mendelson M, Mock J, Erbauch J. Beck depression inventory
(BDI). 1961.
[34] Bastien CH, Vallieres A, Morin CM. Validation of the Insomnia Severity Index as an
outcome measure for insomnia research. Sleep Med. 2001;2(4):297-307.
https://doi.org/10.1016/s1389-9457(00)00065-4.
[35] Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, et al. The Human Brainnetome Atlas:
A New Brain Atlas Based on Connectional Architecture. Cereb Cortex.
2016;26(8):3508-26. https://doi.org/10.1093/cercor/bhw157.
[36] Liu X, Jiao G, Zhou F, Kendrick KM, Yao D, Gong Q, et al. A neural signature for the
subjective experience of threat anticipation under uncertainty. Nat Commun.
2024;15(1):1544. https://doi.org/10.1038/s41467-024-45433-6.
[37] Ceko M, Kragel PA, Woo CW, Lopez-Sola M, Wager TD. Common and stimulus-
type-specific brain representations of negative affect. Nat Neurosci.
2022;25(6):760-70. https://doi.org/10.1038/s41593-022-01082-w.
[38] Vossel S, Geng JJ, Fink GR. Dorsal and ventral attention systems: distinct neural
circuits but collaborative roles. Neuroscientist. 2014;20(2):150-9.
https://doi.org/10.1177/1073858413494269.
[39] Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, et al.
Dissociable intrinsic connectivity networks for salience processing and executive
control. J Neurosci. 2007;27(9):2349-56.
https://doi.org/10.1523/JNEUROSCI.5587-06.2007.
[40] Jones EG. The thalamus. Springer Science & Business Media; 2012.
[41] Squire LR, Cave CB. The hippocampus, memory, and space. Hippocampus.
1991;1(3):269-71. https://doi.org/10.1002/hipo.450010313.
[42] Cardinal RN, Parkinson JA, Hall J, Everitt BJ. Emotion and motivation: the role of
the amygdala, ventral striatum, and prefrontal cortex. Neurosci Biobehav Rev.
2002;26(3):321-52.
[43] Hölzel BK, Hoge EA, Greve DN, Gard T, Creswell JD, Brown KW, et al. Neural
mechanisms of symptom improvements in generalized anxiety disorder following
mindfulness training. NeuroImage: Clinical. 2013;2:448-58.
[44] Santarnecchi E, Egiziano E, D'Arista S, Gardi C, Romanella SM, Mencarelli L, et al.
Mindfulness-based stress reduction training modulates striatal and cerebellar
connectivity. J Neurosci Res. 2021;99(5):1236-52.
https://doi.org/10.1002/jnr.24798.
[45] Prakash RS, Shankar A, Tripathi V, Yang WFZ, Fisher M, Bauer CCC, et al.
Mindfulness Meditation and Network Neuroscience: Review, Synthesis, and
Future Directions. Biol Psychiatry Cogn Neurosci Neuroimaging. 2025;10(4):350-
8. https://doi.org/10.1016/j.bpsc.2024.11.005.
[46] Nolen-Hoeksema S, Wisco BE, Lyubomirsky S. Rethinking Rumination. Perspect
Psychol Sci. 2008;3(5):400-24. https://doi.org/10.1111/j.1745-
6924.2008.00088.x.
[47] Raichle ME. The brain's default mode network. Annu Rev Neurosci. 2015;38:433-
47. https://doi.org/10.1146/annurev-neuro-071013-014030.

[48] King AP, Block SR, Sripada RK, Rauch S, Giardino N, Favorite T, et al. Altered
default mode network (DMN) resting state functional connectivity following a
mindfulness‑based exposure therapy for posttraumatic stress disorder (PTSD) in
combat veterans of Afghanistan and Iraq. Depression and anxiety.
2016;33(4):289-99.
[49] Chou T, Dougherty DD, Nierenberg AA, Deckersbach T. Restoration of default
mode network and task positive network anti-correlation associated with
mindfulness-based cognitive therapy for bipolar disorder. Psychiatry Research:
Neuroimaging. 2022;319:111419.
[50] Chen F, Lv X, Fang J, Li T, Xu J, Wang X, et al. Body–mind relaxation meditation
modulates the thalamocortical functional connectivity in major depressive
disorder: a preliminary resting-state fMRI study. Translational psychiatry.
2021;11(1):546.
[51] Zhang J, Raya J, Morfini F, Urban Z, Pagliaccio D, Yendiki A, et al. Reducing default
mode network connectivity with mindfulness-based fMRI neurofeedback: a pilot
study among adolescents with affective disorder history. Molecular psychiatry.
2023;28(6):2540-8.
[52] Zhang X, Pines A, Stetz P, Goldstein-Piekarski AN, Xiao L, Lv N, et al. Adaptive
cognitive control circuit changes associated with problem-solving ability and
depression symptom outcomes over 24 months. Sci Transl Med.
2024;16(763):eadh3172. https://doi.org/10.1126/scitranslmed.adh3172.
[53] Park C, Rosenblat JD, Lee Y, Pan Z, Cao B, Iacobucci M, et al. The neural systems
of emotion regulation and abnormalities in major depressive disorder.
Behavioural Brain Research. 2019;367:181-8.
https://doi.org/10.1016/j.bbr.2019.04.002.
[54] Rive MM, van Rooijen G, Veltman DJ, Phillips ML, Schene AH, Ruhe HG. Neural
correlates of dysfunctional emotion regulation in major depressive disorder. A
systematic review of neuroimaging studies. Neurosci Biobehav Rev. 2013;37(10
Pt 2):2529-53. https://doi.org/10.1016/j.neubiorev.2013.07.018.
[55] Mochcovitch MD, da Rocha Freire RC, Garcia RF, Nardi AE. A systematic review
of fMRI studies in generalized anxiety disorder: evaluating its neural and cognitive
basis. Journal of affective disorders. 2014;167:336-42.
[56] Wang L, Hermens DF, Hickie IB, Lagopoulos J. A systematic review of resting-
state functional-MRI studies in major depression. Journal of affective disorders.
2012;142(1-3):6-12.
[57] Williams K, Elliott R, McKie S, Zahn R, Barnhofer T, Anderson IM. Changes in the
neural correlates of self-blame following mindfulness-based cognitive therapy in
remitted depressed participants. Psychiatry Res Neuroimaging.
2020;304:111152. https://doi.org/10.1016/j.pscychresns.2020.111152.
[58] Williams K. Neuropsychological mechanisms of mindfulness-based cognitive
therapy for depression. The University of Manchester (United Kingdom); 2017.
[59] Serra-Blasco M, Miquel-Giner N, Vicent-Gil M, Porta-Casteras D, Martinez-
Zalacain I, Cano M, et al. Clinical and Neuroimaging Effects of Mindfulness-Based
Cognitive Therapy for Symptomatic Obsessive-Compulsive Disorder Patients
after First-Line Treatments: A Randomised Controlled Trial. Psychother
Psychosom. 2025:1-17. https://doi.org/10.1159/000548961.

Figure legends

Figure 1. The flow chart of participant recruitment and data analysis. A: The flow

chart of participant recruitment in the healthy population; B: The experiment

procedures for participants in healthy population: all participants took MRI scans and

filled online questionnaires at baseline (T1, pre-intervention); the participants in the

MD group were assigned to take a 4-week mindfulness workshop, while the

participants in the NC group did not take any intervention during this period;

immediately after the 4-week, all participants were invited to take online

questionnaires; at the two months follow-up (T2, post-intervention), all participants

were invited to take MRI scans and fill the same online questionnaires; C: The flow

chart of the recruitment of patients with MDD; D: the experiment procedures for

participants in the MDD population: all participants were invited to take MRI scans

and online questionnaires at the baseline (pre-intervention, T1); participants in the

MD group took part in an 8-week MBCT workshop, while participants in the WL

group did not take any particular intervention during this period; 8 weeks later (post-

intervention, T2), all participants were invited to take MRI scans and online

questionnaires; E: data analysis flow: Brain regions defined by the Brainnetome atlas

were used as regions of interest (ROIs) to extract node-wise average time series.

Edge-wise time series were then generated by computing the element-wise (dot)

product of the time series from each pair of nodes. Pearson correlation coefficients

were calculated between all pairs of edge-wise time series to construct the edge

functional connectivity (eFC) matrix. Sensorimotor network (SMN)–related eFCs

were identified as those containing at least one node within the SMN. Repeated-

measures ANOVA was conducted to identify eFCs showing significant group (MD

vs. WL) × time (T1 vs. T2) interaction effects. These significant eFCs were

subsequently projected onto Yeo's large-scale functional networks, and permutation

tests were performed to determine whether the number of significant eFCs within

each network category exceeded chance levels. Finally, the identified significant eFCs

were used to predict insomnia severity using support vector regression.

Notes: MD: mindfulness; NC: non-treatment; WL: waitlist; ANOVA: analysis of

variance.

Figure 2. The eFC profile in the healthy population showing a significant group ×
time interaction. A\D: the frequency of edges involved in eFCs showing a significant
group × time interaction with $p < 0.001$ (A) or $p < 0.01$ (D); warmer color of edges
indicates higher frequency of engagement; B\E: the frequency of brain regions
involved in eFCs showing a significant group × time interaction with $p < 0.001$ (B) or
$p < 0.01$ (E); C\F: the frequency of eFCs, showing significant interaction with p <

0.001 (C) or p < 0.01 (F), summarized in network categories (of note, the x- and y-

axes are SMN-related edges, i.e., Sub indicates an edge between SMN and subcortical

network); the upper triangular parts display frequencies survived a permutation test

(the eFC categories highlighted in orange squares are healthy population-specific,

while those highlighted in red squares are similar across populations); the lower

triangular parts display frequencies without correction.

Notes: Sub: subcortical network; Vis: visual network; SMN: somatomotor network;

DAN: dorsal attention network; VAN: ventral attention network; Lim: limbic

network; FPN: frontoparietal network; DMN: default mode network.

Figure 3. The eFC profile in the MDD population showing a significant group × time interaction. A\D: the frequency of edges involved in eFCs showing a significant group × time interaction with $p < 0.001$ (A) or $p < 0.01$ (D); warmer color of edges indicates higher frequency of engagement; B\E: the frequency of brain regions involved in eFCs showing a significant group × time interaction with $p < 0.001$ (B) or $p < 0.01$ (E); C\F: the frequency of eFCs, showing significant interaction with $p < 0.001$ (C) or

p < 0.01 (F), summarized in network categories (of note, the x- and y-axes are SMN-

related edges, i.e., Sub indicates an edge between SMN and subcortical network); the

upper triangular parts display frequencies survived a permutation test (the eFC

categories highlighted in blue squares are MDD population-specific, while those

highlighted in red squares are similar across populations); the lower triangular parts

display frequencies without correction.

Notes: Sub: subcortical network; Vis: visual network; SMN: somatomotor network;

DAN: dorsal attention network; VAN: ventral attention network; Lim: limbic

network; FPN: frontoparietal network; DMN: default mode network.

Figure 4. The prediction results of insomnia with eFCs, showing significant group ×

time p < 0.001, in the healthy and MDD populations. A: the association between true

insomnia and predicted insomnia in the healthy population; B: the frequency of edges

involved in eFCs having significant contributions to predicting insomnia (p < 0.005)

in the healthy population; warmer colors indicate higher frequency of edges; C: the

association between true insomnia and predicted insomnia in the MDD population; D:

the frequency of edges involved in eFCs having significant contributions to predicting

insomnia (p < 0.005) in the MDD population; warmer colors indicate higher

frequency of edges.

Notes: MD: mindfulness; NC: non-treatment; WL: waitlist; T1: pre-intervention; T2:

post-intervention; Sub: subcortical network; Vis: visual network; SMN: somatomotor

network; DAN: dorsal attention network; VAN: ventral attention network; Lim:

limbic network; FPN: frontoparietal network; DMN: default mode network.

631 **Table**

633 **Table 1**. Demographic data in the healthy population

Variables	Mindfulness group	Non-treament \Waitlist group	Chi-square/ t-test
Healthy population (Number of participants)	**30**	**21**	
Age (SD)	23.83(3.56)	23.86(2.39)	p = 0.98
Gender(M/F)	5/25	4/17	p = 0.82
Education (U/M/P)	10/14/6	2/14/5	p = 0.14
Ethnicity (Han/Others)	28/2	19/2	--
MDD population (Number of participants)	**16**	**19**	
Age (SD)	30.44(6.17)	29.10(6.56)	*p = 0.54*
Gender(M/F)	4/12	3/16	*p = 0.49*
Education (U/M/P)	2/6/8	4/9/6	*P = 0.52*
Baseline Becker depression	19.47±12.96	21.21±9.94	*P = 0.66*
Ethnicity (Han/Others)	34/0	21/0	----
Age of first onset	26.06±6.96	23.94±6.10	*P = 0.37*
Duration of disease	3.75±3.70	4.06±2.59	*P = 0.78*
Medicine(Yes/No)	13/3	14/5	*P = 0.60*

635 Note: (M/F): Male/Female; (U/M/P): undergraduate/Master/Ph.D. SD: standard

636 deviation.