Title:

8266 SARS-CoV-2 Genomic Assemblies from Asymptomatic Carriers in Japan

Authors and Affiliations:

Hajime Ohyanagi[1†], Junko S Takeuchi[1†], Yuichi Kawanishi[1], Saya Ohi[1], Teiichiro

Shiino[1], Moto Kimura[1], Yusuke Takahashi[2], Shigeru Yoshida[2], Minoru Kato[3], Yukumasa

Kazuyama[3], Masato Ikeda[3], and Wataru Sugiura[1*]

[1]Center for Clinical Sciences, Japan Institute for Health Security, Tokyo, Japan

[2]Department of Clinical Laboratory Science, Health Sciences University of Hokkaido,

Hokkaido, Japan

[3]SB Coronavirus Inspection Center Corp., Tokyo, Japan

[†]These authors equally contributed

*Corresponding author: sugiura.w@jihs.go.jp

Abstract:

In the context of public health, asymptomatic carriers of respiratory infectious diseases

are considered a hidden yet critical factor in the transmission of infection and represent

a key target for efforts to mitigate disease spread. To contribute a unique genomic

resource to the SARS-CoV-2 research community, we collected SARS-CoV-2–positive

samples from asymptomatic individuals at the SB Coronavirus Inspection Center Corp.

during the COVID-19 pandemic in Japan and conducted a comprehensive analysis of

their viral genomes. Using Illumina COVIDSeq technology, we successfully generated

8,266 SARS-CoV-2 genome assemblies, all of which have been made publicly available

to facilitate further research. In this report, we summarize our efforts to collect SARS-

CoV-2–positive samples from asymptomatic individuals and highlight the key features

and accessibility of this genomic dataset.

Background & Summary:

We operate the SB Coronavirus Inspection Center Corp. (SBCVIC) [1] , which has a

key distinguishing feature compared to other medical facilities that conduct testing for

general health concerns: our center collects samples from asymptomatic

individuals. Consequently, the positive samples obtained at our facility are considered

to be from asymptomatic carriers of SARS-CoV-2. Epidemiologically, asymptomatic

carriers are known to have similar transmission capabilities to symptomatic

individuals[2-5]. Therefore, their existence represents a hidden yet critical factor in

controlling the spread of the pandemic[6,7]. From a genomic perspective, samples from

asymptomatic carriers are also a valuable resource for investigating genetic factors in

the viral genome that may be associated with the presence or absence of clinical

symptoms in COVID-19 patients[8].

During the pandemic period in Japan[9,10], spanning from July 2020 to January 2023,

the country experienced eight epidemic waves of SARS-CoV-2 infection. These

included waves dominated by the Alpha (the 4[th] wave), Delta (the 5[th] wave), and

Omicron sublineages BA.1, BA.2, and BA.5 (from the 6[th] wave onward) (**Figure 1A, B**).

Over the course of the study (July 27, 2020 – January 16, 2023), we tested a total of

4,573,575 samples, among which 18,475 tested positive for SARS-CoV-2 using reverse

transcription real-time quantitative polymerase chain reaction (RT-qPCR) (Ct values of

≤ 40) (**Figure 2** and **Table 1**). The overall average positive rate among asymptomatic

individuals was 0.40%. The positive rates varied by wave: 0.05% during the 3[rd] wave,

0.13% in the 4th wave (Alpha), 0.23% in the 5th wave (Delta), 4.5% in the 6th wave,

8.2% in the 7th wave (BA.1/BA.2), and 5.2% in the 8th wave (BA.5) (**Figure 2**). The

numbers of newly confirmed cases in Japan correlates well with the infection rate

among asymptomatic individuals (**Figure 1A**).

Starting October 9, 2020 (approximately six weeks after the initiation of the study),

we began collecting background information on positive cases from 45 prefectures

across Japan, including Shiga, Tokyo, Nagasaki, Osaka, Hokkaido, Fukuoka, Chiba,

Gifu, Kanagawa, Saitama, Hyogo, Aichi, Kyoto, Miyagi, Gumma, Hiroshima, Nagano,

Kumamoto, Ibaraki, Shizuoka, Tochigi, Fukushima, Yamanashi, Nara, Miyazaki,

Okinawa, Mie, Ehime, Iwate, Okayama, Kagawa, Yamagata, Saga, Aomori, Akita,

Niigata, Toyama, Tottori, Ishikawa, Yamaguchi, Oita, Kagoshima, Fukui, Shimane, and

Kochi (**Figure 3**). Among these cases, 51.9% were male, with a median age of 36 years

(interquartile range [IQR]: 27–44) (**Table 1**).

	Number (%)
Tested individuals	4,573,575
Positive cases	18,475
Gender	
Male	6,408 (51.9%)
Female	5,944 (48.1%)
Unknown	6,123
Age (Years)	36 [27 - 44]

Table 1. Statistics of the study participants. Data are presented as median
[interquartile range] for continuous measures and n (%) for categorical measures.

In total, we obtained 18,475 SARS-CoV-2-positive samples from asymptomatic

individuals. We performed whole-genome sequencing on residual saliva samples and

assembled viral genome sequences that met quality thresholds for downstream analysis.

Ultimately, we successfully generated 8,266 genome assemblies from these cases,

which are archived in the SBCVIC genomic dataset and categorized into two groups

(**Table 2** and **Table 3**).

For technical validation, we evaluated the distribution of consensus genome

sequence lengths and examined the lineage composition of the SARS-CoV-2 genomes.

We also assessed the presence of known mutations to confirm the reliability of our

sequencing pipeline for allele typing. In particular, we focused on two adjacent, co-

occurring mutations in the nucleocapsid (N) protein R203K/G204R[11], which are

associated with increased viral infectivity, fitness, and virulence. For further details, see

the **Technical Validation** section.

Methods:

SARS-CoV-2 Testing (Screening of Asymptomatic Cases)

Following previously described protocols[1,12], we conducted SARS-CoV-2 testing

and genome sequencing. The details are summarized below. The SBCVIC provided

routine workplace-based SARS-CoV-2 screenings upon company request, as well as

voluntary screenings requested by local governments. As a result, most positive cases

were identified in asymptomatic individuals [1,12]. All patients gave informed consent via

the opt-out method. Between July 27, 2020, and January 16, 2023, we tested

approximately 4.6 million asymptomatic individuals in Japan. However, individuals

who declined participation in the study were excluded from this count. The study

protocol was approved by the Ethics Committee of the National Center for Global

Health and Medicine (NCGM), Japan (approval number: NCGM-G-003678-00). Each

participant self-collected approximately 2 mL of saliva using the ZEESAN Saliva RNA

Sample Collection Kit (MD-ZSV-001; Zeesan Biotech, Fujian, China), which contained

1 mL of a guanidine-based viral inactivation buffer. The sample was mailed to the

inspection center as a UN 3373 Biological Substance Category B. On the day of arrival,

SARS-CoV-2 RT-qPCR testing was performed using the SARS-CoV-2 Direct Detection

RT-qPCR Kit (Takara Bio, Shiga, Japan). Kits RC30JW and RD003 were used until

April 4 and from April 5, 2021, respectively, according to the manufacturer's protocol.

Because the guanidine-based viral inactivation buffer can inhibit RT-PCR reactions[13],

we modified the pretreatment process. Specifically, 16 µL of the saliva/buffer mixture

was combined with 16 µL of water and 4 µL of pretreatment reagent (Solution A) to

dilute the inhibitor. The resulting 36 µL mixture was incubated at room temperature for

5 minutes, followed by 5 minutes at 95°C. A 2.5 µL aliquot of this treated mixture was

used as the RT-qPCR template. Test results (positive or negative) were determined

based on the protocol, with Ct values ≤ 40 considered positive.

Viral Genome Sequencing

Samples from SARS-CoV-2-positive individuals were subjected to genome

sequencing. The sequencing execution rate was nearly 100% for samples with Ct values

≤ 30 across all periods. However, the rate decreased for higher Ct values (30–40),

particularly from 2022 onward, due to operational constraints (see **Table 2** and **Table 3**).

For RNA extraction, 50 µL of viral nucleic acid was extracted from 200 µL of saliva

samples using MagMAX Viral/Pathogen Nucleic Acid Isolation Kit (Themo Fisher

Scientific, MA, USA) with MVP_Saliva_200_Flex_V1 protocol for KingFisher Flex

System (Themo Fisher Scientific). The eluted RNA (20 µL) was treated with RQ1

RNase-Free DNase (Promega, WI, USA). Subsequent cDNA synthesis, target

amplification using the ARTIC primer set [14], and library preparation were performed

according to the Illumina COVIDseq Test Reference Guide (Illumina Inc., CA, USA).

Sequencing was conducted on an Illumina NextSeq 2000 system. Sequencing data were

analyzed using the Illumina DRAGEN COVID Lineage Pipeline. Consensus sequences

in FASTA format were generated, collected and archived (see **Data Records**). The

ARTIC primer sets and DRAGEN versions used for each sample are documented in the

archive. Sequences ≥ 29,000 nucleotides in length (Category 1 in **Table 2** and **Table 3**)

were analyzed for lineage assignment using Phylogenetic Assignment of Named Global

Outbreak Lineages (PANGOLIN) version 4.3.1 (pangolin-data 1.21) [15].

Control Dataset

To analyze the distribution of SARS-CoV-2 lineages across Japan during the study

period (July 27, 2020 – January 16, 2023), we utilized sequences registered in the

GISAID EpiCov database (https://gisaid.org) as of February 6, 2023. The filtering

criteria were:

• Host: Human

• Location: Asia / Japan

• Collection date: July 27, 2020 – January 16, 2023

• Sequence length: ≥ 29,000 nt

• Passage history: Original

The resulting 539,504 sequences were analyzed using PANGOLIN version 4.3.1

(pangolin-data 1.21) for lineage classification (see **Figure 1B**).

Bioinformatics

The reference genome used was the Wuhan WIV04 strain (hCoV-

19/Wuhan/WIV04/2019, EPI_ISL_402124), downloaded from GISAID. A multiple

using MAFFT [16]. The WIV04 sequence was then removed from the alignment, which

was analyzed using AliView [17] and custom Python scripts to calculate allele frequencies

at the R203K/G204R sites in the N protein.

Data Records:

Of the 18,475 SARS-CoV-2-positive samples, 14,201 were subjected to viral

genome sequencing. By the end of December 2021, sequencing was conducted for

100% of samples with Ct ≤30 and 93.92% of those with Ct value between 30–40. From

2022 onward, the coverage remained high Ct ≤30 samples (for 99.83%) but decreased

to 62.71% for Ct 30–40 samples (**Table 2**).

Year/Group		Positive Samples		Sequenced Samples		Genomes Determined (Category 1 and 2)		Genomes Determined (Category 1)	
- 2021	Ct ≤30	513	**2,025**	513	**1,933**	346	**683**	263	**444**
	Ct 30-40	1,512		1,420		337		181	
2022 -	Ct ≤30	5,258	**16,450**	5,249	**12,268**	4,389	**7,583**	2,473	**2,473**
	Ct 30-40	11,192		7,019		3,194		0	
	total		**18,475**	total	**14,201**	total	**8,266**	total	**2,917**
- 2021	Ct ≤30	100 %	**100** %	100.00 %	**95.5** %	67.45 %	**33.7** %	51.27 %	**21.9** %
	Ct 30-40	100 %		93.92 %		22.29 %		11.97 %	
2022 -	Ct ≤30	100 %	**100** %	99.83 %	**74.6** %	83.47 %	**46.1** %	47.03 %	**15.0** %
	Ct 30-40	100 %		62.71 %		28.54 %		0 %	

Table 2. Summary of viral genome sequencing. The upper section shows actual
sample counts; the lower section shows corresponding percentages relative to the total
number of positive samples. For the definition of Categories 1 and 2, see **Table 3**.

Raw reads and consensus genome sequences for 8,266 SARS-CoV-2 samples were

submitted and are available for download from the International Nucleotide Sequence

Database Collaboration (INSDC) [18]. The corresponding BioSample and other accession

numbers are summarized in **Table 3**. According to operational criteria (based on

genome length and Ct values), sequences were categorized into two groups (see **Table**

3). Assembled genome sequences were also submitted to GISAID [19-21], under the

accession IDs listed.

Category and Criteria	Number of Sequences	BioSample* Accession	Sequence Accession	SRA Run Accession	SRA Experiment Accession	GISAID ID
Category 1: Genome length ≥ 29,000 nucleotides and Ct value ≤ 40 (for - 2021) ≤30 (for 2022 -)	2917	SAMD00738313 - SAMD00741057, SAMD00741082, SAMD00786981 - SAMD00787151	BS007837 - BS010581, BS010689, BS016048 - BS016218	DRR600457 - DRR603201, DRR603222, DRR608552 - DRR608722	DRX581004 - DRX583748, DRX583769, DRX589099 - DRX589269	EPI_ISL_19576276 - EPI_ISL_19582591, EPI_ISL_19582637, EPI_ISL_19575435 - EPI_ISL_19575605
Category 2: Complete Genome not meeting Category 1 criteria and with ≤ 20% unidentified nucleotides	5349	SAMD00741058 - SAMD00741081, SAMD00741083 - SAMD00746626**	BS010669 - BS010688, BS010690 - BS016018	DRR603202 - DRR603221, DRR603223 - DRR608551	DRX583749 - DRX583768, DRX583770 - DRX589098	EPI_ISL_19582592 - EPI_ISL_19582606, EPI_ISL_19582632 - EPI_ISL_19582636, EPI_ISL_19582638 - EPI_ISL_19585391, EPI_ISL_19579847 - EPI_ISL_19580766, EPI_ISL_19578927 - EPI_ISL_19579846, EPI_ISL_19574700 - EPI_ISL_19575434

Table 3. Submission of genomic sequences to public repositories (INSDC/DDBJ

and GISAID)

* BioSample accessions provide metadata describing the biological source materials

used to generate sequencing data.

** 219 BioSample data did not meet Category 2 criteria; sequence data for these were

not submitted.

Technical Validation:

Statistics of Consensus Genome Sequences

We first examined the length distribution of the 8,266 complete consensus genome

sequences across both Category 1 and Category 2 (**Figure 4**). A sufficient number (n =

2,917) of nearly full-length genome sequences ($\geq$ 29,000 nt; Category 1) were included

and exclusively used for SARS-CoV-2 lineage assignment via PANGOLIN, as well as

for subsequent analyses described below.

Epidemiological Distribution of SARS-CoV-2 Variants

Since the SBCVIC dataset represents a SARS-CoV-2 subpopulation within the

broader Japanese population, we anticipated a comparable lineage distribution between

our dataset and that of GISAID. Lineage analysis using PANGOLIN (**Figure 1B** and

1C) revealed a similar distribution of SARS-CoV-2 variants between SBCVIC and

GISAID datasets, supporting the validity of the overall SBCVIC sampling, sequencing,

and analysis workflow.

Detection of R203K/G204R Mutations in the Nucleocapsid Protein

The 2,917 Category 1 genome assemblies were also used for further technical

validation. As a control, we randomly selected 2,917 SARS-CoV-2 genome sequences

from symptomatic individuals of Japanese origin deposited in GISAID during the same

period (July 27, 2020 to January 16, 2023) (**Table 4**). To demonstrate the genotyping

capability of our dataset, we focused on two well-known adjacent co-occurring

mutations R203K/G204R in the N protein of SARS-CoV-2, which are known to affect

viral pathogenicity[11].

Genome sequences were aligned to the SARS-CoV-2 reference genome, and the

allele frequencies at the R203K/G204R sites were calculated for both the GISAID

(symptomatic) and SBCVIC (asymptomatic) groups (see **Method**). Our SBCVIC

dataset exhibited a lower rate of missing allele calls (5.9%) compared to the GISAID

dataset (17.8%), indicating high-quality genotyping across the majority of samples.

Interestingly, the frequency of the mutant allele in the SBCVIC (asymptomatic) group

was comparable to that in the GISAID (symptomatic) group (**Table 4**) ($P = 0.799676$,

G-test), despite our initial expectation that mutation rates would differ between

symptomatic and asymptomatic individuals. While we do not propose a specific

hypothesis here, this allele-level analysis highlights a potential application of the

SBCVIC dataset.

	Wild allele (R203/G204)	Mutant allele (203K/204R)	Allele Not Determined
GISAID (Symptomatic, n=2917)	7	2391	519 (17.8%)
SBCVIC (Asymptomatic, n=2917)	7	2739	171 (5.9%)

Table 4. Number of wild-type and mutant alleles at the nucleocapsid protein mutation

sites (R203K/G204R) in symptomatic and asymptomatic populations.

Usage Note:

The dataset of SARS-CoV-2 samples collected and analyzed from asymptomatic

individuals in Japan is highly distinctive. We demonstrated the validity of this SBCVIC

dataset for analyzing the epidemiological distribution of SARS-CoV-2 variants and for

conducting allele-level genotyping. We believe that this dataset has significant potential

to provide insights into the modulation of SARS-CoV-2 virulence and to enhance our

understanding of viral fitness.

Code Availability

All custom scripts used in this study are available on GitHub

at https://github.com/oyasai55/calc_allele_frequencies/.

Acknowledgements

We thank Mr. Takahiro Sakurai and Mr. Hiroshi Ikawa for their support with

bioinformatics analyses. We also acknowledge Mr. Tsuyoshi Saito, Ms. Natsumi

Miyazaki, Mr. Atsushi Sugi, and Mr. Masanori Tamanaha for their assistance in

managing sample metadata.

Ethics Approval:

All participants provided written informed consent for the collection of saliva

samples, viral genome sequencing, and the use of associated demographic information.

Participation in the study was voluntary, and individuals had the option to opt out via

the study website. The study protocol was approved by the Institutional Review Board

of the National Center for Global Health and Medicine (approval number: NCGM-G-

003678-00). All procedures were conducted in accordance with the principles of the

Declaration of Helsinki, as revised in 2013.

Author contributions

H.O. and J.S.T. designed and performed the analyses and wrote the manuscript.

J.S.T. and Yuichi K. managed the genome sequence data from SBCVIC. S.O. conducted

bioinformatics and statistical analyses. J.S.T. and S.O. conducted figure/table

visualization. T.S. and W.S. supervised the entire analysis process. Moto K. directed the

research contract with the SBCVIC and intellectual properly. S.Y., Minoru K. and

Yukumasa K. was responsible for RT-qPCR testing with the viral genome sequencing at

SBCVIC. M.I. and W.S. conceived an inspection system by SBCVIC and acquired

research funding. All authors critically reviewed and approved the final version of the

manuscript.

Competing interests:

This research was supported by the SB Coronavirus Inspection Center Corp. Moto

Kimura and Wataru Sugiura have received research grants from SB Coronavirus

Inspection Center Corp. Masato Ikeda, Yukumasa Kazuyama, and Minoru Kato are

employees of the SB Coronavirus Inspection Center Corp.

Figures:

In a separate file (pptx).

Figure legends:

Figure 1. Monthly epidemiological distribution of SARS-CoV-2 variants in Japan

(July 27, 2020 - January 16, 2023). (A) Monthly number of newly confirmed COVID-

19 cases in Japan, based on open data from the Ministry of Health, Labour and Welfare

(https://covid19.mhlw.go.jp/en/, accessed May 1, 2023)[22]. (B) All domestic sequences

registered in the Global Initiative on Sharing All Influenza Data (GISAID) EpiCoV

database as of February 6, 2023 (n = 539,504). (C) Positive cases detected in

asymptomatic individuals in this study (n = 2,917). SARS-CoV-2 lineage analysis was

conducted using the Phylogenetic Assignment of Named Global Outbreak Lineages

(PANGOLIN) version 4.3.1. Data visualization was performed using R version 4.3.1.

Figure 2. Number of tested (gray) and SARS-CoV-2-positive (blue) cases in this study. Between July 27, 2020, and January 16, 2023, a total of 4,573,575 saliva samples were tested, of which 18,475 were positive by RT-qPCR testing. Individuals who opted out of the study were excluded from this count.

Figure 3. Geographic distribution of sample collection. This figure was generated using the R package *NipponMap* (https://cran.r-project.org/web/packages/NipponMap)[23,24]. Color intensity indicates the number of samples collected, with darker shades representing higher counts. The top five prefectures with the highest number of samples were Shiga (n = 6,082), Tokyo (n = 3,118), Nagasaki (n = 521), Osaka (n = 429), and Hokkaido (n = 401).

Figure 4. Distribution of SARS-CoV-2 genome consensus sequence lengths obtained in this study (n = 8,266). A histogram shows the distribution of sequence lengths (X-axis) and their frequencies (Y-axis). Category 1 sequences (n = 2,917) are shown in dark purple, and Category 2 sequences (n = 5,349) are shown in green. The dashed line represents the threshold for near-complete genome sequences (29,000 nt).

References:

1   Shiino, T. *et al.* Molecular epidemiology of SARS-CoV-2 genome sentinel surveillance in commercial COVID-19 testing sites targeting asymptomatic individuals during Japan's seventh epidemic wave. *Sci Rep* **14**, 20950 (2024). https://doi.org/10.1038/s41598-024-71953-8

2   Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic

293   review and meta-analysis. *PLoS Med* **17**, e1003346 (2020).
294   https://doi.org/10.1371/journal.pmed.1003346

295 3 Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of
296   asymptomatic and presymptomatic SARS-CoV-2 infections: Update of a living
297   systematic review and meta-analysis. *PLoS Med* **19**, e1003987 (2022).
298   https://doi.org/10.1371/journal.pmed.1003987

299 4 Casey-Bryars, M. *et al.* Presymptomatic transmission of SARS-CoV-2 infection: a
300   secondary analysis using published data. *BMJ Open* **11**, e041240 (2021).
301   https://doi.org/10.1136/bmjopen-2020-041240

302 5 Yang, Q. *et al.* Just 2% of SARS-CoV-2-positive individuals carry 90% of the virus
303   circulating in communities. *Proc Natl Acad Sci U S A* **118** (2021).
304   https://doi.org/10.1073/pnas.2104547118

305 6 Davis, J. T. *et al.* Cryptic transmission of SARS-CoV-2 and the first COVID-19
306   wave. *Nature* **600**, 127-132 (2021). https://doi.org/10.1038/s41586-021-04130-
307   w

308 7 Nabeshima, T. *et al.* COVID-19 cryptic transmission and genetic information
309   blackouts: Need for effective surveillance policy to better understand disease
310   burden. *Lancet Reg Health West Pac* **7**, 100104 (2021).
311   https://doi.org/10.1016/j.lanwpc.2021.100104

312 8 Ferdinand, A. S. *et al.* An implementation science approach to evaluating
313   pathogen whole genome sequencing in public health. *Genome Med* **13**, 121
314   (2021). https://doi.org/10.1186/s13073-021-00934-7

315 9 Kitahara, K., Nishikawa, Y., Yokoyama, H., Kikuchi, Y. & Sakoi, M. An overview of
316   the reclassification of COVID-19 of the Infectious Diseases Control Law in Japan.
317   *Glob Health Med* **5**, 70-74 (2023). https://doi.org/10.35772/ghm.2023.01023

318 10 Takeuchi, J. S. *et al.* Large-scale screening of SARS-CoV-2 variants in Tokyo,
319   Japan: A 3-year and 9-month longitudinal survey. *Glob Health Med* **7**, 151-160
320   (2025). https://doi.org/10.35772/ghm.2025.01004

321 11 Wu, H. *et al.* Nucleocapsid mutations R203K/G204R increase the infectivity,
322   fitness, and virulence of SARS-CoV-2. *Cell Host Microbe* **29**, 1788-1801 e1786
323   (2021). https://doi.org/10.1016/j.chom.2021.11.005

324 12 Terada-Hirashima, J. *et al.* Investigation of the use of PCR testing prior to ship
325   boarding to prevent the spread of SARS-CoV-2 from urban areas to less-
326   populated remote islands. *Glob Health Med* **4**, 174-179 (2022).
327   https://doi.org/10.35772/ghm.2022.01008

13   Katsuno, T. *et al. Diagnostic accuracy of direct reverse transcription-polymerase*
chain reaction using guanidine-based and guanidine-free inactivators for SARS-
CoV-2 detection in saliva samples (Avairable at SSRN (preprint), 2023).
14   *Artic Network SARS-CoV-2*, <https://artic.network/ncov-2019> (
15   O'Toole, A. *et al.* Assignment of epidemiological lineages in an emerging
pandemic using the pangolin tool. *Virus Evol* **7**, veab064 (2021).
https://doi.org/10.1093/ve/veab064
16   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-
780 (2013). https://doi.org/10.1093/molbev/mst010
17   Larsson, A. AliView: a fast and lightweight alignment viewer and editor for large
datasets. *Bioinformatics* **30**, 3276-3278 (2014).
https://doi.org/10.1093/bioinformatics/btu531
18   Karsch-Mizrachi, I. *et al.* The international nucleotide sequence database
collaboration (INSDC): enhancing global participation. *Nucleic Acids Res* (2024).
https://doi.org/10.1093/nar/gkae1058
19   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's
innovative contribution to global health. *Glob Chall* **1**, 33-46 (2017).
https://doi.org/10.1002/gch2.1018
20   Khare, S. *et al.* GISAID's Role in Pandemic Response. *China CDC Wkly* **3**, 1049-
1051 (2021). https://doi.org/10.46234/ccdcw2021.255
21   Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data -
from vision to reality. *Euro Surveill* **22** (2017). https://doi.org/10.2807/1560-
7917.ES.2017.22.13.30494
22   The Ministry of Health, L. a. W., Japan. *Visualizing the data: information on*
COVID-19 infections., <https://covid19.mhlw.go.jp/en/> (2023).
23   NipponMap: Japanese Map Data and Functions. v. 0.2 (2017).
24   R: A language and environment for statistical computing. (2021).

Figure 1

Figure 2

Number of Tested and Positive cases of SARS−CoV−2 in SVCVIC

Figure 3

Figure 4

Histogram of sequence length (n=8,266)