Title:

8,266 SARS-CoV-2 Genomic Assemblies from Asymptomatic Carriers in Japan

Authors and Affiliations:

Hajime Ohyanagi[1†], Junko S Takeuchi[1†], Yuichi Kawanishi[1], Saya Ohi[1], Teiichiro

Shiino[1], Moto Kimura[1], Yusuke Takahashi[2], Shigeru Yoshida[2], Minoru Kato[3], Yukumasa

Kazuyama[3], Masato Ikeda[3], and Wataru Sugiura[1*]

[1]Center for Clinical Sciences, Japan Institute for Health Security, Tokyo, Japan

[2]Department of Clinical Laboratory Science, Health Sciences University of Hokkaido,

Hokkaido, Japan

[3]SB Coronavirus Inspection Center Corp., Tokyo, Japan

[†]These authors equally contributed

*Corresponding author: sugiura.w@jihs.go.jp

Abstract:

In the context of public health, asymptomatic carriers of respiratory infectious diseases

are considered a hidden yet critical factor in the transmission of infection and represent

a key target for efforts to mitigate disease spread. To contribute a unique genomic

resource to the SARS-CoV-2 research community, we collected SARS-CoV-2–positive

samples from asymptomatic individuals at the SB Coronavirus Inspection Center Corp.

during the COVID-19 pandemic in Japan and conducted a comprehensive analysis of

their viral genomes. Using Illumina COVIDSeq technology, we successfully generated

8,266 SARS-CoV-2 genome assemblies, all of which have been made publicly available

to facilitate further research. In this report, we summarize our efforts to collect SARS-

CoV-2–positive samples from asymptomatic individuals and highlight the key features

and accessibility of this genomic dataset.

Background & Summary:

We operate the SB Coronavirus Inspection Center Corp. (SBCVIC) [1] , a large-scale

screening facility specializing in SARS-CoV-2 testing for individuals without symptoms.

The center primarily serves workplace-based routine screening programs and voluntary

screening requests from local governments, and therefore most samples originated from

individuals who did not report symptoms at the time of testing. Epidemiologically,

asymptomatic carriers of SARS-CoV-2 are known to have non-negligible transmission

capabilities comparable to those of symptomatic individuals[2-5]. Therefore, their

existence represents a hidden yet critical factor in controlling the spread of the

pandemic[6,7]. From a genomic perspective, samples from asymptomatic carriers are also

a valuable resource for investigating genetic factors in the viral genome that may be

associated with the presence or absence of clinical symptoms in COVID-19 patients[8].

During the pandemic period in Japan[9,10], spanning from July 2020 to January 2023,

the country experienced eight epidemic waves of SARS-CoV-2 infection[11] (**Figure 1A**).

These included waves dominated by the Alpha (the 4th wave), Delta (the 5th wave), and

Omicron sublineages BA.1, BA.2, and BA.5 (from the 6th wave onward) (**Figure 1B** for

symptomatic cases, and **Figure 1C** for asymptomatic cases). Over the course of the

study (July 27, 2020 – January 16, 2023), we tested a total of 4,573,575 samples, among

which 18,475 tested positive for SARS-CoV-2 using reverse transcription real-time

quantitative polymerase chain reaction (RT-qPCR) (Ct values of ≤ 40) (**Figure 2** and

Table 1). The overall average positive rate among asymptomatic individuals was 0.40%.

The positive rates varied by wave: 0.05% during the 3^{rd} wave, 0.13% in the 4^{th} wave

(Alpha), 0.23% in the 5^{th} wave (Delta), 4.5% in the 6^{th} wave, 8.2% in the 7^{th} wave

(BA.1/BA.2), and 5.2% in the 8^{th} wave (BA.5). The number of newly confirmed cases

in Japan (**Figure 1A**) shows a trend similar to the positive rate among asymptomatic

individuals mentioned above.

Starting October 9, 2020 (approximately six weeks after the initiation of the study),

we began collecting background information on positive cases from 45 prefectures

across Japan, including Shiga, Tokyo, Nagasaki, Osaka, Hokkaido, Fukuoka, Chiba,

Gifu, Kanagawa, Saitama, Hyogo, Aichi, Kyoto, Miyagi, Gumma, Hiroshima, Nagano,

Kumamoto, Ibaraki, Shizuoka, Tochigi, Fukushima, Yamanashi, Nara, Miyazaki,

Okinawa, Mie, Ehime, Iwate, Okayama, Kagawa, Yamagata, Saga, Aomori, Akita,

Niigata, Toyama, Tottori, Ishikawa, Yamaguchi, Oita, Kagoshima, Fukui, Shimane, and

Kochi (**Figure 3**). No positive case was collected from two prefectures (Tokushima and

Wakayama). Among 18,475 positive cases, 6,408 (34.7%) were male, 5,944 (32.2%)

were female, and 6,123 (33.1%) had missing gender information. Age information was

available for 12,544 individuals, among whom the median age was 36 years

(interquartile range [IQR]: 27–44). (**Table 1**).

		Value
Tested individuals		4,573,575
Positive cases		18,475
Gender		
Male	6,408 (34.7%)	
Female	5,944 (32.2%)	
Unknown	6,123 (33.1%)	
Age (Years)		36 [27 - 44]

Table 1. Statistics of the study participants. Gender is shown in n (%), and age is

presented as median [interquartile range] .

In total, we obtained 18,475 SARS-CoV-2-positive samples from asymptomatic

individuals. We performed whole-genome sequencing on residual saliva samples and

assembled viral genome sequences that met quality thresholds for downstream analysis.

Ultimately, we successfully generated 8,266 genome assemblies from these cases,

which are archived in the SBCVIC genomic dataset and categorized into two groups

(**Table 2** and **Table 3**) according to operational criteria (based on genome length and Ct

values).

For technical validation, we evaluated the distribution of consensus genome

sequence lengths and examined the lineage composition of the SARS-CoV-2 genomes.

We also assessed the presence of known mutations to confirm the reliability of our

sequencing pipeline for allele typing. In particular, we focused on two adjacent, co-

occurring mutations in the nucleocapsid (N) protein R203K/G204R[12], which are

associated with increased viral infectivity, fitness, and virulence. For further details, see

the **Technical Validation** section.

Methods:

Participants

Procedures complied with national, institutional committees on human

experimentation and with the Helsinki Declaration (1975, revised in 2008). No

participants were directly recruited for this study. All analyses were performed using

saliva specimens collected by SBCVIC testing laboratories as part of their routine

SARS-CoV-2 testing for asymptomatic individuals. In Japan, explicit opt-in informed

consent is not required for this type of specimen. The institutional observational studies

employed an IRB-approved opt-out process in accordance with national ethical

guidelines. Study information and opt-out procedures were publicly posted on the

institutional websites, and individuals were given the opportunity to decline the

secondary use of their specimens and associated clinical information. Saliva specimens

collected by the testing laboratories were handled under their approved protocol, only

anonymized samples and limited clinical data were provided to our research team. The

study protocol was approved by the Institutional Review Board of the National Center

for Global Health and Medicine (approval number: NCGM-G-004225-00, August 5,

2020).

SARS-CoV-2 Testing (Screening of Asymptomatic Cases)

Following previously described protocols[1,13], we conducted SARS-CoV-2 testing

and genome sequencing. The SBCVIC provided routine workplace-based SARS-CoV-2

screenings upon company request, as well as voluntary screenings requested by local

governments. Each participant self-collected approximately 2 mL of saliva using the

ZEESAN Saliva RNA Sample Collection Kit (MD-ZSV-001; Zeesan Biotech, Fujian,

China), which contained 1 mL of a guanidine-based viral inactivation buffer. The

sample was classified and shipped as UN 3373 Biological Substance, Category B. On

the day of arrival, SARS-CoV-2 RT-qPCR testing was performed using the SARS-CoV-

2 Direct Detection RT-qPCR Kit (Takara Bio, Shiga, Japan). Kits RC30JW and RD003

were used until April 4 and from April 5, 2021, respectively, according to the

manufacturer's protocol. Because the guanidine-based viral inactivation buffer can

inhibit RT-PCR reactions[14], we modified the pretreatment process. Specifically, 16 µL

of the saliva/buffer mixture was combined with 16 µL of water and 4 µL of pretreatment

reagent (Solution A, included in the kit) to dilute the inhibitor. The resulting 36 µL

mixture was incubated at room temperature for 5 minutes, followed by 5 minutes at

95°C. A 2.5 µL aliquot of this treated mixture was used as the RT-qPCR template. Test

results (positive or negative) were determined based on the protocol, with Ct values ≤

40 considered positive.

Viral Genome Sequencing

Samples from SARS-CoV-2-positive individuals were subjected to genome

sequencing. For RNA extraction, 50 µL of viral nucleic acid was extracted from 200 µL

of saliva samples using MagMAX Viral/Pathogen Nucleic Acid Isolation Kit (Thermo

Fisher Scientific, MA, USA) with MVP_Saliva_200_Flex_V1 protocol for KingFisher

Flex System (Thermo Fisher Scientific, MA, USA). The eluted RNA (20 µL) was

treated with RQ1 RNase-Free DNase (Promega, WI, USA). Subsequent cDNA

synthesis, target amplification using the ARTIC[15] primer set (versions used for each

sample are documented in the BioSample archive), and library preparation were

performed according to the Illumina COVIDseq Test Reference Guide (Illumina Inc.,

CA, USA). The sequencing performance rates were 100.00% (-2021) and 99.83%

(2022-) for samples with Ct values ≤ 30 across all periods (see the column **Genome**

Sequencing Performed in **Table 2**). However, the rate decreased for higher Ct values

(30–40), particularly from 2022 onward, due to operational constraints. Sequencing was

conducted on an Illumina NextSeq 2000 system.

Bioinformatics

Sequencing data were analyzed using the Illumina DRAGEN COVID Lineage

Pipeline (https://jp.illumina.com/products/by-type/informatics-products/basespace-

sequence-hub/apps/dragen-covid-lineage.html). Consensus sequences in FASTA format

were generated (see **Data Records** and **Data Overview**). The DRAGEN versions used

for each sample are documented in the BioSample archive. Sequences equal to or longer

than 29,000 nucleotides in length (Category 1 in **Table 2** and **Table 3**) were analyzed

for lineage assignment using Phylogenetic Assignment of Named Global Outbreak

Lineages (PANGOLIN) version 4.3.1 (pangolin-data 1.25.1) [16].

For variant detection, the reference genome used was the Wuhan WIV04 strain

(hCoV-19/Wuhan/WIV04/2019, EPI_ISL_402124), downloaded from GISAID[17-19]. A

multiple sequence alignment, including the WIV04 reference, was generated in FASTA

format using MAFFT (online version 7) [20]. The WIV04 sequence was then removed

from the alignment, which was analyzed using AliView (version 1.28) [21] and custom

Python scripts (Python version 3.12.3[22]) to calculate allele frequencies at the

R203K/G204R sites in the N protein. Python scripts were also used to perform

statistical test (G-test).

To analyze the distribution of SARS-CoV-2 lineages across Japan during the study

period (July 27, 2020 – January 16, 2023), we utilized sequences registered in the

GISAID EpiCov database (https://gisaid.org) as of February 6, 2023. The filtering

criteria were:

•   Host: Human

•   Location: Asia / Japan

•   Collection date: July 27, 2020 – January 16, 2023

•   Sequence length: ≥ 29,000 nt

•   Passage history: Original

The resulting 539,504 sequences were analyzed using PANGOLIN for lineage

classification (see **Figure 1B**).

Data visualization was performed using R version 4.4.0[23] and the R

package *NipponMap*[24].

Data Records:

This dataset includes:

• Raw sequencing reads (SRA accessions under DRP012098)

• Consensus genome assemblies (FASTA format)

• BioSample metadata (BioSample accessions listed in the **Data Overview**

section)

• Lineage annotations (PANGOLIN)

All files follow International Nucleotide Sequence Database Collaboration (INSDC)[25]-

compliant formats. The PANGOLIN lineage annotations are archived exclusively in

GISAID and are not part of the INSDC repositories.

Data Overview:

Of the 18,475 SARS-CoV-2-positive samples, 14,201 were subjected to viral

genome sequencing. By the end of December 2021, sequencing was implemented for

100% of samples with Ct ≤30 and 93.92% of those with Ct value between 30–40. From

2022 onward, the coverage remained high Ct ≤30 samples (for 99.83%) but decreased

to 62.71% for Ct 30–40 samples (**Table 2**). Across all sequencing attempts, a total

of 8,266 SARS-CoV-2 genomes were successfully assembled. Among these, 2,917

sequences met the criteria for Category 1, and the remaining genomes were classified

Year/Group		Positive Samples		Genome Sequencing Performed (Sequencing performance rate)		Genomes Successfully Sequenced, Category 1 and 2 (Sequencing success rate)		Genomes Successfully Sequenced, Category 1 only (Sequencing success rate)	
- 2021	Ct ≤30	513		513 (100.00%)		346 (67.45%)		263 (51.27%)	
			2,025		1,933 (95.46%)		683 (33.73%)		444 (21.93%)
	Ct 30-40	1,512		1,420 (93.92%)		337 (22.29%)		181 (11.97%)	
2022 -	Ct ≤30	5,258		5,249 (99.83%)		4,389 (83.47%)		2,473 (47.03%)	
			16,450		12,268 (74.58%)		7,583 (46.10%)		2,473 (15.03%)
	Ct 30-40	11,192		7,019 (62.71%)		3,194 (28.54%)		0 (0.00%)	
	total	18,475		total 14,201		total 8,266		total 2,917	

Table 2. Summary of viral genome sequencing. The positive samples (18,475 samples
in total) were selectively implemented for genome sequencing (14,201 samples in total).
Eventually, 8,266 samples were successfully sequenced in both quality categories
(Category 1 and Category 2), while Category 1 consists of 2,917 samples. For the
definition of Categories 1 and 2, see **Table 3**.

Raw reads and consensus genome sequences for 8,266 SARS-CoV-2 samples were

submitted and are available for download from the NCBI Sequence Read Archive in the

International Nucleotide Sequence Database Collaboration (INSDC)[25], under the study

accession number DRP012098[26]. While 1,248 of them were used in previous

publication[1], all high-quality genome assemblies are currently available under the single

accession. The corresponding BioSample and other accession numbers are summarized

in **Table 3**. According to operational criteria (based on genome length and Ct values),

sequences were categorized into two groups (see **Table 3)**. Assembled genome

sequences were also submitted to GISAID, under the accession IDs listed in **Table 3**.

Category and Criteria	Number of Sequences	BioSample* Accession	Sequence Accession	SRA Run Accession	SRA Experiment Accession	GISAID ID
Category 1: Genome length ≥ 29,000 nucleotides and Ct value ≤ 40 (for - 2021) ≤30 (for 2022 -)	2917	SAMD00738313 - SAMD00741057, SAMD00741082, SAMD00786981 - SAMD00787151	BS007837 - BS010581, BS010689, BS016048 - BS016218	DRR600457 - DRR603201, DRR603222, DRR608552 - DRR608722	DRX581004 - DRX583748, DRX583769, DRX589099 - DRX589269	EPI_ISL_19576276 - EPI_ISL_19582591, EPI_ISL_19582637, EPI_ISL_19575435 - EPI_ISL_19575605
Category 2: Complete Genome not meeting Category 1 criteria and with unidentified nucleotides ≤ 20%	5349	SAMD00741058 - SAMD00741081, SAMD00741083 - SAMD00746626**	BS010669 - BS010688, BS010690 - BS016018	DRR603202 - DRR603221, DRR603223 - DRR608551	DRX583749 - DRX583768, DRX583770 - DRX589098	EPI_ISL_19582592 - EPI_ISL_19582606, EPI_ISL_19582632 - EPI_ISL_19582636, EPI_ISL_19582638 - EPI_ISL_19585391, EPI_ISL_19579847 - EPI_ISL_19580766, EPI_ISL_19578927 - EPI_ISL_19579846, EPI_ISL_19574700 - EPI_ISL_19575434

Table 3. Submission of genomic sequences to public repositories (INSDC/DDBJ

and GISAID). In total, 8,266 genomic assemblies were submitted to the public

repositories in this dataset.

* BioSample accessions provide metadata describing the biological source materials

used to generate sequencing data.

** 219 BioSample data did not meet Category 2 criteria; sequence data for these were

not submitted.

Technical Validation:

Statistics of Consensus Genome Sequences

We first examined the length distribution of the 8,266 complete consensus genome

sequences across both Category 1 and Category 2 (**Figure 4**). A sufficient number (n =

2,917) of nearly full-length genome sequences (≥ 29,000 nt; Category 1) were included

and exclusively used for SARS-CoV-2 lineage assignment via PANGOLIN, as well as

for subsequent analyses described below.

Epidemiological Distribution of SARS-CoV-2 Variants

Since the SBCVIC dataset represents a SARS-CoV-2 subpopulation within the broader Japanese population, we anticipated a comparable lineage distribution between our dataset and that of GISAID. Lineage analysis using PANGOLIN (**Figure 1B** and **1C**) revealed a similar distribution of SARS-CoV-2 variants between SBCVIC and GISAID datasets, supporting the validity of the overall SBCVIC sampling, sequencing, and analysis workflow.

Detection of R203K/G204R Mutations in the Nucleocapsid Protein

The 2,917 Category 1 genome assemblies were also used for further technical validation. As control datasets, we randomly selected and created three replicates of 2,917 SARS-CoV-2 genome sequences from symptomatic individuals of Japanese origin deposited in GISAID during the same period (July 27, 2020 to January 16, 2023) (**Table 4**). To demonstrate the genotyping capability of our dataset, we focused on two well-known adjacent co-occurring mutations R203K/G204R in the N protein of SARS-CoV-2, which are known to affect viral pathogenicity[12].

Genome sequences were aligned to the SARS-CoV-2 reference genome by MAFFT, and the allele frequencies at the R203K/G204R sites were calculated for both the GISAID (symptomatic) and SBCVIC (asymptomatic) groups (see **Methods**). Our SBCVIC dataset exhibited a lower rate of missing allele calls (5.9%) compared to the GISAID datasets (17.8% for replicate 1, 18.9% for replicate 2, and 18.7% for replicate 3), indicating higher call rate across the majority of samples while not showing the higher genotyping quality itself. Interestingly, the frequency of the mutant allele in the SBCVIC (asymptomatic) group was comparable to that in the GISAID (symptomatic) group (**Table 4**) ($P = 0.799676$ for replicate 1, $P = 0.133389$ for replicate 2, and $P =$

246 0.783957 for replicate 3, G-test), despite our initial expectation that mutation rates

247 would differ between symptomatic and asymptomatic individuals. While we do not

248 propose a specific hypothesis here, this allele-level analysis highlights a potential

249 application of the SBCVIC dataset.

	Wild allele (R203/G204)	Mutant allele (203K/204R)	Allele Not Determined
GISAID (Symptomatic, replicate 1, n=2917)	7	2391	519 (17.8%)
GISAID (Symptomatic, replicate 2, n=2917)	2	2365	550 (18.9%)
GISAID (Symptomatic, replicate 3, n=2917)	7	2365	545 (18.7%)
SBCVIC (Asymptomatic, n=2917)	7	2739	171 (5.9%)

Table 4. Number of wild-type and mutant alleles at the nucleocapsid protein mutation
sites (R203K/G204R) in symptomatic (GISAID, three replicates) and asymptomatic
(SBCVIC, this study) populations.

Usage Note:

The dataset of SARS-CoV-2 samples collected and analyzed from asymptomatic

individuals in Japan is highly distinctive. We demonstrated the validity of this SBCVIC

dataset for analyzing the epidemiological distribution of SARS-CoV-2 variants and for

conducting allele-level genotyping. Possible applications of our dataset are, analysis for

asymptomatic-specific mutations that would be keys for SARS-CoV-2 virulence (see

Technical Validation), training AI classifiers and Genome Large Language Models on

asymptomatic variant data, and building viral fitness model for SARS-CoV-2. We

believe that this dataset has significant potential for epidemiologists, geneticists, and

computational biologists to provide insights into the modulation of SARS-CoV-2

virulence and to enhance our understanding of viral fitness.

Code Availability

All custom scripts used in this study are available on GitHub

at https://github.com/oyasai55/8266_SARS-CoV-2_Genomic_Assemblies_Scripts/.

Acknowledgements

We thank Mr. Takahiro Sakurai and Mr. Hiroshi Ikawa for their support with

bioinformatics analyses. We also acknowledge Mr. Tsuyoshi Saito, Ms. Natsumi

Miyazaki, Mr. Atsushi Sugi, and Mr. Masanori Tamanaha for their assistance in

managing sample metadata.

Author contributions

H.O. and J.S.T. designed and performed the analyses and wrote the manuscript.

J.S.T. and Yuichi K. managed the genome sequence data from SBCVIC. S.O. conducted

bioinformatics and statistical analyses. J.S.T. and S.O. conducted figure/table

visualization. T.S. and W.S. supervised the entire analysis process. Moto K. directed the

research contract with the SBCVIC and intellectual properly. Y.T., S.Y., Minoru K. and

Yukumasa K. was responsible for RT-qPCR testing with the viral genome sequencing at

SBCVIC. M.I. and W.S. conceived an inspection system by SBCVIC and acquired

research funding. All authors critically reviewed and approved the final version of the

manuscript.

Competing interests:

This research was supported by the SB Coronavirus Inspection Center Corp. Moto

Kimura and Wataru Sugiura have received research grants from SB Coronavirus

Inspection Center Corp. Masato Ikeda, Yukumasa Kazuyama, and Minoru Kato are

employees of the SB Coronavirus Inspection Center Corp.

Figures:

In separate files (pdf).

Figure legends:

Figure 1. Monthly epidemiological distribution of SARS-CoV-2 variants in Japan

(July 27, 2020 - January 16, 2023). (A) Monthly number of newly confirmed COVID-

19 cases in Japan, based on open data from the Ministry of Health, Labour and Welfare

(accessed on May 1, 2023). (B) All domestic sequences registered in the Global

Initiative on Sharing All Influenza Data (GISAID) EpiCoV database as of February 6,

2023 (n = 539,504). (C) Positive cases detected in asymptomatic individuals in this

study (n = 2,917).

Figure 2. Number of tested (gray) and SARS-CoV-2-positive (blue) cases in this

study. Between July 27, 2020, and January 16, 2023, a total of 4,573,575 saliva samples

were tested (A), of which 18,475 were positive by RT-qPCR testing (B). Individuals

who opted out of the study were excluded from this count.

Figure 3. Geographic distribution of sample collection. Color intensity indicates the

number of samples collected, with darker shades representing higher counts. The top

five prefectures with the highest number of samples were Shiga (n = 6,082), Tokyo (n =

3,118), Nagasaki (n = 521), Osaka (n = 429), and Hokkaido (n = 401).

Figure 4. Distribution of SARS-CoV-2 genome consensus sequence lengths

obtained in this study (n = 8,266). A histogram shows the distribution of sequence

lengths (X-axis) and their frequencies (Y-axis). Category 1 sequences (n = 2,917) are

shown in dark purple, and Category 2 sequences (n = 5,349) are shown in green. The

dashed line represents the threshold for near-complete genome sequences (29,000 nt).

References:

1   Shiino, T. *et al.* Molecular epidemiology of SARS-CoV-2 genome sentinel
        surveillance in commercial COVID-19 testing sites targeting asymptomatic
        individuals during Japan's seventh epidemic wave. *Sci Rep* **14**, 20950 (2024).
        https://doi.org/10.1038/s41598-024-71953-8

2   Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of
        asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic
        review and meta-analysis. *PLoS Med* **17**, e1003346 (2020).
        https://doi.org/10.1371/journal.pmed.1003346

3   Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of
        asymptomatic and presymptomatic SARS-CoV-2 infections: Update of a living
        systematic review and meta-analysis. *PLoS Med* **19**, e1003987 (2022).
        https://doi.org/10.1371/journal.pmed.1003987

4   Casey-Bryars, M. *et al.* Presymptomatic transmission of SARS-CoV-2 infection: a
        secondary analysis using published data. *BMJ Open* **11**, e041240 (2021).
        https://doi.org/10.1136/bmjopen-2020-041240

5   Yang, Q. *et al.* Just 2% of SARS-CoV-2-positive individuals carry 90% of the virus
        circulating in communities. *Proc Natl Acad Sci U S A* **118** (2021).
        https://doi.org/10.1073/pnas.2104547118

6   Davis, J. T. *et al.* Cryptic transmission of SARS-CoV-2 and the first COVID-19
        wave. *Nature* **600**, 127-132 (2021). https://doi.org/10.1038/s41586-021-04130-
        w

7   Nabeshima, T. *et al.* COVID-19 cryptic transmission and genetic information
        blackouts: Need for effective surveillance policy to better understand disease

346   burden. *Lancet Reg Health West Pac* **7**, 100104 (2021).

347   https://doi.org/10.1016/j.lanwpc.2021.100104

348   8   Ferdinand, A. S. *et al.* An implementation science approach to evaluating

349   pathogen whole genome sequencing in public health. *Genome Med* **13**, 121

350   (2021). https://doi.org/10.1186/s13073-021-00934-7

351   9   Kitahara, K., Nishikawa, Y., Yokoyama, H., Kikuchi, Y. & Sakoi, M. An overview of

352   the reclassification of COVID-19 of the Infectious Diseases Control Law in Japan.

353   *Glob Health Med* **5**, 70-74 (2023). https://doi.org/10.35772/ghm.2023.01023

354   10   Takeuchi, J. S. *et al.* Large-scale screening of SARS-CoV-2 variants in Tokyo,

355   Japan: A 3-year and 9-month longitudinal survey. *Glob Health Med* **7**, 151-160

356   (2025). https://doi.org/10.35772/ghm.2025.01004

357   11   *The Ministry of Health, Labour and Welfare, Japan Infectious Disease Trend*

358   *Visualization System*, <https://covid19.mhlw.go.jp/en/> (2023).

359   12   Wu, H. *et al.* Nucleocapsid mutations R203K/G204R increase the infectivity,

360   fitness, and virulence of SARS-CoV-2. *Cell Host Microbe* **29**, 1788-1801 e1786

361   (2021). https://doi.org/10.1016/j.chom.2021.11.005

362   13   Terada-Hirashima, J. *et al.* Investigation of the use of PCR testing prior to ship

363   boarding to prevent the spread of SARS-CoV-2 from urban areas to less-

364   populated remote islands. *Glob Health Med* **4**, 174-179 (2022).

365   https://doi.org/10.35772/ghm.2022.01008

366   14   Katsuno, T. *et al.* Diagnostic accuracy of direct reverse transcription-polymerase

367   chain reaction using guanidine-based and guanidine-free inactivators for SARS-

368   CoV-2 detection in saliva samples. *J Virol Methods* **326**, 114909 (2024).

369   https://doi.org/10.1016/j.jviromet.2024.114909

370   15   ARTIC Network, SARS-CoV-2 (COVID-19). https://artic.network/viruses/sars-cov-

371   2.

372   16   O'Toole, A. *et al.* Assignment of epidemiological lineages in an emerging

373   pandemic using the pangolin tool. *Virus Evol* **7**, veab064 (2021).

374   https://doi.org/10.1093/ve/veab064

375   17   Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's

376   innovative contribution to global health. *Glob Chall* **1**, 33-46 (2017).

377   https://doi.org/10.1002/gch2.1018

378   18   Khare, S. *et al.* GISAID's Role in Pandemic Response. *China CDC Wkly* **3**, 1049-

379   1051 (2021). https://doi.org/10.46234/ccdcw2021.255

19　Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data -
from vision to reality. *Euro Surveill* **22** (2017). https://doi.org/10.2807/1560-
7917.ES.2017.22.13.30494

20　Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-
780 (2013). https://doi.org/10.1093/molbev/mst010

21　Larsson, A. AliView: a fast and lightweight alignment viewer and editor for large
datasets. *Bioinformatics* **30**, 3276-3278 (2014).
https://doi.org/10.1093/bioinformatics/btu531

22　Python Software Foundation. Python Language Reference, version 3.12.3.
https://www.python.org. (2024).

23　R: A Language and Environment for Statistical
Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-
project.org/. (2024).

24　Tanimura, S. NipponMap: Japanese Map Data and Functions. R package version
0.2, https://cran.r-project.org/web/packages/NipponMap. (2017).

25　Karsch-Mizrachi, I. *et al.* The international nucleotide sequence database
collaboration (INSDC): enhancing global participation. *Nucleic Acids Res* (2024).
https://doi.org/10.1093/nar/gkae1058

26　*NCBI Sequence Read Archive*,
<https://identifiers.org/ncbi/insdc.sra:DRP012098> (2024).

Histogram of sequence length (n=8,266)