Title:

8,266 SARS-CoV-2 Genomic Assemblies from Asymptomatic Carriers in Japan

Authors and Affiliations:

Hajime Ohyanagi[1†], Junko S Takeuchi[1†], Yuichi Kawanishi[1], Saya Ohi[1], Teiichiro

Shiino[1], Moto Kimura[1], Yusuke Takahashi[2], Shigeru Yoshida[2], Minoru Kato[3], Yukumasa

Kazuyama[3], Masato Ikeda[3], and Wataru Sugiura[1*]

[1]Center for Clinical Sciences, Japan Institute for Health Security, Tokyo, Japan

[2]Department of Clinical Laboratory Science, Health Sciences University of Hokkaido,

Hokkaido, Japan

[3]SB Coronavirus Inspection Center Corp., Tokyo, Japan

[†]These authors equally contributed

*Corresponding author: sugiura.w@jihs.go.jp

Abstract:

In the context of public health, asymptomatic carriers of respiratory infectious diseases

are considered a hidden yet critical factor in the transmission of infection and represent

a key target for efforts to mitigate disease spread. To contribute a unique genomic

resource to the SARS-CoV-2 research community, we collected SARS-CoV-2–positive

samples from asymptomatic individuals at the SB Coronavirus Inspection Center Corp.

during the COVID-19 pandemic in Japan and conducted a comprehensive analysis of

their viral genomes. Using Illumina COVIDSeq technology, we successfully generated

8,266 SARS-CoV-2 genome assemblies, all of which have been made publicly available

to facilitate further research. In this report, we summarize our efforts to collect SARS-

CoV-2–positive samples from asymptomatic individuals and highlight the key features

and accessibility of this genomic dataset.

Background & Summary:

We operate the SB Coronavirus Inspection Center Corp. (SBCVIC) [1] , which differs

from typical medical facilities by focusing on testing asymptomatic individuals. As a

result, positive cases detected at our center are considered asymptomatic carriers of

SARS-CoV-2. Epidemiologically, asymptomatic carriers of SARS-CoV-2 are known to

have non-negligible transmission capabilities comparable to those of symptomatic

individuals[2-5]. Therefore, their existence represents a hidden yet critical factor in

controlling the spread of the pandemic[6,7]. From a genomic perspective, samples from

asymptomatic carriers are also a valuable resource for investigating genetic factors in

the viral genome that may be associated with the presence or absence of clinical

symptoms in COVID-19 patients[8].

During the pandemic period in Japan[9,10], spanning from July 2020 to January 2023,

the country experienced eight epidemic waves of SARS-CoV-2 infection[11] (**Figure 1A**).

These included waves dominated by the Alpha (the 4[th] wave), Delta (the 5[th] wave), and

Omicron sublineages BA.1, BA.2, and BA.5 (from the 6[th] wave onward) (**Figure 1B** for

symptomatic cases, and **Figure 1C** for asymptomatic cases). Over the course of the

study (July 27, 2020 – January 16, 2023), we tested a total of 4,573,575 samples, among

which 18,475 tested positive for SARS-CoV-2 using reverse transcription real-time

quantitative polymerase chain reaction (RT-qPCR) (Ct values of ≤ 40) (**Figure 2** and

Table 1). The overall average positive rate among asymptomatic individuals was 0.40%.

The positive rates varied by wave: 0.05% during the 3[rd] wave, 0.13% in the 4[th] wave

(Alpha), 0.23% in the 5th wave (Delta), 4.5% in the 6th wave, 8.2% in the 7th wave

(BA.1/BA.2), and 5.2% in the 8th wave (BA.5). The number of newly confirmed cases

in Japan (**Figure 1A**) shows a trend similar to the positive rate among asymptomatic

individuals mentioned above.

Starting October 9, 2020 (approximately six weeks after the initiation of the study),

we began collecting background information on positive cases from 45 prefectures

across Japan, including Shiga, Tokyo, Nagasaki, Osaka, Hokkaido, Fukuoka, Chiba,

Gifu, Kanagawa, Saitama, Hyogo, Aichi, Kyoto, Miyagi, Gumma, Hiroshima, Nagano,

Kumamoto, Ibaraki, Shizuoka, Tochigi, Fukushima, Yamanashi, Nara, Miyazaki,

Okinawa, Mie, Ehime, Iwate, Okayama, Kagawa, Yamagata, Saga, Aomori, Akita,

Niigata, Toyama, Tottori, Ishikawa, Yamaguchi, Oita, Kagoshima, Fukui, Shimane, and

Kochi (**Figure 3**). No positive case was collected from two prefectures (Tokushima and

Wakayama). Among 18,475 positive cases, 6,408 (34.7%) were male, 5,944 (32.2%)

were female, and 6,123 (33.1%) had missing gender information. Age information was

available for 12,544 individuals, among whom the median age was 36 years

(interquartile range [IQR]: 27–44). (**Table 1**).

	Value
Tested individuals	4,573,575
Positive cases	18,475
Gender	
Male	6,408 (34.7%)
Female	5,944 (32.2%)
Unknown	6,123 (33.1%)
Age (Years)	36 [27 - 44]

Table 1. Statistics of the study participants. Gender is shown in n (%), and age is

presented as median [interquartile range] .

In total, we obtained 18,475 SARS-CoV-2-positive samples from asymptomatic

individuals. We performed whole-genome sequencing on residual saliva samples and

assembled viral genome sequences that met quality thresholds for downstream analysis.

Ultimately, we successfully generated 8,266 genome assemblies from these cases,

which are archived in the SBCVIC genomic dataset and categorized into two groups

(**Table 2** and **Table 3**) according to operational criteria (based on genome length and Ct

values).

For technical validation, we evaluated the distribution of consensus genome

sequence lengths and examined the lineage composition of the SARS-CoV-2 genomes.

We also assessed the presence of known mutations to confirm the reliability of our

sequencing pipeline for allele typing. In particular, we focused on two adjacent, co-

occurring mutations in the nucleocapsid (N) protein R203K/G204R[12], which are

associated with increased viral infectivity, fitness, and virulence. For further details, see

the **Technical Validation** section.

Methods:

SARS-CoV-2 Testing (Screening of Asymptomatic Cases)

Following previously described protocols[1,13], we conducted SARS-CoV-2 testing

and genome sequencing. The SBCVIC provided routine workplace-based SARS-CoV-2

screenings upon company request, as well as voluntary screenings requested by local

governments. As a result, most positive cases were considered asymptomatic

individuals [1,13]. Each participant self-collected approximately 2 mL of saliva using the

ZEESAN Saliva RNA Sample Collection Kit (MD-ZSV-001; Zeesan Biotech, Fujian,

China), which contained 1 mL of a guanidine-based viral inactivation buffer. The

sample was classified and shipped as UN 3373 Biological Substance, Category B. On

the day of arrival, SARS-CoV-2 RT-qPCR testing was performed using the SARS-CoV-

2 Direct Detection RT-qPCR Kit (Takara Bio, Shiga, Japan). Kits RC30JW and RD003

were used until April 4 and from April 5, 2021, respectively, according to the

manufacturer's protocol. Because the guanidine-based viral inactivation buffer can

inhibit RT-PCR reactions[14], we modified the pretreatment process. Specifically, 16 µL

of the saliva/buffer mixture was combined with 16 µL of water and 4 µL of pretreatment

reagent (Solution A, included in the kit) to dilute the inhibitor. The resulting 36 µL

mixture was incubated at room temperature for 5 minutes, followed by 5 minutes at

95°C. A 2.5 µL aliquot of this treated mixture was used as the RT-qPCR template. Test

results (positive or negative) were determined based on the protocol, with Ct values $\leq$

40 considered positive.

Viral Genome Sequencing

Samples from SARS-CoV-2-positive individuals were subjected to genome

sequencing. For RNA extraction, 50 µL of viral nucleic acid was extracted from 200 µL

of saliva samples using MagMAX Viral/Pathogen Nucleic Acid Isolation Kit (Thermo

Fisher Scientific, MA, USA) with MVP_Saliva_200_Flex_V1 protocol for KingFisher

Flex System (Thermo Fisher Scientific, MA, USA). The eluted RNA (20 µL) was

treated with RQ1 RNase-Free DNase (Promega, WI, USA). Subsequent cDNA

synthesis, target amplification using the ARTIC[15] primer set (versions used for each

sample are documented in the BioSample archive), and library preparation were

performed according to the Illumina COVIDseq Test Reference Guide (Illumina Inc.,

CA, USA). The sequencing implementation rates were 100.00% (-2021) and 99.83%

(2022-) for samples with Ct values $\leq$ 30 across all periods (see the column **Sequenced**

Samples in **Table 2**). However, the rate decreased for higher Ct values (30–40),

particularly from 2022 onward, due to operational constraints. Sequencing was

conducted on an Illumina NextSeq 2000 system.

Bioinformatics

Sequencing data were analyzed using the Illumina DRAGEN COVID Lineage

Pipeline (https://jp.illumina.com/products/by-type/informatics-products/basespace-

sequence-hub/apps/dragen-covid-lineage.html). Consensus sequences in FASTA format

were generated (see **Data Records**). The DRAGEN versions used for each sample are

documented in the BioSample archive. Sequences equal to or longer than 29,000

nucleotides in length (Category 1 in **Table 2** and **Table 3**) were analyzed for lineage

assignment using Phylogenetic Assignment of Named Global Outbreak Lineages

(PANGOLIN) version 4.3.1 (pangolin-data 1.25.1) [16].

For variant detection, the reference genome used was the Wuhan WIV04 strain

(hCoV-19/Wuhan/WIV04/2019, EPI_ISL_402124), downloaded from GISAID[17-19]. A

multiple sequence alignment, including the WIV04 reference, was generated in FASTA

format using MAFFT (online version 7) [20]. The WIV04 sequence was then removed

from the alignment, which was analyzed using AliView (version 1.28) [21] and custom

Python scripts (Python version 3.12.3[22]) to calculate allele frequencies at the

R203K/G204R sites in the N protein. Python scripts were also used to perform

statistical test (G-test).

To analyze the distribution of SARS-CoV-2 lineages across Japan during the study

period (July 27, 2020 – January 16, 2023), we utilized sequences registered in the

GISAID EpiCov database (https://gisaid.org) as of February 6, 2023. The filtering

criteria were:

- Host: Human

- Location: Asia / Japan

- Collection date: July 27, 2020 – January 16, 2023

- Sequence length: $\geq$ 29,000 nt

- Passage history: Original

The resulting 539,504 sequences were analyzed using PANGOLIN for lineage

classification (see **Figure 1B**).

Data visualization was performed using R version 4.4.0[23] and the R

package *NipponMap*[24].

Data Records:

Of the 18,475 SARS-CoV-2-positive samples, 14,201 were subjected to viral

genome sequencing. By the end of December 2021, sequencing was implemented for

100% of samples with Ct $\leq$30 and 93.92% of those with Ct value between 30–40. From

2022 onward, the coverage remained high Ct $\leq$30 samples (for 99.83%) but decreased

to 62.71% for Ct 30–40 samples (**Table 2**).

Year/Group		Positive Samples		Sequenced Samples (Sequencing implementation rate)		Genomes Determined, Category 1 and 2 (Sequencing success rate)		Genomes Determined, Category 1 only (Sequencing success rate)	
- 2021	Ct ≤30	513		513 (100.00%)		346 (67.45%)		263 (51.27%)	
			2,025		1,933 (95.46%)		683 (33.73%)		444 (21.93%)
	Ct 30-40	1,512		1,420 (93.92%)		337 (22.29%)		181 (11.97%)	
2022 -	Ct ≤30	5,258		5,249 (99.83%)		4,389 (83.47%)		2,473 (47.03%)	
			16,450		12,268 (74.58%)		7,583 (46.10%)		2,473 (15.03%)

Ct 30-40	11,192	7,019 (62.71%)	3,194 (28.54%)	0 (0.00%)
	total **18,475**	total **14,201**	total **8,266**	total **2,917**

Table 2. Summary of viral genome sequencing. The positive samples (18,475 samples in total) were selectively implemented for genome sequencing (14,201 samples in total). Eventually, 8,266 samples were successfully sequenced in both quality categories (Category 1 and Category 2), while Category 1 consists of 2,917 samples. For the definition of Categories 1 and 2, see **Table 3**.

Raw reads and consensus genome sequences for 8,266 SARS-CoV-2 samples were submitted and are available for download from the NCBI Sequence Read Archive in the International Nucleotide Sequence Database Collaboration (INSDC)[25], under the study accession number DRP012098[26]. While1,248 of them were used in previous publication[1], all high-quality genome assemblies are currently available under the single accession. The corresponding BioSample and other accession numbers are summarized in **Table 3**. According to operational criteria (based on genome length and Ct values), sequences were categorized into two groups (see **Table 3)**. Assembled genome sequences were also submitted to GISAID, under the accession IDs listed in **Table 3**.

Category and Criteria	Number of Sequences	BioSample* Accession	Sequence Accession	SRA Run Accession	SRA Experiment Accession	GISAID ID
Category 1: Genome length ≥ 29,000 nucleotides and Ct value ≤ 40 (for - 2021) ≤30 (for 2022 -)	2917	SAMD00738313 - SAMD00741057, SAMD00741082, SAMD00786981 - SAMD00787151	BS007837 - BS010581, BS010689, BS016048 - BS016218	DRR600457 - DRR603201, DRR603222, DRR608552 - DRR608722	DRX581004 - DRX583748, DRX583769, DRX589099 - DRX589269	EPI_ISL_19576276 - EPI_ISL_19582591, EPI_ISL_19582637, EPI_ISL_19575435 - EPI_ISL_19575605

Category 2: Complete Genome not meeting Category 1 criteria and with unidentified nucleotides ≤ 20%	5349	SAMD00741058 - SAMD00741081, SAMD00741083 - SAMD00746626**	BS010669 - BS010688, BS010690 - BS016018	DRR603202 - DRR603221, DRR603223 - DRR608551	DRX583749 - DRX583768, DRX583770 - DRX589098	EPI_ISL_19582592 - EPI_ISL_19582606, EPI_ISL_19582632 - EPI_ISL_19582636, EPI_ISL_19582638 - EPI_ISL_19585391, EPI_ISL_19579847 - EPI_ISL_19580766, EPI_ISL_19578927 - EPI_ISL_19579846, EPI_ISL_19574700 - EPI_ISL_19575434

Table 3. Submission of genomic sequences to public repositories (INSDC/DDBJ and GISAID). In total, 8,266 genomic assemblies were submitted to the public repositories in this dataset.

* BioSample accessions provide metadata describing the biological source materials used to generate sequencing data.

** 219 BioSample data did not meet Category 2 criteria; sequence data for these were not submitted.

Technical Validation:

Statistics of Consensus Genome Sequences

We first examined the length distribution of the 8,266 complete consensus genome sequences across both Category 1 and Category 2 (**Figure 4**). A sufficient number (n = 2,917) of nearly full-length genome sequences ($\geq$ 29,000 nt; Category 1) were included and exclusively used for SARS-CoV-2 lineage assignment via PANGOLIN, as well as for subsequent analyses described below.

Epidemiological Distribution of SARS-CoV-2 Variants

Since the SBCVIC dataset represents a SARS-CoV-2 subpopulation within the broader Japanese population, we anticipated a comparable lineage distribution between our dataset and that of GISAID. Lineage analysis using PANGOLIN (**Figure 1B** and **1C**) revealed a similar distribution of SARS-CoV-2 variants between SBCVIC and

GISAID datasets, supporting the validity of the overall SBCVIC sampling, sequencing,

and analysis workflow.

Detection of R203K/G204R Mutations in the Nucleocapsid Protein

The 2,917 Category 1 genome assemblies were also used for further technical

validation. As control datasets, we randomly selected and created three replicates of

2,917 SARS-CoV-2 genome sequences from symptomatic individuals of Japanese

origin deposited in GISAID during the same period (July 27, 2020 to January 16, 2023)

(**Table 4**). To demonstrate the genotyping capability of our dataset, we focused on two

well-known adjacent co-occurring mutations R203K/G204R in the N protein of SARS-

CoV-2, which are known to affect viral pathogenicity[12].

Genome sequences were aligned to the SARS-CoV-2 reference genome by MAFFT,

and the allele frequencies at the R203K/G204R sites were calculated for both the

GISAID (symptomatic) and SBCVIC (asymptomatic) groups (see **Methods**). Our

SBCVIC dataset exhibited a lower rate of missing allele calls (5.9%) compared to the

GISAID datasets (17.8% for replicate 1, 18.9% for replicate 2, and 18.7% for replicate

3), indicating higher call rate across the majority of samples while not showing the

higher genotyping quality itself. Interestingly, the frequency of the mutant allele in the

SBCVIC (asymptomatic) group was comparable to that in the GISAID (symptomatic)

group (**Table 4**) ($P = 0.799676$ for replicate 1, $P = 0.133389$ for replicate 2, and $P =$

0.783957 for replicate 3, G-test), despite our initial expectation that mutation rates

would differ between symptomatic and asymptomatic individuals. While we do not

propose a specific hypothesis here, this allele-level analysis highlights a potential

application of the SBCVIC dataset.

	Wild allele (R203/G204)	Mutant allele (203K/204R)	Allele Not Determined
GISAID (Symptomatic, replicate 1, n=2917)	7	2391	519 (17.8%)
GISAID (Symptomatic, replicate 2, n=2917)	2	2365	550 (18.9%)
GISAID (Symptomatic, replicate 3, n=2917)	7	2365	545 (18.7%)
SBCVIC (Asymptomatic, n=2917)	7	2739	171 (5.9%)

Table 4. Number of wild-type and mutant alleles at the nucleocapsid protein mutation
sites (R203K/G204R) in symptomatic (GISAID, three replicates) and asymptomatic
(SBCVIC, this study) populations.

Usage Notes:

The dataset of SARS-CoV-2 samples collected and analyzed from asymptomatic

individuals in Japan is highly distinctive. We demonstrated the validity of this SBCVIC

dataset for analyzing the epidemiological distribution of SARS-CoV-2 variants and for

conducting allele-level genotyping. Possible applications of our dataset are, analysis for

asymptomatic-specific mutations that would be keys for SARS-CoV-2 virulence (see

Technical Validation), training AI classifiers and Genome Large Language Models on

asymptomatic variant data, and building viral fitness model for SARS-CoV-2. We

believe that this dataset has significant potential for epidemiologists, geneticists, and

computational biologists to provide insights into the modulation of SARS-CoV-2

virulence and to enhance our understanding of viral fitness.

Data Availability

The dataset has been deposited to INSDC/DDBJ under the study accession number

DRP012098[26].

Code Availability

All custom scripts used in this study are available on GitHub

at https://github.com/oyasai55/8266_SARS-CoV-2_Genomic_Assemblies_Scripts/.

Acknowledgements

We thank Mr. Takahiro Sakurai and Mr. Hiroshi Ikawa for their support with

bioinformatics analyses. We also acknowledge Mr. Tsuyoshi Saito, Ms. Natsumi

Miyazaki, Mr. Atsushi Sugi, and Mr. Masanori Tamanaha for their assistance in

managing sample metadata.

Ethics Approval:

Procedures complied with national, institutional committees on human

experimentation and with the Helsinki Declaration (1975, revised in 2008). Sample

collection ensured anonymity. Study details and opt-out were provided on the websites

of the participating institutions. The study protocol was approved by the Institutional

Review Board of the National Center for Global Health and Medicine (approval

number: NCGM-G-004225-00, August 5, 2020).

Author contributions

H.O. and J.S.T. designed and performed the analyses and wrote the manuscript.

J.S.T. and Yuichi K. managed the genome sequence data from SBCVIC. S.O. conducted

bioinformatics and statistical analyses. J.S.T. and S.O. conducted figure/table

visualization. T.S. and W.S. supervised the entire analysis process. Moto K. directed the

research contract with the SBCVIC and intellectual properly. Y.T., S.Y., Minoru K. and

Yukumasa K. was responsible for RT-qPCR testing with the viral genome sequencing at

SBCVIC. M.I. and W.S. conceived an inspection system by SBCVIC and acquired

research funding. All authors critically reviewed and approved the final version of the

manuscript.

Competing interests:

This research was supported by the SB Coronavirus Inspection Center Corp. Moto

Kimura and Wataru Sugiura have received research grants from SB Coronavirus

Inspection Center Corp. Masato Ikeda, Yukumasa Kazuyama, and Minoru Kato are

employees of the SB Coronavirus Inspection Center Corp.

Figures:

In separate files (pdf).

Figure legends:

Figure 1. Monthly epidemiological distribution of SARS-CoV-2 variants in Japan

(July 27, 2020 - January 16, 2023). (A) Monthly number of newly confirmed COVID-

19 cases in Japan, based on open data from the Ministry of Health, Labour and Welfare

(accessed on May 1, 2023). (B) All domestic sequences registered in the Global

Initiative on Sharing All Influenza Data (GISAID) EpiCoV database as of February 6,

2023 (n = 539,504). (C) Positive cases detected in asymptomatic individuals in this

study (n = 2,917).

Figure 2. Number of tested (gray) and SARS-CoV-2-positive (blue) cases in this

study. Between July 27, 2020, and January 16, 2023, a total of 4,573,575 saliva samples

were tested, of which 18,475 were positive by RT-qPCR testing. Individuals who opted

out of the study were excluded from this count.

Figure 3. Geographic distribution of sample collection. Color intensity indicates the

number of samples collected, with darker shades representing higher counts. The top

five prefectures with the highest number of samples were Shiga (n = 6,082), Tokyo (n =

3,118), Nagasaki (n = 521), Osaka (n = 429), and Hokkaido (n = 401).

Figure 4. Distribution of SARS-CoV-2 genome consensus sequence lengths

obtained in this study (n = 8,266). A histogram shows the distribution of sequence

lengths (X-axis) and their frequencies (Y-axis). Category 1 sequences (n = 2,917) are

shown in dark purple, and Category 2 sequences (n = 5,349) are shown in green. The

dashed line represents the threshold for near-complete genome sequences (29,000 nt).

References:

1   Shiino, T. *et al.* Molecular epidemiology of SARS-CoV-2 genome sentinel
surveillance in commercial COVID-19 testing sites targeting asymptomatic
individuals during Japan's seventh epidemic wave. *Sci Rep* **14**, 20950 (2024).
https://doi.org/10.1038/s41598-024-71953-8
2   Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of
asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic
review and meta-analysis. *PLoS Med* **17**, e1003346 (2020).
https://doi.org/10.1371/journal.pmed.1003346
3   Buitrago-Garcia, D. *et al.* Occurrence and transmission potential of
asymptomatic and presymptomatic SARS-CoV-2 infections: Update of a living
systematic review and meta-analysis. *PLoS Med* **19**, e1003987 (2022).
https://doi.org/10.1371/journal.pmed.1003987

4   Casey-Bryars, M. *et al.* Presymptomatic transmission of SARS-CoV-2 infection: a
secondary analysis using published data. *BMJ Open* **11**, e041240 (2021).
https://doi.org/10.1136/bmjopen-2020-041240
5   Yang, Q. *et al.* Just 2% of SARS-CoV-2-positive individuals carry 90% of the virus
circulating in communities. *Proc Natl Acad Sci U S A* **118** (2021).
https://doi.org/10.1073/pnas.2104547118
6   Davis, J. T. *et al.* Cryptic transmission of SARS-CoV-2 and the first COVID-19
wave. *Nature* **600**, 127-132 (2021). https://doi.org/10.1038/s41586-021-04130-
w
7   Nabeshima, T. *et al.* COVID-19 cryptic transmission and genetic information
blackouts: Need for effective surveillance policy to better understand disease
burden. *Lancet Reg Health West Pac* **7**, 100104 (2021).
https://doi.org/10.1016/j.lanwpc.2021.100104
8   Ferdinand, A. S. *et al.* An implementation science approach to evaluating
pathogen whole genome sequencing in public health. *Genome Med* **13**, 121
(2021). https://doi.org/10.1186/s13073-021-00934-7
9   Kitahara, K., Nishikawa, Y., Yokoyama, H., Kikuchi, Y. & Sakoi, M. An overview of
the reclassification of COVID-19 of the Infectious Diseases Control Law in Japan.
Glob Health Med **5**, 70-74 (2023). https://doi.org/10.35772/ghm.2023.01023
10   Takeuchi, J. S. *et al.* Large-scale screening of SARS-CoV-2 variants in Tokyo,
Japan: A 3-year and 9-month longitudinal survey. *Glob Health Med* **7**, 151-160
(2025). https://doi.org/10.35772/ghm.2025.01004
11   *The Ministry of Health, Labour and Welfare, Japan Infectious Disease Trend*
Visualization System, <https://covid19.mhlw.go.jp/en/> (2023).
12   Wu, H. *et al.* Nucleocapsid mutations R203K/G204R increase the infectivity,
fitness, and virulence of SARS-CoV-2. *Cell Host Microbe* **29**, 1788-1801 e1786
(2021). https://doi.org/10.1016/j.chom.2021.11.005
13   Terada-Hirashima, J. *et al.* Investigation of the use of PCR testing prior to ship
boarding to prevent the spread of SARS-CoV-2 from urban areas to less-
populated remote islands. *Glob Health Med* **4**, 174-179 (2022).
https://doi.org/10.35772/ghm.2022.01008
14   Katsuno, T. *et al.* Diagnostic accuracy of direct reverse transcription-polymerase
chain reaction using guanidine-based and guanidine-free inactivators for SARS-
CoV-2 detection in saliva samples. *J Virol Methods* **326**, 114909 (2024).
https://doi.org/10.1016/j.jviromet.2024.114909

15 ARTIC Network, SARS-CoV-2 (COVID-19). https://artic.network/viruses/sars-cov-
2.
16 O'Toole, A. *et al.* Assignment of epidemiological lineages in an emerging
pandemic using the pangolin tool. *Virus Evol* **7**, veab064 (2021).
https://doi.org/10.1093/ve/veab064
17 Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's
innovative contribution to global health. *Glob Chall* **1**, 33-46 (2017).
https://doi.org/10.1002/gch2.1018
18 Khare, S. *et al.* GISAID's Role in Pandemic Response. *China CDC Wkly* **3**, 1049-
1051 (2021). https://doi.org/10.46234/ccdcw2021.255
19 Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data -
from vision to reality. *Euro Surveill* **22** (2017). https://doi.org/10.2807/1560-
7917.ES.2017.22.13.30494
20 Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-
780 (2013). https://doi.org/10.1093/molbev/mst010
21 Larsson, A. AliView: a fast and lightweight alignment viewer and editor for large
datasets. *Bioinformatics* **30**, 3276-3278 (2014).
https://doi.org/10.1093/bioinformatics/btu531
22 Python Software Foundation. Python Language Reference, version 3.12.3.
https://www.python.org. (2024).
23 R: A Language and Environment for Statistical
Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-
project.org/. (2024).
24 Tanimura, S. NipponMap: Japanese Map Data and Functions. R package version
0.2, https://cran.r-project.org/web/packages/NipponMap. (2017).
25 Karsch-Mizrachi, I. *et al.* The international nucleotide sequence database
collaboration (INSDC): enhancing global participation. *Nucleic Acids Res* (2024).
https://doi.org/10.1093/nar/gkae1058
26 *NCBI Sequence Read Archive*,
<https://identifiers.org/ncbi/insdc.sra:DRP012098> (2024).

Number of Tested and Positive Cases of SARS–CoV–2 in SBCVIC

Histogram of sequence length (n=8,266)