Development of a behavioral model for the use of psychosocial support services by cancer survivors: A grounded theory pilot study

Short title: A behavioral model for use of psychosocial support services by cancer survivors

Manae Hirayama[1], Daisuke Kasai[1], Ken-ichi Tabei[1]

[1] Graduate School of Industrial Technology, Advanced Institute of Industrial Technology, Tokyo Metropolitan Public University Corporation

Corresponding Author

Ken-ichi Tabei
Graduate School of Industrial Technology, Advanced Institute of Industrial Technology,
Tokyo Metropolitan Public University Corporation
Tokyo, Japan
03-3472-7831
tabei-kenichi@aiit.ac.jp

Abstract

This pilot study explored the factors influencing the theoretical use of psychosocial support services among cancer survivors not currently using those services. It used the Fogg Behavior Model as a theoretical framework. A web-based questionnaire survey was conducted over one month using snowball sampling through social networking services of patient organizations, targeting cancer survivors aged 40 years and older. Free-text responses about support needs from participants who were not using psychosocial support were analyzed using a grounded theory approach. The texts were coded based on the three Fogg Behavior Model elements: "triggers" to initiate behavior, "motivation" to engage in behavior, and "ability" that hinders behavior. The structure serving as the starting point for behavior was modeled. The results revealed that many factors were based on emotional values, such as expectations and hopes for support. "Motivation" indicated the need to build trust with supporters and to organize feelings. "Ability" identified that barriers included self-sufficiency and difficulties verbalizing feelings. "Trigger" required supporters to show human qualities, and for there to be clear pathways to consultation and creative publicity methods. These findings suggest that acquiring psychological values such as reassurance, trust, and hope strongly motivates the use of psychosocial support. Designing appropriate support pathways based on cancer survivors' behaviors and developing effective cognitive strategies are important ways to promote the future use of psychosocial support services.

Keywords: Cancer survivor, Fogg Behavior Model, Psychosocial support, Behavior change, Help-seeking

Introduction

In recent years, advances in cancer drugs and treatments have improved the survival rates for many types of cancer, leading to an increase in the number of cancer survivors [1]. It is estimated that there are more than 50 million cancer survivors worldwide [2], including approximately 3.41 million in Japan [3]. Cancer survivors, regardless of whether they are still receiving treatment, may experience feelings of anxiety or depression, known as emotional distress. Approximately 30% of individuals continue to experience these symptoms even after treatment ends [4]. Emotional distress affects mental health and also has physical and behavioral impacts, leading to an overall decrease in the quality of life [5]. Psychosocial support is considered effective in improving quality of life [6], and providing this support is crucial [7]. In Japan, efforts are being made to promote the use of cancer consultation support centers by providing accurate information [8], but awareness is only 55.1%, and usage remains at 21.1%, indicating that these services are not being fully used [9]. Additionally, patients who need psychosocial support but do not receive it experience greater distress than those who do [10]. According to a survey by an insurance company, about 60% of cancer survivors stated they "hesitate to ask for help" [11]. Other studies have identified stigma attached to psychosocial support, such as the desire to keep personal affairs to themselves, or not wanting to have personal problems managed by others, as well as concerns about whether support is available to meet individual needs [12][13][14]. Another study investigating barriers to availability of psychosocial care services highlighted reasons such as healthcare providers prioritizing treatment or medical procedures over psychosocial care, and a shortage of professionals to provide this support. Even in high-income countries, the psychosocial needs of cancer survivors are generally not fully met [15]. Previous studies have reported low utilization rates, user trends, and the characteristics of those who do

not need psychosocial support. However, it is not clear which specific behavioral factors would

improve utilization rates. This study aimed to identify the factors that would encourage cancer

survivors who do not currently use psychosocial support services to start to do so. We used a

descriptive approach, with the Fogg Behavior Model (FBM) [16] as a theoretical framework to

explain the use of psychosocial support services. The FBM is a theoretical model that indicates

the three necessary elements to trigger behavior: motivation, ability, and trigger [16]. Using this

framework clarifies the specific behavioral factors related to the use of psychosocial support by

cancer survivors, systematizes the design and evaluation of interventions, and facilitates a

common understanding among stakeholders [17]. This makes it easier to examine the

effectiveness of intervention measures.

Several recent studies have applied the FBM to cancer. For example, one study used a design

based on the FBM to make proposals to improve compliance with a liver cancer screening

intervention using a mobile health approach, focused on high-risk groups [18]. A pilot

randomized controlled trial of a mobile app promoting genetic counseling use among ovarian

cancer patients showed improvements in knowledge, self-efficacy, and family communication

[19]. These suggest that applying the FBM to the use of psychosocial support by cancer

survivors is also likely to be effective. This study aimed to clarify the specific motivational,

ability-related, and trigger factors affecting the use of psychosocial support by cancer survivors.

It therefore qualitatively analyzed the factors associated with willingness to use support services

using the FBM, and proposed strategies to improve utilization rates. It is expected to contribute

to the development of effective intervention strategies to improve use of support services.

Methods

Materials

Using Google Forms and snowball sampling, the questionnaire was distributed to cancer survivors aged 40 years or older through the social networking services of multiple patient groups over a one-month period (May 1 to May 31, 2024). The questionnaire included (1) questions about the use of psychosocial support for cancer patients, with three questions for those who answered "yes" and two questions for those who answered "no," and (2) one open-ended question about expectations of psychosocial support.

We adopted a detailed methodological approach based on the FBM and analyzed the survey results. This approach enables a comprehensive understanding of the factors influencing cancer survivors' use of psychosocial support services. The sample size for this pilot study was determined based on the following criteria. For continuous variables [20] [21], 12 participants were required in each group. We also required a total of 36 participants to ensure a 95% confidence interval (0.85–0.95) with a 5% margin of error [22].

Statistical analysis

The basic attributes, use of psychosocial support, and situation from diagnosis to the present were analyzed using descriptive statistics. For the content analysis of psychosocial support needs among those not using counseling support, we used the grounded theory approach [23][24] to structure the psychosocial needs of cancer patients and the actual state of support. The free descriptions were open-coded. We defined code families and concepts based on the three

110   elements of the FBM, motivation, ability, and trigger. Theoretical coding [25] was used to model

111   the structure underlying the behavior (Table 1).

113   **Table 1. Code families used to categorize the requests of cancer survivors not using**

114   **psychosocial support**

Code Family	Concept
Interaction	Empathy and Human Communication
Condition	Qualities and Competencies of Physicians, Medical Staff, and Support Personnel
Barrier	Psychological, Social, and Environmental Barriers
Consequence	Outcomes and Effects Experienced After Use
Means	Methods and Channels That Facilitate Access or Use
Time	Appropriate Timing to Promote Use
Strategy	Promotional Activities and Outreach to Encourage Utilization

The first author was responsible for the survey design and initial coding of free descriptions.

During the analysis, the data were reviewed and discussed multiple times with co-authors

experienced in medicine and patient research to eliminate biased perspectives and unilateral

interpretations of the data. R Version 4.3.2 was used for descriptive statistics, and MAXQDA

was used for the theoretical coding of descriptive data.

Ethical approval

This study was approved by the Research Safety and Ethics Committee of the Tokyo

Metropolitan University of Industrial Technology (Approval No. 23020). The questionnaire

survey was conducted in accordance with the Declaration of Helsinki and relevant guidelines,

and informed consent was obtained from all respondents. All response data were anonymized.

Results

Characteristics of participants

Responses were obtained from 63 participants. The average age was 55.5 years, with 43 women

(68.3%) and 20 men (31.7%) included in the study (Table 2). Overall, 32 people (50.8% said that

they had used psychosocial support services. The reasons for use were as follows: "Had

questions or concerns" (15 people, 46.9%), "Wanted information about medical or welfare

services" (6 people, 18.8%), "Wanted to learn about treatment or care options" (three people,

9.4%), "Needed support from family or those around them" (one person, 3.1%), and "Other"

(seven people, 21.9%). The psychosocial support services used included the Cancer Consultation

Support Center (18 people), cancer salons or patient groups (10 people), telephone services (two

people), apps or online services (one person), and consulting medical professionals (one person).

Table 2. Basic attributes of participants

Patient characteristic	n	%
Age (mean ± SD)	55.5 ± 9.3	
Sex		
Female	43	68.3

Male	20	31.7
Site of cancer		
Breast cancer	18	28.5
GIST	6	9.5
Ovarian cancer/Fallopian tube cancer	6	9.5
Pancreatic cancer	5	7.9
Colorectal cancer	4	6.3
Lung cancer	4	6.3
Glioma	2	3.2
Malignant lymphoma	2	3.2
Stomach cancer	2	3.2
Uterine cancer	2	3.2
Acute lymphoblastic leukemia	1	1.6
Anal cancer	1	1.6
Bladder cancer	1	1.6
Brain cancer	1	1.6
Cancer of unknown primary	1	1.6
Chronic myelogenous leukemia	1	1.6
Colorectal cancer	1	1.6
Kidney cancer	1	1.6
Liver cancer	1	1.6
PNET (Primitive Neuroectodermal Tumor)	1	1.6
Prostate cancer	1	1.6

Vaginal cancer	1	1.6

In contrast, 31 (49.2%) participants said that they did not use these services. The reasons for not

using them were: "I sufficiently understand my current situation" (four people, 12.9%), "I

research and find information myself" (six people, 19.4%), "I do not feel the need for

counselling support" (five people, 16.1%), "The support provided by medical professionals is

sufficient" (four people, 12.9%), and "Other" (13 people, 41.9%) (Table 3).

Table 3. Trends in the use of consultation support

	n	%
Have you ever used cancer counseling support?		
Yes	32	50.8
No	31	49.2
What motivated you to seek cancer counseling support?		
Had questions or concerns	15	46.9
Wanted to know about treatment and care options	3	9.4
Needed support from family or others	1	3.1
Wanted information about medical and social services	6	18.8
Other	7	21.9
What type of cancer counseling support do you use?		
Cancer Consultation Support Center	1	1.6
Cancer salons and patient groups	18	28.6
Telephone services	10	15.9

Mobile apps and online services	2	3.2
Other	1	1.6
Why did you not use counselling support?		
The present situation is fully comprehended.	3	9.7
The present author is conducting independent research and gathering information.	6	19.4
It is not perceived to be necessary to seek counselling or support.	5	16.1
The level of support provided by my healthcare provider is adequate.	4	12.9
Other	13	41.9

Barriers and facilitators to the use of psychosocial support

This section focuses on insights gained from open-ended responses by individuals not using

psychosocial support. We also qualitatively analyzed the specific factors that influenced the use

of support. The results were categorized into themes such as "Motivation", which underpins

behavior, "Ability", which enables or inhibits behavior; and "Trigger", which prompts behavior.

We also included "Interaction", "Conditions", "Outcomes", "Means", "Strategies", "Barriers"

and "Time" (Table 4). Of the 32 open-ended responses, 16 (50.0%) were about motivation, 13

(40.6%) about triggers, and two (6.3%) about ability. Fig 1 shows a model of psychosocial

support use behavior created by adapting the FBM using these results. The model indicates that

when the ability barrier influencing behavior is lower, it is more likely that a trigger will succeed

in driving action. Conversely, when the barrier is higher, it is more likely that a trigger will fail

to drive action. This suggests that when motivation, the driving force behind the behavior, is high,

a trigger is more likely to drive action, whereas when motivation is low, it is more likely to fail

to do so (Fig 1).

Table 4. The coded text and themes

Sex	FBM	Theme	Text
Female	Trigger	Condition	It helps when someone with empathy reaches out to you.
Female	Trigger	Condition	It feels easier to use when someone initiates the conversation.
Female	Trigger	Means	I want support that understands my work and workplace situation.
Female	Trigger	Time	It's helpful if someone listens before the consultation or while waiting.
Female	Trigger	Means	It would be great to have online consultations via Zoom or similar platforms.

Female	Trigger	Means	Support that helps search for necessary information together would be beneficial.
Female	Trigger	Means	I wanted to connect with other patients who have similar experiences.
Female	Trigger	Means	An orientation session right after diagnosis would be very helpful.
Male	Trigger	Condition	It helps to get advice from a trustworthy expert when you're unsure.
Male	Trigger	Strategy	I'd appreciate having booklets and flyers readily available to take home.
Male	Trigger	Strategy	I'd like to see videos or interviews of other patients using consultation services.

Male	Trigger	Means	I think a counseling service could relieve stress by allowing people to share concerns.
Male	Trigger	Strategy	Promoting it in a way that makes me think, "Maybe I should try it," would help.
Female	Ability	Barrier	I've had a vague unease I can't put into words, so I don't know how to seek help.
Male	Ability	Barrier	I tend to rely on looking things up myself and don't seek help from others.
Female	Motivation	Consequence	I want to ask questions I find hard to bring up with my doctor.
Female	Motivation	Consequence	Routine appointments make it hard to raise issues; I want to share even minor concerns.

Female	Motivation	Interaction	I've felt lonely since diagnosis, and I hope someone can support me through it.
Female	Motivation	Consequence	I need help maintaining a positive mindset and emotional balance.
Female	Motivation	Consequence	I wanted to learn how to support my child emotionally while managing my own feelings.
Female	Motivation	Consequence	I'd like referrals or info on hospitals for preventing recurrence or metastasis.
Female	Motivation	Interaction	I appreciate when someone listens closely and helps organize my thoughts and feelings.
Female	Motivation	Interaction	A place where they truly listen to your story is valuable.

Female	Motivation	Interaction	Support should be personalized based on individuals and their family context.
Female	Motivation	Interaction	Support systems should vary to match the patient's stage and specific needs.
Male	Motivation	Interaction	A place where hope for life can be nurtured through counseling is important.
Male	Motivation	Consequence	I want guidance on how to distance myself from people who aren't helpful.
Male	Motivation	Consequence	I'd like to know if remission is possible and how to achieve it.
Male	Motivation	Consequence	I want comprehensive information, including clinical trials.

Male	Motivation	Consequence	I lack information due to limited explanations from my doctor, so I want more details.
Male	Motivation	Consequence	I want a one-stop place to talk about everything—from medical info to daily concerns.
Female	Motivation	Interaction	Rather than qualifications, supporters should be chosen based on their actual abilities.

Fig 1. Behavioral model of psychosocial support use based on Fogg's Behavior Model

The following section discusses each theme in more detail.

Theme 1: Motivation—security and hope brought about by interaction and

outcomes

The primary motivations for using psychosocial support were centered around building trust

through interaction and obtaining a sense of security and hope. For example, one user expressed

expectations about the support provider's character and ability to empathize with users:

"We should select people with ability, rather than just those with qualifications (such as social

workers or nurses) or suitability." (Interaction/Motivation)

Expectations for psychosocial support included the provision of information or assistance with

decision-making and also organizing feelings and gaining a sense of reassurance.

"When I want to ask something that's difficult to bring up with the doctor, I can ask."

(Outcome/Motivation) These points indicate a desire to make a human connection and have

some hope for the future, beyond just the "information" or "skills" provided by psychosocial

support.

Theme 2: Barriers to action (ability): obstacles to use caused by difficulty

articulating feelings and a tendency toward self-reliance

Two statements corresponded to "Ability", and they shared themes such as "difficulty putting

feelings into words" and "hesitation to rely on others".

"I've constantly carried a vague sense of unease that I can't put into words, so I don't know how

to ask for advice or seek support." (Female/Barrier/Ability) *"I tend to think I understand things*

after researching them myself, so I don't really try to rely on others." (Male/Barrier/Ability)

The former expresses frustration at the "difficulty explaining in words", and the latter reveals

hesitation because of a self-reliant attitude. Neither participant rejected psychosocial support

itself, but both were in a psychological state in which it is hard to find an "opportunity" to access

support. This highlights the complexity of "Ability" within the FBM.

Theme 3: Trigger: promoting usage behavior through presentation of conditions, means, and strategies

The trigger to encourage behavior required not only "conditions" and "means", but also

"strategically designed pathways".

• **Conditions**

"*It should be someone with human depth.*" (Condition/Trigger)

"*To receive advice based on expertise from someone trustworthy.*" (Condition/Trigger) The

person who responds and the attitude they take was an important starting point in making

participants feel that it would be okay to consult with that person.

• **Means**

"*It would be good to have a mechanism where orientation is received immediately after*

diagnosis." (Means/Trigger)"*Support where necessary information can be searched together.*"

(Means/Trigger)

Means help to develop a pathway to action, and supportive guidance acts as a trigger.

• **Strategies**

"*If the service was publicized, I might want to use it.*"(Strategy/Trigger)

"*I want to see videos or interviews showing what a consultation support center looks like from*

the patient's perspective." (Strategy/Trigger)

This indicates that "ingenious measures to encourage potential users" are crucial to promote the

use of psychosocial support.

Discussion

Principal results

In this study, we hypothesized that cancer survivors who do not use psychosocial support services experience challenges such as stigma about seeking support and a lack of trust in these services. Based on the FBM, we investigated the factors related to the use of psychosocial support. The analysis revealed that many respondents held emotional values such as expectations and hopes for psychosocial support. However, factors such as "a lack of appropriate pathways" and "an environment that does not create opportunities for action" emerged as barriers to its use in practice.

Comparison with prior work

First and foremost, motivation focused on building trust through interactions and obtaining a sense of security and hope as a result. Women, in particular, emphasized expectations of support to enable "emotional sorting" and developing a sense of security. Previous studies have also reported that female cancer survivors tend to feel anxious [26] and have a strong tendency to seek empathy and support for their feelings [27][28], which is consistent with our results. In contrast, men emphasized the need for information about clinical trials and the remission period of their illness, focusing less on psychological support. Previous studies have also shown that men have a low demand for psychological support [29], which is consistent with our findings. We also obtained insights about barriers to action related to ability, such as difficulties expressing emotions. This highlighted a psychological state of having no opportunity or not wanting to rely on others. The tendency for self-sufficiency may be attributed to the Dunning–

Kruger effect [30]. However, those who self-rate their health literacy as high [31] often do not

realize the inadequacy of their understanding. This tendency has been reported in other studies

[32][33], and was also reflected in our results. The inability to verbalize emotions, known as

inhibited self-disclosure, has been noted as a factor hindering help-seeking behavior in patients

with ovarian cancer [34], which is consistent with our findings.

Triggers included aspects such as the humanity of supporters, clarification of consultation or

counselling procedures, and promotional activities to encourage awareness. These findings

suggest the need to balance a safe access environment with mechanisms that promote action

when introducing psychosocial support. Surveys targeting patients with advanced cancer have

also reported the importance of ensuring accessibility to support [35][36], which our findings

confirm. Previous research also suggests the need for flexible support tailored to conditions and

intentions rather than uniform responses for all patients [37]. Our findings indicate the potential

to promote use of support services more effectively by presenting a combination of elements that

trigger actions. This could serve as concrete guidelines for designing behavior-changing

interventions for psychosocial support.

Strengths and limitations

This study involved several limitations. First, because it was a pilot study conducted via a

web☐based questionnaire, selection bias may have occurred. Future research should address this

issue by employing larger, more representative samples and a variety of data☐collection

methods.

Second, although we outlined a clear framework for discussing limitations, practical implications,

and future research directions, our findings may not generalize to all cancer survivors who do not

use psychosocial support services.

Third, our analysis was restricted to the descriptive content provided in the one-time survey

responses, limiting the depth of insight. Complementary qualitative studies—such as in-depth

interviews or ethnographic observations—are needed to enrich these findings.

Fourth, because the Google Form link was shared on third-party platforms without trackable

URLs, we could not perform click-through analysis or calculate a response rate. Future surveys

should use trackable links to obtain precise engagement metrics.

Finally, building on these results, subsequent research should develop and evaluate practical

interventions—such as tailored counseling services and informational tools—to optimize

psychosocial support for cancer survivors.

Conclusion

In this study, we used the FBM to systematically organize the factors that promote the use of

psychosocial support. This has provided insights that may be useful for understanding the initial

design and intervention points for behavioral change in the future. In particular, psychosocial

support is not merely a place to provide information. Based on the identified importance of

motivation and triggers, it will be necessary to focus on enhancing trust and hope through

individualized communication strategies and clearly defined pathways for initiating support for

patients.

Acknowledgments

We would like to express our deep gratitude to the patient organizations and respondents who

cooperated with the recruitment and survey for this study. We also thank Melissa Leffler, MBA,

from Edanz (https://jp.edanz.com/ac) for editing a draft of this manuscript.

References

References

1. Cancer Information Service, National Cancer Center, Japan. Cancer statistics. National Cancer

Registry, Ministry of Health, Labour and Welfare. 2025 July 3

https://ganjoho.jp/reg_stat/statistics/stat/summary.html

2. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics

2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries.

CA: A Cancer Journal for Clinicians . 2024 May 1;74(3):229–63. https://doi.org/10.3322/caac.21834

3. National Cancer Center Japan. Joint research with IARC identifying associations between lifestyle

risk factors and the prognosis of cancer survivors. 2022 August 4 https://www.iarc.who.int/fr/news-

events/iarc-and-the-national-cancer-center-japan-launch-a-joint-long-term-study-lifestyle-factors/

4. Bower JE, Bak K, Berger A, Breitbart W, Escalante CP, Ganz PA, et al. Screening, assessment and

management of fatigue in adult survivors of cancer: an ASCO clinical practice guideline adaptation.

J Clin Oncol. 2014;32(17): 1840-1850. http://doi.org/10.1200/JCO.2013.53.4495

5. Fujisawa D, Inoguchi H, Shimoda H, Yoshiuchi K, Inoue S, Ogawa A, et al. Impact of depression on

health-utility value in cancer patients. Psychooncology. 2016;25(5): 491-495.

http://doi.org/10.1002/pon.3945

6. Costa ALS, Heitkemper MM, Alencar GP, Damiani LP, da Silva RM, Jarrett ME. Social support

predicts lower stress and better quality of life and resilience in Brazilian patients with colorectal

cancer. Cancer Nurs. 2017;40(5): 352-360. http://doi.org/10.1097/NCC.0000000000000388

7. Ruiz-Rodríguez I, Hombrados-Mendieta I, Melguizo-Garín A, Martos-Méndez MJ. The importance

of social support, optimism and resilience on quality of life in cancer patients. Front Psychol.

2022;13: 833176. http://doi.org/10.3389/fpsyg.2022.833176

8. Ministry of Health, Labour and Welfare. Revision of the Basic Plan for the Promotion of Cancer

Control Measures in the "Living with Cancer" field . 2022 October 27

https://www.mhlw.go.jp/content/10901000/001005572.pdf

9.   National Cancer Center Japan. Report on the Patient Experience Survey 2023 (preliminary) 2025

July 3

https://www.ncc.go.jp/jp/icc/policy-evaluation/project/010/2023/R5_all.pdf

10.   Merluzzi TV, Philip EJ, Yang M, Heitzmann CA. Matching of received social support with need for

support in adjusting to cancer and survivorship. Psychooncology. 2016;25(6): 684-690.

http://doi.org/10.1002/pon.3896

11.   Aflac Life Insurance Co. Survey results on concerns and anxieties of cancer patients: one in two

cancer patients are "lost in cancer information". 2022.Oct 19

https://www.aflac.co.jp/news_pdf/20220819.pdf

12.   Fennell K, Hull M, Jones M, Dollman J. Barriers to accessing services for mental and physical

health conditions in rural Australian adults. Rural Remote Health. 2018;18(1): 4155.

http://doi.org/10.22605/RRH4155

13.   Langbecker D, Ekberg S, Yates P. "Don't need help, don't want help, can't get help": why brain-

tumour patients do not use rehabilitation or psychosocial services. Patient Educ Couns. 2017;100(9):

1744-1750. http://doi.org/10.1016/j.pec.2017.04.004

14.   Yigitbas C, Bulut A. Exploring psychological help-seeking behaviours and stigma perception among

cancer patients: impact on psychosocial adjustment. Support Care Cancer. 2024;32(10): 694.

http://doi.org/10.1007/s00520-024-08870-z

15.   Signorelli C, Lim Høeg B, Asuzu C, Centeno I, Estapé T, Fisher P, et al. International survey of

psychosocial care for cancer survivors in low-/middle- and high-income countries. JCO Glob Oncol.

2024;10: e2300418. http://doi.org/10.1200/GO.23.00418

16.   Fogg BJ. A behavior model for persuasive design. In: Proc 4th Int Conf Persuasive Technology.

New York: ACM; 2009;40: 1-7.

17.   http://doi.org/10.1145/1541948.1541999

18.   Fogg BJ. Fogg Behavior Model. Stanford (CA): Behavior Design Lab, Stanford University; 2019.

Technical report.

19.   Feng GH, Zhao KH, Wang YF, Yue QQ, Chen YS, Huang LL, et al. mHealth interventions to

improve liver-cancer screening among high-risk populations: protocol for an RCT. BMC Public

Health. 2024;24(1): 2501. http://doi.org/10.1186/s12889-024-20025-7

20.   Vogel RI, Niendorf K, Petzel S, Lee H, Teoh D, Blaes AH, et al. A mobile-health app to motivate

genetic-counselling use among women with ovarian cancer: pilot RCT. Gynecol Oncol.

2019;153(1): 100-107. http://doi.org/10.1016/j.ygyno.2019.01.019

21.   Montgomery R. Sample-size justification in feasibility studies: moving beyond published guidance.

Pilot Feasibility Stud. 2025;11: 88. http://doi.org/10.1186/s40814-025-01675-9

22.   Julious SA. Sample size of 12 per group rule of thumb for a pilot study. Pharm Stat. 2005;4(4): 287-

291. http://doi.org/10.1002/pst.185

23.   Kunselman AR. A brief overview of pilot studies and their sample-size justification. Fertil Steril.

2024;121(6): 899-901. http://doi.org/10.1016/j.fertnstert.2024.01.040

24.   Charmaz K. Constructing grounded theory: a practical guide through qualitative analysis. London:

Sage; 2006.

25.   Charmaz K. Grounded theory as an emergent method. In: Morse JM, Stern PN, Corbin J, et al.,

editors. Handbook of emergent methods. New York: Guilford Press; 2008. p. 155-170.

26.   Glaser BG. Theoretical sensitivity: advances in the methodology of grounded theory. Mill Valley

(CA): Sociology Press; 1978.

27.   Matsui T, Hirai K, Gondo Y, Sato S. Help-seeking behaviour and psychosocial-support service use

among Japanese cancer patients. Jpn J Clin Oncol. 2020;50(10): 1175-1181.

http://doi.org/10.1093/jjco/hyaa086

28.   Katayama K, Ishikawa D, Miyagi Y, Takemiya S, Okamoto N, Ogawa A. Cancer telephone

consultations: gender differences in counselling needs. Patient Educ Couns. 2020;S0738-

3991(20)30278-0. http://doi.org/10.1016/j.pec.2020.05.012

29. Feng X, Hu Y, Pfaff H, Liu S, Wang H, Wang Y. Emotional or informational? Help-seeking

behaviours in online health communities. Soc Sci Med. 2025;371: 117902.

http://doi.org/10.1016/j.socscimed.2025.117902

30. Faller H, Koch U, Brähler E, Härter M, Keller M, Schulz H, et al. Satisfaction with information and

unmet information needs in men and women with cancer. J Cancer Surviv. 2016;10: 62-70.

http://doi.org/10.1007/s11764-015-0451-1

31. Kruger J, Dunning D. Unskilled and unaware of it: difficulties in recognising one's own

incompetence. J Pers Soc Psychol. 1999;77(6): 1121-1134. http://doi.org/10.1037/0022-

3514.77.6.1121

32. Nutbeam D. Health promotion glossary. Health Promot Int. 1998;13(4): 349-364.

http://doi.org/10.1093/heapro/13.4.349

33. Peng RX, Shen F. Why fall for misinformation? Information-processing strategies, health

consciousness and overconfidence in health literacy. J Health Psychol. 2024;Epub ahead of print.

http://doi.org/10.1177/13591053241273647

34. Pennycook G, Ross RM, Koehler DJ, Fugelsang JA. Dunning–Kruger effects in reasoning:

implications of failing to recognise incompetence. Psychon Bull Rev. 2017;24(6): 1774-1784.

http://doi.org/10.3758/s13423-017-1242-7

35. Yin H, Zha Y, Zhou Y, Tao H, Zhu D. Barriers and facilitators to help-seeking for symptoms in

ovarian-cancer patients in China: a qualitative study. BMJ Open. 2024;14(11): e087602.

http://doi.org/10.1136/bmjopen-2024-087602

36. Newton JC, O'Connor M, Saunders C, Moorin R, Ali S, Nowak AK, et al. Psychosocial support in

people living with advanced cancer: patients' perspectives. Psychooncology. 2021;30(3): 287-295.

http://doi.org/10.1002/pon.5569

37. Sheridan R, McCaughan D, Hewison A, Roman E, Smith A, Patmore R, et al. Experiences and

preferences for psychosocial support in chronic haematological cancers. BMJ Open. 2023;13(8):

e070467. http://doi.org/10.1136/bmjopen-2022-070467

38. Matsui T, Hirai K, Gondo Y, Sato S. Factors contributing to help-seeking for psychosocial support

among Japanese cancer patients: segmentation approach. Psychooncology. 2020;29(11): 1905-1916.

http://doi.org/10.1002/pon.5523