Title: Staff Perspectives on Implementing Opt-Out Blood-Borne Virus Testing in English Emergency Departments: A Qualitative Study

Authors

Siobhán Allison[1], Tom May[1], Jonathan Roberts[2], Rachel Hill-Tout[3], Stephen Hindle[3], Matthew Hickman[1], Lucy Yardley[1], Rachel Todd[1], Robyn Heath[1], Jeremy Horwood[1]

1: University of Bristol, NIHR Health Protection Research Unit
2: University Hospitals Sussex NHS Foundation Trust, Claude Nicol Centre
3: NHS England

Abstract

Background

A significant challenge to achieving global 2030 elimination goals for Blood-Borne Viruses (BBVs) is identifying undiagnosed individuals and relinking those who are no longer in care. To address this, the UK government has implemented opt-out BBV testing in Emergency Departments (EDs) to increase access to BBV testing in high prevalence areas. All adult ED patients having a routine blood test are automatically tested for HIV, hepatitis B and C, unless they opt-out. This study aimed to identify barriers and facilitators to implementation of ED opt-out BBV testing and provide recommendations for future rollouts.

Method

Semi-structured interviews with 23 staff members across five ED sites in very high HIV prevalence areas were analysed thematically, informed by Normalisation Process Theory.

Results

While there was some variation in staff knowledge and understanding of the programme, overall acceptance of the opt-out testing approach was found to be high. Training impacted positively on the staff's understanding of the purpose of the intervention and the correct process, including the opt-out model. High workloads and competing priorities in EDs were a significant barriers to testing. However, there were specific systems and processes that facilitated the uptake of testing, these

included automation and BBV champions. Giving the programme time to embed into practice and ensuring good feedback loops and flexibility to 'tweak' the process was essential to sustaining the programme.

Conclusion

To embed opt-out testing into emergency care, sites should implement automated test ordering, staff training, clear communication, and dedicated champions, which can help to support earlier diagnosis, reduce inequalities and improve patient outcomes.

Abbreviations

BBV Blood Borne Virus (HIV, Hepatitis B and Hepatitis C)
ED   Emergency Department
EPR Electronic Patient Records
HCA Health Care Assistant
HBV Hepatitis C
HCV Hepatitis B
HIV   Human Immunodeficiency Virus
NPT Normalisation Process Theory
SOP Standardised Operating Procedures

Introduction

The UK government has set a goal to reach both national and World Health Organisation (WHO) targets of zero new HIV transmissions and eliminating Hepatitis C (HCV) and Hepatitis B (HBV) as a public health threat by 2030. A key challenge in reaching these targets is identifying individuals who remain undiagnosed. An estimated 4,700 people in England are living with undiagnosed HIV (1) and 76,820 people remain undiagnosed with hepatitis B virus (HBV)(2). In 2024 approximately 40% of HIV diagnoses in England were made at an advanced stage of HIV disease (CD4 count below 350 cells/mm³)(3), which increases the risk of hospitalisation and mortality, reduces life expectancy, and results in higher treatment costs (4, 5). Prompt HIV diagnosis and initiation of antiretroviral therapy reduces viral load, improves life expectancy (6), and prevents onwards transmission(7). Both HBV and HCV can lead to chronic liver disease and cirrhosis and antiviral treatment can cure most cases of HCV (8).

New and innovative approaches are needed to reach undiagnosed populations. Emergency Department (ED) opt-out blood-borne viruses (BBV) testing (the programme) is one such approach. Opt-out BBV testing involves all adult ED patients who are having a blood test as part of their care, are automatically tested for BBVs unless they opt out. Patients are informed about the testing via posters, leaflets, and video screens.

EDs are well-positioned to implement opt-out BBV testing due to their high patient volumes and their ability to reach people underserved by traditional health services(9). These populations often face significant health inequalities, making robust systems for linkage to care essential (10). The opt-out testing model has the potential to reduce the incidence of missed opportunities of diagnosing BBVs (11). The UK government allocated funding for ED opt-out testing in two waves; Wave 1 in April 2022 to 33 sites in areas with very high HIV prevalence in England (defined as >5 per 1000 in the local population) (across London, Manchester, Blackpool, and Brighton) and Wave 2 in April 2024 to 45 sites in areas with high HIV prevalence across England (defined as >2 per 1000 in the local population).

Evidence from previous ED BBV testing implementation pilots have identified barriers to testing, which are attributed to staff not offering the test rather than

patients actively opting out, despite high public acceptability of BBV testing (16, 17). Traditionally, BBV testing focused on patient risk factors (including country of birth, sexual behaviour and history of injecting drug use). Evidence suggests that healthcare workers' assumptions or unconscious biases may, in some cases, affect whether BBV testing is offered to patients in a routine and equitable way, (12, 13). HIV care has traditionally been provided separately from other medical services, with enhanced privacy and confidentiality (14). This approach emerged in response to the significant stigma faced by people living with HIV, a phenomenon often referred to as the 'exceptionalism' of HIV(15). As a result, many general NHS staff have had limited experience treating individuals living with HIV, or accessing current information about care and treatment, which can contribute to outdated and stigmatising views (14) in contrast, universal opt-out testing removes the need for such judgments, helping to normalise BBV testing and reduce stigma(13). The introduction of changes in practices needs to ensure that staff are well supported and aware of the rationale behind the opt-out approach(16).

There is overlap between populations at risk of HIV and HBV and HCV supporting the public health opportunity and importance of combining the three blood-borne virus (BBV) tests in this programme. Early pilots of opt-out testing in emergency departments focused solely on EDs testing for HIV, missing the opportunity to diagnose viral hepatitis. For example, one study, a single week of testing in an ED revealed that 54 viral hepatitis diagnoses would have been missed if testing had been limited to HIV alone (18).

ED presents unique challenges related to testing implementation, uptake, and connecting patients to ongoing care, compared to other screening programmes in antenatal and prisons. The current study aimed to evaluate the implementation by investigating ED staff views and experiences of opt-out ED BBV testing, as well as the facilitators and barriers to implementation, and provide recommendations to sites.

Methods

Study design

The research employed a qualitative design using semi-structured remote interviews with 23 members of staff involved in the delivery of opt-out BBV testing in Wave 1 sites. Interviews were conducted between April and September 2024.

Participants and recruitment

Staff were recruited from five ED sites in areas with very high HIV prevalence. Sites were selected based on their geographical location, organisational practices, and opt-out testing models/processes. Recruitment involved site Principal Investigators (PIs) informing eligible staff members about the project, either by email or face-to-face, and placing recruitment materials in staff areas (e.g. staff rooms). These materials included a QR code linking to a web page with study information and a section for interested staff to log their contact details. A member of the research team later contacted them to arrange a suitable interview time. Purposive sampling was used to select a diverse range of staff based on their role (e.g. ED staff, laboratory staff, care pathway facilitators) and level of experience (including time in post and current NHS band). Analysis was conducted alongside recruitment, and the data was continuously assessed to ensure that the study objectives were being met and 'information power' had been achieved[19].

Data Collection

Interviews were conducted via telephone or video call. A flexible topic guide was developed informed by Normalisation Process Theory (NPT)[20, 21] and included open-ended questions to elicit views on opt-out testing, including its acceptability, perceived organisational barriers, aspects that worked well and areas for improvement. All interviewees were offered a £10 voucher as a thank you.

Theoretical Framework

NPT is an implementation theory used to understand how new interventions are adopted, embedded and sustained in routine healthcare practices. NPT proposes that the implementation of interventions is dependent on the ability of participants to fulfil four criteria presented in figure 1. NPT informed data collection and analysis by

guiding exploration of how staff made sense of, engaged with, operationalised and appraised the programme.(20, 21).

Figure 1: NPT constructs (21)

Data Analysis

With participant consent, interviews were audio recorded, transcribed, anonymised, and uploaded to NVivo V.12 for analysis. Thematic analysis, using a data-driven inductive approach(22), was applied to identify and examine patterns and themes significant to participants and across the dataset. Three transcripts were initially inductively coded independently by two researchers (TM and SA) who and discussed any new codes of potential significance to the research objective. A coding framework, informed inductive codes and deductively by NPT constructs was developed. The coding framework was then applied to the remaining transcripts by SA. Codes were built into broader categories through comparison across transcripts and higher-level recurring themes were developed informed by the NPT constructs. The data and development of themes were discussed by the multidisciplinary research team to ensure credibility and confirmability, with relevance to the research objectives.

Patient & public involvement

The study team participated in the programmes steering group, which included key individuals from UK Health Security Agency (UKHSA), NHS England (NHSE),

academia, community organisations involved with the care of people living with HIV and Hepatitis and people living with HIV and Hepatitis. These meetings provided a platform for the study team to share regular updates on study progress, discuss emerging challenges, and seek input on key aspects of the research. Study documents (including participant information, posters, and topic guides) were sent to all members of this group for comment and feedback.

Ethics

This study obtained Oxford B Research Ethics Committee, HRA and HCRW approval (IRAS PROJECT ID 327922/REC Reference 23/SC/0357).

Findings

Twenty-three interviews were conducted with staff from five sites involved in the delivery of the programme, including one with a senior NHSE stakeholder overseeing the project. Interviews lasted between 14 and 45 minutes (average 30 minutes). Staff represented a range of roles related to the delivery of the programme including ED Health Care Assistants (n=4), ED Nurses (Band 3-8) (n=5), ED Doctor (n=5), Specialist Nurse (HIV or Hepatitis) (n=5), Specialist Doctor (HIV or Hepatitis) (n=1), Laboratory (n=2), NHS Staff (n=1).

Themes developed from the analysis aligned with four main NPT constructs which are used to structure the findings below, alongside verbatim quotes.

Coherence

Purpose of the intervention (Communal specification)

Many staff understood the purpose of the intervention and the programme's benefits to patients and contributions toward public health elimination goals:

> *I think it's really good. Especially from the point of view of hepatitis C, a lot of people have been found and diagnosed through ED that would otherwise potentially never have been tested… because they're not necessarily within a demographic that would be targeted for testing, so I think it's been really positive (Hep C Nurse, Site 2)*

However, there were some individuals who misunderstood the programme's purpose. For example, one healthcare assistant described the reassurance that opt-out testing provided in the context of potential needle stick injuries:

> *For somebody who's continuously in contact with patients and handling sharps, it is comforting to know that if I accidentally end up with a needle stick injury, most of my patients agree to have the test, and then I know that I'm…if I'm going to be okay or if there's any risks (ED HCA, Site 1)*

New concepts (Differentiation)

Most staff appeared to support the introduction of the programme. However, not all fully understood the concept of opt-out testing, including that patients were informed via posters and verbal consent was not required. Patient-facing staff were therefore required to familiarise themselves with new concepts associated with an opt-out model:

> *And there were a couple of people [ED staff] that said not sure if patient consents to it [testing]… which I suppose there's still a question from staff about what is the consent process? I can't remember exactly how it was phrased but it was that they weren't quite sure if proper consent had been done (ED Doctor, Site 1)*

Some staff believed that patients should be informed they were having a BBV test, and that verbal consent should be sought. This was often due to a lack of familiarity with the opt-out model, including the shift away from requiring verbal consent:

> *I'm not doing [BBV test] without consent– you know – because I dunno what's gonna happen. Is my name on that (HCA, Site 1)*

For others, normal practice involved obtaining verbal consent for all blood tests, including BBVs:

> *Oh of course, we always gain consent when – we gain consent for even before getting bloods, we always gain consent if they're happy for us to take the bloods (ED Nurse, Site 3)*

The need to change previously learned behaviours regarding which regular blood tests they would be requesting posed a barrier to the programme's's delivery, particularly among specific staff groups:

> *And that is a really hard thing to do if people, you know, it's changing someone's learnt behaviour that they always just request their normal bloods and go and the sample. Even though it's such a simple extra it's hard to change that little bit of behaviour, especially in doctors I feel (ED Doctor, Site 1)*

Value in the intervention (Internalisation)

As well as practical benefits, staff also recognised the potential value, benefits and importance of implementing the programme. This was often aligned with existing evidence of its effectiveness or benefit to patients:

> We've been told like there are – like we have had people who have been tested, and they've come back positive, and it's changed their lives in the positive, in the fact that they've got, in some cases, early diagnosis (HCA, Site 1)

For ED staff in particular, it was seen as an opportunity to contribute to population-level preventive medicine, an area they felt was often overlooked in emergency care:

> It's something I don't think A&E does enough of, is preventative medicine. (ED Doctor, Site 4)

Cognitive participation

Key individuals (Initiation)

Implementation demanded considerable time and effort to set up from the service level down to the ward level, including IT and laboratory services. Key individuals were central to ensuring the successful implementation of BBV testing. Some sites had formal roles, such as champions. For others, it was carried out as an informal role on top of their existing workload. Both were key in supporting implementation efforts:

> One of the HIV clinical nurse specialists has really taken a lead (HCV Nurse, Site 2)

These key individuals helped to foster participation across different staff groups:

> We've developed, kind of ED opt-out champions in the emergency department. So we've got a handful of healthcare assistants and nurses who were really almost, kind of empowering to role model the whole service and really encourage it … I think, working more with the BBV champions over there will just make the process more streamlined (HCV Nurse, Site 2)

Buy in (Legitimation)

Education and training were key components in ensuring staff engagement and buy-in, as understanding the purpose of the intervention helped staff to engage with it:

> *But once the kind of process and theory behind it was explained, I think that they [staff members] were very much on board (ED Doctor, Site 1)*

Staff discussed how the intervention and training had shifted their perceptions of BBVs. By making the test opt-out, the focus moved away from individual stereotypical risk, leading to diagnoses in individuals who would typically be considered low risk:

> *I can remember sort of older people in resus having heart attacks, and you take their blood and it … you'd find out later that they're HIV positive and it wasn't the ones you'd expect, you know … the young homosexuals or the drug users ... those people stand out as maybe being HIV positive. So I think it's very useful to think that … anybody could be HIV positive (ED Consultant, Site 4)*

Collective action

Performing intervention tasks (Interactional workability)

The intervention required staff to order, collect and send an additional blood sample to the laboratory. In some sites BBV tests were automatically included in the standard blood set, the electronic ordering system (EOS) automatically included the BBV test in all blood orders. However, staff remained responsible for physically collecting the sample. Initial training was valuable in ensuring staff understood the importance of including the test. To maintain uptake - particularly at sites without automated ordering where staff had to manually request the BBV test via the EOS – staff huddles, post-it notes and other verbal and written reminders proved effective:

> *It's also being done [reminder about testing] by word of mouth … and then they continuously updated in our handover in the early morning before we start our shift (ED Nurse, Site 5)*

It is recommended that patients who return to the ED within a 12-month period do not undergo a repeat BBV test unless there is a clinical reason. Ideally, the ordering systems would automatically 'block' the test being ordered. If blocking is not in place, patients who attend the ED regularly may have unnecessary tests, incurring extra cost and time. Not all sites had this mechanism in place, and staff raised this as a concern:

> *And that would also skew your results, because all the tests that you do, like a chunk of them will be in the same patient within a certain time frame ... So, yeah, I think that needs, because I don't think there's any sort of blocking in A&E (ED Doctor, Site 3)*

Some ED staff highlighted the potential positive adjustments to their day-to-day practice, including the removal of the need to risk-assess patients, with the potential biases associated, which helped to simplify their workflow:

> *I'm not having to double-think it and say ooh, is this the type of person that might have it? Cause, for one, I know that there is no specific type of person, and two, I know that, yeah, that's not expected of me, just send it (ED Doctor, Site 1)*

(Confidence in the Opt-Out Model?) Relational integration

The opt-out model relied on the accessibility of the posters explaining the intervention to patients. Some staff had confidence in these and relied on them to inform patients:

> *We just let the posters do the talking (ED Nurse, Site 1)*

Other staff were concerned that patients had not seen or understood these posters, and therefore sought verbal consent from patients for each individual test:

> *They [patients] could have not seen the poster and they don't know they're being tested for HIV, so it could be that we should mention it when we say, 'We're taking blood. We're also testing for HIV (ED Doctor, Site 3)*

Good working relationships and links between the different departments involved (HIV, Hepatitis, laboratories, etc) ensured the programme continued to run smoothly and allowed issues to be resolved quickly:

> *But I think overall, it works well. I think there's good communication between hepatology, sexual health, the labs. We often sort of meet and chat, and sort of like tweak things to see how we can get them to work better. So, I think that yeah, it is working well (HCV Nurse, Site 2)*

Collaboration across specialist teams also provided support and advice during the early stages, which was useful for responding to any modifications to the programme (e.g. integration of HCV testing):

> *When it came to adding in hep C testing … I introduced our HIV and our hep C leads for testing, so that they could work together, because they had so many commonalities. And, actually, that's been brilliant because er it just means that they can work synergistically and save an awful of time (HIV Doctor, Site 2)*

Consistent messaging and training (Skill-set workability)

There were inconsistencies both between and within sites regarding whether training was provided and who received it. This led to staff not having the confidence to speak to patients about follow-up care they may receive if they had a reactive test:

> *None of us actually know what the next steps are for that patient. So we kind of say to them … that's something if you bring that up to your doctor, your doctor will be able to give you more advice on. But us personally don't know the next steps to be able to advise the patients on what the next steps would be if they was to be tested positive (HCA, Site 1)*

Staff were, at times, only given limited information from senior staff about what was required of them to carry out testing:

You were told in passing [about BBV testing] that you know, to see whether they had anything and it was just a screening to catch people that didn't know (ED Doctor, Site 5)

Workforce (give different name? - Workforce and context (Contextual integration)

EDs experienced high volumes of patients, and staff experienced high workloads, which impacted delivery of BBV testing. Sites reported high staff turnover alongside shift work, with some staff only working night shifts. This led to some staff lacking accurate information about opt-out testing:

It's not going well with doctors because we only see new doctors. When we have locums doctors … they don't know. So I think the doctor needs to have more knowledge about this. Because every week.. someone came to me, what is red for? Oh it's for HIV. Oh, so you have to do it for everyone? Yes you have to do it for everyone? You don't have to ask just do it … so I'm sure most of them they don't really … know about this (HCA, Site 2)

There were some initial concerns about the additional workload associated with introducing another blood test:

Initially, it kind of felt a bit like that [extra work] because it was another label, another blood bottle … But now it's just like I said, automatic (Nurse, Site 4)

Reflexive monitoring

Feedback and impact stories (Systematisation)

When sites provided staff with feedback on BBV tests conducted, new diagnoses, and personal stories, staff found these messages engaging and felt motivated by the positive impact of the tests:

I think it's the patient stories that were more convincing, rather than the numbers, to sort of say here's someone actually who came here, you know, actually a new diagnosis who was started on treatment that never, you know, wouldn't have known without this project, I think that's quite, quite a powerful teaching tool (ED Doctor, Site 1)

Staff reflections (Communal appraisal)

Staff reflected on the effectiveness of the programme and offered recommendations to enhance future rollouts. Key suggestions included comprehensive training, ensuring staff are well-informed about follow-up care and able to address patients concerns and providing regular feedback including patient stories:

> *Yeah, so the training, have somebody, you know, to go through the training with you. But also have somebody who's living with the virus, do a talk and meet some of the staff (HCA, Site 4)*

Others suggested having a champion role, collaborating with experienced sites, fostering teamwork and engagement:

> *I guess what would be really beneficial is linking with other sites that are already doing it, so that you can almost kind of – almost like get a bit of like clinical supervision sessions to, to discuss how it's working, if there are any improvements and, like, buddying up (HCV Nurse, Site 2)*

Overall, staff encouraged new sites to integrate the opt-out testing approach into routine practice:

> *Just do it. You know you're gonna get reluctance by staff. You're gonna get reluctance by patients so just do it. Just do it and just get it over with and it'll become be a routine, like routine so we just crack on. It's automatically done so yes. (ED nurse, Site 1)*

Staff reflected that there was potential to achieve more through the programme. Stigma and lack of knowledge amongst patients were highlighted as an issue. Staff considered if more could be done in educating patients and increasing understanding of BBVs. One participant suggested providing patients with information about BBVs:

> *I think if we come with some, leaflets, to give to them when they're leaving, after the blood test... they can read about HIV, why it's so important having bloods, and then why is the treatment. So I think if we give them to the patient*

> *at the end... they will be more knowledgeable about what we're doing (HCA, Site 2)*

However, this did not seem feasible to some staff. Even though they recognised the benefits, they questioned whether ED was the right environment:

> *I don't know whether A&E is the best place to give education. You know, we have lots of patient leaflets that we hand out and, err, you know, but, but you walk outside the hospital and the bin's full of them (ED Consultant, Site 4)*

Others suggested other forms of health promotion ideas videos using TV screens in the ED and other forms of communication to increase awareness of BBVs and the programme:

> *TV running with, with, some up to date information and maybe some links attached to that (ED Consultant, Site 4)*

Personal Sense of Accomplishment and Purpose (Individual appraisal)

Overall, being part of the programme for some individuals was important and gave them a sense of accomplishment:

> *It's a sense of achievement because that's what we're trying to do, we're trying to reach elimination (HCV Nurse, Site 2)*

Initiating change (Reconfiguration)

ED staff took ownership of identifying issues and making changes to improve the intervention. For example, a HCA identified a problem with staff forgetting to fill the second yellow-top vial and described modifications their site had introduced to address this:

> *With the two yellow you used to get confused before. People, especially the doctors, they used to send one and they thought it's including HIV in the other bloods. So we, we, so yes … so I said, um, I said, can we change, can we do something here so we can void, so they can see one different colour. Which the red I think it's more … um, yes it's red, it's more obvious (HCA, site 1)*

Table 1. Recommendations for Sites

Implementation Planning	1. Foster early connections between hospital management, EDs, labs, HIV and Hepatitis services, IT teams and clinical governance to streamline workflows and resolve issues efficiently. 2. Leverage national and regional NHSE networks by having a designated staff member attend meetings and sharing insights locally. 3. Engage staff early with regular updates, including testing figures and impactful case studies, to build understanding and sustain involvement. 4. Identify and support staff champions to drive momentum and promote programme adoption.
Staff Training and Feedback	5. Provide accessible, practical and knowledge-based training for all staff groups that includes content on the Programme's aims, local processes for test delivery, IT systems, care pathways for reactive results and up-to-date virus knowledge and stigma reduction. 6. Incorporate lived-experience stories in training to enhance staff understanding of the Programme's impact.
Delivery of Testing	7. Share practical solutions to barriers from existing sites, such as modified vials and visual prompts, to improve test coverage (1). 8. Use simple scripts to reinforce the opt-out process and address staff concerns about patient communication.
Reflection	9. Allow time for the programme to embed, enabling staff to adapt to new workflows. 10. Maintain ongoing communication about progress (including numbers of people detected, cost-effectiveness and contributions toward elimination plans) and provide opportunities for staff to share feedback to support long-term sustainability.

Discussion

This study builds upon existing evidence on the acceptability and effectiveness of ED opt-out testing by identifying qualitative insights into the implementation of opt-out testing in EDs(10, 13, 16, 18, 23-28). Findings identified general support and acceptance of BBV testing among staff, despite initial concerns regarding its delivery but also some variation in how staff understood and conceptualised opt-out testing. Staff reported how, over time, training and key individuals helped foster staff buy-in, and the programme became routine once fully implemented and established in their EDs. Collaboration across teams also played a key role in refining the programme and sustaining staff engagement. Regular feedback on results and impact stories promoted staff reflection on the impact of the programme and helped to maintain staff engagement. This study provides further context to the interim data from the Public Health Evaluation of the programme, which indicates promising results for the programme, (29) by showing the acceptability for the programme amongst staff and what barriers may need to be addressed to increase testing.

Overall, acceptance of the opt-out testing approach was found to be high, which aligns with existing evidence on staff acceptability of opt-out testing in EDs with staff particular focus on the value to individuals health(13, 28). Nevertheless, this study identified how the intervention required a change in staff practices as the opt-out model differed from previous blood taking routines. This demonstrates how fostering a strong understanding of the programme among staff through training and regular staff meetings, not only enhances adherence to the model but helps to normalise the process of additional BBV testing(28).

The exceptionalism of HIV testing remains a barrier to testing (16, 24, 30). Some staff in this study still viewed HIV testing as 'different' to other blood tests they take, with the view that explicit consent was needed(16). (24). This ambiguity is echoed in the literature (23), but as the opt-out model becomes more established, staff confidence is expected to grow (25). Training impacted positively on the staff's understanding of the purpose of the intervention and the correct process, including the way patients were informed of the opt-out test. Providing staff with clear, concise language to inform patients particularly in situations where patient awareness is

uncertain may help staff communicate effectively without the need for lengthy consent discussions. However, one study found significant demographic disparities in testing rates when staff used reminders, with testing rates lower in groups that were at higher risk of BBVs (23). To address this issue, training should inform staff about situations (e.g. visual impairment or language barriers) in which the reminders may be helpful and appropriate and when reminders are not needed. It is also essential for sites to ensure that posters and other visual material informing patients about opt-out BBV testing are on display in prominent positions throughout the ED and not lost among other posters(16). This will not only ensure patients are informed but also reassure staff that patients are informed. Sounds good (sorry cant add comments so delete!)

High workloads and competing priorities in EDs were a significant barrier to testing (23, 28). However, there were specific systems and processes that facilitated the uptake of testing. Automation was found to support higher testing uptake, aligning with findings from previous studies (13, 16, 23, 31). By removing the need for staff to manually order tests, automation helped to streamline the process and reduced reliance on individual discretion. One study found that test uptake doubled when automation was introduced, compared to when nurses were required to order tests manually (32). Automation also helped overcome barriers linked to stigma and risk profiling (13) ,contributing to the normalisation of BBV testing. By embedding testing into routine care, automation can help to reduce the influence of staff biases in deciding who should be tested (23).

Healthcare staff stigma towards people living with HIV remains an issue today (33). This programme has the potential to shift perspective and reduce stigma by normalising BBV testing and increasing knowledge through the training provided in the programme. Research has shown that education, including stigma reduction messaging, is a powerful tool for reducing HIV stigma which in turn would engage staff in the intervention (13, 23, 33).

Giving the intervention time to embed into practice and ensuring good feedback loops and the flexibility to modify programme components was essential to sustaining the programme. Champions and key individuals appeared to play an

important role in maintaining staff motivation in the programme. These roles were at times 'official' roles, but often relied on individual's willingness and personal interest to drive the intervention. Key individuals facilitated good communication between staff and created pathways for staff to provide feedback when issues arose, which facilitated adapting the intervention (e.g. a change in blood bottles). Data shows that although sites in the study were not consistently reaching 90% testing targets, staff nonetheless reported that they felt testing was embedded in practice (29). This was in contrast to staff in one study who viewed the test as a 'disruption'(28). Time may be a factor here, as interviews in that study were conducted early in the implementation phase, whereas some sites in the current study had been testing for over two years, highlighting the importance of allowing sufficient time for the intervention to embed (28).

Sites aiming to implement or scale up similar programmes should consider investing in automated test ordering, staff training, clear communication tools and dedicated individuals or champions who can maintain staff momentum and motivation (see Table 1). This can support the implementation of opt-out testing as an accepted part of emergency care that facilitates earlier diagnosis, reduces inequalities and ultimately improves patient outcomes.

Strengths and limitations

The sample in this study included a wide range of staff from HCAs to NHSE staff providing a varied and diverse view on implementation of the programme. The data was collected in 5 of the 32 sites, selected purposively to reflect geographic location, organizational practices and opt-out testing processes, and only evaluated staff views of implementing the programme within ED. The use of NPT allowed for examination of issues with both the design of the intervention and its implementation. Interview data indicated that the NPT constructs of coherence and cognitive participation were well fulfilled for staff with positive consequences for implementation. Some findings may be site-specific and shaped by local characteristics or the length of time the programme had been running. It did not

explore patients views and linkage to care, these will be explored in more detail in the wave 2 evaluation.

Acknowledgments

This research was funded by NHS England and the National Institute for Health Research (NIHR), with support from NIHR Health Protection Research Unit (HPRU) in Behavioural Science and Evaluation (NIHR200877). The views expressed in this article are those of the authors and not necessarily those of NHS England, the NIHR or the Department of Health and Social Care.

Competing Interests

RHT has received sponsorship from Gilead Sciences for presenting on ED opt-out BBV testing. JR has received sponsorship from VIV Health and Gilead Sciences for presenting on ED opt-out BBV testing.
JH has received a grant from Gilead for another study. Gilead Sciences and VIV Health manufacture HIV and hepatitis medications but have not funded ED opt-out BBV testing.
RHT, SH, and JR are all employed by the NHS, which has funded this project

References

1. Schoemig V, Martin V, Shah A, Okumu-Camerra K, Mackay N, Bera S, et al. HIV Action Plan monitoring and evaluation framework 2024 report: Report summarising progress from 2019 to 2023. London: UK Health Security Agency; 2024.

2. Allen E, Balogun K, Burton S, Benson T, Cazachorowski M, Clarke E, et al. Hepatitis B in England: Working to eliminate hepatitis B as a public health threat. UK Health Security Agency; 2023.

3. Kitt H, Shah A, Chau C, Okumu-Camerra K, Bera S, Schoemig V, et al. HIV testing, PrEP, new HIV diagnoses and care outcomes for people accessing HIV services: 2024 report. The annual official statistics data release. London: UK Health Security Agency; 2024.

4. May M, Gompels M, Delpech V, Porter K, Post F, Johnson M, et al. Impact of late diagnosis and treatment on life expectancy in people with HIV-1: UK Collaborative HIV Cohort (UK CHIC) Study. BMJ. 2011;343:d6016.

5. Beck EJ, Mandalia S, Sangha R, Sharott P, Youle M, Baily G, et al. The Cost-Effectiveness of Early Access to HIV Services and Starting cART in the UK 1996–2008. PLOS ONE. 2011;6(12):e27830.

6. Trickey A, Sabin CA, Burkholder G, Crane H, d'Arminio Monforte A, Egger M, et al. Life expectancy after 2015 of adults with HIV on long-term antiretroviral therapy in Europe and North America: a collaborative analysis of cohort studies. Lancet HIV. 2023;10(5):e295-e307.

7. Rodger AJ, Cambiano V, Bruun T, Vernazza P, Collins S, Degen O, et al. Risk of HIV transmission through condomless sex in serodifferent gay couples with the HIV-positive partner taking suppressive antiretroviral therapy (PARTNER): final results of a multicentre, prospective, observational study. The Lancet. 2019;393(10189):2428-38.

8. Fathi H, Clark A, Hill NR, Dusheiko G. Effectiveness of current and future regimens for treating genotype 3 hepatitis C virus infection: a large-scale systematic review. BMC Infectious Diseases. 2017;17(1):722.

9. NHS England. Blood-borne viruses opt-out testing in emergency departments in areas of high and extremely high HIV prevalence: good practice guidance. London NHSE; 2024.

10. Parry S, Bundle N, Ullah S, Foster GR, Ahmad K, Tong CYW, et al. Implementing routine blood-borne virus testing for HCV, HBV and HIV at a London Emergency Department - uncovering the iceberg? Epidemiol Infect. 2018;146(8):1026-35.

11. Guess S, Gormley MA, Moschella P, Roth P, Litwin AH. Missed opportunities for diagnosis of HIV in the emergency department using non-risk-based testing strategy. J Am Coll Emerg Physicians Open. 2023;4(1):e12898.

12. Iott BE, Loveluck J, Benton A, Golson L, Kahle E, Lam J, et al. The impact of stigma on HIV testing decisions for gay, bisexual, queer and other men who have sex with men: a qualitative study. BMC Public Health. 2022;22(1):471.

13. Simmons R, Plunkett J, Cieply L, Ijaz S, Desai M, Mandal S. Blood-borne virus testing in emergency departments - a systematic review of seroprevalence, feasibility, acceptability and linkage to care. HIV Med. 2023;24(1):6-26.

14. Rutland E, Foley E, O'Mahony C, Miller RF, Maw R, Kell P, et al. How normalised is HIV care in the UK? A survey of current practice and opinion. Sex Transm Infect. 2007;83(2):151-4.

15. Kagan D, Seear K, Lenton E, Farrugia A, Valentine K, Mulcahy S, et al. The trouble with normalisation: Transformations to hepatitis C health care and stigma in an era of viral elimination. Sociol Health Illn. 2023;45(7):1421-40.

16. Blakey EG, Fairhead CEL, Rodger AJ, Burns FM, Ralph L, Chadwick DR. Clinical staff attitudes towards opt-out consent for blood-borne virus screening in emergency departments in England. HIV Med. 2025;26(2):326-32.

17. Glew S, Pollard A, Hughes L, Llewellyn C. Public attitudes towards opt-out testing for HIV in primary care: a qualitative study. Br J Gen Pract. 2014;64(619):e60-6.

18. Orkin C, Flanagan S, Wallis E, Ireland G, Dhairyawan R, Fox J, et al. Incorporating HIV/hepatitis B virus/hepatitis C virus combined testing into routine blood tests in nine UK Emergency Departments: the "Going Viral" campaign. HIV Medicine. 2016;17(3):222-30.

19. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies: Guided by Information Power. Qual Health Res. 2016;26(13):1753-60.

20. May CR, Mair F, Finch T, MacFarlane A, Dowrick C, Treweek S, et al. Development of a theory of implementation and integration: Normalization Process Theory. Implement Sci. 2009;4(1):29.

21.   Murray E, Treweek S, Pope C, MacFarlane A, Ballini L, Dowrick C, et al. Normalisation process theory: a framework for developing, evaluating and implementing complex interventions. BMC Med. 2010;8(1):63.

22.   Braun V, Clarke V. Toward good practice in thematic analysis: Avoiding common problems and be(com)ing a knowing researcher. Int J Transgend Health. 2023;24(1):1-6.

23.   Fairhead C, Barber TJ, Okhai H, Durkin R, Hart J, Pinto J, et al. Demographic disparities in blood-borne-virus screening in two London Emergency Departments: a case for implied consent. AIDS Care. 2025;37(5):855-63.

24.   Chadwick DR, Page E, Wilkinson D, Savulescu J. Implied consent for HIV testing in the UK: time for a new approach? Lancet HIV. 2022;9(1).

25.   Crane D, Henderson EJ, Chadwick DR. Exploring the acceptability of a 'limited patient consent procedure' for a proposed blood-borne virus screening programme: a Delphi consensus building technique. BMJ Open. 2017;7(5).

26.   O'Connell S, Lillis D, Cotter A, O'Dea S, Tuite H, Fleming C, et al. Opt-Out Panel Testing for HIV, Hepatitis B and Hepatitis C in an Urban Emergency Department: A Pilot Study. PLoS One. 2016;11(3):e0150546.

27.   Dhairyawan R, O'Connell R, Flanagan S, Wallis E, Orkin C. Linkage to care after routine HIV, hepatitis B & C testing in the emergency department: the 'Going Viral' campaign. Sex Transm Infect. 2016;92(7):557.

28.   Cullen L, Grenfell P, Rodger A, Orkin C, Mandal S, Rhodes T. 'Just another vial...': a qualitative study to explore the acceptability and feasibility of routine blood-borne virus testing in an emergency department setting in the UK. BMJ Open. 2019;9(4):e024085.

29.   UK Health Security Agency. ED BBV opt-out testing 24-month interim report 2024. London: UKHSA 2024.

30.   British HIV Association, British Infection Association, Royal College of Emergency Medicine Joint Working Group. Rapid guidance on opt-out blood-borne virus testing in high-prevalence and extremely high prevalence acute medical settings and emergency department. London 2024.

31.   Smout E, Phyu K, Hughes GJ, Parker L, Rezai R, Evans A, et al. Real-world clinical effectiveness and sustainability of universal bloodborne virus testing in an urban emergency department in the UK. Sci Rep. 2022;12(1):19257.

32.   White DAE, Todorovic T, Petti ML, Ellis KH, Anderson ES. A Comparative Effectiveness Study of Two Nontargeted HIV and Hepatitis C Virus Screening Algorithms in an Urban Emergency Department. Ann Emerg Med. 2018;72(4):438-48.

33.   European Centre for Disease Prevention and Control. HIV stigma in the healthcare setting. Monitoring implementation of the Dublin Declaration on partnership to fight HIV/AIDS in Europe and Central Asia. Stockholm; 2024.