Assessing the repurposing potential of disease-modifying antirheumatic drug targets to reduce Alzheimer's disease risk: a pQTL-based Mendelian randomization and colocalization analysis

Christina N. Kushnir[1], Victoria Taylor-Bateman[1], Neil M. Davies[1,2,3], Emma L. Anderson[1]

Affiliations

[1] Division of Psychiatry, University College London, London WC1E 6BT, UK.

[2] Department of Statistical Science, University College London, London WC1E 6BT, UK.

[3] Department of Public Health and Nursing, Norwegian University of Science and Technology, Norway.

Corresponding author

Christina N. Kushnir MSc

christina.kushnir.23@alumni.ucl.ac.uk

Key words: Drug target Mendelian randomization, DMARD, inflammation, Alzheimer's disease, dementia

Abstract

Background: Current literature has implicated systemic inflammation in the pathogenesis of Alzheimer's disease (AD). However, the viability of anti-inflammatory drug targets as repurposed treatment candidates for AD remains unclear. We utilized two-sample Mendelian randomization (MR) and colocalization analysis to investigate whether disease-modifying antirheumatic drug (DMARD) targets reduce AD risk.

Methods: We investigated 9 DMARD targets, using blood protein quantitative trait loci (pQTLs) from the UK Biobank Pharma Proteomics Project (n = 54,219). The associations of these variants and AD were extracted from the International Genomics of Alzheimer's Project ($n$cases = 21,982, $n$controls = 41,944). We used two-sample MR analyses to estimate the effects of perturbing DMARD targets on AD risk. We also examined the colocalization of target pQTLs and AD signals in the *cis*-regions of target-encoding genes.

Results: Our MR estimates suggested that a 1 SD increase in plasma protein concentration of the etanercept target, *FCGR3B*, increased the risk of AD by 10% (OR 1.10; 95% CI [1.02, 1.19]; p = 0.01). We found little evidence that the remaining DMARD targets reduced AD risk. Further colocalization analysis indicated no evidence of colocalization between encoding genes of DMARD targets and AD, including *FCGR3B*. Though, there was suggestive evidence of a non-colocalized association between *TNF* and AD on chromosome 6 via distinct causal variants (H_3 = 46.5%).

Conclusions: Our findings suggest that the majority of analyzed DMARD targets were unlikely to reduce AD risk, except for *FCGR3B*. Future research in AD drug targeting and repurposing should examine the therapeutic viability of drugs which lower *FCGR3B* protein levels in blood plasma.

Background

Over 50 million people worldwide suffer from dementia, with this number expected to increase nearly threefold by the mid-21st century [1]. Alzheimer's disease (AD), the most common cause of dementia and a leading cause of death in the United States and the United Kingdom, imposes both immense pain on patients and their loved ones as well as a significant socioeconomic burden on healthcare systems, making it imperative to find treatment candidates that do not simply address quality of life but, more urgently, target pathological mechanisms that may prevent or halt disease progression [2, 3].

In recent years, chronic and systemic inflammation has been recognized as a possible modifiable risk factor for AD. Elevated concentrations of cytokines in the central nervous system (CNS) lead to neuronal toxicity and tissue damage which facilitate the susceptibility and development of AD [4, 5]. Additional dysregulation of key pro-inflammatory mediators, such as tumor necrosis factor (TNF-α) and interleukin-6 (IL-6), has been connected to hallmark molecular mechanisms contributing to the pathogenesis of AD [4, 5, 6, 7, 8]. Furthermore, genome-wide association studies (GWAS) have substantiated the role of immune function and inflammation in AD. The largest and most recent AD GWAS to date, conducted by Bellenguez et al. (2022), identified multiple genome-wide significant loci near genes involved in innate immunity and microglial function, such as *TREM2*, *PLCG2*, and *ABI3*, as well as loci near *IL-34*, *TNIP1*, and *GRN*, all of which have roles in immune regulation or inflammatory signaling [9]. These findings provide converging genetic evidence that dysregulated immune responses may contribute to AD pathogenesis, highlighting inflammatory pathways as promising therapeutic targets.

Disease-modifying antirheumatic drugs (DMARDs), a class of anti-inflammatory medications that modulate both cytokine activity and other pro-inflammatory targets, present as potentially viable

drug repurposing candidates for AD prevention and treatment. Importantly, the disease-modifying component of DMARDs makes them encouraging therapeutic candidates to counter the progressive neurodegeneration that characterizes AD. However, observational studies examining the association between prescribed DMARD usage and the risk of AD have yielded mixed and contradictory results [10, 11, 12, 13, 14, 15]. Furthermore, such studies investigate sample populations already impacted by autoimmune and/or rheumatic diseases, making confounding by indication highly likely. While randomized control trials (RCTs) may help clarify the existing research, only two RCTs have investigated the casual effects of DMARDs on the risk of AD, both with small sample sizes (n = 41, n = 15) [16, 17]. These RCTs examined the use of etanercept, a commonly prescribed DMARD, over the course of 6 months with results from both trials suggesting positive trends in favor of a protective effect on AD risk. However, the lack of statistical power limits inference from these studies.

Mendelian randomization (MR) is a gene-based instrumental variable approach that can help to minimize confounding and reverse causation—which are pervasive in observational studies—without the additional logistic and financial constraints of clinical trials [18]. Due to the randomized nature of genetic inheritance, MR can assess the causal effects of risk factors via instrumentation of genetic variants [19]. Additionally, MR can be specifically employed to study the effects of perturbing drug targets [19]. Drug target MR estimates the effects of changing protein levels or expression under the rationale that these traits are likely to mimic the consequences of perturbing these proteins by drugs known to target these mechanisms. The genetic variants are selected based on their association with the drug targets of interest (Fig. 1). Colocalization often accompanies MR studies as a form of sensitivity analysis to reduce the threat that linkage disequilibrium (LD) poses to the assumptions of MR. As neither MR nor colocalization are alone sufficient in establishing causality, the integration of both methods may provide strength to studies in statistical genetics and drug repurposing. Here, we assess whether select DMARD targets are likely to be efficacious therapeutic targets for reducing AD risk, using two-sample MR and colocalization.

Fig. 1. Diagrammatic illustration of the study. MR assumes that the genetic variant (1) is robustly associated with the exposure variable (relevance); (2) has no shared confounders with the outcome (independence); and (3) affects the outcome only through the exposure (exclusion restriction) [20]. Abbreviations: DMARD, disease-modifying antirheumatic drug; *TNF*, tumor necrosis factor; BMI, body mass index

Methods

Selection of DMARD targets

11 DMARDs, totaling 33 target genes, were initially selected for analysis. The rationale for inclusion was based on findings from the current literature regarding associations between DMARD usage and incidence rates of dementia or AD [10, 11, 12, 13, 14, 15]. The drugs identified were methotrexate, sulfasalazine, hydroxychloroquine, leflunomide, adalimumab, etanercept, infliximab, certolizumab, golimumab, tocilizumab, and tofacitinib. Methotrexate and tofacitinib had no available pQTL data and were therefore excluded from analysis. In later stages, the pQTL data for the leflunomide target, *DHODH*, did not meet the required significance threshold; as a result, leflunomide was also excluded from analysis. DMARDs and their corresponding encoding gene targets were identified using the DrugBank database (https://go.drugbank.com) and are listed in Table 1. Note, this table does not

encapsulate all targets of the respective DMARDs but rather the targets with available pQTL data.

Additionally, some targets, such as *TNF*, are shared across multiple drugs in our analysis.

Table 1: Selected DMARDs and their analyzed targets

Drug name	Drug target
Sulfasalazine	*PLA2G1B*
	NFKB1
Hydroxychloroquine	*ACE2*
Etanercept	*LTA*
	FCGR2A
	FCGR2B
	FCGR3B
	TNF
Adalimumab	*TNF*
Infliximab	*TNF*
Certolizumab	*TNF*
Golimumab	*TNF*
Tocilizumab	*IL6R*

pQTL instrument selection

Exposure data for each selected DMARD target were sourced from a large-scale proteogenomic

analysis as part of the UK Biobank Pharma Proteomics Project (UKB-PPP, *n* = 54,219), which provides

detailed mapping of 2,923 proteins and their associations with genetic variants within blood plasma

[21]. We searched for *cis*-acting single nucleotide polymorphisms (SNPs) known to be associated

with circulating protein levels (pQTLs) of encoded DMARD targets (Fig. 2). Lead pQTLs were

identified to proxy the effects of select DMARD target perturbations, based on independence and

strength of association (meeting the genome-wide significance threshold of $p < 5 \times 10^{-8}$). The

following selection criteria were applied for instrumentation: 1) pQTLs within ± 1000 kb of the gene

region (*cis*-acting); and 2) a genome-wide significance threshold of $p < 5 \times 10^{-8}$.

Fig. 2. Flow diagram of the study design. Abbreviations: DMARDs, disease-modifying antirheumatic drugs; UKB-PPP, UK

Biobank Pharma Proteomics Project; pQTL, protein quantitative trait loci; AD, Alzheimer's disease; IGAP, International

Genomics of Alzheimer's Project; GWAS, genome-wide association study; MR, Mendelian randomization

Outcome data

Effect estimates of instrumented pQTLs were extracted from a genome-wide association meta-

analysis of clinically diagnosed AD cases within the International Genomics of Alzheimer's Project

(IGAP) [22]. Stage 1 of the GWAS included 21,982 cases and 41,944 controls, all of whom were of

European ancestry. Risk of late-onset AD (LOAD) was defined as symptom onset at ≥ 65 years of age.

Diagnostic assessment occurred either in clinical evaluation or postmortem autopsy. The mean age

at onset for AD cases ranged from 71.1 to 82.6 years, and mean age at examination (or last follow-

up) for healthy controls ranged from 51.0 to 78.9 years. Further information on quality control can

be found in the original publication.

Statistical analyses

Mendelian randomization analysis

MR analyses were conducted to examine the individual effects of each drug target on the risk of AD.

Due to the outcome GWAS comprising only autosomal data, we did not include SNPs located within

the X chromosome during analysis. To reduce impacts from LD, the instrumented pQTLs underwent

a clumping procedure with a threshold of $r^2 = 0.001$ and a 10,000 kb window to ensure independent

variants. The exposure data was coded using the most recent genome reference consortium human

build 38 (GRCh38), assembly Hg20, while the IGAP GWAS outcome data used GRCh37 with assembly

Hg19. Thus, any instrument pQTLs not found in the outcome dataset were replaced by proxy

variants in high LD from the 1000 Genomes European reference panel ($r^2 > 0.8$). During data

harmonization, palindromic SNPs were excluded if the minor allele frequency was > 0.4. MR was

performed using the inverse variance weighted (IVW) method. Upon completion of analysis, odds

ratios (ORs) and 95% confidence intervals (CIs) for AD outcomes were presented per 1 standard

deviation (SD) increase in blood plasma protein concentration.

Colocalization analysis

Colocalization was performed on *cis*-regions of the encoding target genes of interest (± 500 kb

window) to assess if identified associations with AD resulted from shared causal variants. *ACE2* was

excluded from this portion of the analysis as it is located on the X chromosome and could not be

matched with autosomal outcome data. The following priors were set: SNPs within the default

window and exclusively associated with trait 1 had a probability of $p_1 = 10^{-4}$; SNPs within the default window and exclusively associated with trait 2 had a probability of $p_2 = 10^{-4}$; and SNPs within the default window and associated with both traits 1 and 2 had a probability of $p_{12} = 10^{-5}$. A posterior probability for shared causal variants (PP.H$_4$ ≥ 80%) was considered to demonstrate strong evidence of colocalization. A threshold of PP.H$_4$ ≥ 50% was considered suggestive evidence.

Results

Effects of DMARD targets on the risk of AD

Overall, there was little evidence in our study that DMARD targets affected the risk of AD (Fig. 3). There was little evidence that the sulfasalazine targets, *PLA2G1B* (OR 0.99; 95% CI [0.91, 1.08]; *p* = 0.80) and *NFKB1* (OR 0.89; 95% CI [0.74, 1.07]; *p* = 0.22), affected AD risk. There was little evidence for the hydroxychloroquine target, *ACE2* (OR 1.10; 95% CI [0.97, 1.24]; *p* = 0.14). There was additionally little evidence for the targets of tocilizumab, the TNF inhibitors, and etanercept, including *LTA* (OR 1.03; 95% CI [0.95, 1.12]; *p* = 0.43), *FCGR2A* (OR 1.02; 95% CI [0.94, 1.10]; *p* = 0.63), *FCGR2B* (OR 0.99; 95% CI [0.87, 1.13]; *p* = 0.86), *TNF* (OR 0.92; 95% CI [0.72, 1.16]; *p* = 0.47), and *IL6R* (OR 0.97; 95% CI [0.90, 1.03]; *p* = 0.28). However, there was evidence that increasing plasma levels of *FCGR3B,* an etanercept target, increased the risk of AD (OR 1.10; 95% CI [1.02, 1.19]; *p* = 0.01).

DMARD target	*n* SNPs	Alzheimer's disease	OR (95%CI)	*p* value
ACE2	16		1.10 (0.97, 1.24)	0.14
FCGR2A	15		1.02 (0.94, 1.10)	0.63
FCGR2B	7		0.99 (0.87, 1.13)	0.86
FCGR3B	23		1.10 (1.02, 1.19)	0.01
IL6R	12		0.97 (0.90, 1.03)	0.28
LTA	18		1.03 (0.95, 1.12)	0.43
NFKB1	7		0.89 (0.74, 1.07)	0.22
PLA2G1B	32		0.99 (0.91, 1.08)	0.80
TNF	5		0.92 (0.72, 1.16)	0.47

Odds ratio per 1 SD unit increase in blood plasma concentration

Fig. 3. Estimates of the effects of selected DMARD targets on the risk of developing AD. Error bars correspond to the 95% confidence intervals. Abbreviations: DMARD, disease-modifying antirheumatic drug; SNP, single nucleotide polymorphism; OR, odds ratio; SD, standard deviation

Results of colocalization analysis

Colocalization analyses were consistent with the MR analyses above. There was little evidence of colocalization between any of the targets and AD (PP.H_4 < 10% across all proteins). Despite evidence of a causal effect in the main MR analysis, *FCGR3B* protein and AD did not colocalize in the flanked chromosome 1 region of *FCGR3B* (PP.H_4 = 6.21%). There was also very little evidence of pleiotropy (PP.H_3 = 10.3%). However, there was some evidence of a non-colocalized association between *TNF* protein and AD, potentially through pleiotropic causal variants (PP.H_3 = 46.5%) (Fig. 4).

Fig. 4. Regional genomic plots from colocalization analysis of the flanked *TNF* region on chromosome 6. The LD reference

SNP (rs2229094) was used for plotting. Both traits (*TNF*, AD) had to be compared separately because of the differing builds,

GrCH37/38. No pQTLs in the *TNF* region met the threshold for colocalization with AD.

Discussion

We utilized *cis*-pQTL and GWAS summary statistics to evaluate the causal effects of genetically

perturbed DMARD targets on the risk of AD. To our knowledge, this is the first study to specifically

examine DMARD targets and AD risk using drug target MR and colocalization.

We found limited evidence to support the repurposing potential of select DMARD targets for

reduced AD risk. The MR analyses provided little evidence that 8 out of the 9 selected targets—

ACE2, *FCGR2A*, *FCGR2B*, *IL6R*, *LTA*, *NFKB1*, *PLA2G1B*, and *TNF*—affected the risk of AD. With MR, we

found that a per 1 SD unit increase in plasma *FCGR3B* concentration leads to a 10% greater risk of AD. Though, further analysis of genetic signaling in the flanked region on chromosome 1 revealed that *FCGR3B* did not colocalize with AD and are thus unlikely to share a single causal variant. A possible explanation for this discordance is that the findings from MR may have been driven by distinct variants in LD. Alternatively, the lack of evidence for colocalization may be due to tissue specificity. In support of this, a recent study found evidence of colocalization between *FCGR3B* protein, located in cerebrospinal fluid (CSF), and AD signals [23]. Interestingly, this study observed the opposite direction of effect for *FCGR3B*, with their MR analysis revealing an association between increased CSF *FCGR3B* concentration and a lower risk of AD—although it should be noted that their instruments for *FCGR3B* did not reach genome-wide significance. CSF protein composition is more diverse than that of plasma and may perform as a more accurate biomarker for neurodegenerative diseases [24]. However, plasma, while sub-optimal in this case, has greater availability of proteomic data, rendering it a useful and more accessible tool for analysis. It is possible that the use of a less disease-relevant tissue such as blood plasma may explain the difference in results, though, both sets of findings are indicative of causal effects of *FCGR3B* on the risk of AD.

The evidence in our study suggests that *ACE2*, *LTA*, *FCGR2A*, *FCGR2B*, *and PLA2G1B* are unlikely to be effective candidate targets for AD; these findings should be taken into consideration to reduce the high attrition rate in AD drug development by discouraging trials of irrelevant targets such as those listed above [25]. Additionally, we observed little evidence of an effect for *IL6R,* though our estimate shows a consistent direction of effect with that of a recent MR study reporting significant protective effects of *IL6R* variants on AD risk via downstream C-reactive protein serum levels [26]. As that study utilized data from larger-scale GWASs, including 'proxy' AD cases—as opposed to the present study which only examined clinically diagnosed AD cases—it is possible that with greater power, we may have similarly observed a significant effect of *IL6R*. Alternatively, it is plausible that they observed an effect of *IL6R* due to bias introduced by the inclusion of 'proxy' AD cases. Definitive

conclusions cannot be drawn regarding the efficacy of *NFKB1* and *TNF* due to the imprecision of estimates observed in this study. However, a previously conducted MR found little evidence of a causal effect of *TNF* serum levels on AD [27]. Their instrumentation of pQTLs derived from blood serum raises the concern of tissue relevancy as mentioned earlier in the discussion of *FCGR3B*. Our suggestive evidence of a non-colocalized association between *TNF* and AD signals at the flanked chromosome 6 region highlights the possibility of alternate tissue-specific molecular pathways in which *TNF* and AD associate. Nevertheless, a potential ~28% reduction in disease risk through *TNF* plasma levels, as evidenced by this study, merits further exploration of this target in plasma tissue.

Limitations

Several limitations should be considered. First, this study was limited by the lack of proteomic data available for relevant drug targets. Among 33 initially selected targets from a total of 11 DMARDs, the final MR analyses could only include 9 targets from 8 DMARDs. Notably, methotrexate had to be excluded, which has recently accumulated promising observational evidence for an association with reduced AD risk [28]. Second, clinical evidence has demonstrated differential protein expression according to tissue specificity and it is plausible that this would have impacted our findings, in addition to the sub-optimal relevance of plasma as a biomarker for neurodegenerative processes [29]. Additional analyses that incorporate locus-specific instruments in tissues other than plasma (e.g., CSF) may yield different results. Third, there are several caveats to drug target MR that limit the ability to causally examine broader drug usage and disease outcome associations [30]. As we estimated the effects of perturbing individual drug targets, we cannot infer the likely aggregate effects of specific DMARDs. Furthermore, it is difficult to achieve a complete understanding of the pharmacodynamics of medications more broadly (e.g., understanding the exact molecular pathways in which etanercept affects *FCGR3B* and other targets). Thus, based on the genetic evidence alone, we cannot conclude how prescribed etanercept usage would affect AD risk via *FCGR3B* and other known or unknown targets. Further mechanistic studies and clinical trials are recommended for

comprehensive evaluation of DMARD repurposing. Finally, it is worth noting possible biases which could have further attenuated our estimates. Sourcing instruments from the UK BioBank—which is prone to selective representation—may introduce collider bias [31, 32]. Selected study participants tend to be healthier and may therefore have a lower genetic liability for disease traits, which in turn may potentially impact MR estimates [33]. That said, to date, there is little evidence in the literature specifically linking proteomic data to selection bias.

Conclusions

The findings from our study do not support the majority of selected DMARD targets as efficacious treatment candidates for AD, with exception of the etanercept target, *FCGR3B*. The observed positive effect of increased plasma *FCGR3B* concentration on AD risk warrants further investigation into the viability of therapeutics which lower *FCGR3B* protein levels in blood plasma. Our findings should be considered as part of the current discourse regarding the targeting of inflammatory mechanisms for AD treatment and prevention. Larger-scale MR studies that instrument a greater number of DMARD targets across various tissues are additionally recommended.

Abbreviations

AD: Alzheimer's disease

MR: Mendelian randomization

DMARD: Disease-modifying antirheumatic drug

pQTL: Protein quantitative trait loci

CNS: Central nervous system

TNF-α: Tumor necrosis factor

IL-6: Interleukin-6

GWAS: Genome-wide association study

RCT: Randomized control trial

LD: Linkage disequilibrium

BMI: Body mass index

UKB-PPP: UK Biobank Pharma Proteomics Project

SNP: Single nucleotide polymorphism

IGAP: International Genomics of Alzheimer's Project

LOAD: Late-onset Alzheimer's disease

GRCh: Genome Reference Consortium Human

IVW: Inverse variance weighted

OR: Odds ratio

CI: Confidence interval

SD: Standard deviation

Declarations

Ethics approval and consent to participate: The GWASs included in this study obtained written
informed consent from all participants prior to data collection. Related study protocols were
approved accordingly by the appropriate institutional review boards accordingly. Therefore,
separate ethical approval was not required for the present study.

Consent for publication: Not applicable.

Availability of data and materials: Proteogenomic summary data from the UKB-PPP can be found at
http://ukb-ppp.gwas.eu. Full AD GWAS data can be obtained by requesting access via the National
Institute on Aging Genetics of Alzheimer's Disease Data Storage Site (NIAGADS) data repository, ID:
NG00075. The codes used to generate the results of this study are located at
https://github.com/tkush15/AD_DMARD_MRCOLOC.

Competing interests: The authors declare that they have no competing interests.

Funding: ELA is supported by a UKRI Future Leaders Fellowship (MR/W011581/2). NMD is supported via a Norwegian Research Council 295989.

Authors' contributions: NMD, ELA, and CNK conceptualized the study. ELA provided coding. VTB provided coding and guidance for colocalization. CNK performed all statistical analyses, prepared figures, and drafted the initial manuscript. NMD and ELA guided the interpretation of results and provided written edits to the manuscript draft. All authors contributed to the revision and approval of the final manuscript.

Acknowledgments: Not applicable.

References

1. Nichols E, Steinmetz JD, Vollset SE, Fukutaki K, Chalek J, Abd-Allah F et al. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. Lancet Public Health. 2022;7(2):105-125. https://doi.org/10.1016/S2468-2667(21)00249-8

2. National Center for Health Statistics. Leading Causes of Death. [online]. Centers for Disease Control and Prevention: Hyattsville MD; 2024 [Accessed 2 May 2024]. Available from: https://www.cdc.gov/nchs/fastats/leading-causes-of-death.htm

3. Nandi A, Counts N, Bröker J, Malik S, Chen S, Han R et al. Cost of care for Alzheimer's disease and related dementias in the United States: 2016 to 2060. npj Aging. 2024;10(13). https://doi.org/10.1038/s41514-024-00136-6

4. Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N. A Meta-Analysis of Cytokines in Alzheimer's Disease. Biological Psychiatry. 2010;68(10):930-941. https://doi.org/10.1016/j.biopsych.2010.06.012

5. Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT. Inflammation as a central mechanism in Alzheimer's disease. Alzheimers Dement TRCI. 2018;4(1):575-590. https://doi.org/10.1016/j.trci.2018.06.014

6. Jayaraman A, Htike TT, James R, Picon C, Reynolds R. TNF-mediated neuroinflammation is linked to neuronal necroptosis in Alzheimer's disease hippocampus. BMC ANC. 2021;9(159) https://doi.org/10.1186/s40478-021-01264-w

7. Erta M, Quintana A, Hidalgo J. Interleukin-6, a major cytokine in the central nervous system. Int J Biol Sci. 2012;8(9):1254-1266. https://doi.org/10.7150/ijbs.4679

8. Chang R, Yee K-L, Sumbria RK. Tumor necrosis factor α Inhibition for Alzheimer's Disease. J Cent Nerv Syst Dis. 2017;9. https://doi.org/10.1177/117957351770927

9. Bellenguez C, Küçükali F, Jansen IE, Kleineidam L, Moreno-Grau S, Amin N et al. New insights into the genetic etiology of Alzheimer's disease and related dementias. Nat Genet. 2022;54:412-436. https://doi.org/10.1038/s41588-022-01024-z

10. Chou R, Kane M, Ghimire S, Gautam S, Gui J. Treatment for Rheumatoid Arthritis and Risk of Alzheimer's Disease: A Nested Case-Control Analysis. CNS Drugs. 2016;30:1111-20. https://doi.org/10.1007/s40263-016-0374-z

11. Huang L-C, Chang Y-H, Yang Y-H. Can disease-modifying anti-rheumatic drugs reduce the risk of developing dementia in patients with rheumatoid arthritis? Neurotherapeutics. 2019;16(3):703-9. https://doi.org/10.1007/s13311-019-00715-6

12. Xie W, Hou Y, Xiao S, Zhang X, Zhang Z. Association between disease-modifying antirheumatic drugs for rheumatoid arthritis and risk of incident dementia: a systematic review with meta-analysis. RMD Open. 2024;10(1). https://doi.org/10.1136/rmdopen-2023-004016

13. Sattui S, Navarro-Millan I, Xie F, Rajan M, Yun H, Curtis JR. Incidence of dementia in patients with rheumatoid arthritis and association with disease modifying anti-rheumatic drugs – Analysis of a national claims database. Semin Arthritis Rheum. 2022;57:152083. https://doi.org/10.1016/j.semarthrit.2022.152083

14. Watad A, McGonagle D, Anis S, Carmeli R, Cohen AD, Tsur AM et al. TNF inhibitors have a protective role in the risk of dementia in patients with ankylosing spondylitis: Results from a

nationwide study. Pharmacol Res. 2022;182:106325.

https://doi.org/10.1016/j.phrs.2022.106325

15. Zhou M, Xu R, Kaelber DC, Gurney ME. Tumor Necrosis Factor (TNF) blocking agents are associated with lower risk for Alzheimer's disease in patients with rheumatoid arthritis and psoriasis. PLOS One. 2020;15(3). https://doi.org/10.1371/journal.pone.0229819

16. Butchart J, Brook L, Hopkins V, Teeling J, Pütener U, Culliford D et al. Etanercept in Alzheimer disease: A randomized, placebo-controlled, double-blind, phase 2 trial. Neurology. 2015;84(21):2161-8. https://doi/org/10.1212/WNL.0000000000001617

17. Tobinick E, Gross H, Weinberger A, Cohen H. TNF-alpha modulation for treatment of Alzheimer's disease: a 6-month pilot study. Med Gen Med. 2006;8(2):25.

https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/16926764/

18. Smith GD, Ebrahim S. Mendelian randomization: prospects, potentials, and limitations. Int J Epidemiol. 2004;33(1):30-42. https://doi.org/10.1093/ije/dyh132

19. Gill D, Georgakis MK, Walker VM, Schmidt AF, Gkatzionis A, Freitag DF et al. Mendelian randomization for studying the effects of perturbing drug targets. Wellcome Open Res. 2021;6:16. https://doi.org/10.12688/wellcomeopenres.16544.2

20. Davies NM, Holmes MV, Smith GD. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. BMJ. 2018;362:k601. https://doi.org/10.1136/bmj.k601

21. Sun BB, Chiou J, Traylor M, Benner C, Hsu Y-H, Richardson TG. Plasma proteomic associations with genetics and health in the UK Biobank. Nature. 2023;622(7982):329-338.

https://doi.org/10.1038/s41586-023-06592-6

22. Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC et al. Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing. Nat Genet. 2019;51(3):414-430. https://doi.org/10.1038/s41588-019-0358-2

23. Cruchaga C, Western D, Timsina J, Wang L, Wang C, Yang C et al. Proteogenomic analysis of human cerebrospinal fluid identifies neurologically relevant regulation and informs causal proteins for Alzheimer's disease. Res Sq. 2023. https://doi.org.10.21203/rs.3.rs-2814616/v1

24. Karlsson L, Vogel J, Arvidsson I, Åström K, Janelidze S, Blennow K et al. Cerebrospinal fluid reference proteins increase accuracy and interpretability of biomarkers for brain diseases. Nat Commun. 2024;15. https://doi.org/10.1038/s41467-024-47971-5

25. Cummings J, Morstorf T, Zhong K. Alzheimer's disease drug-development pipeline: few candidates, frequent failures. Alz Res Therapy. 2014;6(37). https://doi.org/10.1186/alzrt269

26. Myserlis EP, Anushree R, Anderson CD, Georgakis MK. Genetically proxied IL-6 signaling and risk of Alzheimer's disease and lobar intracerebral hemorrhage: A drug target Mendelian randomization study. Alzheimers Dement TRCI. 2024;10(3).
https://doi.org/10.1002/trc2.70000

27. Andrews SJ, Goate A. Mendelian randomization indicates that TNF is not causally associated with Alzheimer's disease. Neurobiol Aging. 2019;84.
https://doi.org/10.1016/j.neurobiolaging.2019.09.003

28. Newby D, Prieto-Alhambra D, Duarte-Salles T, Ansell D, Pedersen L, van der Lei J et al. Methotrexate and relative risk of dementia amongst patients with rheumatoid arthritis: a multi-national multi-database case-control study. Alzheimers Res Ther. 2020;12(38).
https://doi.org/10.1186/s13195-020-00606-5

29. Whitehead A, Crawford DL. Variation in tissue-specific gene expression among natural populations. Genome Biol. 2005;6(R13). https://doi.org/10.1186/gb-2005-6-2-r13

30. Anderson EL, Williams DM. Drug target Mendelian randomization: are we really instrumenting drug use? Diabetologia. 2023;66(6):1156-8. https://doi.org/10.1007/s00125-023-05875-x

31. Gkatzionis A, Burgess S. Contextualizing selection bias in Mendelian randomization: how bad is it likely to be? Int J Epidemiol. 2019;48(3):691-701. https://doi.org/10.1093/ije/dyy202

32. Schoeler T, Speed D, Porcu E, Pirastu N, Pingault JB, Kutalik Z. Participation bias in the UK Biobank distorts genetic associations and downstream analyses. Nat Human Behav. 2023;7(7):1216-27. https://doi.org/10.1038/s41562-023-01579-9

33. Munafò MR, Tilling K, Taylor AE, Evans DM, Smith GD. Collider scope: when selection bias can substantially influence observed associations. Int J Epidemiol. 2018;47(1):226-35. https://doi.org/10.1093/ije/dyx206