Occupational Exposure to Heat and Wildfire Smoke in California Correctional Facilities

Rachel Sklar[1*], Samantha Kramer[2], Fred Lurmann[2], Marissa Childs[1], Stefano Bertozzi[3], Elizabeth Noth[4]

1. Department of Environmental and Occupational Health Sciences, University of Washington School of Public Health, Seattle, WA 98195, USA

2. Sonoma Technology, Inc., Petaluma, California

3. Division of Health Policy and Management, School of Public Health, University of California, Berkeley, Berkeley, California, USA; School of Public Health, University of Washington, Seattle, Washington, USA; Instituto Nacional de Salud Pública, Cuernavaca, Mexico

4. Division of Environmental Health Sciences School of Public Health, University of California, Berkeley, Berkeley, California, USA

*Correspondence to rsklar@uw.edu

1. Abstract

Objectives: To assess long-term trends in outdoor heat and wildfire smoke exposure at

410 correctional facilities in California from 2000–2023.

Methods: We used ERA5-Land data to calculate daily wet-bulb globe temperature

(WBGT) and identified days exceeding the NIOSH 28°C threshold. Wildfire smoke

exposure was estimated using NOAA's Hazard Mapping System (HMS). Temporal

trends in daily heat and smoke exposure were modeled using linear regression for each

facility.

Results: Smoke-affected days rose statewide from 4.9% (2006–2010) to 12.5% (2019–

2023), with Northern California experiencing the highest burden and steepest increases.

Days above 28°C WBGT increased from 27.3% to 29.8% statewide, with the Central

Valley and desert regions showing the highest current burden and coastal areas

experiencing the steepest increases. Coastal facilities show emerging heat risks, and some

regions—especially the Central Valley and Sierra foothills—face dual exposure to both

hazards.

Conclusions: Correctional officers and incarcerated workers face rising environmental

risks, with regional variation in exposure. Facilities with concurrent heat and smoke

burdens must balance cooling and filtration needs. Resilience planning and occupational

protections must evolve to meet facility- and region-specific challenges.

2. Introduction

Correctional facilities in California are often located in regions increasingly affected by extreme heat and wildfire smoke. [1,2] As of 2023, over 33,740 correctional officers and 64,788 incarcerated workers were employed within these settings in California. [3,4] Many correctional facilities were not built to withstand modern climate stressors. Their aging, heat-retaining concrete structures and outdated HVAC systems offer limited protection. [5,6] These environmental hazards are compounded by demanding job requirements, making it difficult to mitigate risks for staff and incarcerated individuals alike.

Existing studies have begun to examine heat exposure in U.S. prisons, though the literature remains sparse.[2,7] Research on wildfire smoke exposure in correctional settings remains especially limited. This study explores current and long-term patterns in both heat and wildfire smoke across California correctional facilities, highlighting the need for targeted research on occupational exposures and controls. We focus on working populations given limited occupational research in correctional settings, though incarcerated residents likely face greater exposure due to living continuously in these facilities.

3. Methods

Correctional facilities were identified using North American Industry Classification System (NAICS) codes. Using ArcGIS, we mapped the locations of all 410 identified California correctional facilities spanning federal, state, county, local, and multi-jurisdictional institutions. Facilities included adult and juvenile detention centers with varying security classifications. Heat exposure data were obtained from the ERA5-Land reanalysis dataset spanning 2006-2023, with daily maximum wet-bulb globe temperature (WBGT) values calculated using temperature,

humidity, and solar radiation data at each facility location.[8] For each facility and year, we

calculated the percentage of days exceeding the National Institute for Occupational Safety and

Health (NIOSH) recommended threshold of 28°C WBGT for outdoor work activities. To assess

long-term trends in heat exposure, we fitted linear regression models for each facility, using

percentage of hot days as the dependent variable and year as the independent variable across the

full 24-year study period (2000-2023). We calculated the total change in heat exposure over the

study period by multiplying the annual rate of change (slope coefficient) by 23 years.[2] For

smoke, we calculated the percentage of days each year with detectable smoke from 2006–2023

using NOAA's Hazard Mapping System (HMS) plumes and applied the same linear regression

approach to assess the trend in smoke exposure over a 17-year period. [9]

4. Results

On average, the percentage of smoke-affected days rose from 4.9% (2006–2010) to 12.5%

(2019–2023), reflecting a marked increase in smoke exposure in the state. The steepest increases

and highest current smoke days are concentrated in Northern California, particularly the northern

Sierra foothills and adjacent valleys (Figure 1a). Statewide, the average percentage of heat days

above the 28°C WBGT threshold rose from 27.3% (2006–2010) to 29.8% (2019–2023),

reflecting a more modest increase in outdoor heat exposure. The steepest heat increases appear in

coastal Southern California and the LA basin, while the Central Valley and desert regions

currently experience the highest current heat levels (Figure 1b). Several coastal Southern

California facilities show steep upward trends in heat exposure despite relatively low current

heat levels, indicating emerging thermal risk. Some regions have mismatched burdens such as

the northern part of the state with high smoke but moderate heat, and inland southern areas with

high heat but moderate smoke. The Central Valley and Sierra foothills face a dual burden, with

high and rising levels of both (Figure 1).

Notably, the relationship between current and long-term exposure differs by hazard. For

smoke, high current exposure correlates strongly with the upward trend over the years. However,

for heat, the greatest upward trend is seen in places with low to moderate current burden.

Locations with already high burden may also see increases in the intensity of heat days not

captured in the metric of days over °C WBGT.

5. Discussion

People who live and work in California correctional facilities are facing growing health risks

from heat and wildfire smoke exposure. These risks may be magnified by aging and inadequate

infrastructure.[5,6] Many occupational factors—such as personal protective equipment (PPE) use,

limited environmental control, and extended shifts—may also elevate risk of heat and smoke

exposure for both correctional officers and incarcerated workers. Correctional officers wear

heavy protective equipment that inhibits heat dissipation.[10] They may work overtime shifts due to

staffing shortages which may increase heat exposure especially during emergency incidents. [1]

During wildfire events, officers and incarcerated workers may face a tradeoff between

wearing respiratory protection, which can worsen heat stress, or forgoing masks and risking

smoke exposure.[11] Underlying health conditions and psychotropic medication use further elevate

heat-related illness risk among incarcerated workers.[2]

Regional variability in smoke and heat trends suggests that Southern California facilities may

need to prioritize extreme heat mitigation, while Northern California institutions may face

greater challenges from increasing wildfire smoke, even in the absence of significant heat trends.

A small but important subset of facilities, primarily in the Central Valley and Sierra foothills,

faces concurrent increases in both heat and smoke exposure raising concerns about the health

effects of co-exposure and the challenge of balancing cooling needs with air filtration when

outdoor air intake may worsen indoor air quality. These differences highlight the need for

facility-specific risk assessments and tailored mitigation strategies that account for the distinct

environmental hazards facing each region. Additionally, emerging heat burdens in parts of

coastal California, areas historically considered low risk, raise questions about whether the

facilities in these regions are adequately equipped with the infrastructure and plans needed to

mitigate these exposures.

This study did not directly assess indoor environmental conditions where both free and

incarcerated workers may spend most of their time at work. Future work will examine how these

outdoor estimates translate into occupational risks indoors. Additionally, while the HMS smoke

plume data provide consistent, long-term spatial coverage, they are limited in vertical resolution

and may not accurately capture ground-level concentrations, but are similar to patterns in

estimated ground-level wildfire smoke fine particulate matter.[12] As a result, our estimates may

over- or underestimate smoke days relevant for exposure assessment. Future analyses will

incorporate additional data sources to better characterize surface-level smoke conditions across

correctional settings.

6. Public Health Implications

Environmental hazards like extreme heat and wildfire smoke pose serious threats to the

health, safety, and productivity of both correctional officers and incarcerated workers. These

conditions can lead to more injuries and illnesses, make it harder to perform physical or decision-

intensive tasks, and strain operations and security. For correctional officers, this may mean

increased medical leave, staffing shortages, and financial pressures on agencies to respond to

workers compensation claims. For incarcerated workers, exposure to heat and smoke may limit

participation in job assignments and programming, potentially reducing opportunities for skill-

building and rehabilitation. Addressing these risks through targeted mitigation strategies is

essential to ensure safer, more resilient correctional environments for all. These risks cannot be

addressed through one-size-fits-all solutions, as the correctional facilities explored here differ in

size, infrastructure type, and capacity to implement exposure controls.

7. References

1. LeMasters K, Haber LA. The hidden crisis of incarcerated individuals during wildfires. *The Lancet Regional Health - Americas*. 2025;43:101032. doi:10.1016/j.lana.2025.101032

2. Tuholske C, Lynch VD, Spriggs R, et al. Hazardous heat exposure among incarcerated people in the United States. *Nat Sustain*. 2024;7(4):394-398. doi:10.1038/s41893-024-01293-y

3. Occupational Employment and Wages, May 2023: 33-3012 Correctional Officers and Jailers. U.S. Bureau of Labor Statistics. Accessed May 28, 2025. https://www.bls.gov/oes/2023/may/oes333012.htm#(1)

4. Macomber J, Singh D, Aghakhanian DA, Davison D. *California Prison Industry Board*. California Prison Industry Authority

5. Jay O, Capon A, Berry P, et al. Reducing the health effects of hot weather and heat extremes: from personal cooling strategies to green cities. *The Lancet*. 2021;398(10301):709-724. doi:10.1016/S0140-6736(21)01209-5

6. Sklar R, Noth E, Kwan A, Sear D, Bertozzi S. Ventilation conditions during COVID-19 outbreaks in six California state carceral institutions. *PLOS ONE*. 2023;18(11):e0293533. doi:10.1371/journal.pone.0293533

7. Ovienmhada U, Hines M, Krisch M, Diongue AT, Minchew B, Wood DR. Spatiotemporal Facility-Level Patterns of Summer Heat Exposure, Vulnerability, and Risk in United States Prison Landscapes. *GeoHealth*. 2024;8(9):e2024GH001108. doi:10.1029/2024GH001108

8. Spangler KR, Liang S, Wellenius GA. Wet-Bulb Globe Temperature, Universal Thermal Climate Index, and Other Heat Metrics for US Counties, 2000–2020. *Sci Data*. 2022;9(1). doi:10.1038/s41597-022-01405-3

9.   Vargo JA. Time Series of Potential US Wildland Fire Smoke Exposures. *Front Public*

Health. 2020;8. doi:10.3389/fpubh.2020.00126

10. Hudson S, Ridland L, Blackburn J, Monchuk L, Ousey K. The comfort and functional

performance of personal protective equipment for police officers: a systematic scoping

review. *Ergonomics*. 2024;67(10):1317-1337. doi:10.1080/00140139.2024.2302957

11. Li Y, Tokura H, Guo YP, et al. Effects of wearing N95 and surgical facemasks on heart rate,

thermal stress and subjective sensations. *Int Arch Occup Environ Health*. 2005;78(6):501-

509. doi:10.1007/s00420-004-0584-4

12. Childs ML, Li J, Wen J, et al. Daily Local-Level Estimates of Ambient Wildfire Smoke

PM2.5 for the Contiguous US. *Environ Sci Technol*. 2022;56(19):13607-13621.

doi:10.1021/acs.est.2c02934

Figure 1. Environmental exposure trends at California correctional facilities. *(a) Smoke exposure trends (2006-2023) showing current smoke levels and magnitude of 17-year change across facilities. (b) Heat exposure trends (2000-2023) showing current heat levels and magnitude of 23-year change. The 410 correctional facilities shown span federal, state, county, local, and multi-jurisdictional institutions. Current heat and smoke levels are calculated as an annual average over the last 5 years in the sample (2019-2023).*