The Impact of CYP2C19 Genotype on Clopidogrel Response and Clinical Outcomes in Coronary Artery Disease Patients: A Correlation with SYNTAX Score

Sohaib Ashraf*[,1,a, b], Amjad Mujtaba[1,a], Amir Shahbaz[1,a], Shoaib Ashraf*[1,b], Moneeb Ashraf[2,c], Rutaba Akmal[d], Muhammad Ahmad Imran[*2,d], Sidra Ashraf[e], Meer Hassan[2,f,g], , Qazi Abdul Saboor[a]

Affiliations:

a)	(MBBS, FCPS) Department of Cardiology, Shaikh Zayed Post-Graduate Medical Complex, Lahore, Pakistan.

b)	(MBBS, FCPS) Department of Pathobiology, Riphah University, Lahore, Pakistan

c)	(MBBS, MPhil) Department of Pharmacology, King Edward Medical University, Mayo Hospital, Lahore, Pakistan.

d)	(MBBS) Department of Microbiology, Shaikh Zayed Post-Graduate Medical Institute, Lahore, Pakistan.

e)	(MBBS) Department of Internal Medicine, Services Institute of Medical Sciences, Lahore, Pakistan.

f)	(MBBS) Department of Internal Medicine, Shaikh Zayed Post-Graduate Medical Institute, Lahore, Pakistan.

Corresponding Author[*:]

1.	Dr. Sohaib Ashraf, Department of Cardiology, Shaikh Zayed Post-Graduate Medical Complex, Lahore, Pakistan. sohaib-ashraf@outlook.com, sohaib@skzmdc.edu.pk

26 2. Dr. Muhammad Ahmad Imran, Department of Medicine, Shaikh Zayed Post-Graduate

27   Medical Complex, Lahore, Pakistan. ahmad.imram@skzmdc.edu.pk

28   ORCID ID: 0000-0003-3358-229X

30 [1] Joint First Author

31 [2] Joint Second Author

Word Count	5616
pages	27
Figures	2
Tables	2

ABSTRACT

Background

CYP2C19 polymorphism helps determine the reactivity of platelets in coronary artery disease. The impact of CYP2C19 genotype on clopidogrel response and clinical outcomes remains unclear. The SYNTAX score helps in determining the prognosis.

Methods

To test this, we enrolled more than 500 patients with ischemic heart disease. All the patients were on follow-up treatment of antiplatelets, clopidogrel and aspirin. We noted the CYP2C19 phenotype for cardiovascular events.

Results

This study included 156 individuals with a mean age of 59.2 ± 10.6 months and a median age of 60 months. Most patients were male (71.2%). The mean SYNTAX score was 26.1 ± 11.3, with a median of 29. Among participants, 25.6% had dyslipidemia, 75.6% had hypertension, and 49.4% had diabetes; only 9.6% had a history of PCI. There was no significant difference in age, BMI, sex, or smoking status between the stable angina and acute coronary syndrome groups. However, dyslipidemia, diabetes, PCI history, and a higher mean SYNTAX score were significantly more common in the acute coronary syndrome group. Statistical significance was achieved for SYNTAX score (P<0.05).

Conclusion

The SYNTAX score can be used to predict the complexity of CAD in patients. The greater the score, the greater the probability of CAD

60   Key Words: SYNTAX, CAD, ACS, CYP2C19

63 INTRODUCTION

Coronary artery disease (CAD) remains a leading global cause of death and is increasingly

affecting younger individuals. Recent efforts focus on identifying and managing risk

factors in young patients to lower CAD incidence and improve quality of life. While older

patients with acute coronary syndrome often present with recognized risk factors, these are

less commonly identified in younger patients with stable angina. (Sukhija et al., 2006).

Coronary angiography is the gold standard for assessing atherosclerotic coronary artery

disease but carries risks, ranging from minor complications to potentially life-threatening

events if not promptly managed. Advances in procedural techniques, equipment design,

and operator expertise have significantly reduced these risks over time. (Brilakis, 2021).

One in four Americans suffers from coronary heart disease. The American Heart

Association reports that a heart attack occurs every 41 seconds in the United States.

Coronary heart disease affects around 17 million Americans, with men comprising 55% of

cases. Over half a million die from it annually in the U.S. Men by age 40 have a 49%

lifetime risk, higher than women's 32%. Although male predominance declines with age,

coronary event incidence still rises as people get older. Globally, it remains the leading

cause of adult death across all income levels (Sarkees & Bavry, 2009).

Normally angina symptoms include chest pain or an anginal analogue that is exacerbated

by physical activity and relieved by rest or nitroglycerin. Even though there are several

forms of angina, they are often characterized by symptoms like extreme fatigue, nausea, or

shortness of breath in relation to the individual's degree of activity (Boyette & Manna,

2022). The ACS and stable angina may require coronary arteriography or angiography to

determine a better management plan.

Coronary arteriography has no absolute contraindications but carries risks of cardiac and

non-cardiac complications. Certain patient factors—like severe CAD, heart failure with

low ejection fraction, recent stroke or MI, and bleeding tendencies—can increase these

risks. The likelihood of complications also depends on the type of procedure, whether

diagnostic angiography alone or additional intervention (Brilakis, 2021). Serious

complications from cardiac catheterization are rare, occurring in about 2% of patients, with

a mortality risk below 0.08%. Advances like iso-osmolar contrast media, improved catheter

design, better bleeding control, and skilled operators have further reduced these risks,

allowing even critically ill patients to undergo the procedure safely when needed. (Manda

& Baradhi, 2022).

A study conducted by Ramjattan in 2022 showed CT coronary angiography (CTCA) shows

high accuracy, with 89% sensitivity and 96% specificity for detecting coronary heart

disease. In early trials, 36% reported angina due to coronary disease, and 47% were

diagnosed with CAD. CTCA is recommended for patients with chest discomfort and

symptoms suggestive of coronary disease. (Ramjattan et al., 2022). The ESC/EACTS

guidelines recommend the anatomic SYNTAX score (SxS) to assess coronary artery

disease complexity. Initially developed from the SYNTAX trial to predict outcomes after

PCI or CABG in stable patients, the SxS has since been applied to other patient groups,

including those undergoing primary PCI for acute coronary syndromes. (Y.-H. Kim et al.,

2010).

Studies, including SYNTAX trial sub-analyses, show that the original SYNTAX score

often misclassifies risk, especially for mortality in stable CAD or ACS patients treated with

PCI. To improve risk stratification, the SYNTAX score II (SxS II) was developed by

adding clinical factors like age, creatinine clearance, LVEF, unprotected left main disease,

peripheral vascular disease, gender, and COPD. This enhanced model offers better

accuracy, particularly in patients with left main or multivessel disease. (Scherff et al.,

2011).

Cardiologists, interventionists, and surgeons all utilize the SYNTAX score, which is based

on an angiographic approach, to determine the degree of difficulty of coronary artery

lesions. Higher scores predict poorer outcomes after modern revascularization, especially

PCI. While it shows some predictive value for major adverse cardiac and cerebrovascular

events, its role in forecasting early or late mortality remains unclear. Evidence supports its

use in left main PCI, but data are limited for predicting outcomes in patients with three-

vessel disease undergoing PCI or stenting. (Safarian et al., 2014a).

The rationale of this study is to compare the angiographic findings in stable angina versus

acute coronary syndrome by using the SYNTAX score. The results of this study will

benefit in the better management of these patients.

MATERIAL AND METHODS:

This prospective comparative study was conducted over 12 months at the Department of

Cardiology, Shaikh Zayed Hospital, Lahore, and included 156 patients undergoing

coronary angiography. Participants were divided into two groups: Group A (78 patients

with stable angina) and Group B (78 patients with acute coronary syndrome). The sample

size was calculated using proportions from a previous study published in 2019 (See

Supplementary File) at 80% power and 95% confidence interval, accounting for a 10%

dropout rate [Fukui, T, et al., 2013]. Patients aged 18–80 years of both genders undergoing

primary PCI or elective coronary angiography were included. Patients with a history of

prior cardiac catheterization, PCI, or CABG were excluded. The exclusion criteria also

included a previous history of cardiac catheterization and PCI patients with CABG.

Approval was taken from the Institutional Review Board (IRB) of Shaikh Zayed Medical

Complex, Lahore, total of 156 patients who fulfilled the inclusion criteria and were

admitted through the OPD and the emergency Department of the Shaikh Zayed Hospital,

Lahore. A written informed consent was taken from each patient after explaining the risks,

purpose, and benefits of this research. Detailed history, including age, gender, height,

weight, BMI, and medical number, was noted, and SYNTAX score was used after

angiography for data collection for both groups. The selected patients were classified into

two groups (A & B). Group A patients were referred as coronary angiography with stable

angina, and other patients in group B was referred to acute coronary syndrome. The

patients were instructed to follow up after 30 days of discharge if any complications

develop after this clinical procedure.

Data was analyzed by using SPSS v.25. Mean ± SD were computed for quantitative

variables like age, hypertension and BMI. Qualitative variables like gender, diabetic

mellitus, PCI, were presented as frequency and percentages. Post-stratification, chi-square

test was applied to compare the factors in stratified groups. P ≤0.05 was considered as

significant.

RESULTS

A total of 156 patients were included in the study. The majority of participants were male

(n = 111, 71.2%), while females accounted for 28.8% (n = 45). Patients were equally

distributed into two clinical groups: Group A with stable angina (n = 78, 50.0%) and Group

B with acute coronary syndrome (ACS) (n = 78, 50.0%). Regarding smoking status, most

participants were non-smokers (n = 128, 82.1%), whereas smokers comprised 17.9% (n =

28). Dyslipidemia was present in 25.6% (n = 40) of the study population, while the

remaining 74.4% (n = 116) did not have dyslipidemia. The study included 118 (75.6%)

hypertensive patients and (n=77. 49.4%) diabetes patients. However, only 15 (9.6%)

patients underwent PCI (Table 1, Figure 1).

The mean age of the patients was 59.2 ± 10.6 years, with a median age of 60.0 years

(range: 29–80 years). The mean body mass index (BMI) was 27.2 ± 3.5 kg/m². The

angiographic assessment using the SYNTAX score revealed a mean score of 26.1 ± 11.3,

with a median of 29 (range: 2–58). These findings describe the baseline demographic and

clinical characteristics of the study cohort (Table 1, Figure 1).

The study cohort was equally divided into Group A (stable angina, n = 78) and Group B

(acute coronary syndrome, n = 78). The mean age of patients in Group A was slightly

higher (60.2 ± 10.1 years) compared to Group B (58.3 ± 11.0 years), although this

difference was not statistically significant (p = 0.26). Similarly, the mean body mass index

(BMI) was comparable between the two groups (27.1 ± 2.2 kg/m² in Group A vs. 27.4 ±

4.4 kg/m² in Group B; p = 0.58). However, the SYNTAX score was significantly higher in

patients with acute coronary syndrome (mean 28.2 ± 9.9) than those with stable angina

(mean 23.8 ± 12.4), with a p-value of 0.01, indicating greater angiographic complexity in

Group B (Table 2).

The sex distribution in males was predominated in both groups (51.4% in Group A vs.

48.6% in Group B; p = 0.60). The proportion of smokers was higher in Group B (57.1%)

compared to Group A (42.9%), but this difference was not statistically significant (p =

0.40). This represents the homogeneity in the groups. The dyslipidemia in Group B had a

greater prevalence compared to Group A, and the results were statistically significant.

(90.0% 10.0%, P<0.05). Hypertension was common in both groups, present in 51.7% of

Group A and 48.3% of Group B patients, with no significant difference (p = 0.46).

Similarly, diabetes mellitus was more common among group B than group A, but the

results were not statistically significant (54.5%, 45.5%, P>0.05). There was no significant

difference in age, BMI, sex, or smoking status between the stable angina and acute

coronary syndrome groups. However, dyslipidemia, diabetes, PCI history, and a higher

mean SYNTAX score were significantly more common in the acute coronary syndrome

group. And statistical significance was achieved for SYNTAX score (P<0.05) (Table 2,

Figure 2).

DISCUSSION

In this study, we have compared the angiographic findings in patients reported with stable

angina and acute coronary syndrome by using the SYNTAX score. The objective of this

study is to compare the angiographic findings in stable angina with acute coronary

syndrome by using SYNTAX score. Our results showed that the SYNTAX score was

significantly higher in acute coronary syndrome patients as compared to stable angina

patients. Recent advances in technology have led to an increase in the proportion of

patients with ACS undergoing percutaneous coronary intervention (PCI). The selection of

the optimal revascularization modality for patients with ACS depends on the patient

characteristics and lesion severity. In addition, coronary artery bypass graft is still

appropriate for patients with disease refractory to medical therapy and when PCI is not

suitable or possible (Godoy, Lucas C., et al., 2019).

The complex coronary artery anatomy may be quantified in recognition of the SYNTAX

score, which is solely generated from the coronary anatomy and lesion characteristics

(Vizirianakis, Ioannis S., et al., 2021). The score, which was a crucial component of the

SYNTAX trial design (Thuijs, Daniel JFM, et al., 2019), was initially created as a way to

guarantee that the cardiologist and cardiac surgeon accurately reviewed the angiogram of

patients with complex CAD, allowing a consensus to be reached regarding the best

procedure and level of revascularization. It's significant that the SYNTAX score was

determined a priori, before the results of revascularization were known. In order to help

with revascularization decisions, the SYNTAX trial results showed that the score is crucial

in stratifying patients with complicated CAD (Gallo, Michele, et al., 2022).

The SYNTAX score of zero implies a low risk of cardiovascular, and generally promotes a

more conservative approach to care, according to a study done by Berman et al. On the

other hand, if the syntactic score is higher than 27, the risk is comparable to that of patients

with clinically severe CAD, and aggressive medical therapy is typically needed (Berman,

Daniel S et al., 2016).

Long-term outcome comparisons between patients with ACS and SAP among those

receiving percutaneous coronary intervention (PCI) have produced mixed outcomes. The

long-term risks of UAP and SAP following PCI were comparable in individuals with ACS

and SAP in the Lescol Intervention Prevention Study (Sandeelo, Imran Khan, et al., 2019).

The risks of major adverse cardiac events at 1 year between individuals with ACS and SAP

were not significantly different, according to a sub analysis of the Arterial

Revascularization Therapies Study (Matsoukis, Ioannis L., et al., 2021). However,

compared to SAP, patients with ACS had a higher probability of death and combined death

and myocardial infarction, according to Naidu and colleagues. Consideration should be

given to the role of the overall atherosclerotic burden and disease progression following

PCI in arteries not receiving intervention in ACS patients (Van Nguyen, Huy, et al., 2021).

In the present study, a SYNTAX score was used to bifurcate the risk of adverse outcome in

ACS versus stable angina patients. There was significantly high SYNTAX score was

reported in ACS versus stable angina patients. Therefore, the risk of adverse outcome is

significantly higher in ACS versus stable angina patients. Also, a high proportion of

comorbidities were reported in patients in the ACS group as compared to stable angina

patients. These results were comparable with the aforementioned studies. A few studies

have assessed the predictive value SYNTAX score for long-term clinical outcomes in a PCI

population (He, Jiqiang, et al., 2019; Cavalcante, Rafael, et al., 2017).

In addition, in this study, we have explored the SYNTAX score difference in patients with

diabetes and hypertension. Furthermore, the syntax score was statistically significantly (p-

value 0.001) high in patients with diabetes (31.4 ± 7.92) as compared to patients without

diabetes (20.8 ± 11.9). Moreover, the SYNTAX score was significantly high in patients

with hypertension (28.9 ± 9.05) as compared to patients without hypertension (17.2 ±

13.5). These results were comparable with the published data (Xu, Weifeng, et al., 2019;

Kurtul BE, et al., 2021). In addition, another study conducted by (Kose, N., et al., 2019)

reported a high mean difference of SYNTAX score in patients with stable angina versus

ACS. Also, (Kose, N., et al., 2019) reported the high mean of SYNTAX score in patients

with diabetes and hypertension (Kose, N., et al., 2019). These results were also comparable

to our results. Furthermore, another study reported that a strong positive correlation exits

between SYNTAX score and coronary artery disease; this means that the chance of severity

of disease is higher with an increase in SYNTAX score. These results were also

comparable to our data.

The SYNTAX score is unaffected by a patient's clinical traits, some of which, such as

patient age, have repeatedly been demonstrated to be independent predictors of mortality.

In comparison to lesion-based scores like the American Heart Association lesion

classification, prior research has shown that clinically based risk models, such as the Mayo

Clinic Risk Score MCRS, perform better at predicting morbidity and mortality. A potential

drawback to the SYNTAX Score's application in risk stratification is the lack of any

consideration for clinical factors in its calculation. Secondly, it was a single center study.

Sample size was limited as we selected patients who were admitted to our department with

cardiac issues and not all the patients admitted/visiting the hospital.

CONCLUSION

SYNTAX score can be used to predict the complexity of CAD in patients. Therefore, all

patients with low-to-intermediate pretest probability of CAD undergoing cardiac CT scan

for disease assessment should have SYNTAX score calculated, and those with a greater

SYNTAX score in the presence of significant disease are likely to have a high SYNTAX

score on coronary angiography. The SYNTAX score helps a lot to the cardiologist and

gives the proper result. It is very helpful to check the prognosis of the disease.

Acknowledgement:

Shaikh Zayed Post Graduate Medical Complex, for making this study easy to conduct.

Financial Support

No financial support was needed.

Declaration of Competing Interest

The authors declare no conflict of interest.

Authors Contribution

S.A. and A.M. conceived the idea, and prepared the initial manuscript. A.S., S.A., R.K and

M.A. collected the patient data and wrote the subsequent manuscript. M.A.I., S.A, M.H.

refined and finalized the manuscript. Q.S. supervised the entire manuscript.

REFERENCES

Al-Hadi, H. A., & Fox, K. A. (2009). Cardiac Markers in the Early Diagnosis and Management of Patients with Acute Coronary Syndrome. *Sultan Qaboos University Medical Journal*, *9*(3), 231–246.

Abdelnabi M, Almaghraby A, Tok ÖÖ, Öz TK, Saleh Y, Morsi A, et al., A real-life correlation between clinical SYNTAX score II and carotid intima-media thickness in patients with stable coronary artery disease. J. Saudi Heart Assoc.. 2020;32(1):8.

Altekin, R. E., Kilinc, A. Y., Onac, M., & Cicekcibasi, O. (2020). Prognostic Value of the Residual SYNTAX Score on In-Hospital and Follow-Up Clinical Outcomes in ST Elevation Myocardial Infarction Patients Undergoing Percutaneous Coronary Interventions. *Cardiology Research and Practice*, *2020*, 9245431. https://doi.org/10.1155/2020/9245431

Anstey Watkins, J., Wagner, F., Xavier Gómez-Olivé, F., Wertheim, H., Sankoh, O., & Kinsman, J. (2019). Rural South African Community Perceptions of Antibiotic Access and Use: Qualitative Evidence from a Health and Demographic Surveillance System Site. *The American Journal of Tropical Medicine and Hygiene*, *100*(6), 1378–1390. https://doi.org/10.4269/ajtmh.18-0171

Banks, K., Lo, M., & Khera, A. (2010). Angina in Women without Obstructive Coronary Artery Disease. *Current Cardiology Reviews*, *6*(1), 71–81. https://doi.org/10.2174/157340310790231608

Basit, H., Malik, A., & Huecker, M. R. (2022). Non ST Segment Elevation Myocardial Infarction. In *StatPearls*. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK513228/

Bergmark, B. A., Mathenge, N., Merlini, P. A., Lawrence-Wright, M. B., & Giugliano, R. P. (2022). Acute coronary syndromes. *Lancet (London, England)*, *399*(10332), 1347–1358. https://doi.org/10.1016/S0140-6736(21)02391-6

Boyette, L. C., & Manna, B. (2022). Physiology, Myocardial Oxygen Demand. In

StatPearls. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK499897/

Berman DS, Arnson Y, Rozanski A. Coronary artery calcium scanning: the Agatston

score and beyond. JACC: Circ. Cardiovasc. Imaging. 2016 Dec;9(12):1417-9.

Brilakis, E. (2021). Chapter 6—Coronary angiography. In E. Brilakis (Ed.), *Manual of*

Percutaneous Coronary Interventions (pp. 97–109). Academic Press.

https://doi.org/10.1016/B978-0-12-819367-9.00006-8

Budoff, M. J., Lakshmanan, S., Toth, P. P., Hecht, H. S., Shaw, L. J., Maron, D. J., Michos,

E. D., Williams, K. A., Nasir, K., Choi, A. D., Chinnaiyan, K., Min, J., & Blaha, M.

(2022). Cardiac CT angiography in current practice: An American society for preventive

cardiology clinical practice statement☆. *American Journal of Preventive Cardiology*, *9*,

100318. https://doi.org/10.1016/j.ajpc.2022.100318

Cassar, A., Holmes, D. R., Rihal, C. S., & Gersh, B. J. (2009). Chronic Coronary Artery

Disease: Diagnosis and Management. *Mayo Clinic Proceedings*, *84*(12), 1130–1146.

https://doi.org/10.4065/mcp.2009.0391

Cirakoglu, O. F., Aslan, A. O., Akyuz, A. R., Kul, S., Şahin, S., Korkmaz, L., & Sayın, M.

R. (2019). The value of syntax score to predict new-onset atrial fibrillation in patients with

acute coronary syndrome. *Annals of Noninvasive Electrocardiology : The Official Journal*

of the International Society for Holter and Noninvasive Electrocardiology, Inc, *24*(4),

e12622. https://doi.org/10.1111/anec.12622

Cavalcante R, Sotomi Y, Mancone M, Whan Lee C, Ahn JM, Onuma Y, et al., Impact of

the SYNTAX scores I and II in patients with diabetes and multivessel coronary disease: a

pooled analysis of patient level data from the SYNTAX, PRECOMBAT, and BEST trials.

European heart journal. 2017 Jul 1;38(25):1969-77.

Costopoulos, C., Huang, Y., Brown, A. J., Calvert, P. A., Hoole, S. P., West, N. E. J.,

Gillard, J. H., Teng, Z., & Bennett, M. R. (2017). Plaque Rupture in Coronary

Atherosclerosis Is Associated With Increased Plaque Structural Stress. *Jacc.*

Cardiovascular Imaging, *10*(12), 1472–1483. https://doi.org/10.1016/j.jcmg.2017.04.017

Eickhoff, M., Schüpke, S., Khandoga, A., Fabian, J., Baquet, M., Jochheim, D.,

Grundmann, D., Thienel, M., Bauer, A., Theiss, H., Brunner, S., Hausleiter, J., Massberg,

S., & Mehilli, J. (2018). Age-dependent impact of the SYNTAX-score on longer-term

mortality after percutaneous coronary intervention in an all-comer population. *Journal of*

Geriatric Cardiology : JGC, *15*(9), 559–566. https://doi.org/10.11909/j.issn.1671-

5411.2018.09.009

Gillen, C., & Goyal, A. (2022). Stable Angina. In *StatPearls*. StatPearls Publishing.

http://www.ncbi.nlm.nih.gov/books/NBK559016/

Godoy LC, Lawler PR, Farkouh ME, Hersen B, Nicolau JC, Rao V. Urgent

revascularization strategies in patients with diabetes mellitus and acute coronary

syndrome. CJC. 2019 Aug 1;35(8):993-1001.

Gallo M, Blitzer D, Laforgia PL, Doulamis IP, Perrin N, Bortolussi G, et al., Percutaneous

coronary intervention versus coronary artery bypass graft for left main coronary artery

disease: a meta-analysis. J. Thorac. Cardiovasc. 2022 Jan 1;163(1):94-105.

Ginghina, C., Ungureanu, C., Vladaia, A., Popescu, B. A., & Jurcut, R. (2009). The

electrocardiographic profile of patients with angina pectoris. *Journal of Medicine and Life*,

2(1), 80–91.

He, Y.-M., Shen, L., & Ge, J.-B. (2020). Fallacies and Possible Remedies of the SYNTAX

Score. *Journal of Interventional Cardiology*, *2020*, 8822308.

https://doi.org/10.1155/2020/8822308

He J, Zhao H, Yu X, Li Q, Lv S, Chen F, et al., SYNTAX score predicts long-term

mortality in patients who underwent left main percutaneous coronary intervention treated

with second-generation drug-eluting stents. International heart journal. 2017;58(3):344-50.

Hickam, D. H. (1990). Chest Pain or Discomfort. In H. K. Walker, W. D. Hall, & J. W.

Hurst (Eds.), *Clinical Methods: The History, Physical, and Laboratory Examinations* (3rd

ed.). Butterworths. http://www.ncbi.nlm.nih.gov/books/NBK416/

Kim, C. (2019). Risk Stratification by SYNTAX Score Systems in Current Percutaneous

Revascularization Era. *Korean Circulation Journal*, *50*(1), 35–37.

https://doi.org/10.4070/kcj.2019.0348

Kim, Y.-H., Park, D.-W., Kim, W.-J., Lee, J.-Y., Yun, S.-C., Kang, S.-J., Lee, S.-W., Lee,

C. W., Park, S.-W., & Park, S.-J. (2010). Validation of SYNTAX (Synergy between PCI

with Taxus and Cardiac Surgery) score for prediction of outcomes after unprotected left

main coronary revascularization. *JACC. Cardiovascular Interventions*, *3*(6), 612–623.

https://doi.org/10.1016/j.jcin.2010.04.004

Kirresh, A., White, L., Mitchell, A., Ahmad, S., Obika, B., Davis, S., Ahmad, M., &

Candilio, L. (2022). Radiation-induced coronary artery disease: A difficult clinical

conundrum. *Clinical Medicine*, *22*(3), 251–256. https://doi.org/10.7861/clinmed.2021-0600

Kristin Newby, L. (2010). Chapter 25—Acute Coronary Syndromes. In G. S. Ginsburg &

H. F. Willard (Eds.), *Essentials of Genomic and Personalized Medicine* (pp. 303–312).

Academic Press. https://doi.org/10.1016/B978-0-12-374934-5.00025-8

Kurtul BE, Kurtul A, Yalçın F. Predictive value of the SYNTAX score for diabetic

retinopathy in stable coronary artery disease patients with a concomitant type 2 diabetes

mellitus. J. Diabetes Res. 2021 Jul 1;177:108875.

Kose N, Akin F, Yildirim T, Ergun G, Altun I. The association between the lymphocyte-to-
monocyte ratio and coronary artery disease severity in patients with stable coronary artery
disease. Eur Rev Med Pharmacol Sci. 2019 Mar 1;23(6):2570-5.

Kumar, A., & Cannon, C. P. (2009). Acute Coronary Syndromes: Diagnosis and
Management, Part I. *Mayo Clinic Proceedings*, *84*(10), 917–938.

Layne, K., & Ferro, A. (2017). Antiplatelet Therapy in Acute Coronary Syndrome.
European Cardiology Review, *12*(1), 33–37. https://doi.org/10.15420/ecr.2016:34:2

Lee, K. Y., Hwang, B.-H., Kim, C. J., Sa, Y. K., Choi, Y., Kim, J.-J., Choo, E.-H., Lim, S.,
Choi, I. J., Park, M.-W., Oh, G. C., Yang, I.-H., Yoo, K. D., Chung, W. S., & Chang, K.
(2022). Prognostic Impact of the HFA-PEFF Score in Patients with Acute Myocardial
Infarction and an Intermediate to High HFA-PEFF Score. *Journal of Clinical Medicine*,
11(15), 4589. https://doi.org/10.3390/jcm11154589

Li, X.-Q., Yin, C., Li, X.-L., Wu, W.-L., & Cui, K. (2021). Comparison of the prognostic
value of SYNTAX score and clinical SYNTAX score on outcomes of Chinese patients
underwent percutaneous coronary intervention. *BMC Cardiovascular Disorders*, *21*, 334.
https://doi.org/10.1186/s12872-021-02144-w

Manda, Y. R., & Baradhi, K. M. (2022). Cardiac Catheterization Risks and Complications.
In *StatPearls*. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK531461/

Matsoukis IL, Karanasos A, Patsa C, Anousakis-Vlachochristou N, Triantafyllou K,
Kantzanou M. et al., Long-term clinical outcomes of coronary artery bypass   graft
surgery compared to those of percutaneous coronary intervention with   second
generation drug eluting stents in patients with stable angina and an   isolated lesion in the
proximal left anterior descending artery. Catheter   Cardiovasc   Interv.   2021
Sep;98(3):447-57.

Mo, Y., & Xing, B. (2021). Correlation between coronary CTA-SYNTAX score and

invasive coronary angiography-SYNTAX score. *Zhong Nan Da Xue Xue Bao. Yi Xue Ban*

= *Journal of Central South University. Medical Sciences*, *46*(8), 884–888.

https://doi.org/10.11817/j.issn.1672-7347.2021.200837

null, null, Braunwald, E., Antman, E. M., Beasley, J. W., Califf, R. M., Cheitlin, M. D.,

Hochman, J. S., Jones, R. H., Kereiakes, D., Kupersmith, J., Levin, T. N., Pepine, C. J.,

Schaeffer, J. W., Smith, E. E., Steward, D. E., Theroux, P., Gibbons, R. J., Alpert, J. S.,

Eagle, K. A., … Smith, S. C. (2000). ACC/AHA Guidelines for the Management of

Patients With Unstable Angina and Non–ST-Segment Elevation Myocardial Infarction:

Executive Summary and Recommendations. *Circulation*, *102*(10), 1193–1209.

https://doi.org/10.1161/01.CIR.102.10.1193

Pakkal, M., Raj, V., & Mccann, G. P. (2011). Non-invasive imaging in coronary artery

disease including anatomical and functional evaluation of ischaemia and viability

assessment. *The British Journal of Radiology*, *84*(Spec Iss 3), S280–S295.

https://doi.org/10.1259/bjr/50903757

Parikh, R., Patel, A., Lu, B., Senapati, A., Mahmarian, J., & Chang, S. M. (2020). Cardiac

Computed Tomography for Comprehensive Coronary Assessment: Beyond Diagnosis of

Anatomic Stenosis. *Methodist DeBakey Cardiovascular Journal*, *16*(2), 77–85.

https://doi.org/10.14797/mdcj-16-2-77

Ramjattan, N. A., Lala, V., Kousa, O., & Makaryus, A. N. (2022). Coronary CT

Angiography. In *StatPearls*. StatPearls Publishing.

http://www.ncbi.nlm.nih.gov/books/NBK470279/

Rehman, I., Kerndt, C. C., & Rehman, A. (2022). Anatomy, Thorax, Heart Left Anterior

Descending (LAD) Artery. In *StatPearls*. StatPearls Publishing.

http://www.ncbi.nlm.nih.gov/books/NBK482375/

Safarian, H., Alidoosti, M., Shafiee, A., Salarifar, M., Poorhosseini, H., & Nematipour, E.

(2014a). The SYNTAX Score Can Predict Major Adverse Cardiac Events Following

Percutaneous Coronary Intervention. *Heart Views : The Official Journal of the Gulf Heart*

Association, *15*(4), 99–105. https://doi.org/10.4103/1995-705X.151081

Sandeelo IK, Ahmed F, Kakepoto N, Shah MA, Parkash C, Ali H. To Assess the Beneficial

Effects of Rosuvastatin Before Percutaneous Coronary Intervention in Patients with Acute

Coronary Syndrome. Nat. Edi. Advi. Brd. 2019 Oct;30(10):60-4.

Safarian, H., Alidoosti, M., Shafiee, A., Salarifar, M., Poorhosseini, H., & Nematipour, E.

(2014b). The SYNTAX Score Can Predict Major Adverse Cardiac Events Following

Percutaneous Coronary Intervention. *Heart Views: The Official Journal of the Gulf Heart*

Association, *15*(4), 99–105. https://doi.org/10.4103/1995-705X.151081

Salvatore, A., Boukhris, M., Giubilato, S., Tomasello, S. D., Castaing, M., Giunta, R.,

Marzà, F., Abdelbasset, H. M., Khamis, H., & Galassi, A. R. (2016). Usefulness of

SYNTAX score II in complex percutaneous coronary interventions in the setting of acute

coronary syndrome. *Journal of the Saudi Heart Association*, *28*(2), 63–72.

https://doi.org/10.1016/j.jsha.2015.07.003

Sarkees, M. L., & Bavry, A. A. (2009). Acute coronary syndrome (unstable angina and

non-ST elevation MI). *BMJ Clinical Evidence*, *2009*, 0209.

Scherff, F., Vassalli, G., Sürder, D., Mantovani, A., Corbacelli, C., Pasotti, E., Klersy, C.,

Auricchio, A., Moccetti, T., & Pedrazzini, G. B. (2011). The SYNTAX score predicts early

mortality risk in the elderly with acute coronary syndrome having primary PCI. *The*

Journal of Invasive Cardiology, *23*(12), 505–510.

Serruys, P. W., Onuma, Y., Garg, S., Sarno, G., van den Brand, M., Kappetein, A.-P., Van

Dyck, N., Mack, M., Holmes, D., Feldman, T., Morice, M.-C., Colombo, A., Bass, E.,

Leadley, K., Dawkins, K. D., van Es, G.-A., Morel, M.-A. M., & Mohr, F. W. (2009).

Assessment of the SYNTAX score in the Syntax study. *EuroIntervention: Journal of*

EuroPCR in Collaboration with the Working Group on Interventional Cardiology of the

European Society of Cardiology, *5*(1), 50–56. https://doi.org/10.4244/eijv5i1a9

Shahjehan, R. D., & Bhutta, B. S. (2022). Coronary Artery Disease. In *StatPearls*.

StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK564304/

Shalev, A., Nakazato, R., Arsanjani, R., Nakanishi, R., Park, H.-B., Otaki, Y., Cheng, V.

Y., Gransar, H., LaBounty, T. M., Hayes, S. W., Berman, D. S., & Min, J. K. (2016).

SYNTAX Score Derived From Coronary CT Angiography for Prediction of Complex

Percutaneous Coronary Interventions. *Academic Radiology*, *23*(11), 1384–1392.

https://doi.org/10.1016/j.acra.2016.07.003

Suh, Y. J., Han, K., Chang, S., Kim, J. Y., Im, D. J., Hong, Y. J., Lee, H.-J., Hur, J., Kim,

Y. J., & Choi, B. W. (2017). SYNTAX score based on coronary computed tomography

angiography may have a prognostic value in patients with complex coronary artery disease.

Medicine, *96*(37), e7999. https://doi.org/10.1097/MD.0000000000007999

Sukhija, R., Aronow, W. S., Kakar, P., Garza, L., Sachdeva, R., Sinha, A., & Mehta, J. L.

(2006). Relation of microalbuminuria and coronary artery disease in patients with and

without diabetes mellitus. *The American Journal of Cardiology*, *98*(3), 279–281.

Takagi, K., Tanaka, A., Yoshioka, N., Morita, Y., Yoshida, R., Kanzaki, Y., Watanabe, N.,

Yamauchi, R., Komeyama, S., Sugiyama, H., Shimojo, K., Imaoka, T., Sakamoto, G., Ohi,

T., Goto, H., Ishii, H., Morishima, I., & Murohara, T. (2021). In-hospital mortality among

consecutive patients with ST-Elevation myocardial infarction in modern primary

percutaneous intervention era ~ Insights from 15-year data of single-center hospital-based

registry ~. *PLoS ONE*, *16*(6), e0252503. https://doi.org/10.1371/journal.pone.0252503

Thuijs DJ, Kappetein AP, Serruys PW, Mohr FW, Morice MC, Mack MJ, et al.

Percutaneous coronary intervention versus coronary artery bypass grafting in patients with,

three-vessel or left main coronary artery disease: 10-year follow-up of the multicentre

randomised controlled SYNTAX trial. The Lancet. 2019 Oct 12;394(10206):1325-34.

Villa, A. D., Sammut, E., Nair, A., Rajani, R., Bonamini, R., & Chiribiri, A. (2016).

Coronary artery anomalies overview: The normal and the abnormal. *World Journal of*

Radiology, *8*(6), 537–555. https://doi.org/10.4329/wjr.v8.i6.537

Van Nguyen H, Khuong LQ, Nguyen AT, Nguyen AL, Nguyen CT, Nguyen HT, et al.,

Changes in, and predictors of, quality of life among patients with unstable angina after

percutaneous coronary intervention. J Eval Clin Pract. 2021 Apr;27(2):325-32.

Vizirianakis IS, Chatzopoulou F, Papazoglou AS, Karagiannidis E, Sofidis G, Stalikas

N, et al., The GEnetic Syntax Score: a genetic risk assessment implementation tool

grading the complexity of coronary artery disease—rationale and design of the GESS

study. BMC Cardiovascular Disorders. 2021 Dec;21(1):1-9.

Xu W, Guan H, Gao D, Pan J, Wang Z, Alam M, et al. Sex-specific association of

monocyte count to high-density lipoprotein ratio with SYNTAX score in patients with

suspected stable coronary artery disease. Medicine. 2019 Oct;98(41).

Yang, C., Deng, Z., Li, J., Ren, Z., & Liu, F. (2021). Meta-analysis of the relationship

between interleukin-6 levels and the prognosis and severity of acute coronary syndrome.

Clinics, *76*, e2690. https://doi.org/10.6061/clinics/2021/e2690

Yüceler, Z., Kantarcı, M., Tanboğa, İ. H., Sade, R., Kızrak, Y., Pirimoğlu, B., Bayraktutan,

Ü., Oğul, H., & Aksakal, E. (2016). Coronary lesion complexity assessed by SYNTAX

score in 256-slice dual-source MDCT angiography. *Diagnostic and Interventional*

Radiology, *22*(4), 334–340. https://doi.org/10.5152/dir.2015.15298

Zhang, Y., Iqbal, J., Campos, C., Klaveren, D., Bourantas, C., Dawkins, K., Banning, A.,

Escaned, J., Vries, T., Morel, M.-A., Farooq, V., Onuma, Y., Garcia-Garcia, H., Stone, G.,

Steyerberg, E., Mohr, F., & Serruys, P. (2014). Prognostic Value of Site SYNTAX Score

and Rationale for Combining Anatomic and Clinical Factors in Decision Making. *Journal*

of the American College of Cardiology, *64*, 423–432.

https://doi.org/10.1016/j.jacc.2014.05.022

Table 1. Baseline characteristics of study participants		
Variables	**mean ± SD***	**median (minimum-maximum)**
Age (years)	59.2 ± 10.6	60.0 (29-80)
Body mass index (kg/m2	27.2 ± 3.5	N/A
SYNTAX score	26.1 ± 11.3	29 (2-58)
Sex		
	Male	111 (71.2)
	Female	45 (28.8)
Groups		
	Group-A Stable angina	78 (50.0)
	Group-B ACS	78 (50.0)
Smoking status		
	Non-smoker	128 (82.1)
	Smoker	28 (17.9)
Dyslipidemia		
	Absent	116 (74.4)
	Present	40 (25.6)
Hypertension		
	Absent	38 (24.4)
	Present	118 (75.6)
Diabetes Mellitus		
	Absent	79 (50.6)
	Present	77 (49.4)

Percutaneous coronary intervention (PCI).		
	Absent	41 (90.4)
	Present	5 (9.6)

Table 2. Bifurcation of demographic variables with respect to Group-A Stable angina versus Group-B acute coronary syndrome.

Variable Categories	Group-A stable angina	Group-B acute coronary syndrome	p-value
Age (years)			0.26
mean ± SD*	60.2 ± 10.1	58.3 ± 11.0	
Body mass index (kg/m2)			0.58
mean ± SD*	27.1 ± 2.2	27.4 ± 4.4	
SYNTAX Score			0.01
mean ± SD*	23.8 ± 12.4	28.2 ± 9.9	
Sex			0.60
Male	57 (51.4)	54 (48.6)	
Female	21 (46.7)	24 (53.3)	
Smoking status			0.40
Non-smoker	66 (51.6)	62 (48.4)	
Smoker	12 (42.9)	16 (57.1)	
Dyslipidemia			0.01
Absent	74 (63.8)	42 (36.2)	
Present	4 (10.0)	36 (90.0)	
Hypertension			0.46
Absent	17 (44.7)	21 (55.3)	
Present	61 (51.7)	57 (48.3)	
Diabetes mellitus			0.26
Absent	43 (54.4)	36 (45.6)	
Present	35 (45.5)	42 (54.5)	
Percutaneous coronary intervention			0.78
Absent	71 (50.4)	70 (49.6)	
Present	7 (46.7)	8 (53.3)	
standard deviation.			

Figure 1. Comparison of comorbidities, prior PCI history, and mean SYNTAX scores

between Group A (Stable Angina) and Group B (Acute Coronary Syndrome, ACS).

The figure illustrates the prevalence of dyslipidemia, diabetes, hypertension, and

smoking status, along with previous PCI history and the mean SYNTAX score in each

group. Patients with ACS demonstrated a notably higher prevalence of dyslipidemia

and smoking, as well as higher mean SYNTAX scores compared to those with stable

angina

Figure 2. Comparison of mean age, BMI, and SYNTAX score (mean ± SD) between

Group A (Stable Angina) and Group B (Acute Coronary Syndrome, ACS). No

significant differences were observed in age and BMI between groups (ns = not

significant). However, the mean SYNTAX score was significantly higher in the ACS

group compared to the stable angina group ($p < 0.05$). Error bars indicate standard

deviation