Full title: Optimising supervised machine learning algorithms predicting cigarette cravings

and lapses for a smoking cessation just-in-time adaptive intervention (JITAI)

Short title: Predicting high-risk moments in smoking cessation using machine learning

Corinna Leppin1, Jamie Brown1, Claire Garnett1,2, Dimitra Kale1, Tosan Okpako1, David

Simons3, Olga Perski1,4

1 Department of Behavioural Science and Health, University College London, London, UK

2 School of Psychological Science, University of Bristol, Bristol, UK

3 Department of Anthropology, The Pennsylvania State University, University Park, PA, USA

4Department of Psychology, Stockholm University, Stockholm, Sweden*Corresponding

author: Corinna Leppin, Department of Behavioural Science and Health, University College

London, 1-19 Torrington Place, London WC1E 7HB, UK. corinna.leppin.21@ucl.ac.uk

Statement of competing interests: OP and JB act as unpaid scientific advisors to the

Smoke Free app. They do not receive salaries or any other remuneration for this work. CG,

DK, CL, TO, and DS have no competing interests to declare.

Funding: This analysis is part of a project funded by Cancer Research UK (PRCRPG-

Nov21\100002; https://www.cancerresearchuk.org/funding-for-researchers/our-funding-

schemes/prevention-and-population-research-programme-awards). CG is supported by a

National Institute for Health and Care Research (NIHR) Advanced Fellowship (302923;

https://www.nihr.ac.uk/career-development/research-career-funding-programmes). OP is

supported by HORIZON TMA MSCA Postdoctoral Fellowships - Global Fellowships from the

European Union (Grant agreement number: 101065293; https://ec.europa.eu/info/funding-

tenders/opportunities/portal/screen/opportunities/projects-details/43108390/101065293) and

by the European Research Council (ERC-2025-StG) under Horizon Europe (HORIZON)

(Grant Agreement number: 101219875; https://ec.europa.eu/info/funding-

tenders/opportunities/portal/screen/opportunities/projects-details/43108390/101219875)

Views and opinions expressed are however those of the author(s) only and do not

necessarily reflect those of the European Union. The European Union cannot be held

responsible for them. The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript. There was no additional external

funding received for this study. **CRediT authorship statement:**

Corinna Leppin: Conceptualization, Data curation, Formal analysis, Investigation,

Methodology, Validation, Visualization, Writing - original draft, Writing - original draft

Jamie Brown: Conceptualization, Funding acquisition, Supervision, Writing - review & editing

Claire Garnett: Conceptualization, Supervision, Writing - review & editing

Dimitra Kale: Investigation, Project administration, Writing - review & editing

Tosan Okpako: Investigation, Writing - review & editing

David Simons: Data curation, Visualization, Writing - review & editing

Olga Perski: Conceptualization, Data curation, Investigation, Methodology, Project

administration, Supervision, Validation, Writing - review & editing

Abstract

This study aimed to optimise the balance between participant burden and algorithm performance for predicting high-risk moments in a smoking cessation just-in-time adaptive intervention (JITAI) by systematically varying ecological momentary assessment (EMA) prompt frequency, predictor count, and training data source.

Thirty-seven participants completed 16 EMAs per day for the first 10 days of their smoking cessation attempt, reporting mood, context, behaviour, cravings, and smoking lapses. Random forest algorithms predicting lapses and cravings were evaluated in terms of F1-score and ROC-AUC via mixed effects models accounting for clustering within individuals

Performance across out-of-sample individuals ranged from excellent to poor but was, on average, modest. Lapse prediction outperformed craving prediction, particularly for ROC-AUC (Median F1-score: Lapses 0.436 [IQR 0.180–0.625], Cravings 0.400 [IQR 0.048–0.649]; Median ROC-AUC: Lapses 0.659 [IQR 0.514–0.809], Cravings 0.628 [IQR: 0.510–0.729]). A substantial proportion of configurations fell below commonly used minimum performance thresholds, particularly for F1-score.

Reducing EMA frequency had outcome- and metric-dependent effects. Lapse F1-scores improved with fewer prompts (16 EMAs: 0.254 [IQR 0.081–0.500], 3 EMAs: 0.588 [IQR 0.353–0.667]), while ROC-AUC showed a slight, inconsistent decline (16 EMAs: 0.661 [IQR 0.520–876], 4 EMAs: 0.613 [IQR 0.494–0.786], 3 EMAs: 0.704 [IQR 0.567–0.809]). For cravings, both metrics declined with fewer prompts (F1-score: 16 EMAs: 0.470 [IQR 0.141–0.745]; 3 EMAs: 0.333 [IQR 0.000–0.600]; ROC-AUC: 16 EMAs 0.700 [IQR 0.582–0.811], 3 EMAs 0.544 [IQR 0.421–0.676]). Feature reduction had negligible impact on lapse prediction (F1-score: all features 0.435, selected features 0.441; ROC-AUC: all 0.660, selected 0.657), but slightly reduced craving performance (F1-score: all 0.410 [IQR 0.117-0.646], selected 0.400 [IQR 0.000–0.650]; ROC-AUC: all 0.632, selected 0.622). Including participant-specific data improved lapse F1-scores (None 0.286 [IQR 0.000–0.571], 30pc 0.542 [IQR: 0.329–

0.667]), but did not ROC-AUC (None 0.655 [IQR: 0.512–0.786], 30pc 0.694 [IQR 0.513–

0.852]); and impaired craving ROC-AUC (None 0.650 [IQR: 0.544–0.734], 30pc 0.614 [IQR

0.493–0.730]; F1-score: None 0.424 [IQR 0.143–0.649], 30pc 0.400 [IQR 0.000–0.703]).

Overall, EMA-based machine learning detected lapse risk but showed modest overall

performance and substantial inter-individual variability. Using higher EMA density, larger

predictor sets, and participant-specific training data did not consistently outperform over

more parsimonious approaches. However, machine learning prediction alone is unlikely to

be sufficient for real-world JITAI implementation, and may be best combined with

complementary rules-based approaches.

Introduction

Smoking remains a key cause of morbidity and mortality in the UK (1,2). Improved

interventions are needed to increase smoking cessation rates and improve population health

in line with government targets (3). Traditional behavioural interventions often focus on

general motivation and coping strategies, but may be insufficient to prevent relapse (4,5).

Many people who smoke fail to actively seek out support at critical moments (6,7), and

ensuing lapses often lead to full relapses (8–11). Just-in-time adaptive interventions (JITAIs)

aim to address this gap by delivering timely, tailored support during high-risk moments

(12,13).

JITAIs rely on frequent assessment of tailoring variables (i.e., factors that may precipitate or

shape high-risk moments), actively via ecological momentary assessments (EMAs) or

passively via sensors. Decision rules (i.e., algorithms that specify how to combine data from

tailoring variables) are then used to determine opportune moments for intervention delivery

(13). While early evidence suggests smoking cessation JITAIs are feasible (14–19) and can

be effective (18), there remain questions over how to optimally balance the two (potentially

conflicting) requirements of JITAIs: collecting enough data on psychological and contextual

tailoring variables to effectively detect or predict high-risk moments and receptivity and

creating a monitoring regime that is not too burdensome for users. Balancing these two

requirements can be challenging in practice, as, all else being equal, an algorithm's ability to

detect or predict states of need and receptivity, decipher their nature, and provide

appropriate support is usually (but not always, cf. (20)) improved by more data and more

sophisticated computing techniques. However, the time and effort involved in data collection

and the drain on device battery resulting from passive sensors and complex modelling can

be a burden for users (21,22).

The minimum sampling frequency at which psychological and contextual tailoring variables

need to be assessed to collect data able to predict or detect high-risk moments well enough

to prevent smoking lapses and effectively support smoking cessation is unclear. Given the

frequency of smoking behaviour and the rapidity at which predictor variables may provoke

lapses, hourly or more frequent measurements may be necessary to accurately model

smoking dynamics (23,24). However, previous research indicates that frequent or lengthy

EMAs may reduce compliance and lead to disengagement with the intervention (25,26).

Although intensive data collection can enhance algorithm performance, most users are

unlikely to tolerate more than four to five surveys with around 10 items per day for a

prolonged period of time (16,19,21,27–30). Limiting EMA frequency and item count per EMA

may therefore be key to making a JITAI acceptable and maintaining user engagement.

JITAIs that are adaptive from the first interaction by using group-level algorithms, or

gradually personalised "warm start" hybrid-level algorithms, appear to align better with user

preferences than JITAIs that require extensive pre-quit training before providing tailored

support (31,32). While some previous evidence suggests that hybrid-level algorithms may be

better at predicting high-risk moments during smoking cessation (9,29,33), they also require

iterative retraining of the algorithms, which increases computational demand (and,

particularly if done device-side, drain device battery, and if done server-side, have privacy

risks) and adds complexity to the implementation process. This can limit their feasibility and

acceptability, so a careful trade-off with performance compared to group-level algorithms is

needed. Additionally, this is a relatively new area of research, and there is little guidance on

how long a warm-start period should be, and how the target behaviour, the dynamics of the

outcome, and the intended intervention context influence the optimal length of this warm

start period.

The current study aimed to train and test a random forest algorithm for a future smoking

cessation JITAI. For this purpose, we conducted secondary analyses of previously collected

EMA data (34) to explore how reducing the sampling frequency and the number of variables

and varying the proportion of the test participants' own data in the training set affects the

algorithm's ability to predict high-risk moments, operationalised either as lapses or as

instances of high cravings and predicted in separate algorithms, during a smoking cessation

attempt.

The specific research question was: How does the performance of random forest algorithms

predicting high-risk moments among people during the first 10 days of their smoking

cessation attempt differ according to the frequency of EMA prompts per day, the number of

predictor variables, the proportion of the test participant's own data in the training set, and

whether high-risk moments are operationalised as smoking lapses or cigarette cravings?

Materials and methods

Design

This study comprised secondary analyses of previously collected EMA data from 37 people

who smoke during the first 10 days of their quit attempt (34). The sample size was based on

a simulation informed by the available literature; for more details, see https://osf.io/ywqpv.

Participants received 16 EMAs per day (one every waking hour) to assess their mood, pain

level, activity, social and physical context, cigarette craving, caffeine, alcohol, and nicotine

consumption, and lapse incidence.

Ethical approval was obtained from the UCL Research Ethics Committee (Project ID:

15297.004). Participants were asked for their written informed consent to share their

anonymised research data with other researchers via an open science platform. The

protocol for this study was preregistered on the Open Science Framework prior to the start of

analyses (https://osf.io/tnu72/).

151   Recruitment and participants

Participants were recruited between May 2022 and March 2023, mostly through paid adverts

on Facebook and Instagram. Recruitment commenced on 20 May 2022, with the first

participant's baseline visit taking place on 27 May 2022. The last participant had their

baseline visit on 21 March 2023 and their final follow-up visit on 31 March 2023.

Eligibility requirements included being at least 18 years of age, regularly smoking cigarettes,

residing in or around London, owning a smartphone, and being willing to set a quit date

within seven days of the initial study visit. Smoking cigarettes regularly was defined as

responding "Yes" to the question "Do you smoke cigarettes at all nowadays?" and at least

one (1) to the question "How many cigarettes per day do you usually smoke?". Smoking

status was not bio-verified at baseline. Participants were remunerated a minimum of £20 for

participation, regardless of their EMA completion rate. Participants received £50 if they

completed at least 50% of EMAs on at least 50% of study days and were informed they

would earn an additional £5 for each day their EMA compliance exceeded 70%, up to a

maximum of £100. It was made clear that payment was not tied to their smoking behaviour.

166   Procedure

After completing an online screening survey to confirm eligibility, eligible participants were

invited to attend an in-person baseline visit. At the baseline visit, they completed a baseline

survey, provided written informed consent, and downloaded and received instructions for

how to use the m-Path app (35), through which they received the EMAs throughout the study

period. Participants were asked to set a quit date within the next seven days and

encouraged to set it for the following day to capitalise on their motivation to stop smoking.

Participants also received free access to the premium version of an evidence-based

smoking cessation app (36). At the end of the baseline visit, the in-person follow-up visit was

scheduled.

The 10-day study period began on each participant's selected quit date. During this time,

participants received 16 EMA prompts per day (one every waking hour on the hour or half-

hour) via the m-Path app (signal-contingent sampling), tailored to their usual waking hours.

EMAs needed to be completed within 30 minutes of receipt. Participants were also instructed

to log lapses as they occurred using an in-app button (event-contingent sampling). Each

EMA took around one to three minutes to complete and contained the same 22-25 items

(depending on whether participants chose to add additional items; these additional items are

not considered in this analysis, but more details on them can be found in the previous study

using these data (34)). At the follow-up visit, data from study-related apps were exported, the

m-Path app was deactivated, and participants received payment for their participation.

Participants who reported that they had successfully quit smoking were asked to provide

expired carbon monoxide measures using a Bedfont iCOquit smokerlyzer

(https://www.icoquit.com/) to verify their abstinence (cut-off set at ≤8 parts per million). Full

details on recruitment and procedure are reported in the previous study (34).

Measures

Overall, the machine learning algorithms in this study used data from 34 variables: 12

between-person variables collected at the baseline questionnaire, and 22 within-person

variables collected at every EMA. Full questionnaires and variable lists cand be found in S1

Appendix.

The between-person variables were collected at baseline and comprised two numerical

variables (cigarettes per day; age in years), four binary variables (gender [four levels were

available, but all participants selected either "man" or "woman"); post-16 education; ever-use

of behavioural support for smoking reduction or cessation; ever-use of pharmacological

support for smoking reduction or cessation, and six categorical variables (occupation type;

ethnicity; past quit attempt; time to first cigarette; motivation to stop scale (37); season of

data collection). The within-person variables comprised 14 quasi-continuous variables

measured on 11-point Likert scales (ten variables measuring affect informed by the

circumplex model of affect (38–40): sad, irritable, stressed, anxious, bored, calm, content,

happy, excited, enthusiastic; bodily pain; cigarette cravings; momentary motivation not to

smoke; quitting-related self-efficacy), four binary variables (past-hour caffeine consumption;

past-hour alcohol consumption; past-hour nicotine use other than tobacco use/smoking;

past-hour lapse [i.e., smoking), and four categorical variables (social context; activity;

location; cigarette availability). For training and evaluating the algorithms, categorical

variables were recoded into dummy variables.

In prior formative work using these data, we explored more complex representations of

lapses and other time-varying predictors, including variables that incorporated longer-term

smoking history and cumulative consumption or moving averages over extended windows.

These more complex features did not meaningfully improve algorithm performance relative

to the simpler representations used here and were therefore not pursued further in the

present analyses.

Analyses
216

Data pre-processing
217

We used the pre-processed dataset from the previous study (34). In this pre-processed

dataset, eight out of 46 participants with insufficient EMA adherence (defined as completion

of ≤60% of prompts) were excluded. This adherence threshold was applied to ensure

sufficient data density to support reliable estimation of temporal associations and the training

and evaluation of machine-learning models relying on lagged EMA predictors. This criterion

was also retained for models using fewer prompts/observations, as these were derived from

the same intensive EMA protocol and still required adequate coverage across days and

contexts. Including participants only in reduced-prompt conditions but not in full-prompt

models would have created validity issues in the imputation process, introduced additional

selection biases, and reduced comparability across algorithm specifications. Missing data in

the remaining analytical sample were imputed using the univariate Kalman filter, which is

recommended for univariate time series data imputation (41) and has been previously used

in analysis of intensive longitudinal data on health behaviours (42). As a forward filter that

only uses past and current information and does not consider future states, it is also suitable

for implementation within an eventual JITAI.

Model specification

This study used group- and hybrid-level random forest (RF) algorithms to predict (a) lapses

by or (b) cravings (dichotomised; high ≥ 7 on an 11-point Likert scale from 0 to 10) at one

EMA based on levels of the predictor variables at the previous EMA and the values of the

baseline variables. The two outcomes were predicted using separate algorithms. RF

algorithms are a machine learning method that combines the results of many decision trees

to make more accurate and reliable predictions (43). A decision tree is a flowchart-like model

that splits the data into branches based on values of the predictor variables, aiming to group

similar outcomes together (43). In an RF, each tree is built using a random sample of the

data and a random subset of predictors, which helps reduce overfitting and improves the

model's ability to generalise to new data (43). For every outcome by number of EMAs by set

of predictors combination, the number of candidate predictors at each split and the minimum

number of observations in a terminal node was tuned at the group-level, while the number of

trees in the forest was set at 500, in line with recommendations for training RF algorithms

(44–46). As is standard, the classification threshold was set at 0.5, such that predicted

probabilities ≥0.5 were classified as positive cases/events (lapse or high cravings) and

probabilities < 0.5 as negative cases/non-events (non-lapses or low/no cravings. All

analyses were conducted using R (47), and the models were trained and tested using the

tidymodels framework (48). The complete analysis syntax and data can be found on Github

(https://github.com/OlgaPerski/EMASENS_ML/tree/main) and the Open Science Framework

(https://osf.io/tnu72/).

Selection of algorithm types

As the aim of this study was to train and test an algorithm for use in a future smoking

cessation JITAI, we tested a single algorithm type for pragmatic reasons. The choice of

algorithm architecture was informed by a previous study using the same dataset (34), in

which multiple algorithm architectures (RF, Support Vector Machine, Penalised Logistic

Regression, and XGBoost) were systematically compared. In that earlier work, an RF

architecture showed the most favourable balance performance across area under the

receiver operating characteristic curve (ROC-AUC; overall ability to distinguish between

classes), sensitivity (true positive rate), and specificity (true negative rate). Additionally, RF

algorithms have modest computational demands, are well suited for feature selection, do not

make assumptions about the shape of the relationship between predictors and outcomes,

and can handle a mix of continuous and categorical predictors as well as complex

interactions (43,49–52). Therefore, we decided to take RF algorithms in this study and did

not re-compare different algorithm classes. Instead, we held the algorithm type constant

(RF) and examined how different *specification decisions* (described in more detail in the

following paragraphs) affected performance.

Number and timing of prompts

We fit algorithms based on the data from 3, 4, 5, 6, and all 16 prompts per day. With a view

to JITAI implementation considerations, we used quasi-random prompts evenly spaced

throughout the day. To do this, we randomly selected prompts from blocks within the

participant's daily schedule. The number of prompts to be tested was based on previous

literature (16,19,21,27–30,53). The prompts chosen for each number of prompts per day

were as follows:

• 3: randomly choose one of 1-5, 6-11, 12-16

• 4: randomly choose one of 1-4, 5-8, 9-12, 13-16

• 5: randomly choose one of 1-3, 4-6, 7-9, 10-13, 14-16

• 6: randomly choose one of 1-3, 4-6, 7-8, 9-11, 12-14, 15-16

• All 16 prompts

When selecting the reduced number of prompts, we also included all event-based EMA

reports (i.e., when a participant reported smoking outside of a prompt), regardless of the

number of daily prompts tested.

Outcome definitions

The different outcomes, lapses and high cravings, were predicted using separate algorithms.

We defined lapses as a lapse event that occurred between a participant's current and the

next randomly selected/event-based EMA. The EMA was counted as a lapse if any lapse

event was observed between these prompts. For high cravings, we recorded whether a high

craving event was reported at the next randomly selected prompt. Additionally, we calculated

prior lapses (as a predictor) by checking if a lapse occurred between the previous and

current selected prompts.

High cravings were derived from an 11-point Likert cale (0-10) measuring self-reported

craving, using a cut-off of ≥7. Cravings were dichotomised to align the modelling approach

with the intended downstream use of the algorithm as part of a JITAI, where the primary aim

is to identify discrete high-risk moments at which to deliver support. The cut-off of ≥7 was

chosen for two complementary reasons. Firstly, it is consistent with the cut-off used in the

previous analysis of the same dataset (34), facilitating comparability across studies and

allowing interpretation of results within an established analytic framework. Secondly, a

threshold of ≥7 captured approximately one third of craving observations across prompt

frequencies (more details below), providing a pragmatic balance between identifying

clinically meaningful high-craving states, maintaining sufficient class prevalence for stable

model training and evaluation, and balancing burden with responsiveness.

In line with the previous study (34) and to allow for the calculation of the performance

metrics, only participants with at least one observation in each outcome class (i.e., lapse and

non-lapse for lapse-prediction algorithms, and high craving and low/no craving for craving-

prediction algorithms) were included. When constructing hybrid-level algorithms, participants

were additionally required to have at least one observation in each outcome class in the

testing set.

Number of predictors

We used data-driven feature selection to reduce the number of predictors. Feature selection

describes machine learning techniques that aim to reduce the number of input dimensions

while preserving the algorithm's performance (54). In smoking cessation, filter (55) and

embedded (30) feature selection methods have been previously used to improve the

parsimony of models predicting smoking urges and lapses. The feature selection technique

we used is recursive feature elimination with cross-validation (RFE-CV). RFE is an

embedded feature selection method that is well-suited in this case, as it performs well even

in the presence of correlated predictors (56), can handle both continuous and categorical

predictors together and structured features, and does not have the same bias towards high

cardinality features as some other methods. RFE-CV selects an optimal number of features

without the need for pre-specification (57) and has been shown to have fairly high stability

(58). RFE-CV has shown promise in the epidemiology and healthcare fields (59–61). To

compare algorithms with fewer predictors to those with the full set of 57 predictors, we

compared the performance of algorithms using the optimal set of predictors to that of

algorithms with the full set of predictors. In cases where the optimal set of predictors was the

full set, we compared the performance of algorithms using the top 15 predictors as identified

by RFE-CV with that of algorithms using the full set of predictors. The full list of predictors

used for each algorithm is provided in S2 Appendix.

Proportion of own data in the training set

For this study, group-level algorithms and hybrid-level algorithms were used to predict lapses

and high cravings, with the latter incorporating the initial day or days of the test participant's

own data, along with the data from the other participants, in the training set. Including a

participant's data in the training set means using a portion of their data to help the machine

learning model learn patterns during the training phase, before generating predictions on the

participant's later (unseen) data, which is reserved for testing the model's performance. With

a view to how a JITAI would be implemented in a real-world setting and to retain adequate

amounts of data in the testing set, the specific amount of the participant's own data in the

training set was varied as 10% (first day), 20% (first two days), and 30% (first three days).

Performance

Algorithm performance was evaluated using the F1-score and ROC-AUC as primary metrics

with sensitivity and specificity reported as supplementary metrics. All metrics were calculated

using a fixed probability threshold of 0.5 to convert predicted probabilities into binary class

predictions.

The F1-score summarises binary classification performance as the harmonic mean of

precision and recall (sensitivity), balancing false positives and false negatives: F1-score = 2

× (precision × recall) / (precision + recall)

Precision (positive predictive value) is defined as: Precision = True Positives / (True

Positives + False Positives)

Recall (sensitivity) is defined as: Recall = True Positives / (True Positives + False Negatives)

F1-scores range from 0 to 1, with higher values indicating better performance. Scores below

0.5 are generally not seen as acceptable. Because the F1-score penalises large imbalances

between precision and recall, the F1-score is well suited to imbalanced data.

ROC-AUC reflects the model's overall ability to discriminate between positive and negative

classes across all possible thresholds. It can be interpreted as the probability that a

randomly selected positive observation receives a higher predicted risk than a randomly

selected negative observation. Values range from 0 to 1 with 0.5 indicating chance-level

(and therefore unacceptable) discriminatory ability and 1 indicating perfect discrimination.

Sensitivity and specificity quantify performance at the chosen threshold of 0.5.

Sensitivity (or recall, as mentioned above) is defined as: True Positives / (True Positives +

False Negatives)

Specificity is defined as: True Negatives / (True Negatives + False Positives)

Sensitivity reflects the ability to detect high-risk moments, whereas specificity reflects correct

identification of low-risk moments. For cases such as this, where the detection of high-risk

moments takes precedence, a minimum sensitivity of 0.7 and a minimum specificity of 0.5

are generally considered the boundaries for acceptable performance.

To examine patterns in performance metrics, we used generalised linear mixed models with

link functions appropriate for the distribution of each metric, implemented using the

glmmTMB and lme4 packages in R (62–64). Based on distributional analysis, we used

linear mixed models for ROC-AUC (approximately normally distributed), zero-inflated beta

regression for F1-score (bounded continuous outcome with substantial proportions of zeros),

and zero-one-inflated beta regression for sensitivity and specificity (bounded continuous

outcomes with substantial proportions of boundary values at both zero and one). . Models

included random intercepts for each participant to account for within-subject clustering, and

fixed effects for the number of prompts per day (EMA frequency), outcome type (lapses

versus cravings), use of feature selection (yes/no), and the proportion of own data in the

training set (none, 10%, 20%, 30%). Separate models were fitted for each performance

metric across outcomes, and stratified by outcome (cravings or lapses), with outcome

excluded as a fixed effect in the stratified models. For fixed effects with more than two levels,

marginal effects were estimated using the emmeans package (65) to compute pairwise

comparisons with Tukey's adjustment for multiple comparisons.

Changes from pre-registration

Two changes were made to the pre-registered protocol. The first change was the addition of

group-level algorithms. Originally, we only planned to train and evaluate hybrid-level

algorithms. However, after conducting the initial analyses, we realised that there were a

substantial number of participants who only had lapses in the first day(s) of their quit

attempt, and for whom, consequently, hybrid-level algorithms could not be constructed, and

saw that the performance of hybrid-level algorithms appeared unrelated to the amount of the

participant's own data in the training set. Therefore, we chose to additionally train and

evaluate group-level algorithms and to include the proportion of own data in the training set

as its own research question rather than as a sensitivity analysis. The second change was

related to the method of feature selection. Originally, we planned to only compare the

optimal set of predictors as determined by the RFE-CV algorithms with the full set of

predictors. However, for a substantial number of algorithm configurations, the optimal set of

predictors was all predictors. As that result would not allow us to compare the performance

of algorithms with fewer predictors with that with the full set of predictors, we chose to select

the 15 most important predictors as indicated by the RFE-CV procedure if that was the case

and compare the algorithm with only these predictors with that with the full set of predictors.

Additionally, the preregistration did not specify a primary performance metric, but we focused

on ROC-AUC and the F1-score in the main results for clarity and brevity. Additionally, the

preregistration did not specify how patterns in performance metrics would be analysed.

Given the large number of algorithm configurations (5 prompt frequencies × 2 outcomes × 2

feature selection levels × 4 levels of the test participant's own data in the training set, each

applied to every out-of-sample individual), we used mixed models appropriate for the

distribution of the respective score to summarise performance. This approach offers a more

interpretable and statistically robust overview than could be achieved through visual

inspection or comparisons of specific algorithm configurations alone.

Results

Sample description

409 The analytic sample comprises n = 37 adult people who smoke from in and around London

410 during the first 10 days of their quit attempt. Table 1 outlines participant characteristics.

411 *Table 1 Participant characteristics*

Characteristic	All participants N = 37[1]	Participants with at least one lapse N = 25[1]
Age	42.97 (14.45)	42.48 (13.80)
Gender		
Female	23 (62.2%)	15 (60.0%)
Male	14 (37.8%)	10 (40.0%)
Occupation		
Non-manual	17 (45.9%)	12 (48.0%)
Manual	9 (24.3%)	5 (20.0%)
Other (e.g., student, unemployed, retired)	11 (29.7%)	8 (32.0%)
Post-16 educational qualifications	33 (89.2%)	22 (88.0%)
Ethnicity		
Asian or Asian British (any Asian background)	3 (8.1%)	3 (12.0%)
Black, Black British, Caribbean or African (any Black, Black British or Caribbean background)	1 (2.7%)	0 (0.0%)
Mixed or multiple ethnic groups (e.g., White and Black	2 (5.4%)	2 (8.0%)

Characteristic	All participants N = 37[1]	Participants with at least one lapse N = 25[1]
African, White and Asian)		
Other ethnic group (i.e., Arab)	1 (2.7%)	1 (4.0%)
White (any White background)	30 (81.1%)	19 (76.0%)
Cigarettes per day	13.08 (6.50)	13.48 (6.72)
Time to first cigarette		
Within 5 minutes	9 (24.3%)	7 (28.0%)
6-30 minutes	16 (43.2%)	10 (40.0%)
31-60 minutes	6 (16.2%)	4 (16.0%)
After 60 minutes	6 (16.2%)	4 (16.0%)
Motivation to stop		
I don't want to stop smoking	0 (0.0%)	0 (0.0%)
I think I should stop smoking but don't really want to	2 (5.4%)	2 (8.0%)
I want to stop smoking but haven't thought about when	6 (16.2%)	5 (20.0%)
I really want to stop smoking but don't know when I will	7 (18.9%)	3 (12.0%)
I want to stop smoking and hope to soon	9 (24.3%)	5 (20.0%)
I really want to stop smoking and intend to in the next 3 months	2 (5.4%)	2 (8.0%)
I really want to stop smoking and intend to in the next month	11 (29.7%)	8 (32.0%)
Past-year quit attempt		
No, never	3 (8.1%)	2 (8.0%)
Yes, but not in the past year	16 (43.2%)	9 (36.0%)
Yes, in the past year	18 (48.6%)	14 (56.0%)
Ever use of pharmacological support (e.g., nicotine replacement therapy, varenicline, e-cigarettes)	30 (81.1%)	20 (80.0%)
Ever use of behavioural support (e.g., counselling, telephone support, website, app)	16 (43.2%)	10 (40.0%)
Smoked since quit date		
In the last 24hrs	16 (43.2%)	14 (56.0%)
1-6 days ago	4 (10.8%)	3 (12.0%)
1-3 weeks ago	5 (13.5%)	4 (16.0%)

Characteristic	All participants N = 37[1]	Participants with at least one lapse N = 25[1]
Have not smoked since my planned quit date	12 (32.4%)	4 (16.0%)
% Completed EMAs	76.86 (9.27)	76.38 (9.23)

[1]Mean (standard deviation); n (%); percentages may not always add up to 100% due to rounding

The majority of participants were female (62.2%) and of any White ethnic background

(81.1%). The mean age was 42.97 years (Standard deviation (SD) = 14.45). Prior to quitting,

participants smoked an average of 13.08 cigarettes per day (SD = 6.50). At follow-up, 43.2%

had smoked in the past 24 hours, 10.8% smoked 1-6 days ago, and 32.4% had not smoked

since their planned quit date. The average adherence to the EMAs was 76.86% (SD = 9.27).

Among participants with at least one lapse (n = 25; 67.6% of the entire analytics sample),

demographic and behavioural characteristics were similar: 60.0% were female and 76.0%

identified with any White ethnicity. Their mean age was 42.48 years (SD = 13.80), and they

smoked an average of 13.48 cigarettes per day (SD = 6.72). Despite reporting lapses during

their EMAs, 16.0% reported having remained abstinent since their planned quit date at the

follow-up visit. The average EMA adherence was 76.38% (SD = 9.23).

424 EMA description

Participants in the full analytic sample completed a total of 5,964 EMAs. Table 2 summarises

key descriptives across the different prompt frequencies (16, 6, 5, 4, and 3 prompts per day).

A version of this table restricted to participants with at least one lapse is available in S3

Appendix.

Event-contingent EMAs represented a small fraction of the total, ranging from 0.7% (16

prompts per day) to 4.0% (3 prompts per day). Lapse rates varied by prompt frequency, from

6.9% (16 prompts per day) to 20.8% (3 prompts per day). Due to the difference in

calculation, the proportion of high cravings was higher and more consistent, ranging from

35.8% (16 prompts per day) to 39.9% (3 prompts per day). The proportion of EMAs

recording lapses and high cravings varied substantially across individuals.

Table 2 Description of the observations

Prompts per day	EMAs				Lapses					High cravings				
	Number			Proportion event contingent	Overall proportion	Across individuals			Overall proportion	Across individuals				
	Total	Event contingent	Signal contingent			Median proportion	25th percentile proportion	75th percentile proportion		Median proportion	25th percentile proportion	75th percentile proportion		
16	5,964	44	5,920	0.7%	6.9%	1.2%	0.0%	7.5%	35.8%	29.6%	11.1%	52.8%		
6	2,204	44	2,160	2.0%	12.9%	3.3%	0.0%	18.2%	37.3%	30.9%	15.3%	59.4%		
5	1,742	42	1,700	2.4%	15.2%	3.0%	0.0%	26.0%	38.1%	29.5%	16.9%	53.7%		
4	1,401	41	1,360	2.9%	17.0%	3.7%	0.0%	28.8%	40.1%	29.8%	17.5%	57.7%		
3	1,063	43	1,020	4.0%	20.8%	6.6%	0.0%	32.3%	39.0%	31.2%	16.3%	55.6%		

Overall algorithm performance

Across all algorithm specifications, performance varied markedly between individuals and was centrally modest. For lapse prediction, the median F1-score was 0.436 (mean 0.409; IQR 0.180-0.625), and the median ROC-AUC was 0.659 (mean 0.652; IQR 0.514-0.809). For craving prediction, the median F1-score was 0.400 (mean 0.410; IQR 0.048-0.649), and the median ROC-AUC was 0.628 (mean 0.611; IQR 0.510-0.729). A substantial proportion of configurations fell below commonly used minimum thresholds for acceptable performance, particularly when evaluated using the F1-score (see S4 Appendix to S7 Appendix for details per metric and configuration). . A full summary of key performance measures by predictors, algorithm specifications, and participants can be found in S4 to S7 Appendices.

Research question 1: Prompt frequency

Lapses

Counter to the expectation that denser sampling improves performance, median F1-scores increased monotonically as prompt frequency decreased for lapse prediction: 16 prompts/day (median 0.254; mean 0.308; IQR 0.081-0.500), 6 prompts (median 0.440; mean 0.400; IQR 0.222-0.587), 5 prompts (median 0.444; mean 0.397; IQR 0.143-0.573), 4 prompts (median 0.475; mean 0.432; IQR 0.218-0.640), and 3 prompts/day (median 0.588; mean 0.508; IQR 0.353-0.667) (Table 3).

Zero-inflated beta-regressions of the scores confirmed this pattern: relative to 16 prompts/day, F1-scores were significantly higher at lower prompt frequencies (all p < 0.001) (Table 4). All pairwise contrasts between 16 prompts/day and lower frequencies were statistically significant after adjustment (S8 Appendix).

Table 3 Summary of F1-score by algorithm configurations (lapses)

	Number of algorithms	Median	Mean	Standard deviation	25th percentile	75th percentile	Minimum	Maximum
Lapses overall								
	780	0.436	0.409	0.283	0.180	0.625	0.000	0.986

Prompts per day								
16	158	0.254	0.308	0.274	0.081	0.500	0.000	0.857
6	156	0.440	0.400	0.262	0.222	0.587	0.000	0.945
5	156	0.444	0.397	0.277	0.143	0.573	0.000	0.955
4	156	0.475	0.432	0.285	0.218	0.640	0.000	0.986
3	154	0.588	0.508	0.281	0.353	0.667	0.000	0.962
Feature selection								
All Features	390	0.435	0.404	0.278	0.188	0.620	0.000	0.986
Selected Features	390	0.441	0.413	0.287	0.168	0.632	0.000	0.986
Proportion of own data in training set								
None	242	0.286	0.319	0.286	0.000	0.571	0.000	0.963
10%	198	0.400	0.384	0.282	0.155	0.607	0.000	0.986
20%	170	0.500	0.484	0.238	0.308	0.650	0.000	0.984
30%	170	0.542	0.490	0.278	0.329	0.667	0.000	0.982

461 *Table 4 Zero-inflated beta-regression model (with random intercepts) of F1-Score (lapses)*

Predictor	Estimate (logit scale)	Standard error	95% confidence interval (logit)	z-statistic	p-value	Sig.
Intercept	-0.785	0.203	-1.183, -0.386	-3.859	< 0.001	***
Prompts per day: 6	0.460	0.075	0.313, 0.606	6.162	< 0.001	***
Prompts per day: 5	0.411	0.074	0.266, 0.557	5.537	< 0.001	***
Prompts per day: 4	0.655	0.075	0.508, 0.801	8.753	< 0.001	***
Prompts per day: 3	0.981	0.076	0.833, 1.130	12.957	< 0.001	***
Feature selection: Selected features	0.097	0.047	0.006, 0.188	2.079	0.038	*
Share of own data: 10%	0.091	0.065	-0.037, 0.219	1.395	0.163	
Share of own data: 20%	0.198	0.065	0.072, 0.325	3.071	0.002	**
Share of own data: 30%	0.384	0.067	0.253, 0.514	5.767	< 0.001	***
Reference levels: Prompts per day = 16; Feature selection = All features; Share of own data = None						
Note: Estimate coefficients are on the logit scale (conditional component only).						
This is a zero-inflated beta model with separate processes for zeros and values between 0 and 1.						
For interpretable effect sizes (actual differences in predicted values), see pairwise comparisons on the response scale (0-1) in Appendix S8.						
Significance levels: *** p < 0.001, ** p < 0.01, * p < 0.05						

In contrast, ROC-AUC showed slightly poorer performance with reduced prompting. Median

ROC-AUCs were similar across conditions (16: median 0.661, mean 0.680, IQR 0.520-

0.876; 6: median 0.628, mean 0.637, IQR 0.498-0.791; 5: median 0.660, mean 0.646, IQR

0.510-0.795; 4: median 0.613, mean 0.631, IQR 0.494-0.786; 3: median 0.704, mean 0.665,

IQR 0.567-0.809) (Table 5). Linear mixed models indicated significantly lower ROC-AUC at

6, 5, and 4 prompts/day compared with 16 (p-values between .003 and < .001), while 3

prompts/day did not differ significantly from 16 (p = .226) (Table 6). See S9 Appendix for full

pairwise comparisons.

Table 5 Summary of ROC-AUC by algorithm configurations (lapses)

	Number of algorithms	Median	Mean	Standard deviation	25th percentile	75th percentile	Minimum	Maximum
Lapses overall								
	780	0.659	0.652	0.198	0.514	0.809	0.000	0.998
Prompts per day								
16	158	0.661	0.680	0.214	0.520	0.876	0.052	0.998
6	156	0.628	0.637	0.197	0.498	0.791	0.017	0.968
5	156	0.660	0.646	0.180	0.510	0.795	0.111	0.969
4	156	0.613	0.631	0.206	0.494	0.786	0.028	0.975
3	154	0.704	0.665	0.192	0.567	0.809	0.000	0.967
Feature selection								
All Features	390	0.660	0.656	0.197	0.518	0.811	0.000	0.998
Selected Features	390	0.657	0.648	0.200	0.508	0.808	0.000	0.994
Proportion of own data in training set								
None	242	0.655	0.644	0.199	0.512	0.786	0.017	0.994
10%	198	0.607	0.627	0.207	0.510	0.765	0.000	0.994
20%	170	0.673	0.674	0.181	0.532	0.819	0.258	0.998
30%	170	0.694	0.670	0.201	0.513	0.852	0.206	0.997

474 *Table 6 Linear mixed models of ROC-AUC (lapses)*

Predictor	Estimate	Standard error	95% confidence interval	t-statistic	p-value	Sig.
Intercept	0.677	0.039	0.601, 0.754	17.373	< 0.001	***
Prompts per day: 6	-0.042	0.011	-0.064, -0.020	-3.749	< 0.001	***

Prompts per day: 5	-0.034	0.011	-0.056, -0.012	-2.986	0.003	**
Prompts per day: 4	-0.048	0.011	-0.070, -0.026	-4.288	< 0.001	***
Prompts per day: 3	-0.014	0.011	-0.036, 0.008	-1.212	0.226	
Feature selection: Selected features	-0.009	0.007	-0.023, 0.005	-1.215	0.224	
Share of own data: 10%	-0.016	0.010	-0.036, 0.003	-1.615	0.106	
Share of own data: 20%	0.004	0.011	-0.017, 0.025	0.384	0.701	
Share of own data: 30%	0.001	0.011	-0.020, 0.021	0.057	0.954	
Reference levels: Prompts per day = 16; Feature selection = All features; Share of own data = None						
Significance levels: *** p < 0.001, ** p < 0.01, * p < 0.05						

Cravings

For craving prediction, reduced prompt frequency was associated with consistent performance declines. Median F1-scores decreased from 0.470 (mean 0.465; IQR 0.141-0.745) at 16 prompts/day to 0.447 (mean 0.453; IQR 0.211-0.658) at 6 prompts, 0.368 (mean 0.393; IQR 0.000-0.624) at 5 prompts, 0.361 (mean 0.379; IQR 0.000-0.647) at 4 prompts, and 0.333 (mean 0.358; IQR 0.000-0.600) at 3 prompts/day (Table 7). Zero-inflated beta-regressions indicated significantly lower F1-scores for all reduced-prompt conditions relative to 16 prompts/day (all p ≤ .012) (Table 8).

Table 7 Summary of F1-score by algorithm configurations (cravings)

	Number of algorithms	Median	Mean	Standard deviation	25th percentile	75th percentile	Minimum	Maximum
Cravings overall								
	1,424	0.400	0.410	0.328	0.048	0.649	0.000	0.997
Prompts per day								
16	292	0.470	0.465	0.332	0.141	0.745	0.000	0.997
6	280	0.447	0.453	0.315	0.211	0.658	0.000	0.992
5	274	0.368	0.393	0.323	0.000	0.624	0.000	0.990
4	290	0.361	0.379	0.329	0.000	0.647	0.000	0.987
3	288	0.333	0.358	0.331	0.000	0.600	0.000	0.984
Feature selection								
All Features	712	0.410	0.415	0.322	0.117	0.646	0.000	0.997
Selected Features	712	0.400	0.404	0.334	0.000	0.650	0.000	0.997
Proportion of own data in training set								
None	360	0.424	0.422	0.309	0.143	0.649	0.000	0.997

10%	356	0.419	0.412	0.315	0.123	0.633	0.000	0.997
20%	354	0.333	0.388	0.341	0.000	0.640	0.000	0.996
30%	354	0.400	0.415	0.348	0.000	0.703	0.000	0.996

486 *Table 8 Zero-inflated beta-regression model (with random intercepts) of F1-Score (cravings)*

Predictor	Estimate (logit scale)	Standard error	95% confidence interval (logit)	z-statistic	p-value	Sig.
Intercept	0.278	0.230	-0.173, 0.728	1.208	0.227	
Prompts per day: 6	-0.142	0.056	-0.253, -0.032	-2.523	0.012	*
Prompts per day: 5	-0.398	0.059	-0.514, -0.282	-6.746	< 0.001	***
Prompts per day: 4	-0.346	0.059	-0.461, -0.231	-5.882	< 0.001	***
Prompts per day: 3	-0.381	0.060	-0.499, -0.262	-6.304	< 0.001	***
Feature selection: Selected features	0.021	0.037	-0.053, 0.094	0.555	0.579	
Share of own data: 10%	-0.057	0.053	-0.160, 0.046	-1.085	0.278	
Share of own data: 20%	-0.076	0.054	-0.181, 0.029	-1.423	0.155	
Share of own data: 30%	0.077	0.053	-0.027, 0.181	1.452	0.146	
Reference levels: Prompts per day = 16; Feature selection = All features; Share of own data = None						
Note: Estimate coefficients are on the logit scale (conditional component only).						
This is a zero-inflated beta model with separate processes for zeros and values between 0 and 1.						
For interpretable effect sizes (actual differences in predicted values), see pairwise comparisons on the response scale (0-1) in Appendix S8.						
Significance levels: *** p < 0.001, ** p < 0.01, * p < 0.05						

ROC-AUC showed a similar pattern. Median ROC-AUC declined from 0.700 (mean 0.681;

IQR 0.582-0.811) at 16 prompts/day to 0.652 (mean 0.641; IQR 0.575-0.736), 0.606 (mean

0.598; IQR 0.517-0.696), 0.622 (mean 0.593; IQR 0.500-0.695), and 0.544 (mean 0.541;

IQR 0.421-0.676) at 6, 5, 4, and 3 prompts/day, respectively (Table 9). Linear mixed models

showed significantly lower ROC-AUCs at all reduced prompt frequencies (all p < .001) (Table

10).

Table 9 Summary of ROC-AUC by algorithm configurations (cravings)

	N	Median	Mean	Standard deviation	25th percentile	75th percentile	Minimum	Maximum
Cravings overall								
	1,424	0.628	0.611	0.174	0.510	0.729	0.000	1.000
Prompts per day								
16	292	0.700	0.681	0.165	0.582	0.811	0.122	0.986
6	280	0.652	0.641	0.148	0.575	0.736	0.000	0.983
5	274	0.606	0.598	0.151	0.517	0.696	0.068	0.971
4	290	0.622	0.593	0.164	0.500	0.695	0.025	1.000
3	288	0.544	0.541	0.201	0.421	0.676	0.000	1.000
Feature selection								
All Features	712	0.632	0.624	0.170	0.516	0.744	0.032	1.000
Selected Features	712	0.622	0.598	0.177	0.502	0.717	0.000	0.986
Proportion of own data in training set								
None	360	0.650	0.630	0.160	0.544	0.734	0.025	0.983
10%	356	0.619	0.607	0.178	0.504	0.722	0.000	1.000
20%	354	0.632	0.606	0.177	0.503	0.728	0.032	1.000
30%	354	0.614	0.600	0.178	0.493	0.730	0.000	1.000

496 *Table 10 Linear mixed model of ROC-AUC (cravings)*

Predictor	Estimate	Standard error	95% confidence interval	t-statistic	p-value	Sig.
Intercept	0.714	0.018	0.678, 0.749	39.470	< 0.001	***
Prompts per day: 6	-0.046	0.012	-0.070, -0.023	-3.812	< 0.001	***
Prompts per day: 5	-0.085	0.012	-0.110, -0.061	-6.964	< 0.001	***
Prompts per day: 4	-0.088	0.012	-0.111, -0.064	-7.260	< 0.001	***

Prompts per day: 3	-0.139	0.012	-0.163, -0.115	-11.489	< 0.001	***
Feature selection: Selected features	-0.026	0.008	-0.041, -0.011	-3.403	< 0.001	***
Share of own data: 10%	-0.023	0.011	-0.045, -0.002	-2.156	0.031	*
Share of own data: 20%	-0.023	0.011	-0.044, -0.002	-2.105	0.035	*
Share of own data: 30%	-0.030	0.011	-0.051, -0.008	-2.726	0.006	**
Reference levels: Prompts per day = 16; Feature selection = All features; Share of own data = None						
Significance levels: *** p < 0.001, ** p < 0.01, * p < 0.05p						

Research question 2: Feature selection

Lapses

Feature selection had minimal impact on lapse prediction. Median F1-scores were similar for models using all predictors (median 0.435; mean 0.404; IQR 0.188-0.620) and selected predictors (median 0.441; mean 0.413; IQR 0.168-0.632) (Table 3). Median ROC-AUCs were likewise similar (0.660 vs 0.657; Table 5). CLMMs indicated a statistically significant but small improvement for F1-score with feature selection (p = .038), whereas no significant effect was observed for ROC-AUC (p = .224) (Tables 4 and 6).

Cravings

For cravings, feature selection was associated with slightly worse performance. Median F1-score decreased from 0.410 (mean 0.415; IQR 0.117-0.646) with all predictors to 0.400 (mean 0.404; IQR 0.000-0.650) with selected predictors, and median ROC-AUC decreased from 0.632 (mean 0.624; IQR 0.516-0.744) to 0.622 (mean 0.598; IQR 0.502-0.717) (Tables 7 and 9). Formal modelling confirmed statistically significant decrements for ROC-AUC (p < .001), but not F1-score (p = 0.579) (Tables 10 and 8).

Research question 3: Outcomes

Outcome differences were more apparent for ROC-AUC than F1-score. Median F1-scores (Tables 3 and 7) were similar across outcomes (lapses: median 0.436, mean 0.409, IQR

515 0.180-0.625; cravings: median 0.400, mean 0.410, IQR 0.048-0.649), and the cross-

516 outcome zero-inflated beta model showed no significant difference (p = .061) (Table 11).

517 *Table 11 Pairwise comparison of F1-Score by outcome; based on zero-inflated beta-regression*

Contrast	Estimate	Standard error	z-ratio	p-value
Lapses - Cravings	0.023	0.012	1.875	0.061
Model: Zero-inflated beta regression. Estimates shown are for the conditional component only (values between 0 and 1), excluding the zero-inflation process.				
Note: Estimates are on the response scale (0-1), showing differences in predicted means.				
Positive estimates indicate higher values for the first condition in each contrast; negative estimates indicate higher values for the second condition.				
P-values adjusted using Tukey method.				

519 Across all specifications, lapse prediction achieved higher median ROC-AUC (0.659; mean

520 0.652; IQR 0.514-0.809; Table 5) than craving prediction (0.628; mean 0.611; IQR 0.510-

521 0.729; Table 9). In the cross-outcome linear mixed model (Table 12), this difference was

522 statistically significant (p < .001).

523 *Table 12 Pairwise comparison of ROC-AUC by outcome; based on linear mixed model*

Contrast	Estimate	Standard error	t-ratio	p-value
Lapses - Cravings	0.036	0.008	4.743	< 0.001
Model: Linear mixed model				
Note: Estimates are on the response scale (0-1), showing differences in predicted means.				
Positive estimates indicate higher values for the first condition in each contrast; negative estimates indicate higher values for the second condition.				
P-values adjusted using Tukey method.				

Research question 4: Proportion of own data

Lapses

Median F1-score was lowest when no participant-specific data were included (median 0.286;

mean 0.319; IQR 0.000-0.571), increased at 10% (median 0.400; mean 0.384; IQR 0.155-

0.607), and was highest at 20-30% (medians 0.500-0.542; means 0.484-0.490; IQRs 0.308-

0.667) (Table 3). Zero-inflated beta-regressions indicated that the scores where significantly

higher at 20% and 30% of own data included in the training algorithm, relative to none (p =

0.002 and p < .001, respectively) (Table 4).

Median ROC-AUC increased descriptively with more participant-specific data (none: median

0.655; mean 0.644; IQR 0.512-0.786; 30%: median 0.694; mean 0.670; IQR 0.513-0.852),

but neither linear mixed models nor adjusted pairwise comparisons indicated statistically

significant differences (Table 6; S9 Appendix).

Cravings

For cravings, median F1-score was similar for none and 10% participant-specific data

(medians 0.424-0.419; means 0.422-0.412; IQRs 0.143-0.649), declined at 20% (median

0.333; mean 0.388; IQR 0.000-0.640), and partially recovered at 30% (median 0.400; mean

0.415; IQR 0.000-0.703) (Table 7). Zero-inflated beta-regressions indicated that none of

these differences were statistically significant (all p-values ≥ 0.146) (Table 8).

Median ROC-AUC was highest when no participant-specific data were used (median 0.650;

mean 0.630; IQR 0.544-0.734) and lower when any participant-specific data were included

(Table 9). Linear mixed models indicated significant decrements at 10% (p = .031), 20% (p =

.035), and 30% (p = .006) (Table 10).

Supplementary metrics

Across specifications, sensitivity (S6 Appendix and S10 Appendix) and specificity (S7

Appendix and S11 Appendix) suggested a consistent trade-off across configurations

Sensitivity was typically moderate (often ~0.45-0.55), whereas specificity was higher (~0.75-

0.85) but more variable. For lapses, lower prompt frequencies were associated with

significantly higher sensitivity (all p < .001) alongside substantially lower specificity (again, all

p < .001). For cravings, sensitivity differences across prompt frequencies were smaller and

in the opposite direction (all significant, at p = .032 to p < .001), while specificity was

significantly reduced at 3 and 4 prompts per day, compared to 16 (p = 0.002 and p < .001,

respectively).

Together, these supplementary metrics indicate that some configurations maintained or

increased sensitivity at the expense of specificity, which can inflate certain threshold-

dependent summaries without necessarily improving overall discrimination.

560   Discussion

561   Summary of main findings

This study adds to the growing literature on predicting smoking behaviour from EMA data

using machine learning algorithms (34,55,66,67). It does so by systematically examining the

effects of prompt frequency, feature selection, prediction target (lapse versus craving), and

use of participant-specific data in training on the performance of RF machine learning

algorithms to underpin a future smoking cessation JITAI.

Across a large set of algorithm specifications, predictive performance varied substantially

between individuals and was, on average, modest when evaluated against commonly used

minimum performance thresholds. Median F1-scores for both lapse and craving prediction

were often below 0.5, indicating limited balance between precision and recall, while ROC-

AUC values were generally above chance but rarely reached levels typically considered

strong discrimination. This divergence highlights that many configurations could distinguish

higher-risk from lower-risk moments to some extent, yet, at the threshold of 0.5 at least, still

struggled to reliably identify true positive events without generating substantial false

positives or false negatives. Importantly, these patterns were not driven by a small subset of

poorly performing models but reflected broad variability across participants and

specifications.

Several consistent trends emerged. For lapse prediction, lower EMA prompt frequencies

were associated with higher F1-scores, contrary to expectations that denser sampling would

improve performance, whereas ROC-AUC showed no corresponding monotonic

improvement and, in some cases, favoured higher prompt frequencies. In contrast, craving

prediction showed the expected pattern: reducing prompt frequency led to systematic and

statistically significant declines in both F1-score and ROC-AUC. Feature selection had

minimal impact on lapse prediction and was associated with slightly worse performance for

cravings, suggesting that reducing predictor burden can be achieved with little loss of

accuracy for lapses but not necessarily for craving prediction. Finally, the results indicate that

the effect of incorporating participant-specific data depends strongly on both the outcome

and the metric used to assess performance. For lapses, personalisation improved F1-score

but not ROC-AUC, suggesting gains in precision–recall balance without improved

discrimination. For cravings, incorporating participant-specific data was associated with

poorer performance across both metrics, indicating limited benefit and even potential harm

from warm-start personalisation (hybrid-level algorithms) in this setting. Taken together,

these findings suggest that while EMA-based machine learning can capture meaningful risk

signals, considerations have to be made when it comes to implementing them in real-world

JITAIs.

Practical implicationsTaken together, these findings suggest that it may be *possible*, but not

guaranteed, to deploy algorithms predicting high-risk moments within a smoking cessation

JITAI without imposing excessive user burden, particularly when high-risk moments are

operationalised as lapses rather than cravings. Predictive performance was highly variable

across individuals and, on average, modest, indicating that such algorithms should be

viewed as providing probabilistic risk signals rather than reliable event-level predictions. The

comparatively greater difficulty of predicting cravings may be due to the algorithms

attempting to predict high cravings at the next EMA and lapses by the next EMA. As a

fluctuating measure, the former may simply be inherently more difficult to predict, especially

with longer and more variable gaps between EMAs. Longer and more variable gaps between

EMAs may therefore disproportionately reduce predictive accuracy for cravings compared

with lapses.

Although performance generally declined as EMA prompt frequency was reduced, this

decline was modest for lapse prediction and depended on the metric used. In particular,

lower prompt frequencies were associated with higher F1-scores for lapses, while ROC-AUC

showed no corresponding monotonic improvement and, in some cases, favoured higher

sampling densities. This divergence highlights that conclusions about "acceptable"

performance may depend on which aspect of performance is prioritised. In real-world

implementation, it may therefore be necessary to move beyond a fixed classification

threshold (e.g., 0.5) and instead select thresholds that explicitly trade off sensitivity and

specificity according to intervention goals. This could mean prioritising sensitivity to minimise

missed high-risk moments, even at the cost of more false positives. Such threshold tuning

could materially change how these algorithms perform in practice relative to the results

reported here.

Consistent with the nature of RF models, which intrinsically prioritise the most relevant

features, explicit feature selection had little impact on lapse prediction and was associated

with small performance decrements for craving prediction. Although we cannot conclude a

true absence of an effect, the point estimate of the difference was small. While this does not

imply that feature reduction is universally harmless, the very limited reduction in performance

across metrics and the overall similarity in retained predictors across algorithm specifications

suggests that a relatively small set of psychological and contextual variables captures much

of the predictive signal in these data. Given the substantial benefit of reducing survey

burden, using a more parsimonious model with fewer psychological and contextual tailoring

variables appears to be a justifiable and practical strategy. Nonetheless, concerns about

generalisability remain, discussed further below. Previous research indicates that users of a

smoking cessation JITAI may not be willing to respond to more than four to five surveys with

around 10 items per day (16,19,21,27–30). Therefore, using one of the lapse risk algorithms

using three to six prompts/day and feature selection may enable researchers to develop a

smoking cessation JITAI that has the potential to be both acceptable and effective.

Importantly, despite substantial differences in how well the algorithms performed for different

individuals, algorithms trained with participant-specific data did not consistently outperform

models trained without any of the test participants own data, particularly when looking at

ROC-AUC. This has favourable implications in terms of feasibility as it suggests that

individualised algorithm training is not necessary to maintain acceptable discriminative

ability. However, the benefit of using participant-specific data when evaluating performance

using the F1-score suggests that, when it comes to reliably identifying true positive events

without generating substantial false positives or false negatives, there may still be a benefit

for using hybrid-level algorithms (warm-start personalisation). Additionally,, there may still be

limitations in terms of generalisability (discussed further below), as the algorithms are trained

on individuals who have at least one instance of lapse or high craving and may therefore

work less well on those who have none or very few of these events. With these caveats in

mind, a group-level lapse-risk algorithm using five prompts per day and feature selection

may nevertheless be the best way going forward for a future JITAI based on this research.

Finally, the substantial between-individual variability in performance has important

implications for implementation. For some individuals, algorithm performance was

consistently poor, suggesting that fully automated decision-making may be inappropriate for

all users. One pragmatic approach that may be to combine machine learning outputs with

human-designed decision rules or clinician-informed heuristics, such that algorithm-

generated risk estimates inform intervention delivery as one of multiple algorithms using by a

JITAI. Such hybrid approaches may improve robustness, transparency, and acceptability

while mitigating the risks associated with relying on imperfect predictions in high-stakes

behavioural contexts.

Links to previous research

The predictors most consistently selected through RFE in this study, namely cravings, self-

efficacy, momentary motivation, cigarette availability, and affective states, are largely

consistent with factors identified as important psychological and contextual variables in prior

research identifying predictors of smoking lapses (30,34,67–72). Location, current activity,

and social context were not selected for any feature-reduced algorithms in this study. This

aligns with mixed evidence regarding the predictive utility of these contextual features;

previous research has found variables relating to these concepts to be key predictors in

some models, but not others (34,55,67,68,70,71). A potential reason why these variables

were not selected for any of the feature-reduced algorithms in this study may be how they

were operationalised (the variables in this study were fairly fine-grained) and the fact that

feature selection was done at the group-level (these variables may still be highly relevant for

some individuals). It is also important to note that this study only uses data from the first 10

days of participants' quit attempts. In this period, they may have engaged in non-typical or

unsustainably constrained ways to support their quit attempt and study participation, limiting

the predictive value of these variables. This interpretation is supported by the relatively low

frequency of several physical and social contexts in the data, including "Restaurant/bar,"

"With friend," and "With relative" (see S12 Appendix). Overall, this suggests that while there

is some consistency in the key psychological and contextual predictors of lapse risk, the

relevance of certain factors, such as location and social setting, may be more variable,

potentially depending on the modelling approach, operationalisations, the dataset, and/or

differences between individuals or populations.

The finding that hybrid-level algorithms did not consistently outperform group-level

algorithms, and were indeed outperformed by them on some metrics, contrasts with results

from previous analyses of machine learning algorithms to predict smoking lapses (67),

including the previous study using the same dataset (34). A likely explanation for this

discrepancy is the difference in how the sub-sample of participants' own data that was

included in the training set for the hybrid-level algorithms was selected. In the previous

studies, hybrid-level algorithms were trained with data that included a randomly selected

20% or 40% of each participant's data. In the present study, to simulate how a hybrid-level

algorithm would operate if implemented as part of a real-world smoking cessation

intervention, we used the first 10%, 20%, or 30% of each participant's data chronologically.

This difference in data selection may have affected the representativeness and diversity of

the training data, particularly if, as research using time-varying effect models suggest

(68,69,73–78), the predictors of smoking lapses and cravings early on a quit attempt are

different than the predictors at later points.

694 Strengths, limitations, and directions for future research

This study has several notable strengths. It is, to the authors' knowledge, the first study to

assess how EMA sampling frequency, predictor count, and training data composition affect

algorithm performance for predicting smoking lapses and cravings. These factors are critical

considerations when developing a JITAI, and by systematically examining them, this study

can provide a reference for making informed decisions about the necessary trade-offs.

Additionally, the use of high-frequency EMA data over a critical 10-day period provides

detailed insight into momentary risk factors. Furthermore, the testing of a large number of

algorithm configurations and cross-participant validation allows for a nuanced assessment of

generalisability and individual-level variability in performance. Finally, the preregistered

protocol and open sharing of data and code promote transparency and reproducibility.

There are also several ways for future research to extend and build on the findings from this

study. First, as a secondary analysis of previously collected data, this study used a randomly

selected subset of EMAs to explore the impact of reducing sampling frequency. Future

studies could assess the ecological validity of the results by experimentally varying prompt

frequencies to explore how participants respond to less intensive monitoring regimes and

assess the real-world impact of different prompt frequencies on algorithm performance.

Although visual inspection of the distributions and time series of all time-varying predictors

across participants (see S12 Appendix) did not suggest that reactivity or response blunting

was a major concern in this study, future research that experimentally varies prompt

frequency would also be well placed to directly test whether and how reactivity influences

both response variability and algorithm performance (79). Second, our analyses focused on

short-term prediction, using data from the immediately preceding EMA. Future work could

explore models that account for longer-term behavioural patterns or temporal dynamics

across the quit attempt to explore whether that would improve predictive performance. Third,

the dataset covers only the first 10 days of participants' smoking cessation attempts. While

this period is critical due to the high risk of lapse and relapse, predictors of lapses may

change over time (68,69,73–78). Future research could extend the time period under

investigation.

Although algorithm performance varied substantially across individuals, investigating why

some participants' smoking behaviour was harder to predict than others' would have

required a larger, more diverse sample and is beyond the scope of this study. Future

research could examine potential differential performance and algorithmic bias across

demographic characteristic aim to identify person-level characteristics that influence

algorithm performance to inform the development of more personalised or stratified

approaches. Additionally, as the sample in this study was self-selected, geographically

localised, and skewed toward participants willing to engage in intensive EMA protocols,

future studies should assess algorithm performance in more diverse and representative

populations. This is particularly important given evidence that smoking patterns and lapse

predictors may vary by socioeconomic status (80,81), and given that occupation type

consistently emerged as a key predictor in this study (see S2 Appendix). While the feature

selection procedure identified a relatively stable set of predictors across models, previous

work has shown that feature importance can vary substantially across contexts (30,54).

Further work is needed to evaluate the generalisability of these predictors and to better

understand their interpretability, including the direction and strength of their association with

outcomes. Importantly, relying on a smaller set of predictors may also constrain the range of

intervention targets available to a JITAI at any given moment, particularly when some

predictors are static or non-modifiable, such as baseline smoking and demographic

characteristics. This could limit the diversity of content in the intervention and potentially

affect user engagement over time if similar intervention components are repeatedly selected

and users get bored of them (7,53,82,83). Moreover, using fewer predictors may potential

mean omitting personally salient risk factors which may disproportionately affect algorithm

performance for some individuals. While the results of this study provide evidence that

acceptable predictive performance may be achievable with fewer features under certain

conditions, it is important that feature selection decisions in the context of JITAIs consider

not only predictive accuracy and participant burden, but also the downstream implications for

and consequences of intervention design, including content diversity, sustained user

engagement, and effectiveness.

Finally, the algorithms in this study employed a standard decision threshold of 0.5 to classify

predicted probabilities when distinguishing between lapses/non-lapses and high cravings/no

or low cravings. While intuitively sensible to get accurate predictions and appropriate for

exploratory purposes, future work focused on clinical or real-world implementation may

benefit from adjusting this threshold to achieve higher F1-scores or optimise either sensitivity

or specificity.

758 Conclusion

This study examined how EMA prompt frequency, predictor count, outcome definition, and

training data composition affect the performance of machine learning algorithms for

predicting high-risk moments during smoking cessation. Across specifications, performance

was highly variable between individuals and, on average, modest, with ROC-AUC values

generally above chance but F1-scores frequently below commonly used thresholds at a fixed

decision cut-off. Lapse prediction was more robust than craving prediction, reduced prompt

frequency had modest and metric-dependent effects, explicit feature selection had little

impact on lapse prediction, and incorporating participant-specific data did not consistently

improve performance. Overall, these findings suggest that EMA-based machine learning can

capture meaningful risk signals but should be applied cautiously in JITAIs, with careful

consideration of outcome choice, performance metric, and decision thresholds. A

parsimonious, group-level lapse-risk model using a moderate number of daily prompts may

offer a pragmatic starting point, while substantial individual-level variability highlights the

potential value of flexible thresholding and hybrid algorithm-rule-based decision strategies in

real-world implementation.

Acknowledgements

Thank you to all participants for their invaluable contributions to this research.

References

1. NHS Digital. Statistics on Public Health, England 2023 [Internet]. 2024 [cited 2025 Feb 20]. Available from: https://digital.nhs.uk/data-and-information/publications/statistical/statistics-on-public-health/2023

2. Reitsma MB, Fullman N, Ng M, Salama JS, Abajobir A, Abate KH, et al. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: a systematic analysis from the Global Burden of Disease Study 2015. The Lancet. 2017 May 13;389(10082):1885–906.

3. Song F, Elwell-Sutton T, Naughton F, Gentry S. Future smoking prevalence by socioeconomic status in England: a computational modelling study. Tob Control. 2021 Jul 1;30(4):380.

4. Hartmann-Boyce J, Ordóñez-Mena JM, Livingstone-Banks J, Fanshawe TR, Lindson N, Freeman SC, et al. Behavioural programmes for cigarette smoking cessation: investigating interactions between behavioural, motivational and delivery components in a systematic review and component network meta-analysis. Addiction. 2022 Aug 1;117(8):2145–56.

5. Livingstone-Banks J, Norris E, Hartmann-Boyce J, West R, Jarvis M, Hajek P. Relapse prevention interventions for smoking cessation. Cochrane Database Syst Rev [Internet]. 2019;(2). Available from: ://WOS:000460588000025

6. Brendryen H, Kraft P. Happy Ending: a randomized controlled trial of a digital multi-media smoking cessation intervention. Addiction. 2008 Mar 1;103(3):478–84.

7.   Ubhi HK, Michie S, Kotz D, Wong WC, West R. A Mobile App to Aid Smoking Cessation:

Preliminary Evaluation of SmokeFree28. J Med Internet Res. 2015 Jan 16;17(1):e3479.

8.   Ashare RL, Wileyto EP, Perkins KA, Schnoll RA. The first seven days of a quit attempt

predicts relapse: Validation of a measure for screening medications for nicotine

dependence. J Addict Med. 2013;7(4):249.

9.   Deiches JF, Baker TB, Lanza S, Piper ME. Early Lapses in a Cessation Attempt: Lapse

Contexts, Cessation Success, and Predictors of Early Lapse. Nicotine Tob Res. 2013

Nov 1;15(11):1883–91.

10. Kenford SL, Fiore MC, Jorenby DE, Smith SS, Wetter D, Baker TB. Predicting smoking

cessation: Who will quit with and without the nicotine patch. Jama. 1994;271(8):589–94.

11. Shadel WG, Martino SC, Setodji C, Cervone D, Witkiewitz K, Beckjord EB, et al. Lapse-

induced surges in craving influence relapse in adult smokers: an experimental

investigation. Health Psychol. 2011;30(5):588.

12. Hardeman W, Houghton J, Lane K, Jones A, Naughton F. A systematic review of just-in-

time adaptive interventions (JITAIs) to promote physical activity. Int J Behav Nutr Phys

Act. 2019 Apr 3;16(1):31.

13. Nahum-Shani I, Smith SN, Spring BJ, Collins LM, Witkiewitz K, Tewari A, et al. Just-in-

Time Adaptive Interventions (JITAIs) in Mobile Health: Key Components and Design

Principles for Ongoing Health Behavior Support. Ann Behav Med. 2018 May

18;52(6):446–62.

14. Battalio SL, Conroy DE, Dempsey W, Liao P, Menictas M, Murphy S, et al. Sense2Stop:

a micro-randomized trial using wearable sensors to optimize a just-in-time-adaptive

stress management intervention for smoking relapse prevention. Contemp Clin Trials.

2021;109:106534.

15. Businelle MS, Ma P, Kendzor DE, Frank SG, Vidrine DJ, Wetter DW. An ecological

momentary intervention for smoking cessation: evaluation of feasibility and

effectiveness. J Med Internet Res. 2016;18(12).

16. Hébert ET, Ra CK, Alexander AC, Helt A, Moisiuc R, Kendzor DE, et al. A mobile just-in-

time adaptive intervention for smoking cessation: pilot randomized controlled trial. J Med

Internet Res. 2020;22(3):e16907.

17. Naughton F, Hopewell S, Lathia N, Schalbroeck R, Brown C, Mascolo C, et al. A

Context-Sensing Mobile Phone App (Q Sense) for Smoking Cessation: A Mixed-Methods

Study. JMIR Mhealth Uhealth. 2016 Sep 16;4(3):e106.

18. Naughton F, Hope A, Siegele-Brown C, Grant K, Barton G, Notley C, et al. An

automated, online feasibility randomized controlled trial of a just-in-time adaptive

intervention for smoking cessation (Quit Sense). Nicotine Tob Res. 2023 Jul

1;25(7):1319–29.

19. Yang MJ, Sutton SK, Hernandez LM, Jones SR, Wetter DW, Kumar S, et al. A Just-In-

Time Adaptive intervention (JITAI) for smoking cessation: Feasibility and acceptability

findings. Addict Behav. 2023 Jan 1;136:107467.

20. Meng XL, Xie X. I Got More Data, My Model is More Refined, but My Estimator is

Getting Worse! Am I Just Dumb? Econom Rev. 2014 Feb 10;33(1–4):218–50.

21. Burke LE, Shiffman S, Music E, Styn MA, Kriska A, Smailagic A, et al. Ecological

Momentary Assessment in Behavioral Research: Addressing Technological and Human

Participant Challenges. J Med Internet Res. 2017 Mar 15;19(3):e77.

22. Materia FT, Smyth JM. Acceptability of Intervention Design Factors in mHealth

Intervention Research: Experimental Factorial Study. JMIR Mhealth Uhealth. 2021 Jul

26;9(7):e23303.

23. Chevance G, Perski O, Hekler EB. Innovative methods for observing and changing

complex health behaviors: four propositions. Transl Behav Med. 2021 Feb 1;11(2):676–

85.

24. Ferguson SG, Shiffman S. The relevance and treatment of cue-induced cravings in

tobacco dependence. J Subst Abuse Treat. 2009 Apr 1;36(3):235–43.

25. Dao KP, De Cocker K, Tong HL, Kocaballi AB, Chow C, Laranjo L. Smartphone-delivered

ecological momentary interventions based on ecological momentary assessments to

promote health behaviors: systematic review and adapted checklist for reporting

ecological momentary assessment and intervention studies. JMIR MHealth UHealth.

2021;9(11):e22890.

26. Kabir KS, Kenfield SA, Van Blarigan EL, Chan JM, Wiese J. Ask the Users: A Case

Study of Leveraging User-Centered Design for Designing Just-in-Time Adaptive

Interventions (JITAIs). Proc ACM Interact Mob Wearable Ubiquitous Technol.

2022;6(2):1–21.

27. Crochiere RJ, Zhang F (Zoe), Juarascio AS, Goldstein SP, Thomas JG, Forman EM.

Comparing ecological momentary assessment to sensor-based approaches in predicting

dietary lapse. Transl Behav Med. 2021 Dec 1;11(12):2099–109.

28. Forman EM, Goldstein SP, Zhang F, Evans BC, Manasse SM, Butryn ML, et al. OnTrack:

development and feasibility of a smartphone app designed to predict and prevent dietary

lapses. Transl Behav Med. 2019;9(2):236–45.

29. Goldstein SP, Zhang F, Thomas JG, Butryn ML, Herbert JD, Forman EM. Application of

Machine Learning to Predict Dietary Lapses During Weight Loss. J Diabetes Sci

Technol. 2018 Sep 1;12(5):1045–52.

30. Hébert ET, Suchting R, Ra CK, Alexander AC, Kendzor DE, Vidrine DJ, et al. Predicting

the first smoking lapse during a quit attempt: A machine learning approach. Drug Alcohol

Depend. 2021;218:108340.

31. Ubhi HK, Kotz D, Michie S, van Schayck OCP, West R. A comparison of the

characteristics of iOS and Android users of a smoking cessation app. Transl Behav Med.

2017 Jun 1;7(2):166–71.

32. Cobb CO, Niaura RS, Donaldson EA, Graham AL. Quit Now? Quit Soon? Quit When

You're Ready? Insights About Target Quit Dates for Smoking Cessation From an Online

Quit Date Tool. J Med Internet Res. 2014 Feb 17;16(2):e55.

33. Shiffman S, Hickcox M, Paty JA, Gnys M, Richards T, Kassel JD. Individual differences

in the context of smoking lapse episodes. Smok Cessat Clin Res Dir. 1997 Nov

1;22(6):797–811.

34. Perski O, Kale D, Leppin C, Okpako T, Simons D, Goldstein SP, et al. Supervised

machine learning to predict smoking lapses from Ecological Momentary Assessments

and sensor data: Implications for just-in-time adaptive intervention development. PLOS

Digit Health. 2024 Aug 23;3(8):e0000594.

35. Mestdagh M, Verdonck S, Piot M, Niemeijer K, Tuerlinckx F, Kuppens P, et al. m-Path:

An easy-to-use and flexible platform for ecological momentary assessment and

intervention in behavioral research and clinical practice. PsyArXiv Prepr. 2022;

36. Jackson SE, Perski O, Crane D, Michie S, West R, Brown J. Effectiveness of an offer of

the Smoke Free smartphone application for smoking cessation: protocol for a

randomized controlled trial. Addiction. 2019 Nov 1;114(11):2078–86.

37. Kotz D, Brown J, West R. Predictive validity of the Motivation To Stop Scale (MTSS): A

single-item measure of motivation to stop smoking. Drug Alcohol Depend. 2013 Feb

1;128(1):15–9.

38. Barrett LF, Gross J, Christensen TC, Benvenuto M. Knowing what you're feeling and

knowing what to do about it: Mapping the relation between emotion differentiation and

emotion regulation. Cogn Emot. 2001;15(6):713–24.

39. Posner J, Russell JA, Peterson BS. The circumplex model of affect: An integrative

approach to affective neuroscience, cognitive development, and psychopathology. Dev

Psychopathol. 2005/11/01 edn 2005;17(3):715–34.

40. Russell JA. A circumplex model of affect. J Pers Soc Psychol. 1980;39(6):1161.

41. Moritz S, Sardá A, Bartz-Beielstein T, Zaefferer M, Stork J. Comparison of different

methods for univariate time series imputation in R. ArXiv Prepr ArXiv151003924. 2015;

42. Chevance G, Baretta D, Heino M, Perski O, Olthof M, Klasnja P, et al. Characterizing

and predicting person-specific, day-to-day, fluctuations in walking behavior. PLoS One.

2021;16(5):e0251659.

43. Rhys H. Machine Learning with R, the tidyverse, and mlr. 1st edition. New York:

Manning; 2020.

44. Probst P, Wright MN, Boulesteix AL. Hyperparameters and tuning strategies for random

forest. WIREs Data Min Knowl Discov. 2019 May 1;9(3):e1301.

45. Probst P, Boulesteix AL. To tune or not to tune the number of trees in random forest. J

Mach Learn Res. 2018;18(181):1–18.

46. Scornet E. Tuning parameters in random forests. ESAIM Proc Surv. 2017;60:144–62.

47. R Core Team. R: A Language and Environment for Statistical Computing [Internet].

Vienna, Austria: R Foundation for Statistical Computing; 2024. Available from:

https://www.R-project.org/

48. Kuhn M, Wickham H. Tidymodels: a collection of packages for modeling and machine

learning using tidyverse principles. [Internet]. 2020. Available from:

https://www.tidymodels.org

49. Singh KJ, Meitei AJ, Alee NT, Kriina M, Haobijam NS. Machine learning algorithms for

predicting smokeless tobacco status among women in Northeastern States, India. Int J

Syst Assur Eng Manag. 2022;13(5):2629–39.

50. Strobl C, Malley J, Tutz G. An introduction to recursive partitioning: rationale, application,

and characteristics of classification and regression trees, bagging, and random forests.

Psychol Methods. 2009;14(4):323.

51. Wright MN, Ziegler A, König IR. Do little interactions get lost in dark random forests?

BMC Bioinformatics. 2016 Mar 31;17(1):145.

52. Breiman L. Random forests. Mach Learn. 2001;45(1):5–32.

53. Leppin C, Okpako T, Garnett C, Perski O, Brown J. Exploring perspectives on digital

smoking cessation just-in-time adaptive interventions: A focus group study with adult

smokers and smoking cessation professionals. PLOS Digit Health. 2025 May

2;4(5):e0000705.

54. Tang J, Alelyani S, Liu H. Feature selection for classification: A review. Data Classif

Algorithms Appl. 2014;37.

55. Dumortier A, Beckjord E, Shiffman S, Sejdić E. Classifying smoking urges via machine

learning. Comput Methods Programs Biomed. 2016 Dec 1;137:203–13.

56. Gregorutti B, Michel B, Saint-Pierre P. Correlation and variable importance in random

forests. Stat Comput. 2017 May 1;27(3):659–78.

57. Cheng YC, Chang HT, Lin CY, Chang HY. Predicting Credit Risk in Peer-to-Peer

Lending: A Machine Learning Approach with Few Features. In: 2021 International

Conference on Technologies and Applications of Artificial Intelligence (TAAI). 2021. p.

295–300.

58. Akkaya B. The Effect of Recursive Feature Elimination with Cross-Validation Method on

Classification Performance with Different Sizes of Datasets. In 2021. p. 4–6.

59. Agaal A, Essgaer M. Influence of Feature Selection Methods on Breast Cancer Early

Prediction Phase using Classification and Regression Tree. In: 2022 International

Conference on Engineering & MIS (ICEMIS). 2022. p. 1–6.

60. Choi SG, Oh M, Park D, Lee B, Lee Y ho, Jee SH, et al. Comparisons of the prediction

models for undiagnosed diabetes between machine learning versus traditional statistical

methods. Sci Rep. 2023 Aug 11;13(1):13101.

61. Habenicht R, Fehrmann E, Blohm P, Ebenbichler G, Fischer-Grote L, Kollmitzer J, et al.

Machine Learning Based Linking of Patient Reported Outcome Measures to WHO

International Classification of Functioning, Disability, and Health Activity/Participation

Categories. J Clin Med. 2023;12(17).

62. Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.

glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated

Generalized Linear Mixed Modeling. R J. 2017;9(2):378–400.

63. McGillycuddy M, Warton DI, Popovic G, Bolker BM. Parsimoniously Fitting Large

Multivariate Random Effects in glmmTMB. J Stat Softw. 2025;112(1):1–19.

64. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using

lme4. J Stat Softw. 2015;67(1):1–48.

65. Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means [Internet].

2025. Available from: https://rvlenth.github.io/emmeans/

66. Abo-Tabik M, Costen N, Darby J, Benn Y. Decision Tree Model of Smoking Behaviour.

In: 2019 IEEE SmartWorld, Ubiquitous Intelligence & Computing, Advanced & Trusted

Computing, Scalable Computing & Communications, Cloud & Big Data Computing,

Internet of People and Smart City Innovation,

SmartWorld/SCALCOM/UIC/ATC/CBDCom/IOP/SCI 2019, Leicester, United Kingdom,

August 19-23, 2019 [Internet]. IEEE; 2019. p. 1746–53. Available from:

https://doi.org/10.1109/SmartWorld-UIC-ATC-SCALCOM-IOP-SCI.2019.00311

67. Perski O, Li K, Pontikos N, Simons D, Goldstein SP, Naughton F, et al. Classification of

Lapses in Smokers Attempting to Stop: A Supervised Machine Learning Approach Using

Data From a Popular Smoking Cessation Smartphone App. Nicotine Tob Res. 2023 Jul

1;25(7):1330–9.

68. Koslovsky MD, Hébert ET, Businelle MS, Vannucci M. A Bayesian time-varying effect

model for behavioral mHealth data. Ann Appl Stat. 2020;14(4):1878–902.

69. Koslovsky MD, Hébert ET, Swartz MD, Chan W, Leon-Novelo L, Wilkinson AV, et al. The

Time-Varying Relations Between Risk Factors and Smoking Before and After a Quit

Attempt. Nicotine Tob Res. 2018 Sep 4;20(10):1231–6.

70. Koslovsky MD, Swartz MD, Chan W, Leon-Novelo L, Wilkinson AV, Kendzor DE, et al.

Bayesian Variable Selection for Multistate Markov Models with Interval-censored Data in

an Ecological Momentary Assessment Study of Smoking Cessation. Biometrics. 2018

Jun 1;74(2):636–44.

71. Suchting R, Hébert ET, Ma P, Kendzor DE, Businelle MS. Using Elastic Net Penalized

Cox Proportional Hazards Regression to Identify Predictors of Imminent Smoking Lapse.

Nicotine Tob Res. 2019 Jan 4;21(2):173–9.

72. Gwaltney CJ, Shiffman S, Balabanis MH, Paty JA. Dynamic self-efficacy and outcome

expectancies: prediction of smoking lapse and relapse. J Abnorm Psychol.

2005;114(4):661.

73. Dermody SS, Shiffman S. The time-varying effect of alcohol use on cigarette smoking

relapse risk. Addict Behav. 2020 Mar 1;102:106192.

74. Lanza ST, Vasilenko SA, Liu X, Li R, Piper ME. Advancing the Understanding of Craving

During Smoking Cessation Attempts: A Demonstration of the Time-Varying Effect Model.

Nicotine Tob Res. 2014 May 1;16(Suppl_2):S127–34.

75. Potter LN, Haaland BA, Lam CY, Cambron C, Schlechter CR, Cinciripini PM, et al. A

time-varying model of the dynamics of smoking lapse. Health Psychol. 2021;40(1):40.

76. Shiyko M, Naab P, Shiffman S, Li R. Modeling Complexity of EMA Data: Time-Varying

Lagged Effects of Negative Affect on Smoking Urges for Subgroups of Nicotine

Addiction. Nicotine Tob Res. 2014 May 1;16(Suppl_2):S144–50.

77. Shiyko M, Lanza ST, Tan X, Li R, Shiffman S. Using the Time-Varying Effect Model

(TVEM) to Examine Dynamic Associations between Negative Affect and Self Confidence

on Smoking Urges: Differences between Successful Quitters and Relapsers. Prev Sci.

2012 Jun 1;13(3):288–99.

78. Vasilenko SA, Piper ME, Lanza ST, Liu X, Yang J, Li R. Time-Varying Processes

Involved in Smoking Lapse in a Randomized Trial of Smoking Cessation Therapies.

Nicotine Tob Res. 2014 May 1;16(Suppl_2):S135–43.

79. McCarthy DE, Minami H, Yeh VM, Bold KW. An experimental investigation of reactivity to

ecological momentary assessment frequency among adults trying to quit smoking.

Addiction. 2015;110(10):1549–60.

80. Cambron C, Lam CY, Cinciripini P, Li L, Wetter DW. Socioeconomic Status, Social

Context, and Smoking Lapse During a Quit Attempt: An Ecological Momentary

Assessment Study. Ann Behav Med. 2020 Feb 21;54(3):141–50.

81. Jahnel T, Ferguson SG, Shiffman S, Thrul J, Schüz B. Momentary smoking context as a

mediator of the relationship between SES and smoking. Ambul Assess Addict Disord.

2018 Aug 1;83:136–41.

82. Bendotti H, Lawler S, Ireland D, Gartner C, Hides L, Marshall HM. What Do People Want

in a Smoking Cessation App? An Analysis of User Reviews and App Quality. Nicotine

Tob Res. 2022 Feb 1;24(2):169–77.

83. Zhang M, Wolters M, O'Connor S, Wang Y, Doi L. Smokers' expectations and

perceptions of smoking cessation apps: A systematic review of the qualitative literature.

Int J Med Inf. 2023;175(105069).

1021 Supporting information captions

S1 Appendix Surveys

S2 Appendix List of predictors for each model

S3 Appendix Description of observations lapsers only

S4 Appendix F1-Score performance summaries by predictor specification and participant

S5 Appendix ROC-AUC performance summaries by predictor specification and participant

S6 Appendix Sensitivity performance summaries by predictor specification and participant

S7 Appendix Specificity performance summaries by predictor specification and participant

S8 Appendix F1-Score models and marginal effects

S9 Appendix ROC-AUC models and marginal effects

S10 Appendix Sensitivity models and marginal effects

S11 Appendix Specificity models and marginal effects

S12 Appendix Time-varying variables distributions in full dataset