Long COVID risk by pre-infection symptoms and functional status: A retrospective cohort study

of data from the *All of Us* Research Program.

SHORT TITLE: Long COVID risk by pre-infection symptoms and functional status in the *All*

of Us Research Program.

AUTHORS:

Kristen Kehl-Floberg[1,#a,#b*], Emma Freisberg[2,#c], Aurora Pop-Vicas[2], Ron Gangnon – senior

author[2], Dorothy Farrar Edwards[1,2,3]

[1]Institute for Clinical and Translational Research, University of Wisconsin-Madison, Madison,
Wisconsin, U.S.A.
11
12 [2]Department of Kinesiology, School of Education, University of Wisconsin-Madison, Madison,
13 Wisconsin, U.S.A
14
15 [3]School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin,
16 U.S.A.
17
18 [#a]Geriatric Research, Education, and Clinical Center (GRECC), William S. Middleton Veteran's
19 Memorial Hospital, 2500 Overlook Terrace, Madison, Wisconsin, 53705, U.S.A.
20
21 [#a]Department of Kinesiology, School of Education, University of Wisconsin-Madison, Madison,
22 Wisconsin, U.S.A.
23
24 [#c]Currently unaffiliated.
25

26 * Corresponding author: Kristen E. Kehl-Floberg, kekehl@wisc.edu,

27 Kristen.Kehlfloberg@va.gov

ABSTRACT

IMPORTANCE: Over seven million U.S. adults experience "long COVID", or persistent health issues after COVID-19. Multiple guidelines recommend the inclusion of functional status in long COVID diagnostic criteria, but more evidence is needed to guide this recommendation. This study explores the adjusted odds of developing long COVID by pre-infection symptoms and functional status, and the feasibility of estimating functional status using health records data.

DESIGN: Retrospective cohort study in a multicenter national longitudinal cohort of U.S. adults with history of COVID-19, using health records and survey responses through July 2022 (*All of Us* CDR 7.0).

EXPOSURE(S): Pre-infection (-5 years) incidences of (a) at least one symptom common in long COVID, and (b) functional status, indicated by *All of* Us baseline survey responses and diagnostic/procedure/billing codes. Disease and demographic data covariates were included in the adjusted models.

RESULTS: $N = 65,464$ met inclusion criteria ($n=40,655$ had post-infection occurrences of at least one symptom (long COVID group), while $n=24,809$ had none). Adjusted odds ratios within 99% confidence intervals [99% CI] of developing long COVID increased with lower pre-infection self-reported mental health ("Good" compared to "Excellent" AOR=1.14 [1.04,1.25], $P>0.000$), and more pre-infection symptoms (compared to the median of four, people with zero had much lower odds (AOR=0.15 [0.04,0.61], $P=0.008$). Adjusted odds were not significantly affected by any single pre-infection symptom, self-rated physical ability, or clinical documentation of functional impairment.

CONCLUSIONS. Greater pre-illness symptom burden and lower self-rated mental health

increased the odds of long COVID symptoms, after adjusting for demographics, variant,

functional status, and individual symptoms. Long COVID represents a change from baseline

functioning and health, even in people with pre-infection incident symptoms and functional

impairments. This estimation of pre-infection functional status using harmonized electronic

health records data demonstrated the feasibility of these data in developing the diagnostic utility

of functional status changes in long COVID.

Introduction

Among U.S. adults with a history of SARS-CoV-2 infection, 7.7 to 23 million experience persistent post-acute symptoms [1] termed "long COVID" [2]. About 7.5% of U.S. adults reported long COVID symptoms in June 2024, and about 5.5% reported activity limitations as a result [3]. Long COVID is a patient-generated term for new-onset or exacerbated, and persistent, symptoms and health conditions after COVID-19 infection [2, 4-8]. Common presentations include fatigue, post-exertional malaise, cognitive impairment, headache, breathing difficulty, musculoskeletal pain or weakness, sleep disturbance, and mood disorder [9] among many others [10-12]. Long COVID symptoms affect complex and "chosen" activity patterns (such as productivity and leisure) more dramatically than basic self-care [13, 14] forcing some patients to abandon complex routines and roles in work, family, and community life [6, 15].

Clinical diagnostic definitions and criteria of long COVID are an urgent and ongoing focus of study [2]. Multiple research initiatives have been launched to study recovery from COVID-19 in large samples with the goal of identifying unique signatures of symptoms and biomarkers [10, 16-21], many using electronic health records (EHR) data. These studies have marshalled extraordinary amounts of data into various phenotypes and diagnostic hallmarks [22], providing vital direction on the disease and its impact. However, few have included quantifiable indicators of daily functioning [23], despite early patient reports describing profound impact on daily life [6, 24] and proposed criteria asserting that an impact on daily functioning is a critical diagnostic component [2, 25]. Because most EHR-based phenotyping studies have not included data on how disability manifests or is treated, their resulting diagnostic classifications have excluded this diagnostic feature.

Additionally, although post-infection presentations have been thoroughly investigated,

studies of pre-infection occurrences of these same symptoms, conditions, and functional

impairments are limited [16, 26]. Most studies that explore the emergence of symptoms and

conditions after COVID-19 have not compared this to, nor assessed risk based on, their pre-

infection prevalence [22]. Concerning functional status, pre-infection disability has been found

to be associated with a two- [27] to three-fold [28] increase in long COVID. Because long

COVID includes worsened or exacerbated conditions that were previously well-managed [5],

pre-infection symptoms and functional status may be relevant to both estimating risk of

developing long COVID and tailoring management strategies [22].

The primary aim of this study is to compare pre-infection functional status, pre-infection

occurrences of common long COVID symptoms, and demographics, between participants with

and without long COVID in a diverse population-based sample. Our secondary aim is to estimate

the adjusted odds of developing at least one long COVID condition based on these pre-infection

characteristics.

95 Materials and Methods

96 Design and data source

This is a retrospective cohort study of participants enrolled in the U.S. National Institutes

of Health *All of Us* Research Program through July 2022 (curated data repository version 7.0),

Controlled tier C2022Q4R9 [29] accessed through the *All of Us* Researcher Workbench [30]

between May 1 2023 and June 31 2025. *All of Us* is among the most comprehensive population-

level biobanks in the world [31] enrolling participants from all 50 U.S. States and three

territories and over-recruiting from groups that are historically underrepresented in medical

research. They share baseline function and health data through surveys, biospecimens, and

release of EHR data which is subsequently scraped every three months. The program and data

snapshots can be viewed at https://www.researchallofus.org/data-tools/.

Sample

All of Us participants 18 years of age and older with at least one incidence of any

indicator of COVID-19 illness were included. Diagnostic indicators were (a) laboratory values

for positive PCR, antigen, or antibody test, (b) self-reported illness reported on the 2021-2022

All of Us COVID-19 Participant Experiences (COPE) survey, or (c) the ICD-10 code for

COVID-19 (U07.7) (S1 Appendix A, Table A.1). Informed consent for sharing EHR and survey

data was obtained from all participants by *All of Us* Health Provider Organizations responsible

for participant recruitment and enrollment. IRB oversight was delegated to the *All of Us* Central

IRB. Controlled tier data includes unique participant identifiers and dates; investigators

transformed these (e.g. date of birth to years of age) promptly following import, and collapsed

counts of fewer than 20 to prohibit subgroup membership identification in summary tables.

These steps are in compliance with *All of Us* data use agreement policies.

Long COVID and *Recovered* participant groups

Participants with COVID-19 history were grouped by whether their EHR showed at least

one diagnosis or observation code from a list of 38 long COVID symptom and condition

categories (see below, "Pre- and post-infection incidences of long COVID symptoms", and S1

Appendix A, Table A.2) at least 28 days after first infection. Participants with at least one

symptom were classified in the "long COVID" group; the rest were classified in the "recovered"

group. (S1 Appendix A).

Data sources and variables

Surveys: Demographics, self-reported health, and self-reported function at study enrollment.

All of Us participants provide self-report of their health and daily functional status data through validated self-report measures from the PROMIS battery and *All of Us*-designed surveys at enrollment. This study used responses to "The Basics" (demographics), "Overall Health" (daily functioning and mental health) and "Covid-19 Participant Experience" (COPE) (self-reported COVID-19 symptoms between May 2020 – March 2021) surveys.

Pre- and post-infection incidences of long COVID symptoms and conditions.

Symptoms of long COVID were chosen a-priori based on the Centers for Disease Control and Prevention's list of symptoms [32] and emergency billing code for Post-COVID conditions [33], prior phenotyping and patient-led literature, and the authors' clinical and community research experience. These sources yielded a list of 44 symptoms/conditions ("descendent" codes), which fell under 38 categories ("ancestor" codes) within the *All of Us* data table structure (S1 Appendix A, Table A.2). These 38 ancestor diagnoses were queried for all participants. For each category, the first incidence in the five years prior to infection signified a pre-infection occurrence, and the first incidence 28 days or more after the infection signified a post-infection occurrence (long COVID group only) of any symptom under that category. Because proposed empirical definitions of long COVID include exacerbations or relapses of pre-infection conditions [5, 11, 22], we queried "any mention" of the symptoms/conditions in this list (rather than "first mention"). Symptom persistence was not estimated due to the *All of Us* data structure and privacy policies. S1 Appendix A reports selection process, rationale, condition categories, and exclusions.

Total number of long COVID symptoms/conditions present prior to infection

149   From the pre-infection symptom occurrences, a gross measure of pre-infection symptom

150   burden was computed for each participant by tallying all long COVID symptom categories with

151   observations. This included all symptoms (e.g. headache, joint pain, cough) and conditions

152   (ME/CFS, POTS, and post-viral fatigue syndromes) used in cohort discovery.

Infection variants active at time of first infection

A time-bound variable was created to infer variant exposure at date of first SARS-CoV-2

infection. Periods began and ended around the emergence of major variants as reported by

Markov et al [34]. Our variant periods were "pre-VOC/wild type" from January-November 2020,

"pre-VOC/alpha/beta from November 2020 – April 2021, "alpha/beta/delta" from April-August

2021, "delta" from August-December 2021, "omicron_BA1-BA2" from December 2021 – April

2022, and "omicron_BA2-BA5" from April 2022 – July 2022 (data release cut-off date).

Demographics

Age, sex assigned at birth, race, and ethnicity were included based on evidence that these

factors influence risks for long COVID [35-38] and for functional impairment [39, 40]. Level of

education was included as a proxy for social determinants of health due to its impact on access to

health-supporting societal resources [41], as financial concerns [42] and greater unmet social

needs [43] have been found to be associated with more numerous and persistent long COVID

symptoms (S1 Appendix D, Table D.1).

Pre-COVID daily functioning

Pre-infection functional status was explored using enrollment surveys and functional

status findings (EHR procedure and diagnostic codes) recorded between January 1, 2015 and

four weeks prior to first infection. Survey responses were queried for self-reported performance

of physically demanding daily activities, social roles, and mental/cognitive health. From the

EHR, the incidences of medical billing codes for therapeutic procedures provided by

occupational therapists were tallied [44] (S1 Appendix B). Additionally, procedure codes under

the vocabulary hierarchy "Finding of Functional Performance and Activity" were grouped into

three levels of presumable impairment ("None", "Some", and "Dependent on others"). (S1

Appendix B.)

Quantitative bias analysis of cohort discovery

178   Potential cohort bias was examined using non-parametric propensity score matching.

179 Bias was estimated for participant enrollment before first COVID infection (retained for this

180 analysis) versus after (excluded) and a 90 day symptom start date instead of 28 days (S1

181 Appendix C).

Analysis

Demographics of the long COVID and recovered groups were graphed and tested for

significant difference using Welch's t test (normal), Mann-Whitney U test (non-normal), or X^2

test (categorical). Unadjusted odds ratios were computed for all variables. Logistic regression

was used to model the adjusted odds of developing long COVID based on demographics, variant,

pre-infection symptoms/conditions, and pre-infection functional status (S1 Appendix D, Table

D.1-D.2, Figure D.1). Data access and analysis were performed in the *All of Us* Researcher

Workbench, Jupyter notebooks cloud computing environment, using R [45]. Cohort discovery,

data cleaning and organization, and statistical analyses were completed by the first author using

R packages "stats" [46], "rms" [47], "corrplot" [48], and the "tidyverse" suite [49].

Results

104,992 participants met inclusion criteria of at least one COVID-19 illness. Participants with missing demographic survey responses (n=21,202) were removed. Further removed were by n=16,815 participants who enrolled after their first COVID-19 indicator (allowing use of enrollment questionnaires as pre-infection self-reported baseline functioning and health data); n=1,506 due to non-responses to the social survey questions (permitting these responses to be combined as an ordinal summary scale); n=5 participants with erroneous diagnostic codes (concept name and code was for SARS-CoV-2, but the dates of diagnosis corresponded with previous SARS-type virus outbreaks in the early 2000's); and n=1 due to an extreme and clinically unlikely value of 1286 (or 321 hours) for their 5-year CPT billing code total. The final sample was $N = 65,464$ participants, with 40,655 showing at least one post-infection symptom (long COVID group) and 24,809 showing none (recovered group) (Figure 1).

[Place Figure 1 about here]

Fig. 1. CONSORT diagram of sample flow.

Quantitative bias analysis of cohort discovery

Non-parametric propensity score matching showed a low match rate of 45% between pre- and post-infection enrollees (S1 Appendix C, FigureS C.1-C.2). Changing the classification timeline to 90 days after first infection had a very small impact on the numbers of long COVID versus recovered participants (S1 Appendix C, Figures (C.3-C.4).

Demographics

Demographic distributions, univariate unadjusted odds ratios and multivariate adjusted odds ratios are shown in Table 1. Most participants had their first infection during the pre-variant/wild type period, during which the proportion of long COVID participants outnumbered that of recovered participants by almost 2 to 1 (S1 Appendix E, Figure E.1). The Long COVID participants were older (median 63 (24) versus 60 (28)), and had slightly higher percentages of female and Black participants than the recovered group. Adjusted odds of developing long COVID after infection increased with older age, female sex, Black racial identity, and an earlier variant period of first infection. Groups were effectively equivalent in numbers of participants reporting Hispanic/Latino ethnicity.

Table 1. Demographic, functioning, and long COVID disease characteristics and odds ratios of developing long COVID in *All of Us* participants through July 2022.

	Long COVID Symptoms	Recovered	*p*	OR	AOR
n (total 65,464)	40,655	24,809			
age (mean (SD))	61.01 (16.03)	57.66 (16.82)	< 2.2e-16		
25				0.54 (0.43, 0.69)	0.68 (0.59, 0.77), *p>0.000*
35				0.64 (0.59, 0.70)	0.75 (0.68, 0.83), *p>0.000*
45				0.73 (0.68, 0.79)	0.83 (0.79, 0.89), *p>0.000*
55				0.94 (0.87, 1.02) *p<0.059*	0.93 (0.91, 0.95), *p>0.000*
62				1.00	1.00
75				1.09 (1.02, 1.17)	1.15 (1.10, 1.20), *p>0.000*
85				1.21 (1.12, 1.31)	1.28 (1.18, 1.39), *p>0.000*
Total number of pre-pandemic symptoms of long COVID (mean (SD))	6.11 (3.96)	2.44 (2.93)	< 2.2e-16		
0				0.13 (0.12, 0.14)	0.15 (0.04, 0.61) *p=0.008*
1				0.47 (0.44, 0.51)	0.38 (0.13, 1.11) *p=0.076*

4				1.00	1.00
7				2.12 (1.99, 2.27)	1.61 (0.55, 4.68) p=0.381
23				4.78 (4.44, 5.16)	1.99 (0.00, 1703.58) p=0.842
Variant of Concern			< 2.2e-16		
Pre-VOC/wild type	21120 (51.9)	7415 (29.9)		1.00	1.00
Pre-VOC, alpha, and beta	9890 (24.3)	6306 (25.4)		0.55 (0.52, 0.58)	0.61 (0.57, 0.65), p>0.000
alpha, beta, and delta	3240 (8.0)	1890 (7.6)		0.60 (0.56, 0.65)	0.48 (0.44, 0.53), p>0.000
delta	3462 (8.5)	2878 (11.6)		0.42 (0.39, 0.46)	0.33 (0.30, 0.36), p>0.000
omicron BA1-BA2	2568 (6.3)	3543 (14.3)		0.25 (0.24, 0.27)	0.17 (0.16, 0.19), p>0.000
omicron BA2-BA5	375 (0.9)	2777 (11.2)		0.05 (0.04, 0.06)	0.02 (0.02, 0.03), p>0.000
Sex assigned at Birth (%)			< 2.2e-16		
Female or intersex	26646 (65.5)	15287 (61.6)		1.18 (1.13, 1.24)	1.06 (1.01, 1.13), p=0.005
Race (%)			< 2.2e-16		
Asian	973 (2.4)	984 (4.0)		0.62 (0.55, 0.69),	0.92 (0.79, 1.10), p=0.14
Black or African American	8941 (22.0)	4645 (18.7)		1.20 (1.14, 1.26),	1.12 (1.05, 4.18), , p>0.00
Middle Eastern or North African	285 (0.7)	199 (0.8)		0.89 (0.70, 1.136), 0.21	1.03 (0.76, 1.40), p=0.80
More than one population	783 (1.9)	526 (2.1)		0.93 (0.80, 1.07), 0.18	1.01 (0.84, 1.20), p=0.88
White	39716 (97.7)	18455 (74.4)		1.00	1.00
Ethnicity (%)			0.00018		
Hispanic or Latino	939 (2.3)	689 (2.8)		0.83 (0.73, 0.94)	0.88 (0.75, 1.03), p=0.041
Education (highest grade completed)			< 2.2e-16		
Advanced Degree	9722 (23.9)	7230 (29.1)		0.7 (0.66, 0.73)	0.97 (0.90, 1.05), p=0.343
College Graduate	9654 (23.7)	6557 (26.4)		0.76 (0.72, 0.80)	1.02 (0.94, 1.09) p=0.585
Some College	11771 (29.0)	6104 (24.6)		1.00	1.00
High School/GED	7043 (17.3)	3658 (14.7)		0.99 (0.94, 1.06)	0.96 (0.89, 1.05) p=0.261
Some High School	1592 (3.9)	851 (3.4)		0.96 (0.86, 1.06)	0.85 (0.73, 0.99) p=0.005
Middle School	286 (0.7)	122 (0.5)		1.16 (0.91, 1.48)	1.02 (0.72, 1.46) p=0.866
Elementary	<40 (<0.1)	<40 (<0.1)		1.09 (0.55, 2.17)	0.72 (0.29, 1.77) p=0.342
No Formal Education	<40 (<0.1)	<40 (<0.1)		1.03 (0.25, 4.30)	0.84 (0.13, 5.26) p=0.801
No Answer	545 (1.3)	266 (1.1)		1.11 (0.93, 1.31)	1.04 (0.81, 1.33) p=0.683
Pre-infection functional performance categories**			< 2.2e-16		
No Functional Performance Difficulty	33933 (83.5)	23494 (94.7)		1.00	1.00
Some Functional Performance Difficulty	4106 (10.1)	926 (3.7)		3.07 (2.79, 3.38)	1.01 (0.62, 1.63) p=0.976

Dependent on Others for Care	2616 (6.4)	389 (1.6)		4.65 (4.05, 5.37)	1.38 (0.84, 2.27) *p*=0.096
Pre-infection OT CPT codes billed (median (IQR))	0 (1)	0 (0)	*p* < 2.2e-16		
(mean (SD))	4.45 (17.78)	2.06 (10.22)			
0				1.00	1.00
5				1.85 (1.71, 2.00)	0.99 (0.99, 1.01), 0.479
20				2.14 (1.95, 2.34)	0.99 (0.95, 1.03), 0.479
80				2.29 (2.04, 2.57	0.96 (0.82, 1.12), 0.479
700				3.62 (2.35, 4.65)	0.68 (0.17, 2.76) 0.479
Self-rated ability to complete physical daily activities			< 0.000		
Completely	21740 (52.2)	16608 (66.9)		1	1
Mostly	7715 (19.0)	3821 (15.4)		1.56 (1.47, 1.65)	1.05 (0.97, 1.13) *p*=0.128
Moderately	6415 (15.8)	2542 (10.2)		1.95 (1.83, 2.09)	1.02 (0.93, 1.12) *p*=0.547
A little	4123 (10.1)	1451 (5.8)		2.20 (2.02, 2.39)	0.95 (0.84, 1.07), 0.228
Not at all	661 (1.6)	254 (1.0)		2.01 (1.66, 2.44)	0.83 (0.64, 1.06), 0.046
No Answer	271 (0.7)	133 (0.5)		1.53 (1.20, 2.08)	1.00 (0.72, 1.42), 0.949
Self-rated social role performance and satisfaction (composite measure)			< 2.2e-16		
2	5197 (12.8)	3955 (15.9)		0.91 (0.85, 0.97)	1.04 (094, 1.10) *p*=0.326
3	5259 (12.9)	3921 (15.8)		0.93 (0.86, 0.99) *p*=0.004	1.03 (0.94, 1.10) *p*=0.433
4	8881 (21.8)	6129 (24.7)		1	1
5	6424 (15.8)	3728 (15.02)		1.19 (1.11, 1.37)	1.06 (0.98, 1.20) *p*=0.066
6	6668 (16.4)	3481 (14.03)		1.32, (1.23, 1.42)	1.00 (0.91, 1.10) *p*=0.897
7	3629 (8.9)	1735 (7.0)		1.44 (1.32, 1.57)	0.96 (0.85, 1.10) *p*=0.377
8	2911 (7.2)	1158 (4.7)		1.73 (1.57, 1.92)	1.01 (0.88, 1.20) *p*=0868
9	1106 (2.7)	452 (1.8)		1.69 (1.45, 1.96)	0.93 (0.75, 1.10) *p*=0.345
10	580 (1.4)	250 (1.0)		1.60 (1.31, 1.69)	0.78 (0.59, 1.10) *p*=0.021
Self-rated mental health (cognition and mood)			<0.000		
Excellent	8057 (19.8)	6092 (24.6)		1	1
Very Good	13078 (32.2)	8950 (36.1)		1.10, (1.04, 1.17)	1.04 (0.96, 1.15), *p*=0.215
Good	11924 (29.3)	6286 (25.3)		1.43 (1.35, 1.52)	1.14 (1.04, 1.25), *p*>0.000
Fair	6009 (14.8)	2781 (11.2)		1.63 (1.52, 1.76)	1.02 (0.91, 1.15) *p*=0.609
Poor	1279 (3.1)	545 (2.2)		1.77 (1.55, 2.04)	1.11 (0.89, 1.36) *p*=0.213
No Answer	308 (3.1)	155 (0.6)		1.50 (1.16, 1.95)	0.97 (0.70, 1.33) *p*=0.778

224   Table 1 Legend: Distributions, central tendency, spread, unadjusted odds ratios and adjusted

225   odds ratios of sample demographics, pre-infection symptom totals, and pre-infection functional

226   status indicators.

227   *Run with Yeats' continuity correction.

228   **From the "Finding of functional performance and activity" EHR observations.

Pre-infection daily functioning

EHR-recorded billing code indicators of daily functioning showed higher proportion of

performance difficulty in the long COVID group (10% vs. 4%) and dependence on others for

care (6% vs. 2%). Participants in the long COVID group also more frequently selected lower

ratings on baseline physical activity ability, social score, and mental health. However, when

adjusted for demographics and pre-infection symptom occurrences, only lower self-rated

mental/cognitive health increased the odds of developing long COVID (Table 1).

Pre-infection occurrences of common long COVID symptoms and conditions

The range of pre-infection occurrences of common long COVID symptoms was between

0 and 23. The long COVID group had a higher median number of pre-infection (6(6) vs. 1(4),

Table 1, Figure 2). The proportions of the number of participants with any observations under

these broad categories prior to infection are shown in Table 2 and S1 Appendix E, Figure E.2.

There were fewer than 500 incidences of anosmia with sparse cells by group, so this symptom

was not included in the model. There were no pre-infection observations of POTS, PVFS,

ageusia.

Table 2. Pre-infection prevalences and odds of common post-COVID symptoms by long

COVID group.

Long COVID symptom/condition group	Recovered n(%)	Long COVID cases n(%)	OR (99% CI) P	AOR (99% CI) P
Joint Pain	7626 (30)	25675 (62)	3.74 (3.58, 3.91) 0.00	1.32 (0.82, 2.11) 0.13
Chest Pain	5054 (20)	19400 (47)	3.20 (3.34, 3.68) 0.00	1.03 (0.64, 1.64) 0.89
Abdominal	5237 (21)	18803 (45)	3.17 (3.02, 3.33) 0.00	1.06 (0.66, 1.70) 0.75
Sleep	3902 (15)	18241 (44)	4.29 (4.07, 4.52) 0.00	1.53 (0.96, 2.45) 0.02
Anxiety	4148 (16)	17717 (43)	3.80 (3.61, 3.99) 0.00	1.36 (0.85, 2.18) 0.09
Depression	3634 (14)	16639 (40)	3.99 (3.79, 4.21) 0.00	1.31 (0.81, 2.10) 0.15
Cough	3977 (16)	15833 (38)	3.31 (3.15, 3.49) 0.00	1.08 (0.68, 1.74) 0.66
Dyspnea	3308 (13)	14670 (35)	3.64 (3.44, 3.45) 0.00	1.13 (0.70, 1.81) 0.51
Fatigue	3339 (13)	14172 (34)	3.42 (3.23, 3.61) 0.00	1.11 (0.70. 1.79) 0.54
Rash/Skin	2965 (12)	11488 (28)	2.86 (2.70, 3.04) 0.00	1.27 (0.79, 2.03) 0.19
Headache	2603 (10)	11051 (27)	3.16 (2.97, 3.36) 0.00	0.99 (0.61, 1.59) 0.95
Dizziness	2398 (9)	10388 (25)	3.19 (3.00, 3.40) 0.00	1.03 (0.64, 1.66) 0.86
Diarrhea	2076 (8)	9273 (22)	3.22 (3.01, 3.45) 0.00	1.07 (0.67, 1.71) 0.72
Muscle Pain	1253 (5)	6528 (16)	3.60 (3.31, 3.92) 0.00	1.16 (0.72, 1.86) 0.44
Palpitations	1469 (6)	6308 (15)	2.91 (2.69, 1.71) 0.00	1.14 (0.71, 1.84) 0.48
Fever	1390 (5)	5537 (13)	2.63 (2.45, 2.88) 0.00	0.89 (0.55. 1.43) 0.52
Cognition	1117 (4)	5131 (12)	3.03 (2.78, 3.32) 0.00	1.01 (0.63, 1.63) 0.96
Paraesthesia	1169 (5)	5111 (12)	2.88 (2.64, 3.14) 0.00	1.08 (0.67, 1.74) 0.67
Tachycardia	1225 (5)	4946 (12)	2.68 (2.46, 2.92) 0.00	0.85 (0.53, 1.37) 0.39
Menstrual Disorder	1475 (6)	4304 (10)	1.87 (1.73, 2.03) 0.00	1.07 (0.66, 1.73) 0.71
Musculoskeletal Chest Pain	648 (3)	3220 (8)	3.20 (2.87, 3.60) 0.00	1.09 (0.67, 1.77) 0.66
ME/CFS (Chronic Fatigue)	430 (2)	2228 (5)	3.33 (2.90, 3.83) 0.00	1.03 (0.63, 1.70) 0.87
Sexual Dysfunction	231 (1)	847 (3)	2.26 (1.86, 2.75) 0.00	0.99 (0.59, 1.66) 0.97

Table legend: Percent of participants with versus without long COVID with at least one pre-

infection occurrence of common long COVID symptoms, with unadjusted and adjusted odds

ratios of developing long COVID given each symptom/condition.

[Insert Figure 2 about here]

Fig. 2. Number of long COVID symptoms with at least one pre-infection occurrence, by

group. A histogram with side-by-side light (long COVID group) and dark (recovered group)

bars. Shows the differences in distributions of the total number of long COVID symptoms

experienced in each group *before* infection (between 5 years and 28 days). The *Y* axis plots the

number of participants from zero to over 7,500; the *X* axis plots the range of number of pre-

infection symptoms observed in this cohort, from zero to 23. The long COVID group shows a

right-skewed normal curve (mean=6.11 (3.96)), while the recovered group shows an exponential

distribution (mean=2.44 (2.93)).

Correlations were computed to test the pre-infection symptom categories for collinearity

(S1 Appendix E, Table E.1) and grouped by first principal components analysis (Figure E.3).

Depression and anxiety had the highest correlation (r=0.53). Moderate to low correlations were

found between abdominal pain and both diarrhea (r=0.32) and chest pain (r= 0.32), dyspnea with

both chest pain (r= 0.4) and cough (r=0.33), ME/CFS and fatigue (r=0.34), musculoskeletal and

generic chest pain (r=0.33), and sleep and depression (0.31).

Having zero pre-infection occurrences of any common long COVID symptom/condition

was associated with significantly lower adjusted odds of long COVID (AOR: 0.15, 99% CI:

0.04, 0.61, *p*=0.005). No single symptom/condition category increased the adjusted odds of

developing long COVID. Sleep disturbance was the only symptom category that approached

significance (*p*=0.02).

Discussion

This retrospective cohort study is among the first examinations of long COVID among

participants in the *All of Us* Research Program. To our knowledge, it is also the first population-

level study to explore the prevalence of pre-infection occurrences of long COVID symptoms and

functional status levels, and their associations with developing post-COVID symptoms.

These associative findings contribute several insights into the evolving understanding of

the risks of long COVID. First, adjusting for demographics, pre-infection functional status, and

pre-infection occurrences of common long COVID symptoms resulted in an overall shift in

effect size to smaller or null differences between the long COVID and recovered groups. Nearly

all variables were associated with significant unadjusted odds, suggesting that people with long

COVID symptoms after infection were more likely to be older, female, Black or African

American, have less education, more symptoms, lower pre-infection functional status, and lower

self-rated health and function. However, in the full model and holding all covariates at their

most-frequently observed values, only older age, Black or African American racial identity,

female biological sex, variant period at first infection, and lower self-rated mental health were

associated with increased odds of long COVID. These demographic findings are consistent with

other large studies that have found higher prevalence of long COVID by race [35, 37], older age

[36], and female biological sex [35, 36, 38].

This was the first study exploring the role of pre-infection functional status in a large

dataset. In unadjusted odds, we noted different strengths and directions of associations from the

self-reported versus EHR-derived functional status indicators (with lower pre-infection self-rated

mental health/cognition being associated with developing long COVID, while ability to perform

physical daily activities was not). This was somewhat surprising since many of the concepts

contained in the "Finding of Functional Status and Activity" code hierarchy describe physical

(e.g. motor and mobility) performance, with comparatively few cognitive or behavioral

observations. This discrepancy could be due to different latent constructs being measured, or

under-detection of slight or moderate functional decreases in EHR data [50]. In the adjusted

odds, the only self-rating that retained a significant association with long COVID was a middle

("Good") rating of cognition and/or mood. In contrast, the EHR-derived diagnostic codes for

cognitive and mood disorders were non-significant. Self-reported functional impairments in long

COVID are sometimes dismissed [6] or measured with instruments that haven't been validated

with this population [51, 52]. Exploring change from pre- to post-infection using instruments

with relevance to the patient experience may provide more accurate insights.

This exploration of the pre-infection incidences of symptoms and conditions common in

long COVID showed several insights. First, the distributions of these conditions relative to one

another were remarkably similar between long COVID and recovered participants; with a few

exceptions of slight magnitude, the conditions that were more prevalent in participants who later

developed long COVID were also more prevalent in participants who later recovered. However,

there was a striking difference in the *proportion* of each group experiencing each symptom, often

twice as high or more in those who later developed long COVID. Second, no single symptom

contributed significantly to higher adjusted odds of long COVID; only sleep disorders

approached significance. From a diagnostic perspective, this aligns with patient-led research

findings that long COVID is a change in health status from pre-infection including both new and

exacerbated symptoms and conditions compared to pre-COVID, and that symptom presentation

can vary widely between individuals [5, 12, 22]. Many (but not all) participants who developed

long COVID experienced pre-infection health issues found in long COVID phenotypes, with a

wide range of pre-infection symptom burden. Although the risk of long COVID was lower with

no pre-infection symptoms, it was not eliminated. Additionally, although pre-infection symptom

burden increased the unadjusted odds of long COVID, this association was not significant

(except at zero symptoms) when adjusted for demographics, individual symptoms, and pre-

infection daily functioning. Thus, prior symptom burden was neither necessary nor sufficient to

support a singular causal link. Nevertheless, the assessment of pre-infection health, including the

presence of and interaction between ongoing health needs, should be considered in research and

clinical management of long COVID, as previously effective management strategies may require

adjustment. A future analysis of long COVID participants could explore pre- versus post-

infection changes in symptom and condition distribution, exacerbations versus new conditions,

and dimensional reduction to explore and compare phenotypes.

The finding that earlier periods of first infection increased the likelihood of developing

long COVID (S1 Appendix E) calls for additional context and exploration. Presumably, the

opportunities to develop symptoms should have increased with time. This dataset's cut-off date

of July 2022 left little time for people in the last variant period to develop symptoms, and we did

not control for time from first infection to first symptom. Additionally, there is evidence that

vaccination against COVID-19 has some protective effect against long COVID [16, 53]. Thus,

both the date at first infection and the availability of vaccines starting in 2021 may have

influenced the number of people who developed symptoms. This may be explored in a future

analysis.

Limitations

This analysis contains several limitations. First, retrospective studies using EHR data are

limited by detection bias [50, 54]. Several sources of relevant functional and disease severity

data are unavailable in *All of Us*, and we were unsuccessful in finding proxies for severity of

infection or long COVID symptoms. In studies of COVID-19 there is difficulty verifying that

participants without COVID-19 diagnostic and lab indicators were truly unexposed, particularly

when using retrospective EHR data. It is possible that many participants who were not selected

for this sample based on EHR evidence of COVID-19 infection had, in reality, been ill with the

virus but were not identifiable as such because they did not report their illness or present for care.

Additionally, admitting diagnoses and narrative notes (where functional status is often described)

are suppressed in *All of Us* for privacy, which precluded connections between incidences of

COVID-19 and hospitalizations and the use of text-to-data methods of data extraction. Use of

Remdesivir, used in acute treatment, was explored as proxy for disease severity but was not used.

The cut-off date for this *All of Us* data set predated FDA approval of Remdesivir by six months.

About 800 participants had been prescribed remdesivir, presumably for experimental or off-label

treatment. This relatively small number was insufficient to make inferences across the sample,

and may have been confounded by access to clinical trials. Prospective studies using tools that

measure disease and symptom severity, as well as the patient experience, continue to be critical.

356 Conclusions

357   Long COVID was more likely for participants with self-reported lower mental or

cognitive health prior to infection. It was less likely for participants with no common long

COVID symptoms occurring before infection, however no single long COVID symptom prior to

first infection affected the odds of long COVID after adjusting for demographics, variant, and

functional status. Long COVID represents a change from baseline functioning and health, even

in people with pre-infection incident symptoms and functional impairments. Functional status

indicators exist in harmonized electronic health records data and have the potential to reveal the

functional impacts of long COVID thus improving diagnostic efficiency. Areas of future study

include developing the diagnostic utility of functional status assessment, uncovering the

relationship between long COVID-related functional changes and ageing, and the changes from

pre-infection to post-infection for people with post-COVID symptoms.

368 Acknowledgements

369   We extend gratitude to *All of Us* participants who generously agreed to share their

370   ongoing health data. The first author also thanks the *All of Us* researcher workbench support staff

371   for their expertise and service, the co-authors for their mentorship, Dr. Beth Fields for editing,

372   Dr. Christine Sorkness for respiratory disease consultation, and Dr. Anoop Mayampurath and the

373   University of Wisconsin Data Science Hub for dataset building support.

Funding

This work was supported by a T32 training grant to the first author from 2021

(5TL1TR002375-05) to 2023 (2TL1TR002375-06) through the University of Wisconsin-

Madison School of Medicine and Public Health, and through a Gertrude Gatson Fund grant

PRJ22AW from the University of Wisconsin-Madison Occupational Therapy Program. The All

of Us Research Program is supported by the National Institutes of Health, Office of the Director:

Regional Medical Centers: 1 OT2 OD026549; 1 OT2 OD026554; 1 OT2 OD026557; 1 OT2

OD026556; 1 OT2 OD026550; 1 OT2 OD 026552; 1 OT2 OD026553; 1 OT2 OD026548; 1

OT2 OD026551; 1 OT2 OD026555; IAA #: AOD 16037; Federally Qualified Health Centers:

HHSN 263201600085U; Data and Research Center: 5 U2C OD023196; Biobank: 1 U24

OD023121; The Participant Center: U24 OD023176; Participant Technology Systems Center: 1

U24 OD023163; Communications and Engagement: 3 OT2 OD023205; 3 OT2 OD023206; and

Community Partners: 1 OT2 OD025277; 3 OT2 OD025315; 1 OT2 OD025337; 1 OT2

OD025276.

Data Availability

389   This study used data from the *All of Us* Research Program's Controlled Tier Dataset

390   version 7, available to authorized users on the Researcher Workbench.

Supporting information

S1 File. Appendices A-E. Appendices with expanded description of cohort discovery; disease, symptom, and functional indicators; quantitative analysis of bias; model fitting notes; expanded results.

S1 Table A.1. Identification of cohort - COVID-19 illness indicators. All laboratory observations, COVID-19 Participant Experience (COPE) survey item responses, and diagnostic codes in SNOMED vocabulary indicating either SARS-CoV-2 infection or COVID-19 illness. Generated using queries for lab values, survey item responses, and diagnostic code incidences via the All of Us Researcher Workbench dataset builder.

S2 Table A.2. Long COVID symptoms/conditions used in Cohort discovery and classification. Standard concept names, concept codes, source vocabularies, and included sub-concepts of long COVID symptoms queried via the *All of Us* Researcher Workbench dataset builder. Hierarchical relationships are as organized in the Athena relational database of the Observational Health Data Sciences and Informatics (OHDSI) and queried from the Observational Medical Outcomes Partnership – Common Data Model (OMOP – CDM) table structures.

S3 Table D.1. Variables for model. Demographic, disease, pre-infection symptom, and pre-infection function variables (names, sources, and formats) used as covariates in regression models. Pre-infection variables are based on entry in the medical record between five years and four weeks prior to first infection date. Each variable's intercept value or level is noted.

S1 Fig C.1. Distribution of propensity scores for pre- versus post-infection enrollment. Jitter point horizontal arrays showing the overlapping ranges in propensity for being in the long

COVID group between the unmatched versus matched pre-infection enrollees ('control') and

post-infection enrollees ('treated'; excluded for the present study's analysis). After matching pre-

and post-enrollment participants, about a 45% overlap is seen between these two groups,

indicating that participants may differ in one or more key demographic and disease aspects

relating to when they enrolled in the study.

S2 Fig C.2. Standardized Mean Difference of demographic and disease characteristics by

pre- versus post-infection enrollment.

S3 Fig. C.3. Distribution of propensity scores for 28-day versus 90-day symptom onset date.

Overlaps in propensity score between the participants classified as cases at 28 days ('control')

and those at 90 days ('treated') is visualized in Figure S.C.3. There was no overlap between

cases and controls by the 90-day alternative classification scheme (distance = 1.0).

S4 Fig. C.4. Standardized Mean Difference of demographic and disease characteristics by

28-day versus 90-day symptom onset date

S5 Fig. E.1. Time of first infection, by long COVID group. Histogram of the number of

participants with (darker blue) versus without (lighter blue) long COVID by first infection date

through July 2022. The *Y* axis plots the number of participants ascending from zero (bottom) to

over 1,500 (top); the *X* axis is a timeline from (left to right) January 1 2020 through July 31

2022. Both groups show a similar profile; they begin with a steep spike in infections about

March 2020, a large and prolonged peak between April and September 2020, and briefer peaks

between about October 2020 – January 2021 and in January 2022.

S6 Fig. E.2. Proportion of sample with each pre-infection long COVID symptom, by group.

Bar graph of the percentage of long COVID cases (light bars) and recovered participants (dark

bars) with at least one pre-infection mention of each long COVID symptom/condition within the

five years preceding their first COVID infection. The cases show between three to five times

greater prevalence of every pre-infection symptom/condition, however none of these symptoms

were significant predictors of long COVID when adjusted for demographics, disease factors, and

pre-infection functional status.

S4 Table E.1. Correlation matrix of pre-infection symptoms. Table of Pearsons R correlation

coefficients between pre-infection symptoms.

S7 Figure E.3. Symptom correlations grouped by first principal components analysis. Heat

map of first principle components analysis of pre-infection symptom correlations reported in

Table E.1. Darker green colors indicate stronger correlations, and white corresponds to no

correlation.

References

1. National Center for Health Statistics USCB. Household Pulse Survey - Long COVID. 2024.
2. National Academies of Sciences Engineering and Medicine. A Long COVID Definition: A Chronic, Systemic Disease State with Profound Consequences. Washington, DC: The National Academies Press; 2024.
3. Cohen J, Rodgers YVM. Long COVID Prevalence, Disability, and Accommodations: Analysis Across Demographic Groups. J Occup Rehabil. 2024;34(2):335-49.
4. McCorkell L, G SA, H ED, Wei H, Akrami A. Patient-Led Research Collaborative: embedding patients in the Long COVID narrative. Pain Rep. 2021;6(1):e913.
5. Al-Aly Z, Davis H, McCorkell L, Soares L, Wulf-Hanson S, Iwasaki A, et al. Long COVID science, research and policy. Nat Med. 2024;30(8):2148-64.
6. Ladds E, Rushforth A, Wieringa S, Taylor S, Rayner C, Husain L, et al. Persistent symptoms after Covid-19: qualitative study of 114 "long Covid" patients and draft quality principles for services. BMC Health Serv Res. 2020;20(1):1144.
7. Mendelson M, Nel J, Blumberg L, Madhi SA, Dryden M, Stevens W, et al. Long-COVID: An evolving problem with an extensive impact. S Afr Med J. 2020;111(1):10-2.
8. The Lancet. Facing up to long COVID. Lancet. 2020;396(10266):1861.
9. Crook H, Raza S, Nowell J, Young M, Edison P. Long covid—mechanisms, risk factors, and management. BMJ. 2021;374:n1648.
10. Zhang H, Zang C, Xu Z, Zhang Y, Xu J, Bian J, et al. Data-driven identification of post-acute SARS-CoV-2 infection subphenotypes. Nat Med. 2023;29(1):226-35.
11. Davis EH, Assaf SG, Mccorkell L, Wei H, Low JR, Re'Em Y, et al. Characterizing long COVID in an international cohort: 7 months of symptoms and their impact. eClinicalMedicine. 2021;38:101019.
12. Davis EH, Mccorkell L, Vogel MJ, Topol JE. Long COVID: major findings, mechanisms and recommendations. Nature Reviews Microbiology. 2023;21(3):133-46.
13. Nielsen TB, Leth S, Pedersen M, Harbo HD, Nielsen CV, Laursen CH, et al. Mental Fatigue, Activities of Daily Living, Sick Leave and Functional Status among Patients with Long COVID: A Cross-Sectional Study. International journal of environmental research and public health. 2022;19(22).
14. Carter SJ, Baranauskas MN, Raglin JS, Pescosolido BA, Perry BL. Functional Status, Mood State, and Physical Activity Among Women With Post-Acute COVID-19 Syndrome. Int J Public Health. 2022;67:1604589.
15. Vanichkachorn G, Newcomb R, Cowl CT, Murad MH, Breeher L, Miller S, et al. Post-COVID-19 syndrome (long haul syndrome): Description of a multidisciplinary clinic at Mayo Clinic and characteristics of the initial patient cohort. Mayo Clin Proc. 2021;96(7):1782-91.
16. Brannock DM, Chew FR, Preiss JA, Hadley CE, Mcmurry AJ, Leese JP, et al. Long COVID risk and pre-COVID vaccination: An EHR-based cohort study from the RECOVER program. medRxiv. 20221007 ed2022.
17. Gentilotti E, Górska A, Tami A, Gusinow R, Mirandola M, Rodríguez Baño J, et al. Clinical phenotypes and quality of life to define post-COVID-19 syndrome: a cluster analysis of the multinational, prospective ORCHESTRA cohort. EClinicalMedicine. 2023;62:102107.
18. Mina Y, Enose-Akahata Y, Hammoud DA, Videckis AJ, Narpala SR, O'Connell SE, et al. Deep Phenotyping of Neurologic Postacute Sequelae of SARS-CoV-2 Infection. Neurol Neuroimmunol Neuroinflamm. 2023;10(4).
19. Prabhakaran D, Day G, Munipalli B, Rush B, Pudalov L, Niazi S, et al. Neurophenotypes of COVID-19: risk factors and recovery trajectories. Brain Behav Immun Health. 2022.
20. Reese JT, Blau H, Casiraghi E, Bergquist T, Loomba JJ, Callahan TJ, et al. Generalisable long COVID subtypes: findings from the NIH N3C and RECOVER programmes. EBioMedicine. 2023;87:104413.

21.   Al-Aly Z, Xie Y, Bowe B. High-dimensional characterization of post-acute sequelae of COVID-
19. Nature. 2021;594(7862):259-64.
22.   Thaweethai T, Jolley ES, Karlson WE, Levitan BE, Levy B, Mccomsey AG, et al. Development
of a Definition of Postacute Sequelae of SARS-CoV-2 Infection. JAMA. 2023;329(22):1934.
23.   Hou Y, Gu T, Ni Z, Shi X, Ranney ML, Mukherjee B. Global Prevalence of Long COVID, its
Subtypes and Risk factors: An Updated Systematic Review and Meta-Analysis. medRxiv. 2025.
24.   Callard F, Perego E. How and why patients made Long Covid. Soc Sci Med. 2021;268:113426.
25.   Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-
19 condition by a Delphi consensus. Lancet Infect Dis. 2022;22(4):e102-e7.
26.   Mandel HL, Colleen G, Abedian S, Ammar N, Bailey LC, Bennett TD, et al. Risk of post-acute
sequelae of SARS-CoV-2 infection associated with pre-coronavirus disease obstructive sleep apnea
diagnoses: an electronic health record-based analysis from the RECOVER initiative. Sleep. 2023.
27.   Hall JP, Kurth NK, McCorkell L, Goddard KS. Long COVID Among People With Preexisting
Disabilities. Am J Public Health. 2024;114(11):1261-4.
28.   Kuper H, Smythe T. Are people with disabilities at higher risk of COVID-19-related mortality?: a
systematic review and meta-analysis. Public Health. 2023;222:115-24.
29.   National institutes of Health. All of Us, Curated data repository version 7, Controlled Tier
C2022Q4R9. In: Health NIo, editor.: National Institutes of Health; 2022-2025.
30.   Ramirez AH, Sulieman L, Schlueter DJ, Halvorson A, Qian J, Ratsimbazafy F, et al. The All of
Us Research Program: Data quality, utility, and diversity. Patterns (N Y). 2022;3(8):100570.
31.   Zeng C, Schlueter DJ, Tran TC, Babbar A, Cassini T, Bastarache LA, et al. Comparison of
phenomic profiles in the All of Us Research Program against the US general population and the UK
Biobank. J Am Med Inform Assoc. 2024;31(4):846-54.
32.   Centers for Disease Control and Prevention. Long COVID or Post-COVID Conditions 2023
[Available from: https://www.cdc.gov/covid/long-term-
effects/?CDC_AAref_Val=https://www.cdc.gov/coronavirus/2019-ncov/long-term-effects/.
33.   Centers for Disease Control and Prevention. New ICD-10-CM code for Post-COVID Conditions,
following the 2019 Novel Coronavirus (COVID-19). In: Prevention CfDCa, editor. 2021.
34.   Markov PV, Ghafari M, Beer M, Lythgoe K, Simmonds P, Stilianakis NI, et al. The evolution of
SARS-CoV-2. Nat Rev Microbiol. 2023;21(6):361-79.
35.   Jacobs MM, Evans E, Ellis C. Racial, ethnic, and sex disparities in the incidence and cognitive
symptomology of long COVID-19. J Natl Med Assoc. 2023;115(2):233-43.
36.   Vasilevskaya A, Mushtaque A, Tsang MY, Alwazan B, Herridge M, Cheung AM, et al. Sex and
age affect acute and persisting COVID-19 illness. Sci Rep. 2023;13(1):6029.
37.   Khullar D, Zhang Y, Zang C, Xu Z, Wang F, Weiner MG, et al. Racial/ethnic disparities in post-
acute sequelae of SARS-CoV-2 infection in New York: an EHR-based cohort study from the RECOVER
program. J Gen Intern Med. 2023;38(5):1127-36.
38.   Shah DP, Thaweethai T, Karlson EW, Bonilla H, Horne BD, Mullington JM, et al. Sex
Differences in Long COVID. JAMA Netw Open. 2025;8(1):e2455430.
39.   Ford ND, Slaughter D, Edwards D, Dalton A, Perrine C, Vahratian A, et al. Long COVID and
Significant Activity Limitation Among Adults, by Age — United States, June 1–13, 2022, to June 7–19,
2023. MMWR Morb Mortal Wkly Rep 2023. 2023;72:866-970.
40.   Izaguirre P, Arakaki É, Boero JV, Zalazar Á, Ghirlanda M, Caruso D. Functional Status in Older
Adults Following Hospitalization for Covid-19: A Cohort Study. Ann Geriatr Med Res. 2023.
41.   U.S. Department of Health and Human Services: Office of Disease Prevention and Health
Promotion. Healthy People 2030: Social Determinants of Health 2025 [Available from:
https://odphp.health.gov/healthypeople/priority-areas/social-determinants-health/literature-summaries.
42.   Case KR, Wang CP, Hosek MG, Lill SF, Howell AB, Taylor BS, et al. Health-related quality of
life and social determinants of health following COVID-19 infection in a predominantly Latino
population. J Patient Rep Outcomes. 2022;6(1):72.

43.   Eligulashvili A, Darrell M, Gordon M, Jerome W, Fiori KP, Congdon S, et al. Patients with
unmet social needs are at higher risks of developing severe long COVID-19 symptoms and
neuropsychiatric sequela. Sci Rep. 2024;14(1):7743.
44.   Centers for Medicare and Medicaid Services. Billing and Coding: Therapy Evaluation, Re-
Evaluation and Formal Testing. A53309. Baltimore, MD: DHHS, CMS; 2020 [Available from:
https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=53309.
45.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2022.
46.   R Core Team and contributors worldwide. The R Stats Package. 4.3.0-4.4.3 ed. CRAN: R
Foundation for Statistical Computing; 2025. p. R statistical functions.
47.   Harrell FE. Regression Modeling Strategies. CRAN; 2024.
48.   Wei T. Visualization of a Correlation Matrix (corrplot). 0.92 ed. CRAN: R Foundation for
Statistical Computing; 2021.
49.   Hadley Wickham MA, Jennifer Bryan, Winston Chang, Lucy D'Agostino McGowan, Romain
François, Garrett Grolemund, Alex Hayes, Lionel Henry, Jim Hester, Max Kuhn, Thomas Lin Pedersen,
Evan Miller, Stephan Milton Bache, Kirill Müller, Jeroen Ooms, David Robinson, Dana Paige Seidel,
Vitalie Spinu, Kohske Takahashi, Davis Vaughan, Claus Wilke, Kara Woo, Hiroaki Yutani. Welcome to
the tidyverse. 2.0.0 ed. CRAN: R Foundation for Statistical Computing; 2019.
50.   Deer RR, Rock AM, Vasilevsky N, Carmody L, Rando H, Anzalone JA, et al. Characterizing
Long COVID: Deep Phenotype of a Complex Condition. eBioMedicine. 2021;74:103722.
51.   Gutzeit J, Weiß M, Nürnberger C, Lemhöfer C, Appel KS, Pracht E, et al. Definitions and
symptoms of the post-COVID syndrome: an updated systematic umbrella review. Eur Arch Psychiatry
Clin Neurosci. 2024.
52.   Lemhöfer C, Appel SK, Häuser W, Hettich N, Kohls M, Polidori CM. Post-COVID: Alles eine
Frage der Definition? DMW - Deutsche Medizinische Wochenschrift. 2022;147(21):1391-7.
53.   Mukherjee S, Singer T, Venkatesh A, Choudhury NA, Perez Giraldo GS, Jimenez M, et al.
Vaccination prior to SARS-CoV-2 infection does not affect the neurologic manifestations of long
COVID. Brain Commun. 2025;7(1):fcae448.
54.   Rando HM, Bennett TD, Byrd JB, Bramante C, Callahan TJ, Chute CG, et al. Challenges in
defining Long COVID: Striking differences across literature, Electronic Health Records, and patient-
reported information. medRxiv. 2021.

Figure 1

Figure 2