Prevalence and Determinants of Hypertension Among People Living with HIV on Antiretroviral Therapy: A Cross-Sectional Study at Mwananyamala Hospital, Tanzania.

Meshack Morice[1,2], Jacqueline Mgumia[1]

1.Department of Sociology and Anthropology, University of Dar es Salaam, Dar es Salaam, Tanzania.

2.Department of Community Medicine, Kairuki University, Dar es Salaam, Tanzania.

Co-author: Jacqueline Mgumia, ORCID iD: 0000-0001-6927-7372, P.O. Box 35091, Dar es Salaam, Tanzania. Phone: +255 754 242 323. Email: jhm11@albion.edu.

Corresponding Author: Meshack Morice, ORCID iD: 0009-0007-2236-9852, P.O. Box 65300, Dar es Salaam, Tanzania, Phone: +255 764 910 402, Email: meshackmorice@gmail.com.

ABSTRACT

Background: Hypertension is an increasingly recognized comorbidity among people living with HIV (PLHIV) receiving antiretroviral therapy (ART), contributing substantially to cardiovascular morbidity. As life expectancy among PLHIV improves, understanding both traditional and HIV-related determinants of hypertension is essential. This study assessed the prevalence, associated factors, and hypertension knowledge among PLHIV on ART in Tanzania.

Methods: A hospital-based cross-sectional study was conducted among 296 PLHIV on ART for more than six months at Mwananyamala Regional Referral Hospital. Data were collected using structured questionnaires and clinical records. Blood pressure was measured according to American Heart Association guidelines. Logistic regression analyses were performed to identify factors independently associated with hypertension at $p<0.05$.

Results: The prevalence of hypertension was 32.8%. Knowledge of hypertension was moderate (mean= 53.9%), with only 10.8% correctly identifying normal blood pressure levels. Age ≥ 55 years (aOR = 4.786, $p < 0.001$) and cigarette smoking (aOR = 6.159, $p = 0.020$) were associated with greater risk of hypertension. Long duration since HIV diagnosis (>10 years: aOR = 3.090, $p = 0.011$) was also associated with increased odds of hypertension, whereas shorter ART duration (<5 years: aOR = 0.106, $p = 0.040$) was associated with lower risk.

Conclusions: Hypertension is prevalent among PLHIV on ART in Tanzania, influenced by both traditional and HIV-related factors. Integrating hypertension screening and education within HIV care services is essential. Strengthening community engagement in chronic disease screening can enhance early detection, improve awareness, and promote preventive behaviours, ultimately reducing the dual burden of HIV and non-communicable diseases.

Keywords: HIV, Hypertension, Antiretroviral Therapy, Tanzania.

INTRODUCTION

Hypertension, defined as blood pressure ≥140 mmHg systolic or ≥90 mmHg diastolic or use of antihypertensive medication, is a growing public health concern globally, affecting a quarter of individuals living with HIV in the previous decade and contributing to cardiovascular diseases (CVD), stroke, heart failure, eye problems (blindness) and other health-related issues[1–4] . Among people living with HIV (PLHIV), the advent of antiretroviral therapy (ART) has transformed HIV into a manageable chronic condition, significantly improving survival rates [5–7]. However, this prolonged survival has shifted morbidity patterns from opportunistic infections to non-communicable diseases, notably hypertension [8]. Studies suggest PLHIV on ART face a higher hypertension risk than the general population, potentially due to HIV-induced inflammation, ART side effects, and traditional risk factors such as alcohol drinking and cigarette smoking [9].

In 2020, global prevalence of hypertension among PLHIV on ART was reported to be 23.6% [10], with the prevalence in Sub-Saharan Africa ranging from 19.6% to 53% [11, 12]. The development of hypertension in PLHIV is multifactorial. With traditional risk factors such as age, obesity or overweight, excessive alcohol consumption, cigarette smoking, sedentary lifestyle, and family history of hypertension having been consistently associated with increased hypertension risk [1, 9, 10]. These risk factors are often more prevalent or more impactful among PLHIV due to intersecting socioeconomic, behavioural, and clinical challenges. Beyond traditional risk factors, HIV-related factors have also been associated with changes in blood pressure among people living with HIV (PLHIV). These factors include CD4 cell count, duration of HIV infection, and duration of antiretroviral therapy (ART) use. These blood pressure changes may result from prolonged immune activation and chronic inflammation caused by HIV, as well as ART-induced metabolic changes [9, 12, 13].

In Tanzania, where HIV prevalence declined from 7% in 2003 to 4.4% in 2023 among adults aged 15–49 [14, 15], hypertension among PLHIV remains understudied, yet it represents a growing public health concern even in the general population. Recent nationally representative data from the 2022 Tanzania Demographic and Health Survey indicate that the prevalence of hypertension among adults aged 15-49 years is 11%, and it is strongly associated with increasing age and overweight/obesity [16]. Another study reported a 34.6% prevalence of hypertension among PLHIV, suggesting a substantially higher burden in this population [17]. However, evidence on HIV-related factors (i.e., CD4 count, ART duration, HIV staging, and HIV diagnosis duration) and hypertension risk is inconsistent, with some studies finding no association [11, 18, 19], while others report strong links [12, 20–22]. Knowledge gaps about

hypertension among PLHIV further complicate prevention efforts. Limited awareness of hypertension [22] hinders

behavioural interventions, yet few studies in Tanzania have assessed this. Addressing these gaps is critical, as

hypertension is a primary contributor to CVD, a leading cause of mortality globally [1]. Given the significant burden

of hypertension among PLHIV, this study aimed to evaluate the determinants of hypertension among PLHIV on

ART, examining risk factors and hypertension knowledge to inform targeted interventions.

MATERIALS AND METHODS

Study Design and Setting

This quantitative, hospital-based cross-sectional study was conducted at Mwananyamala Regional Referral Hospital

(MRRH), a major referral facility in Kinondoni District, Dar es Salaam, Tanzania. MRRH's Care and Treatment

Clinic (CTC) records approximately 26,000 total patient visits per year, with an average of 50–100 daily visits,

making it an ideal site for studying hypertension in this population.

Study population

This study enrolled PLHIV aged ≥18 years, on ART for >6 months, attending the CTC during the study period.

Pregnant women and those on contraceptive pills were excluded due to potential blood pressure confounding effects.

Eligibility was confirmed using CTC records.

Sample size and sampling technique

The sample size was calculated using the Cochran's formula, based on a previous prevalence of 26% from

prevalence from a Tanzanian study [22], with adjustments for 10% non-response rate. The final sample size was 329

participants. Simple random sampling was employed: daily CTC patient lists were generated, each patient assigned a

random number, and 329 were selected using a computer-generated random number generator. Of these, 296

participated. Potential recall bias in self-reported data (e.g., smoking, alcohol use, physical activity) was mitigated

by structured, pretested questionnaires with clear, simple questions to improve recall accuracy.

Variables

The dependent variable was hypertension. Independent variables included traditional risk factors (age, gender,

smoking, alcohol use, BMI, family history, physical activity), HIV-related factors (duration of HIV diagnosis, CD4

count, WHO stage, ART duration), and knowledge level. There were no missing data for any variables analysed,

including sociodemographic characteristics, HIV-related factors, and knowledge of hypertension, as all 296

participants provided complete responses and measurements.

Data collection procedures

Data were collected from September 10, 2024 to March 29, 2025, through structured questionnaires administered in Swahili during one-on-one interviews in a private room post-clinic visit. Sociodemographic data (age, gender, education, occupation, smoking, alcohol use, physical activity, family history of hypertension) and hypertension knowledge were self-reported. Hypertension knowledge was assessed using a structured 10-item questionnaire adapted from the Hypertension Knowledge-Level Scale (HK-LS) [23], with some of the questions modified to fit the local study context and population under investigation. The tool included questions on: (1) normal blood pressure values; (2–6) behavioural risk factors including cigarette smoking, alcohol consumption, weight gain, excessive salt intake, and lack of physical exercise; and (7–10) complications of hypertension including heart problems, visual disturbances, stroke, and kidney disease. Each correct response was awarded one point and summed to generate a percentage score (0–100%). Categories of general knowledge levels were low (0–30%), moderate (40–70%), and good (80–100%). HIV-related factors (duration of HIV diagnosis, CD4 count, WHO HIV clinical stage, and ART duration) were extracted from CTC records.

Blood Pressure Measurement

Blood pressure was measured by two registered nurses using a calibrated sphygmomanometer, following American Heart Association guidelines [24]. Participants rested for 5 minutes, seated with feet flat, arm supported at heart level. Two readings, 1–2 minutes apart, were averaged from two separate occasions. Hypertension was defined as ≥140/90 mmHg or current antihypertensive use.

Anthropometric Measurement

Height and weight were measured using a Weight and Height Scale, and BMI was calculated (kg/m²) using a mobile application. BMI categories were: underweight (<18.5), normal (18.5–24.9), overweight (25–29.9), obese (≥30).

Data Analysis

Data were analysed using Statistical Package for the Social Sciences (SPSS) 26. Descriptive statistics (frequencies and crosstabs) were used to summarise participant characteristics and determine the prevalence of hypertension. Knowledge was evaluated through frequencies and composite scores. Bivariate logistic regression analysis was conducted to examine the association between each independent variable and hypertension. Subsequently, variables were included in the multivariate logistic regression model to adjust for potential confounders. Odds ratios (ORs) with 95% confidence intervals (CIs) and p-values <0.05 were considered statistically significant.

RESULTS

Table 1 presents the sociodemographic and clinical characteristics of the study participants by hypertension status. A total of 296 participants were enrolled in the study. The majority were female (66.6%), and more than one-third were aged 45–54 years (34.8%). Most participants had attained primary education (66.9%) and were business owners (48.6%), while one-third were unemployed (33.1%). Regarding lifestyle characteristics, 7.8% participants reported smoking cigarettes, 23.6% reported alcohol consumption, and the majority were either not very active or moderately active (89.8%). A family history of hypertension was reported by 21.3% participants. More than half of the study population was overweight or obese 58.8%.

The overall prevalence of hypertension was 32.8%. Hypertension prevalence increased with age, with the highest proportion observed among participants aged 55 years and above (56.3%). Higher prevalence was also observed among participants with no formal education (59.1%), unemployed individuals (41.8%), those who were physically inactive (39.1%), those with a family history of hypertension (42.9%), and obese participants (45.8%).

Table 1: Participant Characteristics with Hypertension status

Characteristic	Total n (%)	Hypertensive n (%)	Non-hypertensive n (%)
Overall	296 (100)	97 (32.8)	199 (67.2)
Age (years)			
18–24	12 (4.1)	2 (16.7)	10 (83.3)
25–34	32 (10.8)	6 (18.8)	26 (81.3)
35–44	85 (28.7)	21 (24.7)	64 (75.3)
45–54	103 (34.8)	32 (31.1)	71 (68.9)
≥55	64 (21.6)	36 (56.3)	28 (43.8)
Gender			
Female	197 (66.6)	64 (32.5)	133 (67.5)
Male	99 (33.4)	33 (33.3)	66 (66.7)
Level of education			
No formal education	22 (7.4)	13 (59.1)	9 (40.9)
Primary education	198 (66.9)	60 (30.3)	138 (69.7)
Secondary education	58 (19.6)	18 (31.0)	40 (69.0)
High school	4 (1.4)	0 (0.0)	4 (100.0)
College degree	14 (4.7)	6 (42.9)	8 (57.1)
Occupation			
Business owner	144 (48.6)	38 (26.4)	106 (73.6)
Employed	46 (15.5)	15 (32.6)	31 (67.4)
Farmer	8 (2.7)	3 (37.5)	5 (62.5)

Unemployed	98 (33.1)	41 (41.8)	57 (58.2)
Lifestyle factors			
Cigarette smoker	23 (7.8)	3 (13.0)	20 (87.0)
Alcohol drinker	70 (23.6)	25 (35.7)	45 (64.3)
Physical activity			
Not very active	133 (44.9)	52 (39.1)	81 (60.9)
Moderately active	133 (44.9)	38 (28.6)	95 (71.4)
Very active	30 (10.1)	7 (23.3)	23 (76.7)
Family history of hypertension			
Yes	63 (21.3)	27 (42.9)	36 (57.1)
No	233 (78.7)	70 (30.0)	163 (70.0)
BMI (kg/m²)			
Underweight (<18.5)	15 (5.1)	5 (33.3)	10 (66.7)
Normal (18.5–24.9)	107 (36.1)	25 (23.4)	82 (76.6)
Overweight (25–29.9)	102 (34.5)	34 (33.3)	68 (66.7)
Obese (≥30)	72 (24.3)	33 (45.8)	39 (54.2)

Knowledge was moderate (mean 53.85%), with 26.4% scoring low (0–30%), 52% moderate (40–70%), and 21.6%

good (80–100%) (Figure 1).

Figure 1: Histogram showing the distribution of hypertension knowledge scores (%)

Only 10.8% knew normal blood pressure, while awareness of risk factors like exercise (90.9%) and weight gain

(73.0%) was higher than smoking (22.6%). Complications awareness ranged from 46.3% (kidney problems) to

68.6% (heart problems) (Table 2)

Table 2: Hypertension knowledge assessment parameters in PLHIV on ART

	Frequency (n)	Percentage (%)
Normal Blood pressure		
No	264	89.2
Yes	32	10.8
Cigarette smoking		
No	229	77.4
Yes	67	22.6
Alcohol drinking		
No	139	47.0
Yes	157	53.0
Weight gain		
No	80	27.0
Yes	216	73.0
Excessive salt use		
No	113	38.2
Yes	183	61.8
Physical exercise		
No	27	9.1
Yes	269	90.9
Heart problems		
No	93	31.4
Yes	203	68.6
Visual disturbances		
No	120	40.5
Yes	176	59.5
Stroke		
No	142	48.0
Yes	154	52.0
Kidney problems		
No	159	53.7
Yes	137	46.3

Table 3 presents the results of the bivariate and multivariate analyses of factors associated with hypertension among

people living with HIV (PLHIV). In the bivariate analysis, age ≥55 years was significantly associated with increased

likelihood of hypertension (COR 3.173, 95% CI 2.325–6.395, p<0.001). Cigarette smoking increased the odds of

hypertension (COR 3.501, 95% CI 1.014–6.085, p=0.047), while being overweight/obese was associated with lower

odds (COR 0.521, 95% CI 0.312–0.870, p=0.013).

After adjusting for potential confounders in the multivariate analysis, age ≥55 years remained significantly

associated with increased odds of hypertension (aOR 4.786, 95% CI 2.987–7.430, p<0.001). Cigarette smoking was

independently associated with increased odds of hypertension (aOR 6.159, 95% CI 1.338–8.356, p=0.020). Duration

with HIV >10 years was significantly associated with higher odds of hypertension (aOR 3.090, 95% CI 1.291–

7.394, p=0.011) and ART use for less than 5 years was associated with reduced odds of hypertension (aOR 0.106,

95% CI 0.011–0.999, p=0.040). Other variables, including gender, family history of hypertension, alcohol use, BMI,

physical activity level, HIV clinical stage, and CD4 count, were not significantly associated with hypertension

(p>0.05).

Table 3: Bivariate and Multivariate analysis of HIV-related factors associated with HTN (N=296)

	Bivariate		Multivariate	
	COR (95% CI)	P-value	AOR (95% CI)	P-value
Age				
18–34	1		1	
35–44	1.477 (0.594–3.672)	0.402	1.062 (0.355–3.181)	0.914
45–54	2.028 (0.848–4.853)	0.112	2.492 (0.895–6.937)	0.08
≥55	3.173 (2.325–6.395)	**<0.001**	4.786 (2.987–7.430)	**<0.001**
Gender				
Female	1		1	
Male	0.962 (0.576–1.608)	0.884	0.997 (0.463–2.149)	0.994
Family history of hypertension				
No	1		1	
Yes	0.573 (0.323–1.015)	0.056	0.539 (0.273–1.063)	0.074
Cigarrette smoking				
No	1		1	
Yes	3.501 (1.014–6.085)	**0.047**	6.159 (1.338–8.356)	**0.02**
Alcohol use				
No	1		1	
Yes	0.842 (0.479–1.478)	0.548	0.605 (0.286–1.279)	0.188
BMI (kg/m^2)				
Underweight/Normal	1		1	
Overweight/Obese	0.521 (0.312–0.870)	**0.013**	0.716 (0.376–1.361)	0.308
Physical activity level				
Not very active	1		1	
Moderately active	0.474 (0.190–1.184)	0.11	1.003 (0.522–1.928)	0.993
Very active	0.761 (0.301–1.921)	0.563	2.306 (0.692–7.687)	0.174
Duration with HIV (years)				

≤ 5	1		1	
6–10	0.660 (0.337–1.295)	0.227	0.998 (0.429–2.318)	0.996
>10	0.902 (0.479–1.698)	0.749	3.090 (1.291–7.394)	**0.011**
HIV Clinical Staging				
Stage 1	1		1	
Stage 2	0.878 (0.350–2.205)	0.782	0.823 (0.283–2.391)	0.72
Stage 3	0.636 (0.315–1.284)	0.207	0.717 (0.298–1.726)	0.458
Stage 4	0.703 (0.253–1.948)	0.498	0.498 (0.146–1.696)	0.265
CD4 Count (cells/mm³)				
<350	1		1	
350-499	0.904 (0.462–1.769)	0.769	0.680 (0.308–1.501)	0.34
≥500	0.976 (0.519–1.833)	0.939	0.863 (0.402–1.852)	0.706
Duration of ART Use (years)				
>5	1		1	
<5	0.892 (0.476–1.670)	0.072	0.106 (0.011–0.999)	**0.04**

DISCUSSION

This study determined the prevalence of hypertension and its associated factors among people living with HIV

(PLHIV) receiving antiretroviral therapy (ART). The overall prevalence of hypertension in this study was 32.8%,

which is markedly higher than the 11% reported in the general Tanzanian adult population, suggesting a

considerable burden of hypertension among PLHIV [16]. This prevalence is higher than the reported global

prevalence of hypertension among PLHIV, estimated at 23.6% [10], but falls within the wide range reported across

Sub-Saharan Africa (19.6–53%) [11, 12], a region increasingly characterized by the dual burden of infectious and

non-communicable diseases [25]. Compared to previous studies conducted in urban Tanzania, which reported

hypertension prevalence estimates of approximately 25.2% [20] and 34.6% [16], the findings of the present study

indicate a persistently high burden of hypertension in this population. The observed prevalence is comparable to

estimates reported in Ghana (36.7%) [21] and South Africa (53%) [12], yet substantially higher than those reported

in Burundi (17.4%) [13] and Kenya (16%) [18]. These variations across studies may reflect differences in population

age distribution, duration of HIV infection, ART exposure, lifestyle transitions, and the capacity of health systems to

detect and manage non-communicable diseases. Collectively, these findings highlight hypertension among PLHIV

as a significant and emerging public health concern that warrants systematic integration of non-communicable

disease screening and management within HIV care programmes.

Participants' knowledge of hypertension was assessed, revealing moderate overall knowledge levels (53.85%), with

more than one-quarter (26.4%) demonstrating low knowledge. These findings are consistent with previous studies

reporting limited hypertension awareness among PLHIV [11, 26, 27]. This study employed a structured and

validated approach to assessing hypertension knowledge, improving upon prior studies that relied on non-

standardized or vaguely defined measures [22]. By capturing specific domains, including knowledge of normal

blood pressure thresholds, behavioural risk factors, and potential complications, this approach provides a clearer

understanding of existing knowledge gaps. Although this study did not demonstrate a statistically significant

association between knowledge scores and hypertension status, inadequate awareness may still have implications for

long-term prevention, health-seeking behaviour, and treatment adherence. Therefore, strengthening patient education

within HIV clinics remains important, particularly to support integrated prevention and control of non-

communicable diseases within routine HIV care.

Concerning factors associated with hypertension among PLHIV, traditional risk factors such as age and cigarette

smoking have demonstrated a significant association. Increasing age was strongly associated with higher odds of

hypertension, with participants aged ≥55 years having significantly elevated risk of hypertension. This finding is

consistent with extensive epidemiological evidence demonstrating that advancing age remains one of the strongest

predictors of hypertension in both the general population and among PLHIV [16, 17]. Age-related vascular changes,

including arterial stiffness, endothelial dysfunction, and reduced vascular compliance, are well-established

mechanisms contributing to elevated blood pressure [28–31]. Among PLHIV, these age-related processes may be

further compounded by chronic inflammation and immune activation, potentially accelerating vascular ageing [32].

As life expectancy among PLHIV continues to improve due to ART scale-up, the ageing HIV population may

experience a disproportionately higher burden of hypertension and other cardiovascular comorbidities[33]. These

findings underscore the need for early cardiovascular risk assessment within HIV care settings.

Cigarette smoking was also associated with increased odds of hypertension. This finding aligns with previous

studies conducted among PLHIV in Sub-Saharan Africa and other settings [9, 34], where smoking has been

identified as an important modifiable cardiovascular risk factor. The hypertensive effect of smoking may be

mediated through sympathetic nervous system activation, endothelial injury, oxidative stress, and increased arterial

stiffness[35, 36]. In PLHIV, the combined effects of smoking and HIV-related inflammation may synergistically

elevate cardiovascular risk. However, some studies have reported that there is no association between smoking and

hypertension among PLHIV [37, 38], possibly due to underreporting of smoking behaviour, or limited statistical

power to identify such associations, which may be attributed to the small sample size and the low proportion of

patients. Nonetheless, the present findings reinforce the importance of integrating smoking cessation counselling

into routine HIV care as part of comprehensive cardiovascular risk reduction strategies.

HIV-related factors were also significantly associated with hypertension in this study. Longer duration since HIV

diagnosis was independently associated with increased odds of hypertension, consistent with findings from previous

studies [39, 40]. Even though not assessed in this study, biological mechanisms such as chronic immune activation,

persistent inflammation, and endothelial dysfunction have been proposed in the literature to explain the relationship

between long-standing HIV infection and increased risk of vascular damage and elevated blood pressure [41, 42].

However, other studies reported no significant association between HIV duration and hypertension [43, 44], possibly

due to differences in study design and variability in definitions for duration of HIV infection.

ART duration also demonstrated a significant relationship with hypertension risk. Participants with shorter ART

duration (<5 years) had a lower risk of hypertension compared to those on long-term treatment. This finding is

consistent with previous studies indicating that prolonged ART exposure may contribute to hypertension through

mechanisms such as drug-related metabolic toxicity, lipodystrophy, insulin resistance, and dyslipidaemia [45–48].

As ART programs continue to extend life expectancy among PLHIV, the long-term cardiometabolic consequences of

sustained treatment must be proactively addressed through integrated care models. HIV clinical stage was not

significantly associated with hypertension in this study, differing from findings reported by other studies [22, 49].

This may be due to ART's mitigating effects on disease progression.

This study has several limitations. The cross-sectional design limits causal inference between hypertension and

associated factors. But also, a detailed assessment of specific ART regimens was not undertaken. Despite these

limitations, the study provides important insights into the burden and determinants of hypertension among PLHIV.

Conclusions

This study demonstrates that hypertension is a major comorbidity among PLHIV, affecting nearly one-third of

individuals. Although both traditional and HIV-related factors were associated with hypertension risk, the observed

prevalence and key determinants, particularly older age and cigarette smoking, closely mirror patterns documented

in the general urban Tanzanian population. These findings suggest that much of the observed risk may reflect ageing,

accumulated exposure time, and prolonged survival in care rather than HIV-specific biological mechanisms alone.

As life expectancy among PLHIV continues to improve, their cardiovascular risk profile increasingly resembles that

of the broader ageing population within Tanzania's ongoing epidemiological transition. While HIV-related

inflammation and ART-associated metabolic effects remain biologically plausible contributors, the present findings

primarily support an epidemiological explanation grounded in traditional risk accumulation. The presence of

moderate knowledge levels with notable gaps further underscores the need for enhanced patient education within

HIV care settings. Integrating routine hypertension screening, targeted health education, and lifestyle counselling

into HIV care and treatment clinics is essential to improve early detection, strengthen chronic disease management,

and reduce the growing burden of cardiovascular comorbidities among PLHIV in Tanzania.

LIST OF ABBREVIATIONS

- **AHA** – American Heart Association

- **AOR** – Adjusted Odds Ratio

- **ART** – Antiretroviral Therapy

- **BMI** – Body Mass Index

- **CD4** – Cluster of Differentiation 4

- **CI** – Confidence Interval

- **COR** – Crude Odds Ratio

- **CTC** – Care and Treatment Clinic

- **CVD** – Cardiovascular Diseases

- **HIV** – Human Immunodeficiency Virus

- **HK-LS** – Hypertension Knowledge-Level Scale

- **MRRH** – Mwananyamala Regional Referral Hospital

- **PLHIV** – People Living with Human Immunodeficiency Virus

- **REC** – Research Ethics Committee

- **SPSS** – Statistical Package for the Social Sciences

- **WHO** – World Health Organization

DECLARATIONS

Ethical approval and consent to participate

Ethical clearance was obtained from the University of Dar es Salaam Research Ethics Committee (REC) with Ref.No. AB3/12(B), with permissions from regional authorities and MRRH. Written informed consent was obtained from all participants, ensuring confidentiality through coded questionnaires.

Consent for Publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Authors' contributions

Meshack Morice conceptualized the study, collected and analysed data, and wrote the manuscript. Jacqueline Mgumia supervised the study from the beginning to the end and provided critical revisions to the manuscript

Acknowledgments

The author (s) thank Mwananyamala Regional Referral Hospital (MRRH) staff for their support, and participants for their consent and time to participate in this study.

REFERENCES

1. WHO (2025) Cardiovascular diseases (CVDs). https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds). Accessed 7 Jan 2026

2. Fragkou PC, Moschopoulos CD, Dimopoulou D, Triantafyllidi H, Birmpa D, Benas D, Tsiodras S, Kavatha D, Antoniadou A, Papadopoulos A (2023) Cardiovascular disease and risk assessment in people living with HIV: Current practices and novel perspectives. Hellenic J Cardiol 71:42–54. https://doi.org/10.1016/j.hjc.2022.12.013

3.   Ntsekhe M, Baker JV (2023) Cardiovascular Disease Among Persons Living With HIV: New Insights Into
Pathogenesis and Clinical Manifestations in a Global Context. Circulation 147:83–100.
https://doi.org/10.1161/CIRCULATIONAHA.122.057443

4.   WHO (2023) Global report on hypertension: the race against a silent killer - World | ReliefWeb.
https://reliefweb.int/report/world/global-report-hypertension-race-against-silent-killer. Accessed 7 Jan 2026

5.   UNAIDS (2020) UNAIDS report on the global AIDS epidemic shows that 2020 targets will not be met because
of deeply unequal success; COVID-19 risks blowing HIV progress way off course | UNAIDS.
https://www.unaids.org/en/resources/presscentre/pressreleaseandstatementarchive/2020/july/20200706_global
-aids-report. Accessed 7 Jan 2026

6.   Devi SB, Singh TJ, Singh KB, Singh NB, Ningshen R, Singh TB (2020) Survival analysis of people living with
human immunodeficiency virus: a study in a teaching hospital. Int J Adv Med 7:1339–1346.
https://doi.org/10.18203/2349-3933.ijam20203596

7.   Zahra AN, Waluyo A, Yona S, Pakasi TA (2024) Resilience in Relation to Adherence to Antiretroviral Therapy
in People Living With HIV: A Qualitative Study. Glob Qual Nurs Res 11:23333936241233449.
https://doi.org/10.1177/23333936241233449

8.   Bigna JJ, Noubiap JJ (2021) Global burden of hypertension in people living with HIV. BMC Med 19:112.
https://doi.org/10.1186/s12916-021-01981-y

9.   Carter M (2020) A quarter of people living with HIV have hypertension. In: aidsmap.com.
https://www.aidsmap.com/news/jun-2020/quarter-people-living-hiv-have-hypertension. Accessed 7 Jan 2026

10.   Bigna JJ, Ndoadoumgue AL, Nansseu JR, Tochie JN, Nyaga UF, Nkeck JR, Foka AJ, Kaze AD, Noubiap JJ
(2020) Global burden of hypertension among people living with HIV in the era of increased life expectancy: a
systematic review and meta-analysis. J Hypertens 38:1659. https://doi.org/10.1097/HJH.0000000000002446

11.   Derick KI, Khan Z (2023) Prevalence, Awareness, Treatment, Control of Hypertension, and Availability of
Hypertension Services for Patients Living With Human Immunodeficiency Virus (HIV) in Sub-Saharan Africa
(SSA): A Systematic Review and Meta-analysis. Cureus 15:. https://doi.org/10.7759/cureus.37422

12.   Tsuro U, Oladimeji KE, Pulido-Estrada G-A, Apalata TR (2022) Risk Factors Attributable to Hypertension
among HIV-Infected Patients on Antiretroviral Therapy in Selected Rural Districts of the Eastern Cape
Province, South Africa. Int J Environ Res Public Health 19:11196. https://doi.org/10.3390/ijerph191811196

13.   Harimenshi D, Niyongabo T, Preux P-M, Aboyans V, Desormais I (2022) Hypertension and associated factors
in HIV-infected patients receiving antiretroviral treatment in Burundi: a cross-sectional study. Sci Rep 12:1–8.
https://doi.org/10.1038/s41598-022-24997-7

14.   UNAIDS (2020) Country progress report - United Republic of Tanzania, Global AIDS Monitoring 2020.
https://www.unaids.org/sites/default/files/country/documents/TZA_2020_countryreport.pdf. Accessed 7 Jan
2026

15.   THIS 2022-2023 (2023) TANZANIA HIV IMPACT SURVEY. https://www.nbs.go.tz/nbs/takwimu/THIS2022-
2023/THIS2022-2023_Summary_Sheet.pdf. Accessed 7 Jan 2026

16.   Lutambi AM, Mnyagatwa PM, Busunge AJ, Dodo EB, Mwingá GP, Nagai HT, Mkama YJ, Chilongani JE,
Kapyolo EP, Matemba LE, Emidi B (2025) Prevalence and determinants of hypertension among adults of
reproductive age in Tanzania: analysis of a cross-sectional Demographic and Health Survey.
https://doi.org/10.1136/bmjopen-2024-094387

17. Hertz JT, Prattipati S, Kweka GL, Mlangi JJ, Tarimo TG, Mmbaga BT, Thielman NM, Sakita FM, Rubach MP, Bloomfield GS, Manavalan P (2022) Prevalence and predictors of uncontrolled hypertension, diabetes, and obesity among adults with HIV in northern Tanzania. Glob Public Health 17:3747–3759. https://doi.org/10.1080/17441692.2022.2049344

18. Mogaka JN, Sharma M, Temu T, Masyuko S, Kinuthia J, Osoti A, Zifodya J, Nakanjako D, Njoroge A, Otedo A, Page S, Farquhar C (2022) Prevalence and factors associated with hypertension among adults with and without HIV in Western Kenya. PLOS ONE 17:e0262400. https://doi.org/10.1371/journal.pone.0262400

19. Nyangi G, Festo C, Olotu A (2020) The Prevalence of Hypertension and its association with HIV related factors in HIV patients on ART, Bagamoyo district, Eastern Tanzania: Hypertension in adult HIV patients on ART. East Afr J Sci Technol Innov 1:. https://doi.org/10.37425/eajsti.v1i4.72

20. Kato I, Tumaini B, Pallangyo K (2020) Prevalence of non-communicable diseases among individuals with HIV infection by antiretroviral therapy status in Dar es Salaam, Tanzania. PLOS ONE 15:e0235542. https://doi.org/10.1371/journal.pone.0235542

21. Nartey ET, Tetteh RA, Anto F, Sarfo B, Kudzi W, Adanu RM (2023) Hypertension and associated factors among patients attending HIV clinic at Korle-Bu Teaching Hospital. Ghana Med J 57:19–27. https://doi.org/10.4314/gmj.v57i1.4

22. Trifirò S, Cavallin F, Mangi S, Mhaluka L, Maffoni S, Taddei S, Putoto G, Torelli GF (2023) Hypertension in people living with HIV on combined antiretroviral therapy in rural Tanzania. Afr Health Sci 23:129–36. https://doi.org/10.4314/ahs.v23i1.15

23. Alhazmi A, Moafa HN, Kotb M, Sayegh L, Baydhi H, Hazzazi A, Moafa H, Hakami A (2024) Assessing knowledge about hypertension and identifying predictors of inadequate knowledge in Saudi Arabia: A cross-sectional study. PLOS ONE 19:e0299745. https://doi.org/10.1371/journal.pone.0299745

24. AHA (2025) Understanding Blood Pressure Readings. In: www.heart.org. https://www.heart.org/en/health-topics/high-blood-pressure/understanding-blood-pressure-readings. Accessed 7 Jan 2026

25. Modjadji P, Thovhogi N, Sekgala MD, Monyeki KD (2025) Non-Communicable Disease Burden and Dietary Determinants in Women of Reproductive Age in Sub-Saharan Africa: A Scoping Review. Diseases 13:313. https://doi.org/10.3390/diseases13100313

26. Kanyike AM, Nakawuki AW, Akech GM, Kihumuro RB, Kintu TM, Nalunkuma R, Nakandi RM, Mugabi J, Twijukye N, Bwayo D, Katuramu R (2024) Prevalence, Awareness, and Factors Associated with Hypertension Among People Living with HIV in Eastern Uganda. A Multicentre Cross-Sectional Study. HIVAIDS - Res Palliat Care 16:325–335. https://doi.org/10.2147/HIV.S477809

27. Manavalan P, Madut DB, Hertz JT, Thielman NM, Okeke NL, Mmbaga BT, Watt MH (2022) Hypertension among adults enrolled in HIV care in northern Tanzania: comorbidities, cardiovascular risk, and knowledge, attitudes and practices. Pan Afr Med J 41:. https://doi.org/10.11604/pamj.2022.41.285.26952

28. Herzog MJ, Müller P, Lechner K, Stiebler M, Arndt P, Kunz M, Ahrens D, Schmeißer A, Schreiber S, Braun-Dullaeus RC (2025) Arterial stiffness and vascular aging: mechanisms, prevention, and therapy. Signal Transduct Target Ther 10:282. https://doi.org/10.1038/s41392-025-02346-0

29. Çelik MC, Kalçık M, Birgün A, Yetim M, Bekar L, Karavelioğlu Y (2025) Endothelial dysfunction and vascular stiffness: molecular drivers of cardiovascular aging. Explor Cardiol 3:101279. https://doi.org/10.37349/ec.2025.101279

30. Jensen LJ (2024) Functional, Structural and Proteomic Effects of Ageing in Resistance Arteries. Int J Mol Sci 25:2601. https://doi.org/10.3390/ijms25052601

31. Boutouyrie P, Chowienczyk P, Humphrey JD, Mitchell GF (2021) Arterial Stiffness and Cardiovascular Risk in
Hypertension. Circ Res 128:864–886. https://doi.org/10.1161/CIRCRESAHA.121.318061

32. Prakoeswa FRS, Maharani F, Bestari RS, Aisyah R, Ichsan B, Nursanto D, Listiansyah R, Tuanaya MRN
(2025) Aging and HIV: Recent Findings in Contributing Factors. AIDS Res Treat 2025:8814760.
https://doi.org/10.1155/arat/8814760

33. Prakash P, Swami Vetha BS, Chakraborty R, Wenegieme T-Y, Masenga SK, Muthian G, Balasubramaniam M,
Wanjalla CN, Hinton AO, Kirabo A, Williams CR, Aileru A, Dash C (2024) HIV-Associated Hypertension:
Risks, Mechanisms, and Knowledge Gaps. Circ Res 134:. https://doi.org/10.1161/CIRCRESAHA.124.323979

34. Uwanyirigira D, Biracyaza E, Uzabakiriho I, Omolo J, Hakizayezu F, Nzayirambaho M (2023) Prevalence and
Factors Associated with Hypertension Among HIV Positive Patients on Antiretroviral Therapy: A Hospital-
Based Cross-Sectional Study in Rwanda. Vasc Health Risk Manag 19:857–870.
https://doi.org/10.2147/VHRM.S442108

35. Hahad O, Schmitt VH, Arnold N, Keller K, Prochaska JH, Wild PS, Schulz A, Lackner KJ, Pfeiffer N,
Schmidtmann I, Michal M, Schattenberg JM, Tüscher O, Daiber A, Münzel T (2023) Chronic cigarette
smoking is associated with increased arterial stiffness in men and women: evidence from a large population-
based cohort. Clin Res Cardiol 112:270–284. https://doi.org/10.1007/s00392-022-02092-1

36. Gokmen O, Ozgur G (2023) The Effects of chronic smoking on retinal vascular densities and choroidal
thicknesses measured by optical coherence tomography angiography. Eur J Ophthalmol 33:455–462.
https://doi.org/10.1177/11206721221124650

37. Assogba CO, Ale BM, Youm E, Houehanou YC, Sinsin AS, Assiongbon SJE, Ale OAJ, Noubiap JJ, Amidou
SA, Houinato DS (2024) Prevalence and associated factors of hypertension among people living with
HIV/AIDS on antiretroviral therapy in Benin's National Teaching Hospital. Sci Rep 14:17035.
https://doi.org/10.1038/s41598-024-64212-3

38. Mbuthia GW, Magutah K, McGarvey ST (2021) The Prevalence and Associated Factors of Hypertension
among HIV Patients. Int J Hypertens 2021:5544916. https://doi.org/10.1155/2021/5544916

39. Tegegne KD, Adela GA, Kassie GA, Mengstie MA, Seid MA, Zemene MA, Feleke SF, Dejenie TA, Abebe EC,
Anley DT, Dessie AM, Gesese MM, Yimer N, Gebeyehu NA (2023) Prevalence and factors associated with
hypertension among peoples living with HIV in East Africa, a systematic review and meta-analysis. BMC
Infect Dis 23:724. https://doi.org/10.1186/s12879-023-08679-x

40. Melo ES, Antonini M, Costa CRB, Sorensen W, Gir E, Reis RK (2020) Evaluation of cardiovascular risk
factors in people living with HIV in São Paulo, Brazil. J Infect Dev Ctries 14:89–96.
https://doi.org/10.3855/jidc.11326

41. Wei S, Evans PC, Strijdom H, Xu S (2025) HIV infection, antiretroviral therapy and vascular dysfunction:
Effects, mechanisms and treatments. Pharmacol Res 217:107812. https://doi.org/10.1016/j.phrs.2025.107812

42. Perkins MV, Joseph SB, Dittmer DP, Mackman N (2023) Cardiovascular Disease and Thrombosis in HIV
Infection. Arterioscler Thromb Vasc Biol 43:175–191. https://doi.org/10.1161/ATVBAHA.122.318232

43. Godongwana M, De Wet-Billings N (2021) Time to hypertension development among people living with HIV
in South Africa: A longitudinal analysis of the National Income Dynamics Survey (NIDS). Heliyon 7:e07750.
https://doi.org/10.1016/j.heliyon.2021.e07750

44. Denu MKI, Revoori R, Buadu MAE, Oladele O, Berko KP (2024) Hypertension among persons living with
HIV/AIDS and its association with HIV-related health factors. AIDS Res Ther 21:5.
https://doi.org/10.1186/s12981-023-00576-2

45. Thet D, Siritientong T (2020) Antiretroviral Therapy-Associated Metabolic Complications: Review of the
Recent Studies. HIVAIDS - Res Palliat Care 12:507–524. https://doi.org/10.2147/HIV.S275314

46. Ergin HE, Inga EE, Maung TZ, Javed M, Khan S, Ergin HE, Inga EE, Maung TZ, Javed M, Khan S (2020)
HIV, Antiretroviral Therapy and Metabolic Alterations: A Review. Cureus 12:.
https://doi.org/10.7759/cureus.8059

47. Prakash P, Swami Vetha BS, Chakraborty R, Wenegieme T-Y, Masenga SK, Muthian G, Balasubramaniam M,
Wanjalla CN, Hinton AO, Kirabo A, Williams CR, Aileru A, Dash C (2024) HIV-Associated Hypertension:
Risks, Mechanisms, and Knowledge Gaps. Circ Res 134:e150–e175.
https://doi.org/10.1161/CIRCRESAHA.124.323979

48. Acharya S, Karanjkar V, Goyal M, Deshpande PV, Anonymous, Setia MS, Rathod D, Kannan P, Krishna A,
Pathni AK, Palkar A, Harshana A, Anonymous (2025) Prevalence of Hypertension and Its Associated Factors
Among Adult People Living with HIV/AIDS at Anti-Retroviral Treatment (ART) Centers in Mumbai, India.
HIVAIDS - Res Palliat Care 17:105–119. https://doi.org/10.2147/HIV.S499508

49. Kamiński M, Prymas P, Konobrodzka A, Filberek P, Sibrecht G, Sierocki W, Osińska Z, Bogdański P (2020)
Blood pressure in relation to the World Health Organization AIDS clinical staging among adults living in rural
Kenya. HIV AIDS Rev 19:93–98. https://doi.org/10.5114/hivar.2020.96242