Title: Modulation of Corticospinal Excitability and Muscle Synergies During Complex

Locomotor Task in Individuals With and Without Cerebral Palsy: A TMS and EMG Study

Authors: Yosra Cherni[1,2,3], Marco Ghislieri[4,5], Laurent Bouyer[1,2], Catherine Mercier[1,2*]

Affiliations:

(1) Centre interdisciplinaire de recherche en réadaptation et intégration sociale (Cirris), CIUSSS
de la Capitale-Nationale, Québec, Québec, Canada

(2) École des sciences de la réadaptation, Université Laval, Québec, Québec, Canada

(3) Current address : École de Kinésiologie et sciences de l'activité physique, Montréal, Québec,
Canada; Institut de génie biomédical, Montréal, Québec, Canada; Centre de recherche Azrieli du
CHU Sainte Justine, Montréal, Québec, Canada

(4) BIOLAB, Département de l'électronique et des télécommunications, École polytechnique de
Turin, Turin, Italie

(5) Polito[BIO]Med Lab, École polytechnique de Turin, Turin, Italie

Corresponding author: Catherine Mercier, email: catherine.mercier@rea.ulaval.ca

Running title: Corticospinal excitability and muscle synergies during gait in cerebral palsy

Abstract:

Introduction: Studies employing transcranial magnetic stimulation and electromyography suggest that disrupted functional corticospinal connectivity significantly contributes to locomotor impairments. In individuals with Cerebral Palsy (CP), the corticospinal tract has therefore been identified as a potential target for improving gait control. However, it remains uncertain whether this pathway can be further activated given the damage caused by the brain lesion. Moreover, muscle synergies, a cooperative activation of groups of muscles, play an essential role in efficient and adaptive locomotion. Understanding and improving the modulation of these synergies could lead to better rehabilitation strategies for individuals with CP. The objective of this study was to assess whether a complex walking task promotes an increase in corticospinal excitability and a modulation of muscle synergies compared to a simple walking task in individuals with CP.

Methods: Fourteen individuals with CP and fourteen control subjects took part in this study. Each participant performed a simple walking task and a complex walking task (i.e., stepping onto virtual targets) at comfortable speed, in counterbalanced order. Motor evoked potentials (MEPs) in the tibialis anterior muscle were induced using transcranial magnetic stimulation. Muscle synergies were extracted from surface electromyography signals acquired from six key lower-limb muscles during both tasks.

Results: MEPs were elicitable in 11/14 participants for the CP group and 14/14 for control group. In the complex task, MEPs increased by 59.4% in the CP group (MEP_simple task=1.89 [1.00-3.09] vs. MEP_complex task=2.70 [1.59-4.80] mV/s^2; $p \leq 0.01$) and 113.8% in the control group (MEP_simple task=1.95 [0.99-2.72] vs. MEP_complex task =2.91 [1.97-3.66] mV/s^2; $p \leq 0.01$). An increase in the number of synergies was observed during complex task in CP group (p=0.018).

Conclusions: These results suggest that performing a complex walking task allows to enhance the corticospinal excitability in both individuals with CP and control subjects. Moreover, CP individuals showed that either the number or the structure of synergies are modulated by the complex task, in comparison to control subjects. Longitudinal studies are recommended to assess the impact of the integration of complex tasks in gait rehabilitation interventions.

Keywords: Cerebral palsy, Locomotion, Gait, Motor control, EMG, Motor modules

1. Introduction:

Cerebral palsy (CP) is the most common motor disability in childhood, often resulting from a defect or lesion in the developing brain(1). While the etiology of CP varies, damage to the corticospinal tract (CST), which connects the cerebral cortex to the spinal cord and plays a key role in voluntary motor control, is a common feature of CP (2,3). This damage affects the signals that are sent from the motor cortex to the muscles, leading to a range of motor deficits. Studies based on non-invasive brain stimulation techniques, such as transcranial magnetic stimulation (TMS), and on surface electromyography (sEMG) showed that individuals with CP often exhibit reduced corticospinal excitability (4,5). This means that the descending drive from the motor cortex to the muscles is weaker, leading to difficulties in initiating and controlling voluntary movements. This may be reflected in a reduction in performance in daily living tasks, including walking (6–8).

Muscle synergy theory has been recently proposed to quantitatively and non-invasively assess the motor control system both in physiological and pathological conditions (9,10). This theory suggests that instead of independently controlling each muscle, the central nervous system may coordinate the activation levels and timing of all muscles involved in a specific movement using a small number of neural commands. Specifically, the muscle synergy theory can be derived from sEMG data using specific factorization methods, enabling the study of underlying motor control patterns and strategies that are believed to reflect neural function (10). Even if the neural origin of muscle synergies is still a topic of discussion among the scientific community (11), several studies have demonstrated the usefulness of the muscle synergy theory in describing motor control system of patients with orthopedical and neurological diseases affecting movements (12–15).

From this perspective, muscle synergies may represent a valuable tool for studying motor control alterations in CP patients who are expected to exhibit altered corticospinal excitability. In fact, altered corticospinal excitability is commonly associated with motor impairments, such as spasticity and weakness (16,17), and changes in this variable are believed to reflect neuroplasticity (18). Therefore, corticospinal excitability alterations may disrupt both the composition and the number of muscle synergies (19,20), leading to less coordinated and more stereotyped movements. In some cases, individuals may also rely on pathological movement

patterns to compensate for the motor deficits. Human locomotion is a widely studied movement for assessing motor control. It is one of the most important activities of daily living and can be easily performed by patients who can walk independently without aid or external support. In daily-life activities, humans often must adapt their gait to avoid obstacles or speed changes. Unpredictable disturbances, such as any deviation from regular walking, can lead to falls and injuries (21,22). Much of the existing research has focused on the automatic control of walking by studying repetitive movements in highly controlled conditions (e.g., walking on a straight-line path or treadmill). These "simple" walking tasks predominantly engage automatic neural circuits[3], like the spinal structures, and depend less on voluntary control circuits, such as the CST (23), which are crucial for daily-life adaptations. Measuring corticospinal excitability and sEMG data during walking conditions mainly involving voluntary control circuits (i.e., the corticospinal tract) may provide insights into how the CNS controls lower-limb muscles in a more ecological environment. This is crucial for understanding the neural mechanisms of gait, which is a complex and fundamental task of daily living. In addition, although it is commonly accepted that muscle synergies are controlled at the spinal cord level(24), cortical contributions may be essential for ensuring a flexible and goal-directed motor control. In control subjects, evidence suggests that fundamental muscle synergy structures (synergy weights or vectors) remain consistent despite external environmental changes during walking (25,26), such as varying gait speeds (27,28), different slopes (29), or distinct walking tasks (e.g., nordic or beam walking) (30,31). However, in children with CP, Kim et al. (32) showed that muscle synergies are more sensitive to changes during walking tasks (e.g., self-selected speed, variable speed, and restricted width) than in typically developing children. Investigating both corticospinal excitability and muscle synergies during regular and complex walking tasks may improve our knowledge about locomotor control in individuals with CP and refine rehabilitation intervention.

The objectives of this study were to: (1) assess whether a complex walking task promotes an increase in corticospinal excitability and a modulation of muscle synergies compared to simple walking task in individuals with CP and controls and (2) to evaluate whether the change is different between the two groups. We hypothesized that an increase in the corticospinal excitability (i.e., complex task) might further involve the corticospinal tract and, thus, determine a reorganization of the locomotor muscle synergies (in terms of number and robustness of the muscle synergies). Moreover, it was expected that control subjects would show a greater increase

109 in corticospinal excitability compared to individuals with CP. In terms of muscular synergy, we
110 expected a modulation of the number and structures.

2. Materials and Methods

2.1. Participants

CP and control groups were recruited for this study. For both groups, the inclusion criteria were:
(1) to be aged between 12 and 45 years; (2) to be able to walk without walking aid for
approximately 3 minutes. To be eligible to perform the corticospinal excitability assessment part
of the study, participants must not have any exclusion criteria for TMS as detailed by Rossi et al.
(33). Inclusion criteria specific to the CP group required participants to have a confirmed
diagnosis of unilateral or bilateral cerebral palsy and an ability to understand simple instructions
(e.g., walks normally, walks on targets, etc). For both groups, the exclusion criteria were any
unstable health condition that could affect task performance. Participants with contraindications
to TMS performed part of the protocol, which involved only sEMG measurements without TMS.
Prior to taking part in the study, all participants provided their informed consent for the
experimental procedure in accordance with the Declaration of Helsinki. The study was approved
by the Quebec City Rehabilitation Institute Ethics Review Board (2021-2209, RIS).

2.2. Experimental protocol

Individuals with CP eligible for TMS were assessed during two evaluation sessions. The first
session involved Magnetic Resonance Imaging (MRI) to obtain the patient-specific brain
anatomical structure, which was then used to guide the experimenter in positioning the TMS coil.
During the second session, CP patients underwent neurophysiological measurements (i.e., motor
evoked potential (MEP) and sEMG) while performing two consecutive treadmill walking tasks at
a comfortable self-selected speed: *a*) a Simple Walking (SW) task and *b*) a Complex Walking
(CW) task. As developed in our previous study (34), the CW task involves walking on a treadmill
and placing the feet on virtual targets presented at different intervals (80%, 100%, 120% of the
normal step length) on a screen facing the subject. The real-time positions of the feet were also
displayed on the screen as feedback, with each foot represented by a small sphere centered at the
heel position. The distance between visual targets in the sagittal plane randomly varied with each
step. During the SW task, instead, no visual targets were displayed on the screen facing the
participants. Control participants attended only the second evaluation session (i.e.,

neurophysiological measurements), a template brain being used for the positioning of the TMS
coil.

2.3. Data acquisitions

Corticospinal excitability. MEPs from Tibialis Anterior (TA) muscle were assessed using
single-pulse TMS (Magstim Rapid2; custom "batwing" coil (34,35)) applied to the motor cortex
(contralateral to the most-affected lower limb in CP subjects and the non-dominant lower limb in
control subjects). The hotspot and active motor threshold were determined while the participant
was in a seated position, with knees flexed at 90 degrees and maintaining approximately 20
degrees of dorsiflexion against gravity to facilitate MEPs. The hotspot was identified as the zone
that evokes MEPs at the lowest stimulation intensity (36). The active motor threshold (aMT) was
determined at the hotspot as the lowest stimulation intensity required to evoke a MEP in 5/10
trials (37). Then, single-pulse stimulations at 110% of the aMT were administered during
walking, with 25 stimulations per condition. These stimulations occurred around 40% of the
walking cycle, specifically in the late stance phase when the activity of the TA muscle is
minimal, to ensure consistency in comparisons across different conditions (34). The timing
corresponding to 40% of the gait cycle was determined in real time using the treadmill force
plates. A neuronavigation system (Brainsight, Rogue research, Montreal, Canada) guided the
experimenter while positioning the coil above the experimentally determined hotspot. The
position of the coil at the moment of application of each pulse was recorded to allow offline
rejection of off-target stimuli and comparison of coil positioning across conditions. Additionally,
the position of each foot was tracked using 3 photo-reflective markers and visualized in real-time
using an optoelectronic system equipped with 8 infrared cameras (Vicon Motion Systems). To
avoid patrticipants' anticipation, stimulations were provided on random gait cycles. On average, a
single pulse stimulation was provided every 5 strides with a range of 3-7 strides. Participants not
eligible for TMS took part only in the protocol assessing muscle synergies, described in the
following section.

Muscle activity. sEMG signals of the most affected (CP group) or non-dominant (control group)
lower limb were recorded by focusing on the muscles involved in the walking tasks. In particular,
the following 6 lower-limb muscles were monitored: TA, Soleus (SOL), Gastrocnemius lateralis

(GL), Rectus femoris (RF), Vastus lateralis (VL), and Semitendinosus (ST). The ground
electrode was positioned on the patella. Ag/AgCl electrodes (Kendall Medi-trace 200, Covidien)
were placed on the volunteers' skin over the belly of each muscle. Prior to electrode placement,
the skin was shaved and cleaned to minimize the impedance between the skin and the electrode.
Each electrode location was determined according to the recommendations of Surface EMG for
Non-Invasive Assessment of Muscles (SENIAM) (38). The ground electrode was positioned on
the patella of the homolateral lower limb. All signals were sampled at 1000 Hz.
178
179 **2.4. Data analysis**
180 **2.4.1. MEP analysis**

181 To quantify corticospinal excitability, the area under the curve was calculated for each rectified
182 MEP and then averaged for each task and participant(34). In each group, the CST excitability
183 change between the two tested conditions (i.e., MEP modulation) was measured as follows:

$$\left(\frac{(\text{MEPs complex task } - \text{ MEPs simple task})}{\text{MEPs simple task}} \right) x \ 100$$

The error in TMS coil location was measured through the neuronavigation software as the
distance between the location of each stimulation and the hotspot (expressed in millimeters).
Finally, the mean background sEMG amplitude was measured (considering 50-ms windows
preceding every TMS stimulation) and compared across task conditions. All data were analyzed
offline by using custom MATLAB routines (MATLAB r2018a, The MathWorks Inc., Natick,
MA, USA).

2.4.2. Muscle synergy extraction

Raw sEMG signals were band-pass filtered between 10 Hz and 450 Hz by means of a zero-lag
4th-order Butterworth filter(39). Thereafter, filtered sEMG signals were full-wave rectified and
the sEMG envelopes were obtained using a 4-th order low-pass Butterworth filter with a cut-off
frequency set at 10 Hz (40). Then, separately for each muscle, the resulting envelopes were
amplitude-normalized by the maximum of each sEMG envelope obtained during the SW task. A
5% body weight vertical cut-off force threshold was applied to the ground reaction forces to
identify gait events. Gait cycles duration was normalized to 1000 samples to avoid biases due to
different gait cycle durations.

The Non-Negative Matrix Factorization (NNMF) was used to extract locomotor muscle

synergies during both walking conditions(41–43). The NNMF algorithm involves an iterative

factorization of the sEMG envelope matrix E, which has dimensions $t \times m$. It decomposes E

into two matrices: the activation coefficient matrix C with dimensions $t \times s$ and the weight

vector W with dimensions $s \times m$. This can be expressed as follows:

$$E = W \times C + residuals$$

The iterations continue until the Frobenius norm of the residuals is minimized. In this context, t

represents the number of time instants, m represents the number of muscles, and s represents the

final number of muscle synergies. For further details, please refer to Hug et al.(44).

In the present study, matrix E contains the time- and amplitude-normalized sEMG envelopes It

has dimensions $(N \times t) \times m$, with N denoting the number of gait cycles, t representing the

number of time instants ($t = 1000$), and m representing the number of acquired muscles ($m = 6$).

Different NNMF solutions were tested by running several times the algorithm on the same sEMG

data changing the muscle synergy number (s) between 1 and 6 (i.e., the total number of muscles

acquired). For each number of muscle synergies, the reconstruction quality was assessed using

the Variance Accounted For (VAF) parameter. Consequently, the final number of muscle

synergies required to properly reconstruct the original sEMG signals was defined by choosing the

smallest number of synergies granting a VAF value equal or greater than 90% (45). Muscle

synergies were extracted through custom MATLAB routines (MATLAB r2018a, The

MathWorks Inc., Natick, MA, USA).

2.5. Statistical analysis

For the groups' characteristics, parameter estimates are represented as median and

interquartile. In all the analyses, the significance level (α) was set to 0.05. Mann-Whitney U-tests

were used to test differences in anthropometric characteristics between groups (i.e., CP and

control).

To test our first objective related to corticospinal excitability, the nparLD test (46), a non-

parametric test equivalent to a repeated-measures ANOVA, was applied to MEP results (i.e.,

MEP under the curve area, EMG background and coil error), with the within-subject factor

Condition (i.e., SW and CW tasks) and the between-subject factor *Group* (i.e., CP and control).

In the presence of a significant interaction, *post-hoc* analyses were carried out. NparLD analysis

was carried out using RStudio software. Between groups' MEP modulations were compared

using a Mann-Whitney U test.

To test our second objective related to muscle synergies, a Wilcoxon test was used to compare

the final number of muscle synergies between SW and CW tasks in CP subjects. Note that there

is no group factor in this analysis as the control group served as a reference. The number of

synergies was determined for each condition in the control group. Specifically, the N synergies

identified in the control group during the SW were used as a reference for comparison with the

synergies of the CP group performing the same task. Similarly, the N synergies identified in the

control group during the CW served as a reference for comparison with the synergies of the CP

group in that task. To assess differences in motor control strategies between individuals with CP

and control participants, the average activation coefficients ($\overline{C}$) and weight vectors ($\overline{W}$) extracted

during the walking tasks of control individuals were used as a reference. Then, the Cosine

Similarity (S) was used to assess the similarity between the muscle synergies of each CP

individual and the reference (i.e., muscle synergies averaged over control group) considering both

C and W vectors. The S value may range between 0 (totally dissimilar) and 1 (totally

similar)(47). Muscle synergies characterized by a CS value below 0.8 were classified as *subject-*

specific muscle synergies, whereas muscle synergies characterized by a S value equal to or

higher than 0.8 were classified as *shared* (or *common*) muscle synergies. Muscle synergies were

compared using custom MATLAB scripts.

3. Results

3.1. Participants

Fourteen individuals with CP and fourteen control subjects were included in the present study. Groups' characteristics are reported in **Table 1**. Individuals with CP had Gross Motor Function Classification System (48) (GMFCS) scores of I (n=3), II (n=8) and III (n=3). Three participants had contraindication to TMS due to a history of epilepsy. Consequently, corticospinal excitability measurements were conducted on 11 CP patients and 14 healthy controls. The characteristics of the CP individuals and controls included in the study are detailed in **Table 1**.

Table 1. Characteristics of CP and control groups.

	Sex	Age (years)	Mass (kg)	Heigh (cm)	aMT (%)
CP (*n* =14)	5♀	27 [16-35]	63 [56-80]	166 [162-175]	65 [64-70]*
CTL (*n* = 14)	8♀	25 [23-28]	64 [58-79]	170 [165-178]	59 [55-64]
p-value	0.276	0.322	0.662	0.433	***0.005***

Parameter estimates are represented as mean ± standard deviation over the group. P-values represent the results of the comparison between groups using Mann-Whitney U-tests. CP: Cerebral Palsy participants; CTL: Control participants; aMT: active motor threshold; *only 11/14 CP participants had TMS assessments. Significant statistical differences are highlighted in ***italic bold***.

3.2. Corticospinal excitability

Figure 1 shows results related to corticospinal excitability (i.e., MEP under curve area and MEP modulation), presented as median and interquartile range across each group. The NparLD results showed a significant effect of the condition (SW *vs.* CW; $p < 0.0001$), but not of the group (CP *vs* Control; p=0.905) nor a condition x group interaction (p=0.332). Based on the Mann-Whitney U test, MEP modulation did not differ significantly between groups (p=0.396, Figure 2B). An effect of the condition was observed in the sEMG background (p=0.015), without any effect of the group or condition x group interaction (p=0.691 and 0.626, respectively). Regarding the coil placement during walking tasks, the target errors of the included stimulation were less than 5 mm and no significant effects of group, condition, or group x condition interaction were observed (p>0.05).

280
281 **Figure 1.** (A) Corticospinal excitability during Simple Walking (SW, blue) and Complex
282 Walking (CW, purple) tasks in cerebral palsy (CP, ■) and Control (CTL, ■) groups. NparLD p-
283 value for the condition effect (SW vs. CW) is displayed. No significant effect for the group factor
284 (CP vs Control) nor at the interaction level (condition x group) are observed. (B) Modulation (%)
285 between the two tasks in terms of motor evoked potential (MEP) and EMG background.
286

3.3. Muscle synergy analysis

On average, four muscle synergies were required to properly reconstruct the sEMG envelopes of

control participants during both walking conditions (i.e., SW and CW), with an average VAF

value of $93 \pm 2\%$ during both SW and CW tasks, respectively. For individuals with CP, an

increase in the number of muscle synergies ($p < 0.05$) was observed during the CW task (n=4 [4-

4]) compared to the SW task (n=3 [3-4]). An average VAF of $93\% \pm 1\%$ and $92 \pm 2\%$ were

obtained during SW and CW tasks, respectively.

Reference muscle synergies, obtained by averaging the muscle synergies of the control

participants, are shown in **Figure 2**. **Figure 3** and **Figure 4** show the weight vectors (W) and the

activation coefficients (C) of each CP patient during both walking tasks with the indication of the

shared and specific muscle synergies.

Figure 2. Reference muscle synergies extracted from control participants during the SW (blue) and the CW (purple) task. Muscle abbreviations: TA Tibialis Anterior, SOL Soleus, GL Lateral Gastrocnemius, RF Rectus Femoris, VL Vastus Lateralis, and ST Semitendinosus.

Results showed the presence of *subject-specific* synergies (i.e., considered to be dissimilar to those of CTL participants) in 11 out of 14 CP participants. Only the three CP participants showing a GMFCS score of I (CP09, CP13, CP14) presented 4 *shared* muscle synergies (i.e., similar to those of CTL participants, with CS scores ranging from 0.87 to 0.99).

Figure 3. Synergies' weight vectors of CP participants during simple (SW) and complex (CW) tasks. Shared weight vectors are shown in blue and purple colors, while patient-specific weight vectors are shown in pink. For each weight vector, the cosine similarity (CS) value between the specific vector and the corresponding reference synergy is represented. Muscle abbreviations: TA Tibialis Anterior, SOL Soleus, GL Lateral Gastrocnemius, RF Rectus Femoris, VL Vastus Lateralis, and ST Semitendinosus.

318
Figure 4. Synergies' activation coefficients of CP participants during simple (SW) and complex
(CW) tasks. Shared activation coefficients are shown in blue and purple colors, while patient-
specific activation coefficients are shown in pink. For each activation coefficient, the cosine
similarity (CS) value between the specific vector and the corresponding reference synergy is

represented. Muscle abbreviations: TA Tibialis Anterior, SOL Soleus, GL Lateral
Gastrocnemius, RF Rectus Femoris, VL Vastus Lateralis, and ST Semitendinosus.
325

4. Discussion

The objective of this study was to assess the effect of a CW task on CST excitability and muscle synergies in individuals with and without CP. The two main findings are that: 1) compared to the SW task, the CW task induced a significant increase in corticospinal excitability and a modulation of muscle synergy structure in individuals with CP; and 2) both groups showed increased CST excitability during the CW task, no significant differences between groups were observed for this metric. This study is the first to investigate corticospinal excitability during walking in individuals with CP, providing essential information on the neuromuscular adaptations in this population.

Corticospinal excitability

Partially consistent with our first hypothesis, both groups increased their CST excitability under the CW task. However, no differences in terms of MEP increase between groups were observed. Although the relationship between movement complexity and MEP size has not yet been established in the literature, our results align with several previous studies highlighting that complex motor tasks increase CST excitability in healthy subjects during manual (23,49,50) or locomotor tasks (34,51). For example, Barthélémy *et al.* (51) observed an increase in corticospinal excitability when participants walked against resistive forces. These results support the idea that CST excitability is modulated in a task-dependent manner, and may suggest that complex walking tasks can serve as powerful stimuli for motor system engagement. Moreover, it is acknowledged that MEP amplitude can be sensitive to cognitive processes such as precision demands(52) or decision-making (53). Thus, it is reasonable to speculate that heightened cognitive processing during the preparation of our visually guided task (*i.e.,* CW) may lead to an increase in CST excitability. Even if differences in MEPs amplitudes and shape were demonstrated between stroke survivors and healthy controls (54), our results revealed no differences in MEP area under the curve and increase between individuals with CP and controls. This is likely because the CP individuals included in this study were only mildly affected, as the majority have a GMFCS level of I or II. Then, the lack of group difference in MEP modulation may suggest that individuals with mild CP retain the capacity to modulate CST excitability in response to increased task demands, comparable to controls. However, the CP group showed a higher aMT than the control participants. This suggests that the higher stimulation intensity used

relative to aMT in the CP group may have compensated for reduced corticospinal excitability,
and then masking potential differences between the groups. Thus, similar MEP responses do not
necessarily reflect equivalent CST reactivity (55), but may rather indicate reduced recruitment
efficiency in CP, requiring greater stimulation to evoke comparable outputs. In conclusion, these
preliminary results support that MEP may serve as valuable as a dynamic marker of CST function
during functional tasks such as walking. It might be preferable to measure MEPs when people are
moving rather than when they are at rest as this provides a better picture of their ability to
function and adapt to their motor impairments.

Muscle synergy modulation

Overall, individuals with CP recruited fewer muscle synergies than healthy controls during the
SW task, indicating a general reduction in the complexity of their motor control system, which is
often linked to poorer motor performance (9). In motor control systems with fewer muscle
synergies, the central nervous system's ability to independently control synergies responsible for
different biomechanical functions is significantly compromised (14,45,56). The reduction in
motor control system complexity can be observed through increased muscle coactivation, where
weight vectors show the activation of a larger number of muscles, and decreased independence of
neural control signals, indicated by activation coefficients that are more similar to each other and
more overlapping throughout the task duration.

When moving from the SW task to the CW task, results demonstrated a reorganization in CP
muscle synergies, in general revealing an increased number of muscle synergies, lower cosine
similarity values compared to the reference muscle synergies, and an increased number of
subject-specific muscle synergies. In other words, individuals with CP require a reorganization of
their motor control strategies (in terms of both activation timing and muscle enrollment
throughout the task duration) to cope with the new, more complex walking tasks. It aligns with
studies suggesting that motor modules in individuals with neurological impairments can be
modulated in response to environmental or biomechanical demands (57,58). CP group showed
that either the spatial (i.e., weight vectors) or temporal (i.e., activation coefficients) components
of the muscle synergies differed when compared to reference synergies of controls, who
maintained highly consistent muscle synergies across walking conditions.

Clinical implications

The findings of the current study highlight the potential for enhancing corticospinal excitability during a CW task. The results of this study showed a significant increase in CST excitability in both CP and control participants. These results suggest that individuals can increase their CST excitability to facilitate precise movement control in CW scenarios that demand accurate foot placement. Regarding muscle synergies, our results are in line with the literature highlighting that individuals with CP often employ simpler motor control strategies during gait compared to their typically developing peers (9). However, to obtain effective motor control during more complex locomotor tasks, individuals with CP required modulation of their muscle synergies by increasing motor control complexity (i.e., increasing the number of muscle synergies). Overall, the parallel increase in CST excitability and modulation of muscle synergies during the CW task in individual with CP suggest that task complexity may act as a powerful driver of neural and muscular adaptations. These results underline the value of implementing cognitively and sensorimotor challenging walking tasks in rehabilitation.

Limitations

Some limitations of this study need to be highlighted. First, the sample size is relatively small sample size, which increases the risk of type 2 error and may impact the generalizability of the findings. Second, it is important to consider that the level of attention during the two walking tasks was not controlled, potentially affecting CST excitability. Third, the study did not include an analysis of the biomechanical changes and gait spatiotemporal parameters associated with the reduced number of muscle synergies in individuals with CP. Future studies should consider incorporating these elements to provide a more comprehensive understanding of the gait dynamics and functional implications in CP population.

5. Conclusion

This study has contributed to elucidating the mechanisms of gait control in CP. Additionally, our results clarified the increased involvement of the CST and the reorganization of motor control strategies in CP patients during more challenging walking tasks. These results pave the way for improved gait rehabilitation in CP by targeting complex tasks. Although our study measured CST excitability only in the TA muscle, future research should also investigate the SOL and RF muscles, as they play significant roles in walking.

REFERENCES

1. Bax M, Goldstein M, Rosenbaum P, Leviton A, Paneth N, Dan B, et al. Proposed definition and classification of cerebral palsy, April 2005. Dev Med Child Neurol. août 2005;null(08):571‑6.

2. Kuo HC, Ferre CL, Carmel JB, Gowatsky JL, Stanford AD, Rowny SB, et al. Using diffusion tensor imaging to identify corticospinal tract projection patterns in children with unilateral spastic cerebral palsy. Dev Med Child Neurol. 2017;59(1):65‑71.

3. Cassidy JM, Carey JR, Lu C, Krach LE, Feyma T, Durfee WK, et al. Ipsilesional motor-evoked potential absence in pediatric hemiparesis impacts tracking accuracy of the less affected hand. Res Dev Disabil. déc 2015;47:154‑64.

4. Brouwer B, Ashby P. Altered corticospinal projections to lower limb motoneurons in subjects with cerebral palsy. Brain J Neurol. juin 1991;114 (Pt 3):1395‑407.

5. Riquelme I, Cifre I, Muñoz MA, Montoya P. Altered corticomuscular coherence elicited by paced isotonic contractions in individuals with cerebral palsy: a case-control study. J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol. déc 2014;24(6):928‑33.

6. Duncan PW. Stroke disability. Phys Ther. mai 1994;74(5):399‑407.

7. Palmer JA, Needle AR, Pohlig RT, Binder-Macleod SA. Atypical cortical drive during activation of the paretic and nonparetic tibialis anterior is related to gait deficits in chronic stroke. Clin Neurophysiol Off J Int Fed Clin Neurophysiol. janv 2016;127(1):716‑23.

8. Cherni Y, Bouyer L, Bretheau F, Mercier C. Effect of a complex walking task on corticospinal excitability and muscle activity in individuals with cerebral palsy: A multiple-case study. Gait Posture. 1 sept 2022;97:S324‑5.

9. Bekius A, Bach MM, van der Krogt MM, de Vries R, Buizer AI, Dominici N. Muscle Synergies During Walking in Children With Cerebral Palsy: A Systematic Review. Front Physiol. 2 juill 2020;11:632.

10. Bizzi E, Cheung VCK. The neural origin of muscle synergies. Front Comput Neurosci [Internet]. 29 avr 2013 [cité 25 févr 2020];7. Disponible sur: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3638124/

11. Cheung VCK, Seki K. Approaches to revealing the neural basis of muscle synergies: a review and a critique. J Neurophysiol. 1 mai 2021;125(5):1580‑97.

12. Steele KM, Rozumalski A, Schwartz MH. Muscle synergies and complexity of neuromuscular control during gait in cerebral palsy. Dev Med Child Neurol. déc 2015;57(12):1176‑82.

13. Cherni Y, Hajizadeh M, Begon M, Turpin NA. Effects of Body Weight Support and Guidance Force Settings on Muscle Synergy during Lokomat Walking. En préparation. 2020;

14. Ghislieri M, Lanotte M, Knaflitz M, Rizzi L, Agostini V. Muscle synergies in Parkinson's disease before and after the deep brain stimulation of the bilateral subthalamic nucleus. Sci Rep. 28 avr 2023;13(1):6997.

15. Roh J, Rymer WZ, Perreault EJ, Yoo SB, Beer RF. Alterations in upper limb muscle synergy structure in
chronic stroke survivors. J Neurophysiol. 1 févr 2013;109(3):768‑81.

16. Condliffe EG, Jeffery DT, Emery DJ, Treit S, Beaulieu C, Gorassini MA. Full Activation Profiles and
Integrity of Corticospinal Pathways in Adults With Bilateral Spastic Cerebral Palsy: Neurorehabil
Neural Repair [Internet]. 29 déc 2018 [cité 11 nov 2020]; Disponible sur:
https://journals.sagepub.com/doi/10.1177/1545968318818898

17. Hussain AW, Onambele GL, Williams AG, Morse CI. Muscle size, activation, and coactivation in adults
with cerebral palsy: Properties of Spastic Hemiplegia. Muscle Nerve. janv 2014;49(1):76‑83.

18. Dai W, Nakagawa K, Nakajima T, Kanosue K. Determinants of Neural Plastic Changes Induced by
Motor Practice. Front Hum Neurosci [Internet]. 2021 [cité 5 janv 2024];15. Disponible sur:
https://www.frontiersin.org/articles/10.3389/fnhum.2021.613867

19. Duclay J, Pasquet B, Martin A, Duchateau J. Specific modulation of corticospinal and spinal
excitabilities during maximal voluntary isometric, shortening and lengthening contractions in
synergist muscles. J Physiol. 1 juin 2011;589(Pt 11):2901‑16.

20. Carson RG, Riek S, Mackey DC, Meichenbaum DP, Willms K, Forner M, et al. Excitability changes in
human forearm corticospinal projections and spinal reflex pathways during rhythmic voluntary
movement of the opposite limb. J Physiol. 1 nov 2004;560(Pt 3):929‑40.

21. Ataullah AHM, De Jesus O. Gait Disturbances. In: StatPearls [Internet]. Treasure Island (FL):
StatPearls Publishing; 2023 [cité 23 oct 2023]. Disponible sur:
http://www.ncbi.nlm.nih.gov/books/NBK560610/

22. Shumway-Cook A, Patla AE, Stewart A, Ferrucci L, Ciol MA, Guralnik JM. Environmental demands
associated with community mobility in older adults with and without mobility disabilities. Phys Ther.
juill 2002;82(7):670‑81.

23. Kennefick M, Burma JS, van Donkelaar P, McNeil CJ. Corticospinal excitability is enhanced while
preparing for complex movements. Exp Brain Res. mars 2019;237(3):829‑37.

24. Desrochers E, Harnie J, Doelman A, Hurteau MF, Frigon A. Spinal control of muscle synergies for
adult mammalian locomotion. J Physiol. janv 2019;597(1):333‑50.

25. Dussault-Picard C, Havashinezhadian S, Turpin NA, Moissenet F, Turcot K, Cherni Y. Age-related
modifications of muscle synergies during daily-living tasks: A scoping review. Clin Biomech. 1 mars
2024;113:106207.

26. Gonzalez-Vargas J, Sartori M, Dosen S, Torricelli D, Pons JL, Farina D. A predictive model of muscle
excitations based on muscle modularity for a large repertoire of human locomotion conditions. Front
Comput Neurosci. 2015;9:114.

27. Buurke TJW, Lamoth CJC, van der Woude LHV, den Otter AR. Synergistic Structure in the Speed
Dependent Modulation of Muscle Activity in Human Walking. PloS One. 2016;11(4):e0152784.

28. Kibushi B, Hagio S, Moritani T, Kouzaki M. Speed-Dependent Modulation of Muscle Activity Based on
Muscle Synergies during Treadmill Walking. Front Hum Neurosci. 2018;12:4.

29. Saito A, Tomita A, Ando R, Watanabe K, Akima H. Similarity of muscle synergies extracted from the
lower limb including the deep muscles between level and uphill treadmill walking. Gait Posture. janv
2018;59:134‑9.

30. Sawers A, Allen JL, Ting LH. Long-term training modifies the modular structure and organization of
walking balance control. J Neurophysiol. déc 2015;114(6):3359‑73.

31. Boccia G, Zoppirolli C, Bortolan L, Schena F, Pellegrini B. Shared and task-specific muscle synergies of
Nordic walking and conventional walking. Scand J Med Sci Sports. mars 2018;28(3):905‑18.

32. Kim Y, Bulea TC, Damiano DL. External walking environment differentially affects muscle synergies in
children with cerebral palsy and typical development. Front Hum Neurosci [Internet]. 23 sept 2022
[cité 16 août 2024];16. Disponible sur: https://www.frontiersin.org/journals/human-
neuroscience/articles/10.3389/fnhum.2022.976100/full

33. Rossi S, Antal A, Bestmann S, Bikson M, Brewer C, Brockmöller J, et al. Safety and recommendations
for TMS use in healthy subjects and patient populations, with updates on training, ethical and
regulatory issues: Expert Guidelines. Clin Neurophysiol Off J Int Fed Clin Neurophysiol. janv
2021;132(1):269‑306.

34. Dambreville C, Neige C, Mercier C, Blanchette AK, Bouyer LJ. Corticospinal Excitability Quantification
During a Visually-Guided Precision Walking Task in Humans: Potential for Neurorehabilitation.
Neurorehabil Neural Repair. nov 2022;36(10‑11):689‑700.

35. Cai W, George JS, Chambers CD, Stokes MG, Verbruggen F, Aron AR. Stimulating deep cortical
structures with the batwing coil: How to determine the intensity for transcranial magnetic
stimulation using coil–cortex distance. J Neurosci Methods. 15 mars 2012;204(2):238‑41.

36. Wassermann EM. Variation in the response to transcranial magnetic brain stimulation in the general
population. Clin Neurophysiol Off J Int Fed Clin Neurophysiol. juill 2002;113(7):1165‑71.

37. Cavaleri R, Schabrun SM, Chipchase LS. The number of stimuli required to reliably assess
corticomotor excitability and primary motor cortical representations using transcranial magnetic
stimulation (TMS): a systematic review and meta-analysis. Syst Rev. 6 mars 2017;6(1):48.

38. Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. Development of recommendations for SEMG
sensors and sensor placement procedures. J Electromyogr Kinesiol Off J Int Soc Electrophysiol
Kinesiol. oct 2000;10(5):361‑74.

39. De Luca CJ. The Use of Surface Electromyography in Biomechanics. J Appl Biomech. mai
1997;13(2):135‑63.

40. Ranaldi S, Gizzi L, Severini G, De Marchis C. Optimal Identification of Muscle Synergies From Typical
Sit-to-Stand Clinical Tests. IEEE Open J Eng Med Biol. 2023;4:31‑7.

41. Lee DD, Seung HS. Learning the parts of objects by non-negative matrix factorization. Nature. oct
1999;401(6755):788‑91.

42. Turpin NA, Uriac S, Dalleau G. How to improve the muscle synergy analysis methodology? Eur J Appl
Physiol. 1 avr 2021;121(4):1009‑25.

43. Rabbi MF, Pizzolato C, Lloyd DG, Carty CP, Devaprakash D, Diamond LE. Non-negative matrix
factorisation is the most appropriate method for extraction of muscle synergies in walking and
running. Sci Rep. 19 mai 2020;10(1):8266.

44. Hug F, Turpin NA, Couturier A, Dorel S. Consistency of muscle synergies during pedaling across
different mechanical constraints. J Neurophysiol. juill 2011;106(1):91‑103.

45. Clark DJ, Ting LH, Zajac FE, Neptune RR, Kautz SA. Merging of Healthy Motor Modules Predicts
Reduced Locomotor Performance and Muscle Coordination Complexity Post-Stroke. J Neurophysiol.
févr 2010;103(2):844‑57.

46. Noguchi K, Gel YR, Brunner E, Konietschke F. nparLD: An R Software Package for the Nonparametric
Analysis of Longitudinal Data in Factorial Experiments. J Stat Softw. 18 sept 2012;50:1‑23.

47. Oliveira ASC, Gizzi L, Kersting UG, Farina D. Modular organization of balance control following
perturbations during walking. J Neurophysiol. oct 2012;108(7):1895‑906.

48. Wood E, Rosenbaum P. The Gross Motor Function Classification System for Cerebral Palsy: a study of
reliability and stability over time. Dev Med Child Neurol. mai 2000;42(5):292‑6.

49. Abbruzzese G, Trompetto C, Schieppati M. The excitability of the human motor cortex increases
during execution and mental imagination of sequential but not repetitive finger movements. Exp
Brain Res. oct 1996;111(3):465‑72.

50. Roosink M, Zijdewind I. Corticospinal excitability during observation and imagery of simple and
complex hand tasks: implications for motor rehabilitation. Behav Brain Res. 12 nov
2010;213(1):35‑41.

51. Barthélemy D, Alain S, Grey MJ, Nielsen JB, Bouyer LJ. Rapid changes in corticospinal excitability
during force field adaptation of human walking. Exp Brain Res. mars 2012;217(1):99‑115.

52. Bonnard M, Galléa C, De Graaf JB, Pailhous J. Corticospinal control of the thumb-index grip depends
on precision of force control: a transcranial magnetic stimulation and functional magnetic resonance
imagery study in humans. Eur J Neurosci. févr 2007;25(3):872‑80.

53. Hadar AA, Rowe P, Di Costa S, Jones A, Yarrow K. Motor-evoked potentials reveal a motor-cortical
readout of evidence accumulation for sensorimotor decisions. Psychophysiology.
2016;53(11):1721‑31.

54. Kang YJ, Ku J, Kim H, Park H. Facilitation of Corticospinal Excitability According to Motor Imagery and
Mirror Therapy in Healthy Subjects and Stroke Patients. Ann Rehabil Med. 1 déc 2011;35:747‑58.

55. Oudega M, Perez MA. Corticospinal reorganization after spinal cord injury. J Physiol. 15 août
2012;590(Pt 16):3647‑63.

56. Ambrosini E, De Marchis C, Pedrocchi A, Ferrigno G, Monticone M, Schmid M, et al. Neuro-
Mechanics of Recumbent Leg Cycling in Post-Acute Stroke Patients. Ann Biomed Eng. nov
2016;44(11):3238‑51.

57. McGowan CP, Neptune RR, Clark DJ, Kautz SA. Modular control of human walking: Adaptations to
altered mechanical demands. J Biomech. 10 févr 2010;43(3):412.

58. Janshen L, Santuz A, Arampatzis A. Muscle Synergies in Patients With Multiple Sclerosis Reveal
Demand-Specific Alterations in the Modular Organization of Locomotion. Front Hum Neurosci
[Internet]. 27 janv 2021 [cité 3 juill 2025];14. Disponible sur:
https://www.frontiersin.org/journals/human-
neuroscience/articles/10.3389/fnhum.2020.593365/full