A mega-analysis of low frequency resting-state measures in mood and psychosis-

spectrum disorders

Maya L. Foster[1], Milana Khaitova[2], Saloni Mehta[2], Jean Ye[3], Dustin Scheinost[1-6]

1) Department of Biomedical Engineering, Yale University

2) Department of Radiology & Biomedical Imaging, Yale School of Medicine

3) Interdepartmental Neuroscience Program, Yale School of Medicine

4) Department of Statistics & Data Science, Yale University

5) Yale Child Study Center, Yale School of Medicine

6) Yale Biomedical Imaging Institute, Yale School of Medicine

Highest Academic Degrees Obtained by Co-Authors:

● Maya L. Foster – M.S./M.Phil.
● Milana Khaitova – B.A.
● Saloni Mehta – M.B.B.S.
● Jean Ye – M.S.
● Dustin Scheinost – Ph.D.
18

19   All correspondence and offprint requests should be sent or mailed to Maya L. Foster

20   (maya.foster@yale.edu), Magnetic Resonance Research Center (MRRC), 300 Cedar St,

21   New Haven, CT 06519.

23   Word count: 2,999 (excluding key points, headers, abstract, references, legends, and

24   tables)

Key Points

Question: How do the amplitude and complexity of low-frequency oscillations in fMRI signals associate with psychosis-spectrum and mood disorders?

Findings: Findings suggest that the amplitude and complexity of low-frequency oscillation are associated with mood and psychosis-spectrum disorders in a wide-spread and global manner. Complexity, which is well-studied in EEG but not fMRI—emerged as a promising measure for further research. Our results also support a frequency-based behavioral encoding system operating in the BOLD signal.

Meaning: Our findings indicate that mood and psychosis-spectrum disorders differ from healthy controls in a mechanistically distinct way, as well as from one another across the whole brain. They also suggest that differences in low-frequency oscillations may represent shared and distinct markers of mental health and may help in developing treatment strategies that target these disruptions.

ABSTRACT

Objective: Conduct a mega-analysis of two complementary measures of resting-state

functional magnetic resonance imaging (rsfMRI) dynamics—amplitude of low-frequency

fluctuation (ALFF) and low-frequency spectral entropy (lfSE)—in mood and psychosis-

spectrum disorders to evaluate group differences and clinical symptom associations.

Design: ALFF and lfSE were calculated at the node-level by filtering data from 0.01 Hz

to 0.08 Hz, regressing demographic variables, and harmonizing sites. Group differences

were assessed using the Wilcoxon signed test. Symptom associations were evaluated

with Spearman's rho. Analyses were conducted at both whole-brain and network levels,

with sensitivity analyses to evaluate the impact of frequency brands.

51
52 **Setting**: Four independent open-source case-control datasets with resting-state

53 functional magnetic resonance imaging were used: the Center for Biomedical Research

54 Excellence, the Human Connectome Project for Early Psychosis, the Strategic

55 Research Program for Brain Sciences, and the UCLA Consortium for Neuropsychiatric

56 Phenomics.

57
Participants: Included participants had a mood disorder (bipolar, dysthymia, or major

depressive disorder, n=228, aged 38.31 ± 12.56 years), a psychosis-spectrum disorder

(early psychosis, schizophrenia spectrum disorder, or mood disorder with psychotic

symptoms, n=318, aged 29.8 ± 13.21 years), or a healthy control (n=535, aged 39.89 ±

15.3 years).

Main outcomes and Measures: To identify group differences and symptom
associations in mood and psychosis-spectrum disorders using ALFF and lfSE.

Results: ALFF in psychosis-spectrum was significantly lower than mood disorders and
controls (q's<0.001) at the whole-brain and network levels. lfSE in controls was
significantly lower than both psychosis-spectrum and mood disorders at the whole-brain
and network levels (q's<0.001). Whole-brain ALFF is positively associated with mood
symptoms ($rho=0.27$, $p<0.05$). Whole-brain lfSE is negatively associated with positive
($rho=-0.13$, $p<0.05$) and mood ($rho=-0.38$, $p<0.01$) symptoms. A greater sensitivity of
group differences and symptom associations to frequency ranges was observed in
mood disorders. ALFF is sensitive to medication.

Conclusions and Relevance: Widespread, global differences in ALFF and lfSE
underly psychosis-spectrum and mood disorders. lfSE may be applicable for wider use
in fMRI. Differences in spectral measures of brain dynamics may represent shared and
distinct markers of mental health.

INTRODUCTION

Mood (e.g., major depressive disorder and bipolar disorder) and psychosis-

spectrum (e.g., first-episode psychosis and schizophrenia) disorders cause significant

disability globally[1–4] and current treatment outcomes remain suboptimal. These

disorders also have shared and unique genetic risks[5–11]. Thus, continuing to advance

knowledge of their shared and unique mechanisms will be important for improving

differential diagnosis[11,12] and symptom management for affected individuals. Widely

associated with cognition and mental health[13–16], changes in brain dynamics may be

linked to the pathophysiology of mood and psychosis-spectrum disorders. However,

while applications of brain dynamics methods are becoming increasingly popular, there

is a scarcity of work exploring their simultaneous application to mood and psychosis-

spectrum disorders, raising questions of whether linked dynamics signatures are shared,

distinct, or both.

Spectral measures succinctly characterize brain dynamics across frequency

bands[17–19]. One widely used metric is the amplitude of low-frequency fluctuations

(ALFF), which quantifies the signal strength in the 0.01-0.08 Hz range and is well-

established in functional neuroimaging[20–24]. A comparable measure is low-frequency

spectral entropy (lfSE)—a frequency-bounded derivative of spectral entropy that

quantifies the complexity or unpredictability of fMRI signal fluctuations in the 0.01–0.08

Hz range. Spectral entropy has been broadly used in EEG[24], but not in fMRI studies.

ALFF and lfSE provide distinct, but complementary insights (amplitude versus

complexity) into intrinsic brain dynamics that may help biologically differentiate disorders

and mediate individual differences in symptom severity.

Prior work has identified ALFF differences across major functional networks in

mood[25–27] and psychosis-spectrum[28–30,27] disorders. In schizophrenia, studies found

reduced ALFF in the default mode (DMN), salience (SAL), sensorimotor, parietal, and

visual networks compared to healthy controls[31]. Early-stage schizophrenia has been

marked by increased ALFF in subcortical and visual networks and decreased ALFF in

the DMN and parietal networks[31]. In contrast, chronic schizophrenia includes

widespread increases in ALFF across frontal, SAL, temporal, and limbic networks,

alongside reductions in sensorimotor, parietal, and occipital areas[31]. In depression,

studies report increased ALFF in several regions—including lateralized subcortical

areas, occipital lobe, limbic areas, frontoparietal network (FPN), and the DMN—

compared to matched healthy controls[26,32–35]. Decreased ALFF is also observed in

somatosensory cortex[25,36,32], limbic areas, and cerebellum[37,38] for major depressive

disorder (MDD) compared to controls. In bipolar disorder, increased ALFF in the right

caudate and putamen[39], bilateral insula, medial prefrontal cortex, and decreased ALFF

in the left cerebellum[40] have been observed. In contrast, lfSE has not been studied in

mood and psychosis-spectrum disorders using fMRI. Moreover, to our knowledge no

previous work has comprehensively contrasted ALFF with lfSE in mood and psychosis-

spectrum disorders in one study.

Mega-analyses (e.g., a method that pools together the raw, individual level data

from multiple studies), are a promising approach for finding shared and distinct

biological indices across disorders. By pooling data from several smaller studies, mega-

analyses maximize population variance, increase generalizability, and improve

statistical power[41,42]. ALFF and lfSE are ideal for these analyses because they are

computationally tractable in large samples, easy to pool across diverse datasets due to

their conceptual simplicity, and neurobiologically interpretable at different spatial scales

(i.e., regional, network, or whole brain levels). By better accounting for psychiatric

heterogeneity, mega-analyses analyses can support the development of treatments that

target shared and distinct clinical mechanisms[43–45].

Here, we perform a mega-analysis in over 1000 individuals to compare ALFF and

lfSE in individuals with mood disorders, individuals with psychosis-spectrum disorders

and healthy controls. A secondary objective was to assess the utility of lfSE compared

to ALFF. Our study expands on previous empirical findings by adding a complexity

measure (i.e., lfSE)[46], using a large-scale sample size (n=1081), and comparing

measures across a wider range of symptoms. We also assess their association with

relevant symptom measures. Based on findings from previous works, we hypothesized

that ALFF and lfSE case-control group differences will be in the FPNI[47,48], SAL[47], and

DMN[47,49].

METHODS

Participants. We used four publicly available resting-state fMRI datasets (see Table

S1 for participant demographics), the Center for Biomedical Research Excellence[50]

(COBRE, n=99), the Human Connectome Project Early Psychosis[51] (HCP-EP, n=169),

the Strategic Research Program for Brain Sciences[52] (SRPBS, n=609) Multi-disorder

Connectivity Dataset, and the University of California Los Angeles (UCLA) Consortium

for Neuropsychiatric Phenomics[53] (CNP, n=204). The sample included 535 healthy

controls (HCs), 228 mood disorder (major depressive disorder, bipolar disorder, and

dysthymia), and 318 psychosis-spectrum disorder (early psychosis and schizophrenia

spectrum disorder) participants. All patients were diagnosed according to the DSM-5[54].

Demographics, symptom scores, medication information, and diagnosis breakdowns

are summarized in the Supplement methods and tables S1, S5, and S6.

Symptom Measures. Psychosis symptoms were assessed with the Positive and

Negative Syndrome Scale[55] (PANSS), which measures the presence and severity of

positive, negative, and general psychopathology for an individual within the past week.

The scale is widely used in clinical psychosis studies[56] and has demonstrated reliability

in assessing psychopathology of schizophrenia populations[57]. 230 participants had

PANSS scores (HCP-EP=107, SRPBS=123). Mood symptoms in individuals (n=62) with

a mood disorder (SRPBS=39) and health controls (SRPBS=23) were assessed with the

Beck's Depression Inventory[58] (BDI-II) . The BDI-II measures hallmark symptoms of

depression and has high internal consistency[58,59].

Data Preprocessing. The acquisition and imaging parameters for the datasets are

detailed elsewhere[50,51,60,52]. However, an abridged version is available in the

Supplement (see Table S2). Images were motion and slice time corrected using

SPM12[61]. BioImage Suite was used to perform mean white matter regression, cerebral

spinal fluid regression, and gray matter time, removing linear trends, and low-pass

filtering[62]. The Shen-268 atlas[61] was warped from Montreal Neurological Institute (MNI)

space into single-subject space. Next, the mean-time course of each node was

calculated by averaging its constituent voxels' time-series data.

Quality Control. Participants were excluded if they had an average frame-to-frame

displacement exceeding 0.2 mm **(**n=28), insufficient quality of linear or nonlinear

registration (n=16), had >100 missing nodes (n=10) or had less than 75% coverage of

relevant nodes for a given network or whole-brain analysis (n=15, see Supplement for

more details on data preservation process). This left a final sample of 1081 subjects for

further analysis: 823 for whole-brain analyses (averaging across all nodes) and varying

sample sizes for network analyses (Table S3-4).

ALFF Calculation and Analysis. First, the power spectral density of each node's time

series was calculated using periodogram function from the SciPy [63] signal processing

API in Python 3.9. This calculation entailed converting each participant's node time-

series into the frequency domain via the fast Fourier transform (FFT). Then, the power

spectral density was integrated over the low-frequency range (0.01 Hz ≤ frequency ≤

0.08 Hz) using SciPy's trapezoid integration method to calculate total power. Next, we

took the square root of that operation to quantify ALFF.

$$\text{ALFF} = \sqrt{\int_{f=0.01\,Hz}^{0.08\,Hz} P(f)\,\mathrm{df}}$$

Note: f=frequency and df=delta frequency

ALFF in this study is considered a proxy for the intensity of neural-associated brain

activity or the energy content of the low-frequency band. A high ALFF signifies greater

oscillation intensity while a low ALFF signifies lower oscillation intensity in the low-

frequency range. ALFF was assessed group-wise at the whole-brain by averaging all

nodes and network-levels using 10 canonical networks from the Shen 268 atlas.

Low-frequency Spectral Entropy Calculation and Analysis. lfSE was also calculated

at the node-level. Individual node time-series underwent a band-pass filter in the low-

frequency range (0.01 Hz ≤ frequency ≤ 0.08 Hz) using SciPy's butter and filtfilt

functions (order = 2)[21]. The signal was transformed into the frequency domain using

FFT, followed by power spectral density calculation and normalization to a probability

density function. Then, lfSE was calculated as the Shannon entropy of the power

spectral density was calculated using AntroPy[64] package in Python 3.

$$\hat{S}_{spectral} = \frac{-\sum_{f=0}^{Fs/2} P(f)\log_2 P(f)}{\log_2(num_f)}, \quad 0.01\,Hz < f < 0.08\,Hz$$

Note: f=frequency, Fs=sampling frequency, num$_f$= number of frequencies derived by the FFT

201
As a complexity measure, high lfSE (flatter power spectral density spectrum) represents

low complexity while low lfSE (power spectral density distribution with more peaks)

represents high complexity. lfSE was assessed group-wise at the whole-brain by

averaging all nodes and network-levels using 10 canonical networks from the Shen 268

atlas.

Statistical Analyses. Statistical analyses were constrained to the whole-brain and

network levels to capture macroscale brain dynamics. Broader scale such as these

have greater power than granular levels of inference[65]. Group comparisons of each low-

frequency brain measure were evaluated at $p<0.05$ using the Wilcoxon two-sided rank

sum test while correcting for multiple comparisons with false discovery rate (FDR). We

controlled for self-reported age, motion, and sex as variables of non-interest. First,

these factors were regressed[66] using a generalized linear model on ALFF and lfSE

values at the node level. Second, we applied the *neuroCombat*[67] R package to

harmonize data across sites. For empirical ranges, please refer to table S5). Where

relevant, we report corrected (q) and uncorrected (p) results for transparency but only

interpret significant corrected results ($q<0.05$).

Secondary analyses. We correlated symptoms scores (PANSS and BDI-II) and

covariate-regressed ALFF and lfSE values using Spearman's correlation at the whole-

brain and network levels. Participants for the PANSS analyses came from multiple sites

and underwent site harmonization using *neuroCombat*[67]. Participants for the BDI-II

analyses came from one site, so harmonization was not applied. Network-level

associations were considered statistically significant at $q<0.05$ after FDR correction.

Additionally, we evaluated similarity between the ALFF and lfSE by correlating these

values across participants ($p<0.05$) using Spearman's rho at the whole-brain and

network levels. We also calculated these similarities independently for each group.

These correlations were Fisher transformed and compared across groups using a two-

sample z-test.

Sensitivity Analysis. We used different frequency cutoffs (0.01-0.045 Hz and 0.035-

0.09 Hz) to assess if finer spectral divisions were driving the identified trends and

associations[68]. We also examined the effect of medication exposure by including current

medication status (binary: 1=on, 0=off) as an additional covariate.

RESULTS

Demographic differences between clinical groups. Significant group differences

were found for self-reported sex (χ^2=18.081, df = 2, p<0.001), age (W=75.48, df=2,

p<0.001) and average frame displacement (W= 8.599, df=2, p=0.014). Participants with

psychosis-spectrum (29.8 ± 13.21 years) were significantly younger than mood

disorders (38.31 ± 12.56 years) and controls (39.89 ± 15.3 years). Participants with

mood disorders also had significantly higher motion frame displacement (0.087 ± 0.045

mm) compared to matched HCs (0.083 ± 0.046 mm) and those with psychosis-

spectrum disorder (0.075 ± 0.037 mm).

Group differences in ALFF. At the whole-brain level, mood disorders exhibited the

greatest ALFF, followed by controls and then psychosis-spectrum (Figure 2A, Table 1).

Psychosis-spectrum ALFF values were significantly lower than both mood disorders and

controls (p's<0.001). No differences in whole-brain ALFF were detected between mood

disorders and controls. Similar differences were observed at the network level, where all

networks showed significant (q's<0.05, FDR-corrected) differences when comparing

psychosis-spectrum to mood disorders or health controls (Table 1). Results were similar

when potential outliers were removed (Table S8) and when removing psychosis-

spectrum patients with known affective disorders (Table S9).

Group differences in IfSE. At the whole-brain level, psychosis-spectrum had the

highest IfSE, followed by mood disorders and then HCs (Figure 2B). Controls were

significantly lower than both psychosis-spectrum and mood disorders at the whole-brain

(p's<0.001). No differences in whole-brain lfSE were detected between psychosis-

spectrum and mood disorders. At the network level, all networks exhibited significant

differences between health controls and psychosis-spectrum (q's<0.05, FDR-corrected).

The medial frontal, FPN, motor, subcortical, and cerebellar networks exhibited

significant differences between HCs and mood disorders (q's<0.05, FDR-corrected).

While no differences between psychosis-spectrum and mood disorders existed at the

whole-brain level, every network (other than the cerebellar network) exhibited significant

between-group differences (q's<0.05, FDR-corrected), likely due to the increases

sample size used in the network analyses (Table 2). Results were similar when

removing psychosis-spectrum patients with affective disorders (Table S10).

The psychosis-spectrum lfSE distribution was bimodal. Since the second (higher)

peak in the distribution may influence the results, we repeated the analyses after

removing these high-leverage points (see Supplemental Methods, Figure S4). Results

were similar (Table S11); however, mood disorders now showed the highest lfSE, and

only visual networks remained significantly different between psychosis-spectrum and

mood disorders (q's<0.05, FDR-corrected).

Associations with psychosis and mood symptoms. Whole-brain ALFF is positively

associated with BDI-II scores (rho=0.36, p= 0.0046; Figure 2C, Table S12) but had no

association with PANSS scores (Table S13). Additionally, whole-brain lfSE is negatively

associated with positive PANSS (rho=-0.13, p=0.0418, Table S14) and BDI-II scores

(rho=-0.38, p=0.002; see Figure 2D-E, Table S15). No other significant associations

with ALFF or lfSE at the whole-brain or network-level were observed (Tables S12-15).

Correlation of ALFF and lfSE. At all scales of analysis, ALFF and lfSE were negatively

correlated, such that individuals with higher ALFF values had lower lfSE values (rho's =

-0.39 to -0.22, Table S16-17). This relationship was also observed in our group

comparison, where ALFF was consistently greater in mood disorder compared to

psychosis-spectrum across scales and lfSE was consistently greater in psychosis-

spectrum compared to mood disorder across scales.

Sensitivity analyses of finer-scale frequency ranges. Results at the finer-scale

frequency ranges were broadly similar to the main results, with wide-spread differences

observed between groups (Tables S18-25). However, results involving the mood

disorder group exhibited the most changes in finer-scale frequency ranges. For example,

the lfSE of the DMN was significantly different between controls and mood disorders at

0.035-0.090 Hz, but not 0.01-0.045 Hz or the full frequency range. Additionally, whole-

brain ALFF was positively associated with BDI-II scores at 0.01-0.045 Hz but negatively

associated at 0.035-0.09 Hz (Table S22).

Medication differentially impacts ALFF and lfSE. Results after controlling for

medication were broadly similar for lfSE, but not ALFF (Table 3, Table S26). Only

comparisons involving the mood disorder group were significant for ALFF. The ALFF

comparisons between controls and psychosis spectrum were not significant.

DISCUSSION

In this study, we conducted a mega-analysis using two spectral measures—

oscillation intensity (ALFF) and complexity (lfSE) —to compare mood and psychosis-

spectrum disorders. Our results highlight extensive differences between these two

clinical conditions that span multiple dimensions of brain dynamics.

Notably, ALFF and lfSE uniquely differentiated clinical groups, providing distinct

insights into the biological underpinnings of mood and psychosis-spectrum disorders.

Psychosis-spectrum participants exhibited larger, widespread differences in lfSE

compared to mood disorder participants and controls, while mood disorder participants

exhibited more focal differences in lfSE compared to controls. Also, in line with several

studies [25,26,32–38], we observed significant differences between mood disorder

participants and controls in ALFF after controlling for medication. In some cases, ALFF

and lfSE results even converged. For example, despite our initial hypotheses, there is

widespread overlap in the functional networks that had significant group differences

between mood and psychosis-spectrum disorders.

Our sensitivity analyses revealed that ALFF results in mood disorders vary by

frequency band. In the narrower range of 0.035 to 0.090 Hz, larger group differences

and significant associations with mood symptoms were discovered. Moreover, there

was a reversal in the sign of significant mood symptom associations in the two finer-

scale frequency bands (0.01-0.045 Hz and 0.035-0.090 Hz). Aligned with these findings,

a previous study showed that associations between depressive symptoms and spectral

measures in the subgenual gyrus differ between frequency bands[69]. These diverging

associations are evidence of the functional specificity of frequency bands in the blood-

oxygen level dependent (BOLD) signal[70–72], a phenomena that has largely been

attributed to EEG studies of mental health[73]. As such, the notion that different

frequency bands reflect physiology occurring at a range of intrinsic time scales[73–76] is

likely a universal principle of brain function. Under this consideration, the finer bands

represented in our work comprise multiple physiological processes that contribute to

mood disorder pathology. If too broad a frequency range is adopted, interactive effects[69]

can dominate and obscure significant results in a competitive or mutually constructive

way[77]. Continued investigations of the functional relevance of specific frequency bands

in mood disorders and other psychiatric conditions are needed in fMRI research.

Out of the two measures, IfSE performed the best in detecting significant group

differences and symptom associations. It was also robust to common sources of

variance, which is crucial for mega-analyses, as they often encompass real-world

conditions (e.g., use of different scanners, participants with varying medication histories).

IfSE may more effectively capture changes in brain dynamics because as a measure of

complexity and irregularity[78], it considers the entire energy distribution across a

frequency band. Previous work employing similar methods in other imaging modalities

have identified signal irregularity and high signal variance as key features of the

psychosis-spectrum[79–82]. Our results, alongside broader literature findings, position

IfSE—a standard measure in the EEG literature[24] and related modalities[78,83–85]—as a

valuable index for increased application in fMRI. Future research should apply the IfSE

metric under different conditions and test its robustness against different covariates to

fully assess its mechanistic insights and resilience to non-interest variance sources.

In contrast, we observed a change in the significance of two group difference

pairs in ALFF analyses with and without medication (Table 3). This inconsistency may

indicate that ALFF is more susceptible to medication effects, or more specifically,

medication effects in mood and psychosis-spectrum disorders. Future studies should

further probe ALFF's sensitivity to medication in other psychiatric populations to confirm.

This study has several strengths. We combined distinct diagnoses into broader

groupings (e.g., mood and psychosis-spectrum disorders), which may better reflect the

dimensional nature of disorders and allow datasets with different diagnostic criteria to

be combined in a mega-analysis. We also used two spectral measures, including lfSE

(which is uncommon in fMRI analysis). It also has several limitations. First, participants

were assumed to exclusively have mood or a psychosis-spectrum disorder. However,

comorbidities are common, and individuals could belong in multiple groupings [86].

Second, we focus on measures that compress brain dynamics into a single number.

More complex summaries of dynamic measures[87] may prove to be more sensitive and

is a natural next step. Third, although we identify associations with symptoms scores,

causality cannot be established.

Through a mega-analysis, we investigated group differences using spectral

measures of brain dynamics in mood and psychosis-spectrum disorders. Our approach

revealed distinct brain signal properties in addition to shared characteristics between

the two patient groups. Future studies should continue to elucidate common

denominators as well as diagnosis-, symptom-, and individual-specific brain deviations

that build on top of disorder commonalities. Better characterization of shared brain

features, and disorder specific deviations could pave the way towards plausible

prevention and tailored treatment efforts.

Data and Code Availability. Data are publicly available through the Center for

Biomedical Research Excellence Phase I grant[50], the Human Connectome Project Early

Psychosis[51], the Strategic Research Program for Brain Sciences[52] and the UCLA

Consortium for Neuropsychiatric Phenomics LA5c Study[88].

(https://openneuro.org/datasets/ds000030/versions/00016). Preprocessing scripts are

available at https://www.nitrc.org/projects/bioimagesuite.

Acknowledgements. We are grateful to the participants who dedicated their time and

energy to participate in these large-scale data consortiums as well as the study team

who collected and collated the data. We also would like to thank Dr. Michelle Hampson

and Dr. Nicha Dvornek for initial input that guided the evolution of this research project

and Dr. Margaret L. Westwater for critically insightful feedback on the manuscript.

Funding Support. This publication was made possible by CTSA Grant Number UL1

TR001863 from the National Center for Advancing Translational Science (NCATS), a

component of the National Institutes of Health (NIH). Its contents are solely the

responsibility of the authors and do not necessarily represent the official views of NIH.

This work was also supported by NIH grants R01MH126133-03 (to MF), F31AA032179

(to JY), R01MH121095 (to DS and SM), and 5R01DK136623-02 (to DS and MK).

Author Contributions and Assistive Grammar Tools Acknowledgement. MF had full

access to the data and takes on full responsibility for accuracy of the data analysis. MF

conceived and designed the study. MF participated in the acquisition, preprocessing,

analysis, and interpretation of data. MF and DS participated in drafting the manuscript.

MF and DS performed statistical analyses. SM and JY provided critical review of the

content and provided clinical interpretations of the analyses. MK participated in quality

control of data. JY participated in acquisition and preprocessing of some of the data.

Portions of the writing in this manuscript were edited with artificial intelligence (AI)-

based tools to improve clarity and grammar. The authors critically reviewed and edited

all AI-assisted content to ensure accuracy and originality. No AI tools were used in the

generation of scientific content, data analysis, or interpretation. All intellectual content,

interpretations, and conclusions are our own.

REFERENCES

1. Ayano G, Demelash S, yohannes Z, et al. Misdiagnosis, detection rate, and associated factors of severe psychiatric disorders in specialized psychiatry centers in Ethiopia. *Ann Gen Psychiatry*. 2021;20:10. doi:10.1186/s12991-021-00333-7

2. Calabrese J, Al Khalili Y. Psychosis. In: *StatPearls*. StatPearls Publishing; 2024. Accessed March 31, 2024. http://www.ncbi.nlm.nih.gov/books/NBK546579/

3. Sekhon S, Gupta V. Mood Disorder. In: *StatPearls*. StatPearls Publishing; 2025. Accessed February 14, 2025. http://www.ncbi.nlm.nih.gov/books/NBK558911/

4. Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet Psychiatry*. 2022;9(2):137-150. doi:10.1016/S2215-0366(21)00395-3

5. International Schizophrenia Consortium, Purcell SM, Wray NR, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009;460(7256):748-752. doi:10.1038/nature08185

6. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *The Lancet*. 2013;381(9875):1371-1379. doi:10.1016/S0140-6736(12)62129-1

7. Ruderfer DM, Ripke S, McQuillin A, et al. Genomic dissection of bipolar disorder and schizophrenia including 28 subphenotypes. *Cell*. 2018;173(7):1705-1715.e16. doi:10.1016/j.cell.2018.05.046

8. Fusar-Poli P, Solmi M, Brondino N, et al. Transdiagnostic psychiatry: a systematic review. *World Psychiatry*. 2019;18(2):192-207. doi:10.1002/wps.20631

9. Bigdeli TB, Fanous AH, Li Y, et al. Genome-Wide Association Studies of Schizophrenia and Bipolar Disorder in a Diverse Cohort of US Veterans. *Schizophr Bull*. 2020;47(2):517-529. doi:10.1093/schbul/sbaa133

10. Xie C, Xiang S, Shen C, et al. A shared neural basis underlying psychiatric comorbidity. *Nat Med*. 2023;29(5):1232-1242. doi:10.1038/s41591-023-02317-4

11. Brand SJ, Möller M, Harvey BH. A Review of Biomarkers in Mood and Psychotic Disorders: A Dissection of Clinical vs. Preclinical Correlates. *Curr Neuropharmacol*. 2015;13(3):324-368. doi:10.2174/1570159X13666150307004545

12. Kas MJH, Penninx BWJH, Knudsen GM, et al. Precision psychiatry roadmap: towards a biology-informed framework for mental disorders. *Mol Psychiatry*. 2025;30(8):3846-3855. doi:10.1038/s41380-025-03070-5

13. Newson JJ, Thiagarajan TC. EEG Frequency Bands in Psychiatric Disorders: A
Review of Resting State Studies. *Front Hum Neurosci.* 2019;12.
doi:10.3389/fnhum.2018.00521

14. Raut RV, Snyder AZ, Mitra A, et al. Global waves synchronize the brain's functional
systems with fluctuating arousal. *Sci Adv.* 2021;7(30):eabf2709.
doi:10.1126/sciadv.abf2709

15. Xu Y, Long X, Feng J, Gong P. Interacting spiral wave patterns underlie complex
brain dynamics and are related to cognitive processing. *Nat Hum Behav.*
2023;7(7):1196-1215. doi:10.1038/s41562-023-01626-5

16. Foster M, Scheinost D. Brain states as wave-like motifs. *Trends in Cognitive*
Sciences. Published online 2024. Accessed April 29, 2024.
https://www.cell.com/trends/cognitive-sciences/abstract/S1364-6613(24)00057-3

17. Duff EP, Johnston LA, Xiong J, Fox PT, Mareels I, Egan GF. The power of spectral
density analysis for mapping endogenous BOLD signal fluctuations. *Human Brain*
Mapping. 2008;29(7):778-790. doi:10.1002/hbm.20601

18. Elble RJ, Ondo W. Tremor rating scales and laboratory tools for assessing tremor. *J*
Neurol Sci. 2022;435:120202. doi:10.1016/j.jns.2022.120202

19. Zang Z, Qiao Y, Yan S, Lu J. Reliability and Validity of Power Spectrum Slope
(PSS): A Metric for Measuring Resting-State Functional Magnetic Resonance
Imaging Activity of Single Voxels. *Front Neurosci.* 2022;16:871609.
doi:10.3389/fnins.2022.871609

20. Yang H, Long XY, Yang Y, et al. Amplitude of low frequency fluctuation within visual
areas revealed by resting-state functional MRI. *Neuroimage.* 2007;36(1):144-152.
doi:10.1016/j.neuroimage.2007.01.054

21. Zou QH, Zhu CZ, Yang Y, et al. An improved approach to detection of amplitude of
low-frequency fluctuation (ALFF) for resting-state fMRI: Fractional ALFF. *Journal of*
Neuroscience Methods. 2008;172(1):137-141. doi:10.1016/j.jneumeth.2008.04.012

22. Zuo XN, Di Martino A, Kelly C, et al. The Oscillating Brain: Complex and Reliable.
Neuroimage. 2010;49(2):1432-1445. doi:10.1016/j.neuroimage.2009.09.037

23. Li MT, Zhang SX, Li X, et al. Amplitude of Low-Frequency Fluctuation in Multiple
Frequency Bands in Tension-Type Headache Patients: A Resting-State Functional
Magnetic Resonance Imaging Study. *Front Neurosci.* 2021;15.
doi:10.3389/fnins.2021.742973

24. Yen C, Lin CL, Chiang MC. Exploring the Frontiers of Neuroimaging: A Review of
Recent Advances in Understanding Brain Functioning and Disorders. *Life.*
2023;13(7):1472. doi:10.3390/life13071472

25. Wang L, Dai W, Su Y, et al. Amplitude of Low-Frequency Oscillations in First-
Episode, Treatment-Naive Patients with Major Depressive Disorder: A Resting-State
Functional MRI Study. *PLOS ONE*. 2012;7(10):e48658.
doi:10.1371/journal.pone.0048658

26. Chen Q, Bi Y, Zhao X, et al. Regional amplitude abnormities in the major depressive
disorder: A resting-state fMRI study and support vector machine analysis. *Journal of*
Affective Disorders. 2022;308:1-9. doi:10.1016/j.jad.2022.03.079

27. Chen C, Zhang B, Qin X, et al. Altered resting-state brain activity of the superior
parietal cortex and striatum in major depressive disorder and schizophrenia. *Asian*
Journal of Psychiatry. 2024;102:104303. doi:10.1016/j.ajp.2024.104303

28. Turner JA, Damaraju E, Van Erp TGM, et al. A multi-site resting state fMRI study on
the amplitude of low frequency fluctuations in schizophrenia. *Front Neurosci*. 2013;7.
doi:10.3389/fnins.2013.00137

29. Wang P, Yang J, Yin Z, et al. Amplitude of low-frequency fluctuation (ALFF) may be
associated with cognitive impairment in schizophrenia: a correlation study. *BMC*
Psychiatry. 2019;19:30. doi:10.1186/s12888-018-1992-4

30. Li X, Liu Q, Chen Z, et al. Abnormalities of Regional Brain Activity in Patients With
Schizophrenia: A Longitudinal Resting-State fMRI Study. *Schizophrenia Bulletin*.
2023;49(5):1336-1344. doi:10.1093/schbul/sbad054

31. Gong J, Wang J, Luo X, et al. Abnormalities of intrinsic regional brain activity in first-
episode and chronic schizophrenia: a meta-analysis of resting-state functional MRI.
J Psychiatry Neurosci. 2020;45(1):55-68. doi:10.1503/jpn.180245

32. Wu Z, Luo Q, Wu H, et al. Amplitude of Low-Frequency Oscillations in Major
Depressive Disorder With Childhood Trauma. *Front Psychiatry*. 2021;11:596337.
doi:10.3389/fpsyt.2020.596337

33. Zhang B, Qi S, Liu S, Liu X, Wei X, Ming D. Altered spontaneous neural activity in
the precuneus, middle and superior frontal gyri, and hippocampus in college
students with subclinical depression. *BMC Psychiatry*. 2021;21(1):280.
doi:10.1186/s12888-021-03292-1

34. Zhang Z, Bo Q, Li F, et al. Increased ALFF and functional connectivity of the right
striatum in bipolar disorder patients. *Progress in Neuro-Psychopharmacology and*
Biological Psychiatry. 2021;111:110140. doi:10.1016/j.pnpbp.2020.110140

35. Luo G, Zhou J, Liu L, Song X, Peng M, Zhang X. Abnormal ReHo and ALFF values
in drug-naïve depressed patients with suicidal ideation or attempts: Evidence from
the REST-meta-MDD consortium. *Progress in Neuro-Psychopharmacology and*
Biological Psychiatry. 2025;136:111210. doi:10.1016/j.pnpbp.2024.111210

36. Liu F, Guo W, Liu L, et al. Abnormal amplitude low-frequency oscillations in
medication-naive, first-episode patients with major depressive disorder: a resting-
state fMRI study. *J Affect Disord*. 2013;146(3):401-406.
doi:10.1016/j.jad.2012.10.001

37. Liu J, Ren L, Womer FY, et al. Alterations in amplitude of low frequency fluctuation
in treatment-naïve major depressive disorder measured with resting-state fMRI.
Human Brain Mapping. 2014;35(10):4979-4988. doi:10.1002/hbm.22526

38. Zhou M, Hu X, Lu L, et al. Intrinsic cerebral activity at resting state in adults with
major depressive disorder: A meta-analysis. *Progress in Neuro-*
Psychopharmacology and Biological Psychiatry. 2017;75:157-164.
doi:10.1016/j.pnpbp.2017.02.001

39. Zhang Z, Bo Q, Li F, et al. Increased ALFF and functional connectivity of the right
striatum in bipolar disorder patients. *Prog Neuropsychopharmacol Biol Psychiatry*.
2021;111:110140. doi:10.1016/j.pnpbp.2020.110140

40. Gong J, Wang J, Qiu S, et al. Common and distinct patterns of intrinsic brain activity
alterations in major depression and bipolar disorder: voxel-based meta-analysis.
Transl Psychiatry. 2020;10(1):353. doi:10.1038/s41398-020-01036-5

41. Costafreda SG. Pooling fMRI data: meta-analysis, mega-analysis and multi-center
studies. *Front Neuroinform*. 2009;3. doi:10.3389/neuro.11.033.2009

42. Bigdely-Shamlo N, Touryan J, Ojeda A, Kothe C, Mullen T, Robbins K. Automated
EEG mega-analysis I: Spectral and amplitude characteristics across studies.
NeuroImage. 2020;207:116361. doi:10.1016/j.neuroimage.2019.116361

43. Zambrano-Vazquez L, Levy HC, Belleau EL, et al. Using the research domain
criteria framework to track domains of change in comorbid PTSD and SUD. *Psychol*
Trauma. 2017;9(6):679-687. doi:10.1037/tra0000257

44. McQuaid RJ. Transdiagnostic biomarker approaches to mental health disorders:
Consideration of symptom complexity, comorbidity and context. *Brain Behav Immun*
Health. 2021;16:100303. doi:10.1016/j.bbih.2021.100303

45. Holmes AJ, Patrick LM. The Myth of Optimality in Clinical Neuroscience. *Trends in*
Cognitive Sciences. 2018;22(3):241-257. doi:10.1016/j.tics.2017.12.006

46. Tenev A, Markovska-Simoska S, Müller A, Mishkovski I. Entropy, complexity, and
spectral features of EEG signals in autism and typical development: a quantitative
approach. *Front Psychiatry*. 2025;16. doi:10.3389/fpsyt.2025.1505297

47. Menon V. Large-scale brain networks and psychopathology: a unifying triple network
model. *Trends Cogn Sci*. 2011;15(10):483-506. doi:10.1016/j.tics.2011.08.003

48. Foster ML, Ye J, Powers AR, Dvornek NC, Scheinost D. Connectome-based
predictive modeling of early and chronic psychosis symptoms.
Neuropsychopharmacol. Published online February 27, 2025:1-9.
doi:10.1038/s41386-025-02064-9

49. Schilbach L, Hoffstaedter F, Müller V, et al. Transdiagnostic commonalities and
differences in resting state functional connectivity of the default mode network in
schizophrenia and major depression. *NeuroImage: Clinical.* 2016;10:326-335.
doi:10.1016/j.nicl.2015.11.021

50. COBRE Phase 3 | The Mind Research Network (MRN). Accessed February 14,
2025. https://www.mrn.org/common/cobre-phase-3

51. Lewandowski KE, Bouix S, Ongur D, Shenton ME. Neuroprogression across the
Early Course of Psychosis. *J Psychiatr Brain Sci.* 2020;5:e200002.
doi:10.20900/jpbs.20200002

52. Tanaka SC, Yamashita A, Yahata N, et al. A multi-site, multi-disorder resting-state
magnetic resonance image database. *Sci Data.* 2021;8(1):227. doi:10.1038/s41597-
021-01004-8

53. Poldrack RA, Congdon E, Triplett W, et al. A phenome-wide examination of neural
and cognitive function. *Sci Data.* 2016;3(1):160110. doi:10.1038/sdata.2016.110

54. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental
Disorders.* Fifth Edition. American Psychiatric Association; 2013. Accessed March
26, 2025. https://psychiatryonline.org/doi/book/10.1176/appi.books.9780890425596

55. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS)
for schizophrenia. *Schizophr Bull.* 1987;13(2):261-276. doi:10.1093/schbul/13.2.261

56. Shafer A, Dazzi F. Meta-analysis of the positive and Negative Syndrome Scale
(PANSS) factor structure. *Journal of Psychiatric Research.* 2019;115:113-120.
doi:10.1016/j.jpsychires.2019.05.008

57. Kay SR, Opler LA, Lindenmayer JP. Reliability and validity of the positive and
negative syndrome scale for schizophrenics. *Psychiatry Res.* 1988;23(1):99-110.
doi:10.1016/0165-1781(88)90038-8

58. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring
depression. *Arch Gen Psychiatry.* 1961;4:561-571.
doi:10.1001/archpsyc.1961.01710120031004

59. Beck AT, Steer RA, Carbin MG. Psychometric properties of the Beck Depression
Inventory: Twenty-five years of evaluation. *Clinical Psychology Review.*
1988;8(1):77-100. doi:10.1016/0272-7358(88)90050-5

60. Gorgolewski KJ, Durnez J, Poldrack RA. Preprocessed Consortium for
Neuropsychiatric Phenomics dataset. *F1000Res*. 2017;6:1262.
doi:10.12688/f1000research.11964.2

61. Shen X, Tokoglu F, Papademetris X, Constable RT. Groupwise whole-brain
parcellation from resting-state fMRI data for network node identification. *Neuroimage*.
2013;82:403-415. doi:10.1016/j.neuroimage.2013.05.081

62. Shen X, Finn ES, Scheinost D, et al. Using connectome-based predictive modeling
to predict individual behavior from brain connectivity. *Nat Protoc*. 2017;12(3):506-
518. doi:10.1038/nprot.2016.178

63. Virtanen P, Gommers R, Oliphant TE, et al. SciPy 1.0: fundamental algorithms for
scientific computing in Python. *Nat Methods*. 2020;17(3):261-272.
doi:10.1038/s41592-019-0686-2

64. antropy.app_entropy — antropy 0.1.9 documentation. Accessed July 24, 2025.
https://raphaelvallat.com/antropy/build/html/generated/antropy.app_entropy.html

65. Noble S, Mejia AF, Zalesky A, Scheinost D. Improving power in functional magnetic
resonance imaging by moving beyond cluster-level inference. *Proceedings of the
National Academy of Sciences*. 2022;119(32):e2203020119.
doi:10.1073/pnas.2203020119

66. Schöttner M, Bolton TAW, Patel J, Nahálka AT, Vieira S, Hagmann P. Exploring the
latent structure of behavior using the Human Connectome Project's data. *Sci Rep*.
2023;13(1):713. doi:10.1038/s41598-022-27101-1

67. Package neuroCombat details | Neuroconductor. Accessed August 25, 2025.
https://neuroconductor.org/package/neuroCombat#citation

68. Zheng R, Chen Y, Jiang Y, et al. Dynamic Altered Amplitude of Low-Frequency
Fluctuations in Patients With Major Depressive Disorder. *Front Psychiatry*. 2021;12.
doi:10.3389/fpsyt.2021.683610

69. Song X, Hu X, Zhou S, et al. Association of specific frequency bands of functional
MRI signal oscillations with motor symptoms and depression in Parkinson's disease.
Sci Rep. 2015;5(1):16376. doi:10.1038/srep16376

70. Caplan JB, Madsen JR, Raghavachari S, Kahana MJ. Distinct Patterns of Brain
Oscillations Underlie Two Basic Parameters of Human Maze Learning. *Journal of
Neurophysiology*. 2001;86(1):368-380. doi:10.1152/jn.2001.86.1.368

71. Ries A, Hollander M, Glim S, Meng C, Sorg C, Wohlschläger A. Frequency-
Dependent Spatial Distribution of Functional Hubs in the Human Brain and
Alterations in Major Depressive Disorder. *Front Hum Neurosci*. 2019;13:146.
doi:10.3389/fnhum.2019.00146

72. Bonnefond M, Kastner S, Jensen O. Communication between Brain Areas Based on
Nested Oscillations. *eNeuro*. 2017;4(2). doi:10.1523/ENEURO.0153-16.2017

73. Attar ET. Review of electroencephalography signals approaches for mental stress
assessment. *Neurosciences (Riyadh)*. 2022;27(4):209-215.
doi:10.17712/nsj.2022.4.20220025

74. Buzsáki G, Draguhn A. Neuronal oscillations in cortical networks. *Science*.
2004;304(5679):1926-1929. doi:10.1126/science.1099745

75. Motivation, emotion, and their inhibitory control mirrored in brain oscillations -
PubMed. Accessed August 7, 2025. https://pubmed.ncbi.nlm.nih.gov/17145079/

76. Liu ZZ, Qu HJ, Tian ZL, et al. Reproducibility of frequency-dependent low frequency
fluctuations in reaction time over time and across tasks. *PLOS ONE*.
2017;12(9):e0184476. doi:10.1371/journal.pone.0184476

77. Guo X, Li J, Su Q, et al. Transcriptional correlates of frequency-dependent brain
functional activity associated with symptom severity in degenerative cervical
myelopathy. *NeuroImage*. 2023;284:120451.
doi:10.1016/j.neuroimage.2023.120451

78. Helakari H, Kananen J, Huotari N, et al. Spectral entropy indicates
electrophysiological and hemodynamic changes in drug-resistant epilepsy – A
multimodal MREG study. *NeuroImage: Clinical*. 2019;22:101763.
doi:10.1016/j.nicl.2019.101763

79. Yang GJ, Murray JD, Repovs G, et al. Altered global brain signal in schizophrenia.
Proceedings of the National Academy of Sciences. 2014;111(20):7438-7443.
doi:10.1073/pnas.1405289111

80. Bachiller A, Lubeiro A, Díez Á, et al. Decreased entropy modulation of EEG
response to novelty and relevance in schizophrenia during a P300 task. *Eur Arch*
Psychiatry Clin Neurosci. 2015;265(6):525-535. doi:10.1007/s00406-014-0525-5

81. Akar SA, Kara S, Latifoğlu F, Bilgiç V. Analysis of the Complexity Measures in the
EEG of Schizophrenia Patients. *International Journal of Neural Systems*. Published
online February 21, 2016. doi:10.1142/S0129065716500088

82. Karanikolaou M, Limanowski J, Northoff G. Does temporal irregularity drive
prediction failure in schizophrenia? temporal modelling of ERPs. *Schizophr*.
2022;8(1):23. doi:10.1038/s41537-022-00239-7

83. Poza J, Hornero R, Abásolo D, Fernández A, Escudero J. Analysis of spontaneous
MEG activity in patients with Alzheimer's disease using spectral entropies. *Annu Int*
Conf IEEE Eng Med Biol Soc. 2007;2007:6180-6183.
doi:10.1109/IEMBS.2007.4353766

84. Gómez C, Hornero R. Entropy and Complexity Analyses in Alzheimer's Disease: An
MEG Study. *Open Biomed Eng J*. 2010;4:223-235.
doi:10.2174/1874120701004010223

85. Ferdinando H, Moradi S, Korhonen V, Helakari H, Kiviniemi V, Myllylä T. Spectral
entropy provides separation between Alzheimer's disease patients and controls: a
study of fNIRS. *Eur Phys J Spec Top*. 2023;232(5):655-662.
doi:10.1140/epjs/s11734-022-00753-w

86. McGrath JJ, Lim CCW, Plana-Ripoll O, et al. Comorbidity within mental disorders: a
comprehensive analysis based on 145 990 survey respondents from 27 countries.
Epidemiol Psychiatr Sci. 2020;29:e153. doi:10.1017/S2045796020000633

87. Hu L, Chen H, Su W, et al. Aberrant static and dynamic functional connectivity of the
executive control network in lung cancer patients after chemotherapy: a longitudinal
fMRI study. *Brain Imaging and Behavior*. 2020;14(3):927-940. doi:10.1007/s11682-
020-00287-6

88. Gorgolewski KJ, Durnez J, Poldrack RA. Preprocessed Consortium for
Neuropsychiatric Phenomics dataset. *F1000Res*. 2017;6:1262.
doi:10.12688/f1000research.11964.2

 FIGURE HEADINGS

 Figure 1. *Schematic of analysis pipeline at node and network levels.* This figure

 illustrates the study design and measurement calculation process. Four large-scale

 datasets were used in this study. A) First, the power spectrums were taken from

 preprocessed node-wise data, and the two low-frequency measures were calculated.

 Then, covariate regression of nuisance variables such as age, mean frame

 displacement, and sex were regressed using a generalized linear model. Where

 relevant, harmonization was performed to account for site effects. Then, the appropriate

 means were calculated for whole-brain (average over all nodes) and network (average

 over all member network nodes were taken for each of the 10 functional networks)

 analyses. B) Finally, group differences and symptom associations at whole-brain and

 network levels were calculated. FFT=Fast Fourier transform, ALFF=Amplitude of Low-

 Frequency Fluctuation, lfSE=low-frequency spectral entropy. $\bar{x}$=mean.

 Figure 2. *Group Differences at Whole-brain Level and Symptom Associations.* Panels A

 and B feature a violin plot of the results for ALFF and lfSE respectively. Dots represent

 the participants (n=823; controls=366, mood=139, psychosis=318), colors represent the

 groups (controls, mood, psychosis). Lines of fit are based on linear model (for

 visualization). Correlations are reported as Spearman's ρ. N.S. = not significant.

 ***=p<0.001. Panels C-E depict significant ALFF and lfSE symptom associations. Dots

 represent participants with a symptom score. A) Psychosis disorders had significantly

 lower ALFF than controls and mood disorders. B) Controls had significantly lower lfSE

 than mood and psychosis disorders. C) Significant BDI-II symptom associations with

 whole-brain ALFF (n=62). D) Significant positive PANSS symptom associations with

694   whole-brain lfSE (n=230). E) Significant BDI-II symptom associations with whole-brain

695   lfSE (n=62).

 TABLES: MAIN ANALYSIS

Scale	n	Clinical Group			Wilcoxon Rank Pairwise p		
		Control	Mood	Psychosis	C vs M	C vs P	M vs P
Whole-brain	823	0.008 ± 0.021	0.01 ± 0.019	-0.014 ± 0.039	p = 0.399 q = 0.399	**p = 5.88e-15** **q = 1.77e-14**	**p = 1.83e-11** **q = 2.74e-11**
Medial Frontal	846	0.008 ± 0.025	0.011 ± 0.025	-0.014 ± 0.041	p = 0.149 q = 0.149	**p = 4.28e-13** **q = 1.28e-12**	**p = 2.68e-11** **q = 4.02e-11**
Frontoparietal	852	0.007 ± 0.022	0.01 ± 0.022	-0.013 ± 0.04	p = 0.146 q = 0.146	**p = 5.49e-12** **q = 1.65e-11**	**p = 7.03e-11** **q = 1.05e-10**
Default Mode	882	0.007 ± 0.026	0.011 ± 0.025	-0.015 ± 0.045	p = 0.0889 q = 0.0889	**p = 3.77e-11** **q = 1.13e-10**	**p = 9.94e-11** **q = 1.49e-10**
Motor	881	0.008 ± 0.025	0.011 ± 0.024	-0.015 ± 0.042	p = 0.212 q = 0.212	**p = 2.34e-14** **q = 7.01e-14**	**p = 4.57e-12** **q = 6.86e-12**
Visual I	1081	0.004 ± 0.028	0.008 ± 0.029	-0.013 ± 0.043	p = 0.0671 q = 0.0671	**p = 3.63e-08** **q = 5.44e-08**	**p = 9.89e-09** **q = 2.97e-08**
Visual II	953	0.006 ± 0.03	0.008 ± 0.031	-0.013 ± 0.045	p = 0.636 q = 0.636	**p = 1.87e-09** **q = 5.6e-09**	**p = 1.74e-07** **q = 2.61e-07**
Visual Association	906	0.008 ± 0.025	0.011 ± 0.029	-0.016 ± 0.044	p = 0.142 q = 0.142	**p = 5.94e-14** **q = 1.78e-13**	**p = 9.27e-12** **q = 1.39e-11**
Salience	936	0.006 ± 0.024	0.01 ± 0.025	-0.014 ± 0.039	p = 0.0987 q = 0.0987	**p = 2.48e-12** **q = 3.72e-12**	**p = 2.14e-12** **q = 3.72e-12**
Subcortical	906	0.007 ± 0.026	0.012 ± 0.028	-0.016 ± 0.041	p = 0.212 q = 0.212	**p = 2.03e-14** **q = 6.09e-14**	**p = 1.79e-12** **q = 2.68e-12**
Cerebellar	823	0.011 ± 0.033	0.012 ± 0.029	-0.017 ± 0.053	p = 0.998 q = 0.998	**p = 6.45e-13** **q = 1.93e-12**	**p = 1.17e-08** **q = 1.75e-08**

697 **Table 1.** *Amplitude of low-frequency (ALFF, 0.01-0.08 Hz) value distributions by whole-*

698 *brain, network and clinical group.* Reported values have had covariates linearly

699 regressed and undergone site harmonization. n=sample size. Bolded table entries

700 represent significant comparisons after FDR correction.

Scale	n	Clinical Group			Wilcoxon Rank Pairwise p		
		Control	Mood	Psychosis	C vs M	C vs P	M vs P
Whole-brain	823	-0.006 ± 0.012	0 ± 0.015	0.007 ± 0.023	**p = 1.52e-05** **q = 2.28e-05**	**p = 2.5e-13** **q = 7.49e-13**	p = 0.141 q = 0.141
Medial Frontal	846	-0.006 ± 0.014	-0.002 ± 0.015	0.008 ± 0.028	**p = 0.00509** **q = 0.00706**	**p = 8.25e-13** **q = 2.48e-12**	**p = 0.00706** **q = 0.00706**
Frontoparietal	853	-0.005 ± 0.014	-0.001 ± 0.019	0.007 ± 0.025	**p = 0.00107** **q = 0.00161**	**p = 2.39e-10** **q = 7.18e-10**	**p = 0.0463** **q = 0.0463**
Default Mode	882	-0.004 ± 0.016	-0.004 ± 0.019	0.007 ± 0.03	p = 0.218 q = 0.218	**p = 7.27e-08** **q = 2.18e-07**	**p = 0.00191** **q = 0.00287**
Motor	881	-0.006 ± 0.014	-0.003 ± 0.019	0.008 ± 0.028	**p = 0.00545** **q = 0.00817**	**p = 4.66e-11** **q = 1.4e-10**	**p = 0.00848** **q = 0.00848**
Visual I	1081	-0.002 ± 0.017	-0.009 ± 0.03	0.009 ± 0.026	p = 0.0793 q = 0.0793	**p = 1.95e-09** **q = 3.37e-09**	**p = 2.25e-09** **q = 3.37e-09**
Visual II	954	-0.003 ± 0.018	-0.004 ± 0.027	0.007 ± 0.027	p = 0.164 q = 0.164	**p = 1.41e-08** **q = 4.24e-08**	**p = 0.000741** **q = 0.00111**
Visual Association	906	-0.005 ± 0.016	-0.006 ± 0.027	0.01 ± 0.028	p = 0.123 q = 0.123	**p = 7.11e-13** **q = 2.13e-12**	**p = 8.02e-05** **q = 0.00012**
Salience	846	-0.005 ± 0.013	-0.004 ± 0.021	0.009 ± 0.027	p = 0.0632 q = 0.0632	**p = 4.85e-10** **q = 1.46e-09**	**p = 0.000724** **q = 0.00109**
Subcortical	853	-0.005 ± 0.013	-0.005 ± 0.025	0.009 ± 0.026	**p = 0.00158** **q = 0.00237**	**p = 4.02e-12** **q = 1.2e-11**	**p = 0.0309** **q = 0.0309**
Cerebellar	882	-0.007 ± 0.019	0.002 ± 0.02	0.007 ± 0.034	**p = 5.31e-06** **q = 7.96e-06**	**p = 2.65e-08** **q = 7.96e-08**	p = 0.641 q = 0.641

701 **Table 2.** *Low-frequency spectral entropy (lfSE, 0.01-0.08 Hz) value distributions by*

702 *whole-brain, network and clinical group.* Reported values have had covariates linearly

703 regressed and undergone site harmonization. n=sample size. Bolded table entries

704 represent significant comparisons after FDR correction.

Scale	n	Clinical Groups (median +/- std)			Wilcoxon Rank Pairwise p		
		C	M	P	C vs M	C vs P	M vs P
Whole-brain	554 (C=232 M=105 P=217)	-0.004 ± 0.021	0.026 ± 0.03	-0.008 ± 0.039	**p = 2.31e-18** **q = 6.93e-18**	p = 0.937 q = 0.937	**p = 1.56e-13** **q = 2.34e-13**
Medial Frontal	554 (C=232 M=105 P=217)	-0.004 ± 0.025	0.025 ± 0.032	-0.008 ± 0.038	**p = 7.97e-15** **q = 2.39e-14**	p = 0.788 q = 0.788	**p = 1.91e-12** **q = 2.86e-12**
Frontoparietal	554 (C=232 M=105 P=217)	-0.005 ± 0.026	0.022 ± 0.026	-0.005 ± 0.038	**p = 2.82e-16** **q = 8.45e-16**	p = 0.539 q = 0.539	**p = 1.98e-11** **q = 2.97e-11**
Default Mode	554 (C=232 M=105 P=217)	-0.004 ± 0.026	0.026 ± 0.034	-0.008 ± 0.042	**p = 1.52e-14** **q = 4.56e-14**	p = 0.946 q = 0.946	**p = 6.04e-12** **q = 9.05e-12**
Motor	555 (C=232 M=106 P=217)	-0.004 ± 0.024	0.025 ± 0.032	-0.008 ± 0.039	**p = 3.87e-15** **q = 1.16e-14**	p = 0.755 q = 0.755	**p = 1.88e-12** **q = 2.82e-12**
Visual I	558 (C=232 M=109, P=217)	-0.004 ± 0.027	0.019 ± 0.031	-0.006 ± 0.04	**p = 4.76e-11** **q = 1.43e-10**	p = 0.92 q = 0.92	**p = 6.96e-08** **q = 1.04e-07**
Visual II	556 (C=232 M=107 P=217)	-0.004 ± 0.028	0.026 ± 0.039	-0.008 ± 0.043	**p = 7.82e-14** **q = 2.35e-13**	p = 0.645 q = 0.645	**p = 1.12e-10** **q = 1.68e-10**
Visual Association	555 (C=232 M=106 P=217)	-0.004 ± 0.025	0.025 ± 0.034	-0.008 ± 0.041	**p = 3.55e-15** **q = 1.07e-14**	p = 0.722 q = 0.722	**p = 1.55e-12** **q = 2.33e-12**
Salience	555 (C=232 M=106 P=217)	-0.004 ± 0.023	0.024 ± 0.031	-0.007 ± 0.038	**p = 2.49e-15** **q = 7.47e-15**	p = 0.871 q = 0.871	**p = 1.21e-11** **q = 1.81e-11**
Subcortical	554 (C=232 M=105 P=217)	-0.004 ± 0.024	0.024 ± 0.032	-0.008 ± 0.04	**p = 4.28e-13** **q = 1.28e-12**	p = 0.416 q = 0.416	**p = 1.29e-10** **q = 1.94e-10**
Cerebellar	554 (C=232 M=105 P=217)	-0.005 ± 0.03	0.03 ± 0.042	-0.009 ± 0.055	**p = 5.36e-14** **q = 1.61e-13**	p = 0.643 q = 0.643	**p = 2e-09** **q = 3e-09**

Table 3. *Effect of Medication Exposure on ALFF Group Differences.* A subsample of the

patient group had known exposure to a psychiatric medication and a sensitivity analysis

was performed on this subsample. Mood-psychosis effects remained consistent with

analyses that did not consider medication exposure. However, case-control differences

for mood disorders became significant, while those for psychosis-spectrum disorders

became insignificant. Motor and visual I networks were analyzed using the mean only

harmonization algorithm (see Supplementary methods).

A. Node Level

For each subject:

Time Series

Power Density Spectrum

Low-frequency Measures

FFT

PSD   0.01 0.08   Hz

PSD   0.01 0.08   Hz

PSD   0.01 0.08   Hz

ALFF

$$A_{low\,freq} = -\sum_{f_{min}}^{f_{max}} \sqrt{\overline{P(f)df}}$$

$$0.01\,Hz < f < 0.08\,Hz$$

lfSE

$$\hat{S}_{spectral} = \frac{-\sum_{f=0}^{Fs/2} P(f)\log_2 P(f)}{\log_2(num_{freq})}$$

$$0.01\,Hz < f < 0.08\,Hz$$

Covariate Regression
+
Site Harmonization

all nodes $\quad \bar{X} \quad$ network member nodes

B. Whole-brain & Network Levels

Group differences

Symptom associations