Title: Effects of ketamine and esketamine on death, suicidal behaviour, and suicidal ideation in psychiatric disorders: A systematic review and meta-analysis

Running head: Ketamine/esketamine and suicidality

Martin Plöderl [1] (0000-0003-4659-9314), Ruth Cooper [2] (0000-0002-9735-4731), Tom Walker [3] (0009-0004-0886-5428), Vibha Shah [4], Mark Horowitz [5] (0000-0003-1318-2029), Constantin Volkmann [6] (0000-0002-5359-4338), Joanna Moncrieff [7] (0000-0003-1214-6974)

Roles and Affiliations

[1] Clinical Psychologist, Center for Inpatient Psychotherapy and Crisis Intervention, University Clinic for Psychiatry/Psychotherapy/Psychosomatics, Paracelsus Medical University, 5020 Salzburg, Austria. m.ploederl@salk.at

[2] Research Associate, Department of Health Service & Population Research, David Goldberg Centre, 18 De Crespigny Park, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK, SE5 8AF.

[3] Senior Trainee in Psychiatry, West London NHS Foundation Trust, UB2 4SA London, UK. thomas.walker6@nhs.net

[4] Consultant Psychiatrist, North East London Foundation Trust, Goodmayes Hospital, 157 Barley Lane, Goodmayes, Ilford IG3 8XJ, UK. vibha.shah@nelft.nhs.uk

[5] Clinical Research Fellow in Psychiatry, Research and Development Department, North East London Foundation Trust, Goodmayes Hospital, Ilford IG3 8XJ, UK. mark.horowitz@nelft.nhs.uk

[6] Consultant Psychiatrist, Department of Psychiatry, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, 10117 Berlin, Germany. 8

[7] Professor of critical and social psychiatry, Division of Psychiatry, University College London, London WC1E 6BT, and North East London Foundation Trust, Goodmayes Hospital, Essex, IG3 8XJ. j.moncrieff@ucl.ac.uk

Correspondence to: Martin Plöderl, Christian Doppler Clinic, Paracelsus Medical University Salzburg, Ignaz-Harrer-Str. 79, 5020 Salzburg, Austria. m.ploederl@salk.at

ABSTRACT

Background: Ketamine and esketamine have been claimed to possess anti-suicidal effects and potentially to transform suicide prevention. This study provides an updated overview of evidence from clinical trials.

Methods: We conducted a systematic review and meta-analysis to establish whether ketamine or esketamine reduce death, suicides, suicide attempts or suicidal ideation, compared to active or inert placebo among people with psychiatric disorders. We searched EMBASE, PubMed and PsycINFO until 09.01.25 for randomised controlled trials which compared the effect of ketamine or esketamine with placebo for the treatment of psychiatric disorders. Risk of bias was assessed using the Cochrane Risk of Bias Tool. Data were synthesised with meta-analysis, including methods for double-zero events.

Results: We included 66 trials with a total of 5103 participants. Rates of suicidal behavior were 0.91% for ketamine/esketamine and 1.23% for placebo, with the 95% confidence/credible intervals including the null-effect, OR = 0.86 [0.46 to 1.54] (Bayesian Analysis) and with low heterogeneity. There were 21 (0.77%) suicide attempts, 4 (0.14%) suicides and 6 (0.21%) deaths with ketamine/esketamine compared to 26 (1.18%), 1 (0.05%) and 2 (0.09%) on placebo. Ketamine/esketamine significantly reduced suicide ideation up to 2 weeks (standardized mean differences [SMD] at 12h to 24h of -0.35 [-0.51 to -0.19], I^2 = 12%), but had small effects (SMD < 0.20, I^2 ≤ 25%) thereafter. Quality assessment revealed unreliable blinding and selective reporting.

Discussion: There is insufficient evidence for a preventive effect of ketamine/esketamine for suicidal behavior. The observed immediate but short-term effect on suicide ideation may be overestimated due to unblinding bias. Our review is the most comprehensive on suicidality, however, more evidence is needed to draw conclusions on suicidal behavior.

Other: No specific funding was involved. The protocol was registered with PROSPERO (CRD42023364156).

KEY MESSAGES

What is already known on this topic

- Ketamine and esketamine have been considered to tranform suicide prevention with evidence from multiple studies suggesting they may rapidly alleviate depression and suicide ideation.

- There have been numerous calls for studies to investigate if these findings translate to suicidal behavior. With the many new studies that have emerged only recently, such a review is overdue, together with an update on suicide ideation.

What this study adds

- Despite including studies up to January 2025, evidence on whether ketamine/esketamine can effectively reduce suicidal behavior remains inconclusive.

- A small to medium short-term effect on suicide ideation within the first three days was observed for ketamine but not for esketamine and unblinding bias is likely involved.

- Repeated doses did not show superiority over single administrations in reducing suicide ideation.

How this study might affect research, practice, or policy

- More studies are needed to judge if ketamine/esketamine can reduce suicidal behavior, and how unblinding effects can explain the short-term effect on suicide ideation.

- The practice of repeated dosings should be questioned and needs further investigation.

INTRODUCTION

Suicide is a major public health concern across the world. Overall, 1 in 100 people die by suicide and among people aged 15-29 it is the fourth leading cause of death worldwide [1]. In the UK suicide is the most common cause of death among both men and women aged 20 to 34 [2]. Despite a worldwide decline in the rate of suicide since 2000, rates are increasing in the US and some European countries, including the UK [2,3]. A broad range of psychiatric disorders is associated with in increase of suicides and suicide attempts [4].

Ketamine and its S-enantiomer, esketamine, are increasingly used for the treatment of treatment-resistant depression and also investigated for a broader range of mental health problems such as bipolar depression, postpartum depression, substance use disorders and anxiety disorders (Table 1). Furthermore, they have been suggested to have the potential to offer rapid relief from suicidal thoughts and to help prevent suicide [5,6] and Esketamine was licensed in the US for people with treatment resistant depression and major depressive disorder with suicidal thoughts or actions [7]. Previous summaries of the literature have claimed promising results for both agents [8–10], suggesting that esketamine is "the only FDA-approved antidepressant that works in hours not weeks – thus potentially transforming treatment of suicidal patients" [8]. There is also some evidence that ketamine reduces suicide ideation independent of depression symptoms [11]. On the other hand, post marketing data suggests that esketamine may be associated with an increase in suicidal ideation and suicides [12] and chronic, recreational ketamine misuse is associated with depression [13–15]. While previous systematic and non-systematic reviews have generally suggested that ketamine and esketamine reduce suicidal ideation and behaviour, at least short-term [16–19], they also highlight methodological limitations of the research and a lack of data on medium or long-term effects [17]. However, these limitations have not been explored in detail, and no existing reviews have considered the outcome of deaths by suicide or suicide attempts, even though reducing suicide is the ultimate aim of this intervention. Moreover, there is the repeated request for using harder outcomes in psychiatric research, such as suicide attempts or suicide, because softer outcomes (reduction in symptom scores) are more prone to bias [20]. Reviews on suicidal behavior are lacking partly due to the difficulty of conducting meta-analysis of rare events, but statistical methods to manage this situation have been developed recently [21,22]. Furthermore, existing reviews, even those of otherwise good quality, had significant limitations, for example, mixing outcomes at different time-points, or errors in data extraction [18,23,24]. Finally, many studies appeared recently and an update is required.

In the current systematic review and meta-analysis we aimed to answer the following question: does ketamine or esketamine reduce suicidal behavior (suicides, suicide attempts), deaths of any cause (as additional outcome because suicidal intent is sometimes unknown), and suicidal ideation for people with primary psychiatric disorders? We examined short-term and longer-term outcomes. We also explored methodological limitations of the existing studies.

METHODS

This systematic review and meta-analysis was conducted following guidance from the Preferred Reporting Items for Systematic Reviews and Meta- Analyses (PRISMA) guidelines [25]; see Supplement for the PRISMA Checklist.

Eligibility criteria

All randomised controlled trials (RCTs), comparing ketamine or esketamine with inert placebo or active placebo in individuals with a primary diagnosis of any major psychiatric disorder, were eligible for inclusion in this review. Data from the first period of cross-over trials were also included (to protect against the carry-over effect). Trials comparing ketamine or esketamine in combination with an antidepressant, psychotherapy or other intervention were included as long as other interventions were equally available to those in the control group. Studies were excluded if the drugs under investigation were intended as an analgesic and/or anaesthetic agent. In addition, although not specified in the protocol due to being unforeseen at the time, studies were excluded if data on suicidality were not available despite contacting the authors.

Information sources and search strategy

We searched EMBASE, PubMed and PsycINFO from inception until 01.10.2022 using the strategy outlined in Table S1. The search was updated on 09.01.25. We had no language restrictions. We also searched citations of included studies and previous reviews.

Selection process

Reviewers (TW, VS, MP and JM) screened titles and abstracts and subsequently full texts. Inclusion criteria for selected studies were double checked by RC, JM, and MP. Discrepancies were discussed and resolved within the team.

Data collection process

All authors extracted summary statistics from an assigned portion of selected studies using a standardised form. Uncertainties were resolved by discussion. In cases of missing data or areas of uncertainty, the corresponding author of the paper was contacted, and we looked up the data on trials registers using the study registration number. Summary statistics from Janssen's esketamine trials were calculated based on individual patient data accessed via the Yale Open University Data Access (YODA) project. All data used in this review were double checked by MP and RC.

Data items and outcome measures

We extracted the following data: trial design, length of follow-up, participant number and characteristics, intervention, control, and Conflicts of Interest (COIs).

Outcomes consisted of death by any cause, suicides, suicide attempts and suicidal ideation. We extracted outcome data at seven time-points, similar to a previous review [18]: within 4 hours, >4 to ⩽12 hours, >12 to ⩽24 hours, >24 to ⩽72 hours, >72 hours to ⩽2 weeks, >2 to ⩽4 weeks, and >4 weeks. If there were more than one measure within a specific time frame, we used the one closest to the end of the time frame.

Suicidal behavior was made up of a composite of suicides and suicide attempts. We checked the trial registers for suicidal behavior and mortality, and we assumed that no such events occurred when the paper reported that "no serious adverse events occurred" or when tabulated adverse events did not include suicidal behavior/mortality.

For suicidal ideation, we used results based on assessments with suicide-specific scales such as the Beck Scale for Suicidal Ideation (BSI) or the Columbia Suicide Severity Rating Scale (C-SSRS). If this was not available, we used the suicidal ideation item of a depression rating scale in this hierarchy: the authors designated primary outcome and if this was not specified the Montgomery-Åsberg Depression Rating Scale (MADRS), followed by the Beck Depression Inventory (BDI), Hamilton Depression Rating Scale (HDRS), and other scales such as the Quick Inventory of Depressive Symptomatology (QIDS).

When examining the trials, it became clear that there were distinct "treatment phases" and "follow-up" phases, although these were not defined consistently in the original studies. Results from follow-up periods were included as long as randomization was maintained, in order to include data

from the same time point in studies with different treatment durations. Data from non-randomized open-label treatment phases were excluded.

Quality assessment

The risk of bias in the included studies was assessed using the Cochrane Risk of Bias tool [26]. This measure rates studies as being at high, low or unclear risk of bias according to various domains. Criteria for the rating of each domain were established by consensus within the research team in accordance with the Cochrane guidelines. Quality assessments were carried out by MP, MH, JM, SA, RC, TW, and VS and each was independently checked by JM and VS.

Synthesis methods

We reported results for suicide attempts, suicides, and mortality according to whether they occurred in the treatment phase, the follow-up phase, and both phases combined, since further information about timing was generally not available. We included studies without any events to avoid biased results, following recent recommendations and different random effects meta-analytic approaches suggested by Xu et al. (2021) These authors also recommended using more than one method to confirm that results are not dependent on a specific method. Therefore, as outlined in our protocol, we used a random effects two-stage exact method [22] and a one-stage Bayesian methods as in a previous publication [28] which was based on a binomial-normal hierarchical model with a weak prior (a normal distribution with mean = 0 and variance = 2.82, with a limitation to delta=250, meaning that odds ratios larger than 250 or smaller than 1/250 are considered unlikely, and a half-normal prior for the heterogeneity tau=0.5). We also used a Bayesian model with a more informative prior (delta = 15). In addition, we used one-stage frequentist beta-binomial and GLMM methods [29] and the following two stage models: Peto's method, Mantel-Haenszel methods with continuity corrections, and two methods based on risk differences (Mantel-Haenszel and arcsine transformation methods). All other methods were based on odds-ratios as measures of effect-size. Heterogeneity was assessed with I^2 and τ^2 if the meta-analytic model allowed for this (with R's default parameters). We combined drug-arms for studies with different doses as recommended [30].

For suicide ideation as outcome, we ran random effects meta-analysis using R's "meta" package, with Hedges g as measure of effect size, I^2 and τ^2 as measures of heterogeneity (using Sidik & Jonkman's method to estimate heterogeneity). Meta-analyses were based on drug-placebo differences in suicide ideation at specific time-points and not on differences in change from baseline, except for one study where only change-scores were available. Publication bias was explored with Egger's

regression test. For the few studies which only reported medians we used the procedure suggested by Wan et al. to calculate means and standard deviations [31]. For one study [32] no information about the standard deviations were available and we imputed these with those from other studies with the same assessment instruments, as recommended [30]. We combined results from multiple arms with different doses following the recommendation by Cochrane [30] or, if possible, by using individual patient data. We planned to conduct various pre-specified subgroup or subset analyses, where sufficient data permitted, namely: inclusion or exclusion of people at significant risk of suicide, different primary psychiatric disorders (including substance misuse), esketamine or ketamine, method of administration (e.g., intravenous, oral), sample size, inert or active placebo, different trial designs (RCT or single blinded trial or open label study) and clinician or self-report measures of suicidal ideation. We also conducted a subgroup analysis of single versus repeated doses which was not included in the protocol. We did not distinguish between the treatment-phase and follow-up phase in the analysis for suicide ideation because this was the only way that longer-term effects from single-administrations could be compared to those of repeated administrations.

We used R 4.3.0 for all analysis [33]. The protocol was developed and registered prior to data extraction (PROSPERO CRD42023364156).

Patient and public involvement

No patient and public involvement happened at the stage of planning the analysis. However, one member of the review-team has experiences of mental health difficulties and treatments. During the data-analysis, a representative of Austria's patient groups who is also a member of the expert panel in Austria's national suicide prevention program SUPRA (Suicide Prevention Austria) was informed about the aims and methods of the review and asked to provide feedback after discussing the study with other representatives of patient groups. This did not result in practical suggestions about the study aims or methodology. Patient representatives felt it was important that the review fulfils quality criteria and will be publicly available. They expressed concerns about the validity of the assessment of suicidality and about potential addiction or dependence problems with longer-term use of ketamine or esketamine. We include these points in the discussion section and intended to publish this review open access.

RESULTS

Study selection and characteristics

The search identified 2668 titles, of which 164 and additional 18 titles via reference checking were subjected to full-text screening (Figure 1). A total of 117 studies were excluded following full-text screening (Table S2). A total of 66 studies were identified that met eligibility criteria, including 50 involving ketamine, 15 esketamine, and 1 arketamine. Twelve of the ketamine trials were cross-over trials and the rest were parallel group trials. A single dose was used in 35 trials (31 ketamine trials). In repeated dose trials, doses were administered between weekly and two times daily and treatment lasted between three days and ten weeks. Follow-up lasted between 24 hours to 12 months. The trials involved 5103 participants in total. Further characteristics are detailed in Table 1.

Risk of bias in studies

Full quality ratings are shown in Figure S2. These revealed that patients and study personnel were rarely likely to be adequately blinded. Only 15 studies tested the integrity of the double blind, of which 12 showed evidence of unblinding among participants and or assessors (Table S3). These included seven of the eight studies that employed an active placebo agent such as midazolam or remifentanil, with some of these finding rates of correct guesses by participants and assessors between 80 and 100% [34,35]. Only three out of these 15 studies were considered adequately blinded, two of which involved administering ketamine along with an anaesthetic [36,37]. Furthermore, we rated 20 (30%) of studies as high risk of bias due to selective reporting of outcomes, and one trial publication did not report suicidality as an outcome despite this being planned (Dwyer. et al. 2021). Overall 56 (85%) trials were rated as being at high risk of bias in some domain.

Suicidal Behavior and Deaths

For suicides and suicide attempt combined, and for both the treatment and follow-up phase combined, there were 25 (0.91%) incidents in 2742 patients randomized to ketamine or esketamine compared to 27 (1.23%) incidents in 2201 patients randomized to placebo. The drug-placebo differences were not statistically significant in any of the meta-analyses. For example, the credible interval in the Bayesian meta-analysis with weak priors (which we use for reporting throughout this section) overlapped with the null-effect (OR = 1), OR = 0.86 [0.45 to 1.56] (Table 2). In the treatment-period, there were 9 (0.33%) incidents with ketamine/esketamine and 9 (0.41%) with placebo, and for the follow-up period there were 16 (1.06%) and 18 (1.42%) incidents, respectively. None of the meta-analytic results were statistically significant (Table 2, forest-plots with results for

individual studies in Supplement). The number of deaths of any cause in combined treatment and follow-up phases was 6 (0.21%) in ketamine or esketamine and 2 (0.09%) in the placebo groups and the credible interval in the meta-analysis overlapped with the null-effect, OR = 1.64 [0.5 to 5.27] (Table S13). For suicides, there were 4 (0.14%) incidences with ketamine or esketamine compared to 1 with placebo (0.05%), OR = 1.53 [0.47 to 5.55] (Table S10). For suicide attempts, there were 21 (0.77%) incidences with ketamine or esketamine, compared to 26 (1.18%) with placebo and again the difference overlapped with the null-effect, OR = 0.75 [0.39 to 1.38] (Table S7). Results were comparable with other meta-analytic models and for different treatment phases (Tables S4-S15). None of the drug-placebo differences for deaths, suicides, and suicide attempts were statistically significant. There was no indication of reporting bias in any of the analysis for suicidal behaviour and deaths.

Suicide Ideation

Suicidal ideation was statistically significantly reduced among people allocated to ketamine/esketamine compared to placebo, with a small to medium effect size up to 24 hours after randomisation. At subsequent follow-ups differences were small (SMD < 0.20) and by 2 weeks follow-up they were not statistically significant (Table 3, Figure 2). An end of treatment phase analysis (combining results from studies of all durations) showed a significant effect, SMD = -0.24 [-0.35 to -0.13]. Heterogeneity was only statistically significant for results in the first 4 hours. There was no indication of publication bias in any of the analysis for suicidal ideation.

Subgroup analysis

Deaths, suicides and suicide attempts were too rare to do subgroup analyses. For suicidal ideation, subgroup analysis showed that the effects of ketamine were statistically significant up until 2 weeks, whereas the effects for esketamine were close to zero and not statistically significant after 24 hours (Table 4). The effects for esketamine were consistently smaller and statistically significantly smaller for >24h to 72, and >2weeks to 4 weeks, and end of treatment. There were no significant differences between trials where suicidality was an inclusion criterion (vs. suicidality as exclusion criteria or unspecified) except for the time >24h – 72h and end of treatment, where trials with suicidal patients had significantly larger effect than other trials. There were no significant differences between trials with active placebos (defined as an active drug), or inert placebos (including those with a bittering agent) except at the end of treatment where trials with active placebos showed larger effects. There were no significant differences between trials where suicide ideation was assessed by a clinician

compared with by patients. There was insufficient data for comparing different diagnoses (most studies involved MDD), trial design, or low risk of bias (or post-hoc, exploratory subgroup analysis for age-groups, doses, etc.). Repeated administrations were numerically comparable and not statistically significantly superior to a single dose at any time point (we ran analysis after 72h since repeated dosing usually starts after this time point).

DISCUSSION

The evidence that ketamine or esketamine prevent suicidal behavior or mortality is inconclusive. Numerically, there were fewer suicide attempts and more suicides and deaths in the drug arms than in the placebo arms, but there is much uncertainty in the estimations of differences. Medium-sized protective effects on suicide ideation were only observed up to three days, and these were generally smaller for esketamine than ketamine. Repeated dosings were not superior to single doses.

In general, drawing conclusions is hampered by the lack of long-term data, insufficient statistical power, and selective reporting. Another problem is the acute psychoactive effects of ketamine/esketamine. These may influence suicidal ideation and mood in their own right, and may also undermine the double blind design. This was confirmed by the few trials which tested blinding and found that it was mostly not successful, even for active placebos [39,40]. In two trials in which ketamine was given during an anaesthetic procedure and blinding was successful, the drug-placebo differences were close to zero (see subgroup analysis in Supplement). In one of these, people who guessed they had had ketamine showed substantially lower symptoms at follow-up than people who guessed they had had placebo (p=0.001), despite there being no effect of the actual substance [37].

Furthermore, it remains unknown if the evidence base for ketamine/esketamine is affected by problems seen in antidepressant research where suicidal behavior was sometimes reclassified as emotional worsening or "drop-out" [41,42]. In addition, it is often not clear if suicidal behavior was assessed by blinded raters, which is known to result in lower efficacy compared to non-blinded raters [43]. Whereas assessment of suicide ideation was often part of a separate and systematic assessment process, suicidal behavior was often only identified during the assessment of spontaneously reported adverse events. Consequently, our risk of bias assessment provides a conservative estimation of the likely biases due to rater blinding.

Esketamine is approved by the FDA and increasingly used but appears less effective than ketamine. This was also reported in other reviews with depression symptoms as an outcome [17,23,44,45] and may reflect greater amplified placebo effects of ketamine, which is usually administered intravenously. More head-to-head comparisons are needed for clarification.

A challenge to the clinical practice of repeated dosings for maintaining efficacy [23] is our finding that this is not superior to single doses. This has not been systematically explored in placebo controlled trials, to our knowledge, but is in line with a recent finding of controlled and uncontrolled trials based on pre-post differences [46]. The diminishing effect could be explained by decreasing expectancy effects over time or tolerance effects. This result challenges claims that ketamine or esketamine work by stimulating neurogeneration or other processes that evolve over time.

Although numbers of suicides and deaths of any cause were small, the fact that they were higher in the ketamine/esketamine arms than in the placebo arms may represent a signal of concern, given that post-marketing surveillance data on esketamine suggests it may increase the incidence of suicidal ideation and behaviour [12]. Of note, two additional suicides and one other death occurred in drug-arms of enrichment or open label phases of studies that were excluded from our analysis [47,48]. However, suicide attempts were numerically lower in the drug arms. Concerns about potential addiction problems from longer-term use were also pointed out by our patient representatives and are well known [49,50].

Strengths of this review include the inclusion of suicidal behavior and mortality and statistical methods to handle these rare events, thus avoiding biases resulting from excluding trials with zero events. Previous reviews were either outdated or did not report on suicidal behavior as outcome, especially not on suicides and deaths of any cause [16–18]. Furthermore, in contrast to some previous reviews, results for suicide ideation were aggregated for different time points and the review is up to date by including studies up to January 2025. The larger number of included studies replicated the findings of a short-term effect on suicide ideation and the weaker effect for esketamine compared to ketamines [17,19,23,45], but is more precise. Limitations include potential biases resulting from our assumption that no suicidal behavior occurred when none were mentioned in the reported adverse events. Some authors did not respond to data requests and this may introduce bias if non-response is systematically related to published outcomes. Additional limitations also affecting other reviews include the use of means and standard deviations when suicide ideation has a skewed distribution or when suicide ideation was measured on a Likert-scale with few response

options, and that studies did not report results separately by sex/gender, and that follow-up time-frames substantially varied (and that we had no further information about the exact timing of events to be able to restrict time-frames).

In conclusion, there is insufficient data to ascertain whether ketamine or esketamine reduce suicide and suicide attempts. Short-term effects for ketamine on suicide ideation are detected, but these are not sustained and may be the product of acute psychoactive effects or unblinding.

ACKNOWLEDGEMENTS

We thank the authors who responded to our requests for data. We also thank Janssen and Yale for make patient-level data available via the YODA platform (YODA statement: "This study, carried out under YODA Project [2024-0612], used data obtained from the Yale University Open Data Access Project, which has an agreement with JANSSEN RESEARCH & DEVELOPMENT, L.L.C.. The interpretation and reporting of research using this data are solely the responsibility of the authors and does not necessarily represent the official views of the Yale University Open Data Access Project or JANSSEN RESEARCH & DEVELOPMENT, L.L.C.."). We thank Silvi Mühringer, patient representative of SUPRA Austria for discussions and suggestions for our study. We thank Simone Amendola for assistance in designing the study and data-extraction.

Research ethics and consent

This is a systematic review of already published studies.

Contributor and guarantor information

Protocol development: MP, RC, TW, MH, CV, JM; literature search: TC, VS, MP, JM; data extraction, data management, data-cleaning: MP, RC, TW, VS, MH, CV, JM, statistical analysis: MP, drafting: MP, RC, JM; reviewing and revising: MP, RC, TW, VS, MH, CV, JM. Responsible for overall content and guarantor: MP and JM. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Copyright/licence for publication

*T*he Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display

and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above.

Competing interests

All authors have completed the ICMJE uniform disclosure form at http://www.icmje.org/disclosure-of-interest/ and declare: CV, MP, RC, TW, VS no support from any commercial organisation for the submitted work and have no other conflicts of interest to declare; JM has received royalties for books about psychiatric drugs; MH received support from a clinical research fellowship at NELFT (NHS), royalties from the Maudsley Deprescribing Guidelines, and is a co-founder of Outro Health, a digital clinic in the US where he receives consulting fees.

Data sharing

The data and statistical code are available on the OSF https://osf.io/xzj7k/

Caption to Figure 1

Figure 1. PRISMA flowchart.

Caption to Figure 2

Figure 2. Meta-analytic results for effects of ketamine and esketamine on suicide ideation at different time-points. The y-axis refers to the standardized mean-difference. Negative values correspond to increased reduction in suicide ideation in the drug arms compared to the placebo arms. The horizontal dashed lines are two thresholds for statistical significance at SMD = -0.3 and SMD = -0.5.

Table 1. Study characteristics

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Abbar 2022 [44], France, NCT02299440	156 (73/83)	Adults with Bipolar disorder, depression, other disorders; current suicidal ideation	Double blind RCT	**TR**: 2 days **TR+FU**: 6 weeks **TR+FU for analysis**: 6 weeks	IV Ketamine (0.5mg/kg), 2 doses given at baseline and 24 hours.	Placebo: saline	BSS – clinician	Yes, directly Timing of suicide attempts clarified with author	Yes
Abdallah 2022 [51], USA, NCT02655692	158 (104/54)	Adult (18 to 70) veterans; PTSD; no suicidality meriting crisis intervention	Double blind RCT	**TR**: 4 weeks **TR+FU**: 8 weeks **TR+FU for analysis**: only 4-week TR because TR-nonresponders were offered open label ketamine in FU	IV Ketamine (0.2 and 0.5mg/kg), twice weekly, 4 weeks	Placebo: saline	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported on trial register	Yes
Ahmed 2023 [52](30) Egypt NCT04101474	36 (18/18)	Adults (18 to 65 years); MDD; current suicidal risk	Double blind RCT	**TR**: 2 weeks **TR+FU**: 3 weeks **TR+FU for analysis**: 3 weeks	IV Ketamine (0.5mg/kg), 2 doses over 2 weeks	Placebo: saline	SPS, using the post-dose assessments	Yes, directly Not reported but author confirmed that no events occurred	No
Anderson 2017, [36,53] UK, ISRCTN 14689382	79 (40/39)	Adults with Uni- or bipolar depression; suicidality not specified in eligibility criteria	Double blind RCT	**TR**: varied – max 10 weeks. **TR+FU**: max. 26 weeks. **TR+FU for analysis**: max. 26 weeks	IV Ketamine (0.5mg/kg) as an adjunct to ECT anaesthetic + ECT course. Twice weekly, average 11 ECT treatments (in both arms)	Placebo: saline + ECT course	MADRS-SI, data from Witt et al. 2019 who requested data from authors	Yes, directly Timing of suicide attempts assumed to coincide with reported detainments in EUDRA information	No
Burger 2016 [32], USA, none reported	10 (3/7)	Adults (18 to 65 years, on active military duty) with depression and suicidality; current suicidal ideation	Double blind RCT	**TR**: 24 hours (single dose) **TR+FU**: max. 3 weeks (max. 1 week inpatient treatment with 2 weeks follow up **TR+FU for analysis**: max 3 weeks	IV Ketamine (0.2 mg/kg), 1 dose	Placebo: saline	BSS – patient, M's from figure, SD's imputed with other studies using patient-BSS	Yes, indirectly No events in AE reported in paper	Not reported
Canuso 2018, [54], USA, NCT02133001	68 (36/32)	Adults (19 to 64 years) with MDD; current suicidal ideation	Double blind RCT	**TR** : 25 days **TR+FU**: 81 days **TR+FU for analysis**: 81 days	Intranasal Esketamine, 84mg (max total dose), twice weekly for 4 weeks	Placebo with bittering agent[a]	SIBAT Module 8, data via YODA	Yes, directly	Yes

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Chen 2023, [55], USA + China, NCT03434041	252 (126/126)	Adults (18 to 64 years) with MDD, no current suicide ideation with intent	Double blind RCT	**TR**: 4 weeks **TR+FU**: 12 weeks **TR+FU for analysis**: only 4-week TR because in some patients moved to other trial afterwards	Intranasal Esketamine, 56mg up to 84mg, twice weekly for 4 weeks	Placebo	C-SSRS, data via YODA	Yes, directly	Yes
Colla 2024 [56], Switzerland SNCTP Nr. 000002938	30 (20/10)	Adults (18 to 70 years); moderate to severe depression; exclude suicidality	Double blind RCT	**TR**: 2 weeks **TR+FU**: 2 weeks (no FU) **TR+FU for analysis**: 2 weeks	Ketamine, 2 dose groups: 160mg or 240mg per day for 14 days	Placebo	SSF	Yes, indirectly No events in AE reported in paper	Yes
Dakwar 2019 [57], USA, NCT01535937	55 (27/28)	Adults under age 70; SUD (Cocaine); nothing mentioned regarding suicidality	Double blind RCT	**TR**: 24h (single dose) plus 5 week mindfulness based therapy **TR+FU**: 31 weeks **TR+FU for analysis**: 31 weeks	Intravenous Ketamine 0.5mg/kg, 1 dose + mindfulness behavioral modification	Active control: Midazolam, 0.025mg/kg + mindfulness behavioral modification	Not assessed	Yes, indirectly No events in AE reported in the trial register	Yes
Dakwar 2020 [58], USA, NCT02539511	40 (17/23)	Adults under age 70; SUD (Alcohol); excluded when current suicide risk or past year suicide attempt (according to protocol)	Double blind RCT	**TR**: 24h (single dose) in second week of 5 week psychological treatment **TR+FU**: 31 weeks **TR+FU for analysis**: 30 weeks	Intravenous Ketamine 0.71mg/kg, 1 dose + motivational enhancement therapy	Active control: Midazolam, 0.025mg/kg + motivational enhancement therapy	Not assessed	Yes, indirectly No events in AE reported in the trial register	Yes
Daly 2018 [59] (36) USA, Belgium NCT01998958	Panel A: 67 (34/33) (phase 1) Panel B: 41 (20/21) (phase 1)	Adults (20 to 64 years) with MDD and TRD; excluded current suicidal ideation	Double blind, doubly randomised, delayed start RCT Phase 1: participants randomised. Phase 2: placebo group with moderate/severe symptoms	**TR**: 7 days (phase 1), 7 days (phase 2). Open label phase: 59 days **TR+FU**: 130 days **TR+FU for analysis**: 7 days (phase 1) – pre-crossover and pre-open label phase (randomization only upheld in phase 1)	Panel A: Intranasal Esketamine, 3 dose groups: 28, 56, or 84mg twice weekly for each of the 2 weeks (double blind phase) Panel B: Intranasal Esketamine, 2 dose groups: 14 or 56mg twice weekly	Placebo with bittering agent[a]	C-SSRS, data via YODA	Yes, directly	Yes

Study, country, registration number	Total N randomised (interventio n/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
			re-randomise d to intervention.						
Domany 2019 [60], Israel NCT02037503	41 (22/19)	Adults (18 to 75 years); MDD; suicidality: not reported	Double blind RCT	**TR**: 21 days **TR+FU**: 28 days **TR+FU for analysis**: 28 days	Oral Ketamine (1mg/kg) three times a week	Placebo: saline	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in publication	Not reported
Domany 2020 [61], USA, NCT01887990	18 (9/9)	Adults (18 to 65) with MDD, bipolar depression, depression, dysthymia; current suicidal ideation	Double blind RCT	**TR**: 24 hours (single dose) **TR+FU**: 14 days **TR+FU for analysis**: 14 days	IV Ketamine (0.2mg/kg), 1 dose	Placebo: saline	BSS – patient, from figure	Yes, indirectly No events in AE reported in trial register	No
Domany 2021 [62], USA NCT02183272	30 (15/15)	Adults (18 to 65 years); eligible participants – required hospitalisation due to suicidal risk	Double blind RCT	**TR**: 24 hours (single dose) **TR+FU**: 28 days **TR+FU for analysis**: 28 days	IN Ketamine, 40mg, single dose	Placebo: saline	BSS – patient, from figure	No Only specific AEs reported, no information on the trial register, no response from author	Not reported
Duek 2023 [63] USA NCT02727998	28 (14/14)	Adults; PTSD; current suicide risk excluded (protocol: ongoing serious suicidal risk as assessed by evaluating investigator or by scoring 5 or more on item-10 of the MADRS)	Double blind RCT	**Length of treatment phase**: 24 hours (single dose) **End of treatment FU**: 90 days **Total trial duration**: 90 days **FU used in analysis**: 90 days	IV Ketamine (0.5mg/kg) Single dose + Psychoptherapy (Retrieve and reprocess exposure therapy)	Active control: Midazolam + Psychopthera py (Retrieve and reprocess exposure therapy)	BDI-SI, raw-data available online	Yes, indirectly No events in AE reported in trial register	Yes
Dwyer 2021 [38] USA NCT02579928	17 (6/11)	Adolescents (13 to 17 years); MDD; excluded active suicidal ideation	Double blind crossover RCT	**TR**: 24 hours (single dose), cross-over 2 weeks later **TR+FU**: 28 days (both cross-over phases) **TR+FU for analysis**: 14 days	IV Ketamine (0.5 mg/kg) single dose	Active control: Midazolam	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in trial register	Yes

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Fava 2020 [35] USA, NCT01920555	99 (80/19)	Adults with MDD and TRD (18 to 70 years); suicidality not specified in eligibility criteria	Double blind RCT	**TR**: 24 hours (single dose) **TR+FU**: 30 days **TR+FU for analysis**: 30 days	IV Ketamine, 1 dose, 3 dose groups: 0.1, 0.2, 0.5, 1.0 mg/kg	Active control: Midazolam	MADRS-SI, data unavailable	Yes, directly	Yes
Feder 2014 [64], USA, NCT00749203	41 (22/19)	Adults (18 to 55 years) with chronic PTSD, excluded when active suicidality	Double blind crossover RCT	**TR**: 24 hours (single dose), cross-over 2 weeks later **TR+FU***: 4 weeks (both cross-over phases) **TR+FU for analysis**: 2 weeks	IV Ketamine (0.5 mg/kg) single dose	Active control: Midazolam (0.045mg/kg)	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in trial register	Yes
Feder 2021 [65], USA, NCT02397889	51 (15/15)	Adults (18 to 70 years) with chronic PTSD, excluded when active suicidal ideation	Double blind RCT	**TR**: 2 weeks **TR+FU**: max 7.5 months **TR+FU for analysis**: only 2-week TR because only responders were followed up	IV Ketamine (0.5 mg/kg) thrice weekly for two weeks	Active control: Midazolam (0.045mg/kg)	C-SSRS, data unavailable	Yes, directly Paper reports that there was no suicidal behavior, and no deaths in AE in trial register	Yes
Fedgchin 2019 [66], America, Europe NCT02417064	346 (233/113)	Adults (18 to 64 years) with MDD and TRD; excluded suicidality (some ideation possible)	Double blind RCT	**TR**: 4 weeks **TR+FU**: 28 weeks **TR+FU for analysis**: only 4-week TR because responders could enter other trial	Intranasal Esketamine 56, 84mg twice weekly for 4 weeks	Placebo with bittering agent[a]	C-SSRS, data via YODA	Yes, directly	Yes
Fineberg 2023 [39], USA NCT03395314	22 (10/12)	Borderline Personality Disorder; excluded current acute suicide risk	Double blind RCT	**TR**: 24 hours (single dose) **TR+FU**: 28 days **TR+FU for analysis**: 28 days	IV Ketamine (0.5 mg/kg), single dose	Active control: Midazolam	BSS – clinician, data from figure in appendix	Yes, indirectly No events in AE reported in trial register	Yes
Fu 2020 [35], USA, Europe, Asia, South Africa, NCT03039192	226 (114/112)	Adults (18 to 64 years), MDD; current suicidal ideation	Double blind RCT	**TR**: 25 days **TR+FU**: 90 days **TR+FU for analysis**: 90 days	Intranasal Esketamine, 84mg, twice weekly for 4 weeks	Placebo with bittering agent[a]	CGI-SS-r, data via YODA	Yes, directly	Yes
George 2017 [67], Australia, NCT01441505	16 (12/4)	Older adults (≤60 years), MDD, Bipolar disorder; high suicide risk excluded.	Crossover RCT	**TR**: 5 weeks with repeated cross-over. Remitters could exit the treatment phase after 1 week. Open label phase: 8 weeks **TR+FU**: 65 weeks	Subcutaneous Ketamine, weekly over 5 weeks: 0.1, 0.2, 0.3, 0.4, 0.5 mg/kg (blinded treatment phase)	Active control: Midazolam, single dose randomly inserted in first 3	MADRS-SI, data from Witt et al. 2019 who obtained data from author	Yes, indirectly No events in AE reported in publication	Yes

Study, country, registration number	Total N randomised (interventio n/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
				TR+FU in analysis: only first week (before first cross-over)		treatment sessions			
Grabski 2022 [68], UK, NCT02649231	96 (48/48)	Adults (18 to 65 years); Alcohol use Disorder; excluded suicidal ideation	Double blind RCT	**TR:** 1 week-3 weeks **TR+FU:** 6 months from first treatment **TR+FU for analysis:** 26-28 weeks	2 groups: IV Ketamine (0.8 mg/kg) 3 times over max 3 weeks + psychological therapy (group 1) or + alcohol education (group 2)	Placebo: saline + psychological therapy (group 1) or + alcohol education (group 2)	BDI-SI, data provided by author	Yes, indirectly No events in AE reported in trial register	Yes
Glue 2024b [40] New Zealand ACTRN12619000 311156	29 (17/8 completers)	Adults (18 to 50 years); MDD; excluded significant suicidal ideation	Double blind crossover RCT	**TR:** 24 hours (single dose), first crossover after 1 week **TR+FU:** 3 weeks **TR+FU for analysis:** 1 week	IV Ketamine (2 groups: 0.5 or 1 mg/kg), single doses	Active control: fentanyl	MADRS-SI, data provided by author	Yes, indirectly No events in AE reported in publication	Yes
Grunebaum 2017 [69], USA, none reported	16 (7/9)	Adults (18 to 65 years), Bipolar Disorder and major depressive episode; current suicidal ideation	Double blind RCT with open label phase	**TR:** 24 hours (single dose) **TR+FU:** 6 weeks **TR+FU used for analysis: only** 24 hours TR because open label treatment with ketamine (single dose) for non-responders in the control group after 24 hours	IV Ketamine (0.5 mg/kg), single dose	Active control: Midazolam	BSS – clinician, data from Witt et al. 2019 but, in contrast to Witt, only up to 24h	Yes, directly	Yes
Grunebaum 2018 [70], USA, NCT01700829	82 (41/41)	Adults (18-65 years); MDD; current suicidal ideation	Double blind RCT with open label phase	**TR:** 24 hours (single dose) **TR+FU:** 6 weeks **TR+FU used for analysis: only** 24 hours TR because open label treatment with ketamine for non-remitters afterwards	IV Ketamine (0.5 mg/kg), single dose	Active control: Midazolam	BSS – clinician, data from Witt et al. 2019 but, in contrast to Witt, only up to 24h	Yes, directly Reported overdoses in Table S6 were suicide attempts, as confirmed by author	Yes
Hu 2016 [71], China, none reported	30 (15/15)	Adults (18-60 years); MDD; excluded if had suicide attempt during current episode	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 4 weeks **TR+FU used for analysis:** 4 weeks	IV ketamine (0.5mg/kg) single dose + escitalopram	Saline + escitalopram	QIDS-SR	No Only specific AEs reported, no information on the trial register, no response from author	Yes

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Ionescu 2019 [72], USA, NCT01582945	26 (13/13)	Adults (18 to 65 years); MDD; current suicidal ideation	Double blind RCT	**TR:** 3 weeks **TR+FU:** 15 weeks **TR+FU for analysis:** 15 weeks	IV ketamine (0.5mg/kg) 6 times to 2 infusions a week, for 3 weeks	Saline	C-SSRS, n's assuming no drop to out, exact time of infusion not given thus results for 72h unavailable	No Cannot be clarified with information in publication or trial register, author did not respond	Yes
Ionescu 2020 [73], Argentina, various European countries, Turkey, USA. NCT03097133	230 (115/115)	Adults (18 to 64 years); MDD; current suicidal ideation	Double blind RCT	**TR:** 25 days **TR+FU:** 90 days **TR+FU used for analysis:** 90 days	Intranasal esketamine 84mg twice weekly for 4 weeks	Placebo with bittering agent[a]	CGI-SS-r, data via YODA	Yes, directly	Yes
Jones 2024 [74] USA NCT03539887	29 (18/11)	Adults (21 to 60 years); MDD, bipolar depression, with/without comorbid alcohol use disorder; current suicidal ideation	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 48 hours **TR+FU for analysis:** 48 hours	Intranasal Ketamine, 50mg, single dose	Placebo: saline	BSS – clinician, data from figure	Yes, indirectly No events in AE reported in trial register	Yes
Lapidus 2014 [75], USA, NCT01304147	20 (10/10)	Adults (21 to 65 years), MDD chronic or recurrent without psychotic features, excluded when high risk of suicide	Double blind crossover RCT	**TR:** 24 hours (single dose), crossover after one week **TR+FU:** 14 days **TR+FU for analysis:** 7 days	Intranasal Ketamine, 50mg, single dose	Placebo: saline	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in trial register	Yes
Leal 2023 [76] Brazil UMIN000038347	10 (5/5)	Adults (18 to 65 years); MDD; suicidality not specified in eligibility criteria.	Double blind RCT with crossover	**TR:** 24 hours (single dose), crossover after one week **TR+FU:** 2 weeks **TR+FU for analysis:** 1 week	IV Arketamine (0.5 mg/kg) single dose	Placebo: saline	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in publication	Yes
Lii 2024 [37], USA, NCT03861988	40 (20/20)	Adults (18 to 80 years, mean age 51 years), MDD, excluded when	Triple blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 14 days **TR+FU for analysis:** 14 days	IV Ketamine (0.5 mg/kg), single dose	Placebo: saline	MADRS-SI, data from author	Yes, indirectly No suicidal behavior but one death reported in trial register	Yes

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
		"subject is at significant risk for suicidal behavior during participation in the study"							
Lijffijt 2022 [77] USA NCT02556606	33 (20/13)	Adults (>55 years); MDD; excluded when serious or imminent suicide risk according to publication on study methods	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 7 days, 28 days for responders at day 7 **TR+FU for analysis:** 7 days	IV Ketamine (3 groups: 0.1, 0.25, 0.5 mg/kg), single dose	Active control: Midazolam	C-SSRS, data unavailable	Yes, directly	Yes
Loo 2016 [78], Australia, none reported	15 (all participants took intervention /control dose)	Adults (18+ years); MDD; clinician interview; not specified	Double blind RCT with crossover	**TR:** 5 weeks, crossovers at least one week separated **TR+FU:** max. 31 weeks **TR+FU for analysis:** 1-2 week (afterwards those on midazolam may have been randomized to ketamine)	Ketamine (0.1, 0.2, 0.3, 0.4, 0.5 mg/kg) 3 routes of administration: IV, intramuscular, subcutaneous. Patients assigned to receive 1 route	Active control: Midazolam, single dose randomly inserted into the first 3 treatment sessions	MADRS-SI, data from Witt et al. 2019 who obtained data from authors	Yes, directly Confirmed by author that no suicidal behavior and deaths occurred	Yes
Loo 2023 [79], Australia, New Zealand ACTRN12616001096448	181 (89/92)	Adults (≥ 18 years); MDD; suicidality not specified in eligibility criteria.	Double blind RCT	**TR:** 4 weeks **TR+FU:** max. 26 weeks (for those who entered open label phase) **TR+FU for analysis:** only 4-week TR	Ketamine, subcutaneously, First cohort with fixed dose: 0.5mg/kg. Second cohort flexible dose: 0.5- 0.9 mg/kg, twice weekly for 4 weeks	Active control: Midazolam	MADRS-SI, data yet unavailable	Yes, directly timing of suicide attempts clarified with author (both were in the 4-week treatment phase)	Yes
Milak 2020 [80], USA, NCT01558063	43 (38/5)	Adults (18 to 59 years), MDD; excluded actively suicidal patients.	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 24 hours **TR+FU for analysis:** 24 hours	Intravenous Ketamine (0.1, 0.2, 0.3, 0.4, or 0.5 mg/kg) single dose	Placebo	HDRS-SI, data unavailable	Yes, directly No events in AE reported in trial register, confirmed by author	No
Murrough 2013, [81], USA, NCT00768430	73 (48/25)	Adults (21 to 80 years); MDD; excluded actively suicidal patients.	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 7 days (non-responders), 4 weeks or until relapse (responders) **TR+FU for analysis:** 7 days **Follow up used in analysis:** 7 days	IV Ketamine (0.5 mg/kg), single dose	Active control: Midazolam	MADRS-SI, data from Witt et al. 2019 who obtained data from author	Yes, directly	Yes

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Murrough 2015, [82], USA, NCT01507181	24 (12/12)	Adults (18 to 80 years); suicidal ideation (>4 on MADRS item 10), had a range of mood and anxiety disorders; current suicidal ideation	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 7 days and 5-week safety assessment **TR+FU for analysis:** 7 days (no data beyond this available)	Ketamine (0.5 mg/kg), single dose	Active control: Midazolam	BSS – patient, data from Witt et al. 2019 who obtained data from author (erroneously reported as MADRS)	Yes, indirectly No events in AE reported in trial register	Yes
Ochs-Ross 2020 [83], USA, Europe, Brazil, South Africa, NCT02422186	138 (72/66)	Older adults (≥65 years); MDD; excluded suicidal ideation with intent to act	Double blind RCT with open label phase	**TR:** 4 weeks **TR+FU:** 6 weeks **TR+FU for analysis:** only 4-week TR because afterwards patients could participate in an open label study	Intranasal Esketamine (28, 56, or 84 mg) twice weekly for 4 weeks.	Placebo with bittering agent	C-SSRS, data via YODA	Yes, directly	Yes
Ohtani 2024 [84], Japan jRCTs031210124	34 (17/17)	Adults (20 to 59 years); MDD; suicidality not specified in eligibility criteria.	Double blind RCT	**TR:** 14 days **TR+FU:** max. 28 days (max. 14 days after the last dose) **FU used for analysis:** max. 28 days	IV Ketamine (0.5 mg/kg), twice weekly for 2 weeks	Placebo: saline	MADRS-SI, data provided by author	Yes, directly No events in AE reported in trial register, confirmed by author	Yes
Pattanaseri 2024 [85], Thailand NCT05026203	21 (11/10)	Adults (18 to 64 years); MDD; suicidality not specified in eligibility criteria.	Double blind RCT	**TR:** 3 days **TR+FU:** 31 days **TR+FU for analysis:** 31 days	IV Ketamine (0.5 mg/kg), 3 doses over 3 days	Active control: Midazolam	MADRS-SI, data provided by author	Yes, indirectly No events in AE reported in trial register	No
Phillips 2019 [86], Canada, NCT01945047	43 (22/21)	Adults (18 to 65) with treatment resistant depression, suicidality not specified in eligibility criteria	Double blind crossover RCT	**TR:** 24 hours (single dose), crossover after min. 7 days **TR+FU:** min. 14 days **TR+FU for analysis:** 7 days Trial had two more open label phases which cannot be used	IV Ketamine (0.5 mg/kg), single dose	Active control: Midazolam 30 microgram/kg	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in publication	Yes
Popova 2019 [87], USA, Europe, NCT02418585	227 (114/109)	Adults (18 to 64 years); Depression or MDD; excluded suicidal ideation with intent to act or suicidal behaviour	Double blind RCT	**TR:** 4 weeks **TR+FU:** max. 28 weeks **TR+FU for analysis:** only 4-week TR because responders entered other trial	Intranasal esketamine (56 or 84mg) twice weekly for 4 weeks	Placebo with bittering agent[a]	C-SSRS, data via YODA	Yes, directly	Yes

Study, country, registration number	Total N randomised (interventio n/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Price 2022 [88], USA NCT03237286	154 (103/51)	Adults (18 to 60 years); MDD; acute suicidality excluded	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 1 year **TR+FU for analysis:** 1 year **FU used in analysis:** 1 year 1-year FU reported in Price (2023)	IV Ketamine (0.5 mg/kg), single dose Group 1: + Automated Self-Association Training (ASAT). Group 2: + sham ASAT	Placebo: saline + ASAT	MADRS-SI; QIDS-SR, data provided by author including data for 1-year FU (Price 2023)	Yes, directly No events in AE reported in trial register and confirmed by author	Yes
Ray-Griffith 2017, (66), USA, none reported	16 (8/8)	Adults (18+ years); Uni- or bi-polar depression; suicidality – not specified	Double blind RCT	**TR:** varied, average: 4.9 treatments (range 2-6, thus max. 2 weeks. **TR+FU:** TR + 90 days **TR+FU for analysis:** up to 5 follow-ups. These were the time points reported in the paper as only 6 participants completed follow-ups beyond these time points.	Ketamine (initially 1 mg/kg, uptitrated until anesthesia) + ECT (thrice weekly) Number of treatments varied, average 5.1 treatments	Methohexital (initially 1mg/kg, uptitrated until anesthesia+ ECT.	HDRS-SI and BDI-SI, data from Witt et al. 2019	Yes, directly No events as confirmed by author	Yes
Rodriguez 2013 [89], USA, NCT01100255	15 (8/7)	Adults (18 to 55 years) with OCD, excluded when "prominent suicide ideation"	Double blind crossover RCT	**TR:** 24 hours (single dose), crossover after min. 7 days **TR+FU:** min. 14 days **TR+FU used for analysis:** 7 days	Intravenous ketamine (0.5 mg/kg), single dose	Placebo: saline	HDRS-SI, data unavailable	Yes, indirectly No events in AE reported in publication and on trial register	Yes
Schiweck 2021 [90], Belgium, EudraCT 2016-001715-21	28 (7/21) according to information on trial register	Adults (18 to 65 years) with MDD	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 4 days (6 days for responders, protocol and supplement says day 21) **TR+FU for analysis:** 4 days	Intravenous ketamine (0.5 mg/kg), single dose	Placebo: saline	HDRS-SI, data unavailable	Yes, indirectly No events in AE reported in trial register	Yes
Shiroma 2020 [91], USA, NCT02360280	58 (28/30)	Adults (18 to 75 years) with treatment resistant MDD, excluded when imminent risk of suicide ideation and/or behavior with intent	Double blind RCT	**TR:** 10 days **TR+FU:** 6 months **TR+FU for analysis:** only 10-day TR can be used because afterwards both groups received ketamine	Intravenous ketamine (0.5mg/kg), thrice weekly	Active control: Midazolam	C-SSRS, data unavailable	Yes, indirectly No events in AE reported in trial register	No
Singh 2016a [92], USA, NCT01627782	68 (35/33) Twice weekly: 35 (18/17)	Adults (18 to 64 years); MDD; excluded clinically significant suicidal ideation	Double blind RCT with open label phase	**TR:** 4 weeks **TR+FU:** up to 13 weeks (optional open label phase after TR) **TR+FU for analysis:** only 15 days of TR, afterwards non-responders could	Intravenous ketamine (0.5 mg/kg) two or three times weekly for 4 weeks	Placebo: Sodium Chloride	C-SSRS, data via YODA	Yes, directly	Yes

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
	Thrice weekly: 33 (17/16)			leave study and open label phase was optional					
Singh 2016b [93], Europe, NCT01640080	30 (20/10)	Adults (18 to 64 years); MDD: excluded acutely suicidal	Double blind, double randomisation, RCT	**TR:** 1 week **TR+FU:** 5 weeks **TR+FU for analysis:** day 4 of TR, afterwards placebo non-responders received esketamine	IV Esketamine (2 groups: 0.20 mg/kg or 0.40 mg/kg), 2 doses over 4 days	Placebo	C-SSRS, data via YODA	Yes, directly	Yes
Smith-Apeldoorn 2024 [94], Netherlands NTR6161	113 (57/56)	Adults (18 to 80 years); MDD; excluded active suicidal intent	Double blind RCT followed by open label phase	**TR:** 6 weeks **TR+FU:** 16 weeks **TR+FU used for analysis:** 10 weeks (TR + 4-week washout phase but not 6 week open label phase)	Esketamine (30-90 mg/day) for 6 weeks	Placebo	HDRS-SI, data obtained from authors	Yes, directly	Yes
Sos 2013 [95], Czech Republic, EudraCT Nr. 2009-010625-39	30 (11/19) (pre-cross-over)	Adults (18 to 65 years); MDD; excluded any suicidal risk	Double blind cross-over RCT	**TR:** 24 hours (single dose), cross-over at day 7 **TR+FU:** 2 weeks **TR+FU for analysis:** 7 days – pre-cross-over period	IV Ketamine (0.27 mg/kg), single dose	Placebo: saline	MADRS-SI, data from Witt et al. 2019 who obtained data from authors	Yes, indirectly No events in AE reported in publication	No
Su 2017 [96], Taiwan, UMIN000016985	71 (47/24)	Adults, MDD; suicidality – not specified	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 28 days **TR+FU for analysis:** 28 days	IV Ketamine, 2 groups: 0.2 or 0.5 mg/kg, single dose	Placebo: saline	HDRS-SI, data from Witt et al. (2019) who obtained data from author	Yes, indirectly No events in AE reported in publication	Yes
Su 2023 [97], Taiwan UMIN000033916 and UMIN000033760	84 (42/42)	Adults (20 to 64 years); MDD; included suicidal ideation	Double blind RCT	**Length of treatment phase:** 24 hours (single dose) **End of treatment FU:** 14 days **Total trial duration:** 14 days **FU used in analysis:** 14 days	IV Ketamine (0.5 mg/kg), single dose	Active control: Midazolam	C-SSRS, M's from figure, SD's imputed with the mean SD of other studies	Yes, directly (Note: suicide attempts reported but not in the study publication)	No
Sumner 2020 [34], New Zealand, ACTRN12615000 573550	30 (17/16)	Adults; MDD; imminent risk of suicide excluded	Double blind cross-over RCT	**TR:** 24 hours (single dose), washout period min 3 weeks (4 weeks for women) then crossover **TR+FU:** min. 5 weeks **TR+FU for analysis:** 3 weeks	IV Ketamine (0.44 mg/kg), single dose	Active placebo: remifentanil	MADRS-SI, data provided by author	Yes, directly No information in publication but author confirmed that no events occurred.	No

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
Takahashi 2021, [98], Japan, NCT02918318	202 (122/80)	Adults (20 to 64 years); MDD; excluded suicidal ideation with intent to act or suicidal behaviour	Double blind RCT with open label phase	**TR:** 4 weeks **TR+FU:** 8-28 weeks **TR+FU for analysis:** only 4-week TR, afterwards open-label treatment for responders	Esketamine nasal spray, 3 groups: 28, 56 or 84mg twice weekly for 4 weeks	Placebo	C-SSRS, data via YODA	Yes, directly	Yes
Taylor 2018 [99], USA, NCT02083926	18 (9/9)	Adults, Social anxiety disorder; suicidality – not specified	Double blind crossover RCT	**TR:** 24 hours (single dose), 28 day washout then crossover **TR+FU:** 42 days **TR+FU for analysis:** 28 days	IV Ketamine (0.5 mg/kg), single dose	Placebo: saline	HDRS-SI, data unavailable	Yes, indirectly No events in AE reported in trial register	Yes
Tiger 2020 [100], Sweden, none reported	30 (20/10)	Adults (20 to 80 years); MDD; excluded suicidality	Double blind RCT with open label phase	**TR:** 24 hours (single dose), open label phase **TR+FU:** 72 hours **TR+FU for analysis:** only 24-hour TR because afterwards all patients were treated open label	Ketamine (0.5 mg/kg) single dose	Placebo: saline	MADRS-SI, data unavailable	Yes, directly. Not reported but author confirmed that no events occurred.	No
Wang 2024 [101], China NCT04414943	364 (182/182)	Adults (≥18 years); prenatal depression, suicidality – not specified	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 42 days **TR+FU for analysis:** 42 days	IV Esketamine (0.2 mg/kg), single dose (given after childbirth)	Placebo: saline	HDRS-SI or EPDS-SI, data unavailable	Yes, indirectly No events in AE reported in publication	No
Xiao 2023 [102], China ChiCTR2100050335	30 (15/15)	Adults (18 to 60 years); MDD; excluded suicidality	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 6 weeks **TR+FU for analysis:** 6 weeks	IV Esketamine (0.2 mg/kg), single dose	Active control: Midazolam	MADRS-SI, data unavailable	Yes, indirectly No events in AE reported in publication	No
Yang 2023 [103], China, ChiCTR-ROC-2000039069	300 (200/100)	Women (over 18 years); Postpartum depressive symptoms; suicidality – not specified	Double blind RCT	**TR:** 24 hours (single dose) **TR+FU:** 42 days **TR+FU for analysis:** 42 days	IV Esketamine (2 groups: 1 or 2mg/kg), single dose, received after giving birth	Placebo: saline	EPDS-SI but data unavailable	Yes, indirectly No events in AE reported in publication	No
Zarate 2006 [104], USA, NCT00088699	18 (9/9)	Adults (18 to 65 years) with treatment resistant	Doube blind crossover RCT	**TR:** 24 hours (single dose), crossover after 7 days **TR+FU:** 14 days	Ketamine (0.5 mg/kg) single dose	Placebo: saline	HDRS-SI, data provided by author	Yes, directly No events occurred as confirmed by authors	No

Study, country, registration number	Total N randomised (intervention/control)	Sample (population, diagnosis; suicidality present)	Randomised Controlled Trial (RCT) design	Lengths of Treatment Phase (TR), TR+Follow-up Phase (TR+FU), and TR+FU used for analysis	Intervention, number of treatments, length of treatment	Control	Suicide ideation measures and data availability	Suicidal behavior and deaths: information available (direct/indirect)?	Conflict of interest? Yes/No/not reported
		MDD without serious suicide risk		**TR+FU for analysis:** 7 days, pre-cross-over					
Zarate 2012 [105], USA, NCT00088699	Publication: 15 Full data-set: 44 (21/23)	Adults (18 to 65 years); Bipolar Affective Disorder; suicidality – not specified	Double blind cross-over RCT	**TR:** 24 hours (single dose), cross-over after 2 weeks **TR+FU:** 4 weeks **TR+FU for analysis:** 14 days, pre-cross over	IV Ketamine (0.5 mg/kg), single dose	Placebo: saline	MADRS-SI, data provided by author for full sample	Yes, directly No events in AE reported in trial register, confirmed by author	Yes
Zhou 2024 [106], China, ChiCTR2000041232	54 (27/27)	Adolescents (13 to 18 years); MDD; eligibility criteria included presence of suicidal ideation	Double blind RCT	**TR:** 6 days **TR+FU:** 33 days **TR+FU for analysis:** 33 days	IV Esketamine (0.25 mg/kg), 3 times over 5 days	Active control: MIdazolam	C-SSRS, change from baseline, n's from Fig. 2E, data for endpoints unavailable	Yes, directly One event ("suicidal behavior") in Midazolam, assumed that this was a suicide attempt	No

References are to be found in the online supplement.

AE: adverse events; BHS: Beck Hopelessness Scale; BDI-SI: suicide ideation item from the Beck Depression Inventory (BDI); BSS: Beck Scale for Suicide Ideation; C-SSRS: Columbia-Suicide Severity Rating Scale; CGI-SS-r: Clinical Global Impression of Severity of Suicidality Revised version; EPDS-SI: suicide ideation item from the Edinburgh Postpartal Depression Scale (EPDS); HDRS-SI: suicide idation item from the Hamilton Depression Rating Scale (HDRS); IN: intranasal; IV: Intravenous; M: mean; MADRS-SI: Montgomery- Åsberg Depression Rating Scale – suicidal ideation item; MDD: Major Depressive Disorder; QIDS-SR: Quick Inventory of Depressive Symptomatology – Self-Report; SD: standard deviation; SIBAT: Suicide Ideation and Behavior Assessment Tool; SPS: Suicide Probability Scale; SSF: Suicide Status Form; SSI: Scale for Suicidal Ideation; SUD: substance use disorder; TRD: Treatment Resistant Depression; n: sample size; YODA: Yale University Open Data Access Project

In most trials, patients were continued on their antidepressant and we only reported other adjunct treatments.

a. Bittering agent simulates esketamine taste
b. Direct: suicidal events/deaths reported in publication or the trial register or via corresponding author; indirect: AE reported but no suicidal events/deaths seem to have occurred

Table 2: Meta-analysis of suicides and suicide attempts (Treatment + Follow-Up Phases combined)

Analysis	OR [95% CI]	p	I^2 [95% CI]	τ^2 [95%CI]	number of studies
Exact (Liu et al. 2012)	0.97 [0.48 to 1.97]	..	..	..	64
Bayesian - Uninformative prior (delta = 250)	0.86 [0.45 to 1.56]	..	..	1.34 [1.01 to 2.59]	64
Bayesian - Informative prior (delta = 15)	0.86 [0.46 to 1.54]	..	..	1.33 [1.01 to 2.55]	64
Binomial-normal GLMM, fixed intercept	0.91 [0.52 to 1.60]	..	..	0.00	64
Binomial-normal GLMM, fixed intercept, alternative parameterization	0.91 [0.51 to 1.60]	..	..	0.00	64
Binomial-normal GLMM, random intercept	0.89 [0.51 to 1.56]	..	..	0.00	64
Binomial-normal GLMM, random intercept, alternative parameterization	0.89 [0.53 to 1.57]	..	..	0.00	64
Hypergeometric-normal GLMM	0.92 [0.53 to 1.58]	..	..	0.00	64
Conditional binomial GLMM	0.92 [0.26 to 1.58]	..	..	0.00	64
Beta-binomial model	0.66 [0.52 to 1.66]	..	..		64
Peto	0.88 [0.49 to 1.61]	0.69	19 [0 to 57]	0.27 [0.00 to 2.10]	13
Mantel-Haenszel - with continuity correction	0.88 [0.59 to 1.30]	0.52	0 [0 to 29]	0.00 [0.00 to 0.00]	64
Mantel-Haenszel - with continuity/treatment arm correction	0.99 [0.66 to 1.48]	0.97	0 [0 to 29]	0.00 [0.00 to 0.00]	64
Mantel-Haenszel - risk difference	Risk difference 0.00 [-0.01 to 0.01]	0.99	0 [0 to 29]	0.00 [0.00 to 0.00]	64
Arcsine square root transformed risk difference	Risk difference 0.00 [-0.03 to 0.02]	0.78	0 [-]	0.00 [-]	64

Note: OR: odds-ratio, CI: confidence interval or credible interval for Bayesian analysis, p: p-value of odds-ratio, I^2 and τ^2: measures of heterogeneity, GLMM: generalized linear mixed models.

All methods are appropriate except Peto's method, because there are both single-arm-zero-events studies and double-arm-zero-events studies and the total event count in each arm is non-zero [27].

Table 3: Meta-analysis of suicidal ideation

Result	up to 4h	>12h up to 24h	>24h up to 72h	>72h up to 2w	>2w up to 4w	>4w	End of Treatment
SMD [95%-CI]	-0.40 [-0.63 - -0.16]	-0.35 [-0.51 to -0.19]	-0.16 [-0.28 to -0.04]	-0.16 [-0.26 to -0.06]	-0.14 [-0.27 to 0.00]	-0.01 [-0.20 to 0.17]	-0.24 [-0.35 to -0.13]
p	< 0.01	< 0.001	< 0.01	< 0.01	0.05	0.87	< 0.001
Heterogeneity	I^2=52%, τ^2=0.2, p<0.01	I^2=12%, τ^2=0.06, p=0.29	I^2=17%, τ^2=0.04, p=0.21	I^2=0%, τ^2=0.03, p=0.90	I^2=25%, τ^2=0.04, p=0.16	I^2=8%, τ^2=0.02, p=0.37	I^2=24%, τ^2=0.05, p=0.10
Number of trials	23	25	29	36	19	9	38
Number of observations	1261	1435	2529	2858	2159	792	2827
Small study effect p-Value of Egger's regression test	0.63	0.75	0.17	0.50	0.88	too few studies	0.02

Note: SMD: standardized mean difference, CI: confidence interval, I^2 and τ^2: measures of heterogeneity, p: p-value

Table 4: Meta Analytic Results, Suicide Ideation - Subgroup Analyses

Subgroup analysis	up to 4h	>12h up to 24h	>24h up to 72h	>72h up to 2w	>2w up to 4w	>4w	End of Treatment
Drug							
—Esketamine	-0.31 [-0.53 to -0.09]*	-0.23 [-0.43 to -0.02]*	-0.05 [-0.16 to 0.07]	-0.10 [-0.21 to 0.01]	-0.06 [-0.18 to 0.05]	0.04 [-0.27 to 0.35]	-0.06 [-0.18 to 0.05]
—Ketamine	-0.41 [-0.69 to -0.13]*	-0.38 [-0.58 to -0.19]*	-0.32 [-0.52 to -0.12]*	-0.22 [-0.38 to -0.07]*	-0.26 [-0.54 to 0.02]	-0.10 [-0.32 to 0.13]	-0.41 [-0.56 to -0.25]*
— test of difference (p-value)	0.09	0.05	< 0.001	0.08	< 0.01	0.33	< 0.001
Suicidality exclusion or inclusion criterion							
—Exclusion/unknown	-0.32 [-0.69 to 0.06]	-0.24 [-0.48 to -0.00]*	-0.06 [-0.19 to 0.07]	-0.14 [-0.27 to -0.01]*	-0.12 [-0.30 to 0.07]	0.10 [-0.18 to 0.37]	-0.15 [-0.29 to -0.02]*
—Inclusion	-0.44 [-0.74 to -0.13]*	-0.42 [-0.61 to -0.23]*	-0.31 [-0.52 to -0.10]*	-0.20 [-0.37 to -0.03]*	-0.17 [-0.38 to 0.04]	-0.09 [-0.32 to 0.13]	-0.37 [-0.56 to -0.18]*
— test of difference (p-value)	0.58	0.32	< 0.01	0.31	0.25	0.21	< 0.01
Doses							
—Repeated	..	..	..	-0.15 [-0.26 to -0.03]*	-0.11 [-0.23 to 0.01]	-0.02 [-0.23 to 0.18]	..
—Single	..	..	..	-0.19 [-0.39 to 0.01]	-0.17 [-0.57 to 0.22]	0.02 [-0.40 to 0.44]	..
— test of difference (p-value)	..	..	..	0.53	0.16	0.88	..
Rating							
—Clinician	-0.38 [-0.59 to -0.17]*	-0.35 [-0.51 to -0.19]*	-0.14 [-0.25 to -0.03]*	-0.16 [-0.27 to -0.05]*	-0.15 [-0.30 to -0.01]*	-0.09 [-0.27 to 0.10]	-0.24 [-0.37 to -0.11]*
—Patient	-0.64 [-1.48 to 0.20]	-0.47 [-0.94 to -0.01]*	-0.44 [-1.00 to 0.11]	-0.18 [-0.50 to 0.13]	0.05 [-0.45 to 0.56]	-0.01 [-0.41 to 0.40]	-0.33 [-0.63 to -0.03]*
— test of difference (p-value)	0.82	0.49	0.15	0.79	0.42	0.74	0.43
Placebo							
—Active	-0.24 [-0.62 to 0.14]	-0.43 [-0.70 to -0.16]*	-0.24 [-0.50 to 0.03]	-0.19 [-0.48 to 0.10]	0.00 [-0.56 to 0.56]	-0.15 [-0.59 to 0.30]	-0.48 [-0.74 to -0.22]*
—Non-active	-0.48 [-0.77 to -0.19]*	-0.30 [-0.48 to -0.11]*	-0.15 [-0.29 to -0.01]*	-0.15 [-0.26 to -0.05]*	-0.14 [-0.28 to -0.01]*	0.01 [-0.19 to 0.21]	-0.17 [-0.28 to -0.05]*
— test of difference (p-value)	0.74	0.09	0.30	0.73	0.78	0.48	< 0.001

Note: Results are standardized mean differences with their 95%-confidence intervals in brackets

* $p < 0.05$

REFERENCES

World Health Organization. *Suicide in the world. Global health estimates.* World Health Organization 2019.

UK Parliament. Suicide Statistics. UK: House of Commons Library 2025.

Ilic M, Ilic I. Worldwide suicide mortality trends (2000-2019): A joinpoint regression analysis. *World J Psychiatry*. 2022;12:1044–60. doi: 10.5498/wjp.v12.i8.1044

Xu YE, Barron DA, Sudol K, *et al.* Suicidal behavior across a broad range of psychiatric disorders. *Mol Psychiatry*. 2023;28:2764–810. doi: 10.1038/s41380-022-01935-7

Price RB, Mathew SJ. Does Ketamine Have Anti-Suicidal Properties? Current Status and Future Directions. *CNS Drugs*. 2015;29:181–8. doi: 10.1007/s40263-015-0232-4

Johns Hopkins Medicine. Esketamine for Treatment-Resistant Depression. 2023. https://www.hopkinsmedicine.org/health/treatment-tests-and-therapies/esketamine-for-treat ment-resistant-depression (accessed 2 April 2025)

Johnson & Johnson. SPRAVATO® (esketamine). Official Patient Website. https://www.spravato.com/what-is-spravato/ (accessed 2 April 2025)

Hochschild A, Grunebaum MF, Mann JJ. The rapid anti-suicidal ideation effect of ketamine: A systematic review. *Preventive Medicine*. 2021;152:106524. doi: 10.1016/j.ypmed.2021.106524

McIntyre RS, Rosenblat JD, Nemeroff CB, *et al.* Synthesizing the Evidence for Ketamine and Esketamine in Treatment-Resistant Depression: An International Expert Opinion on the Available Evidence and Implementation. *Am J Psychiatry*. 2021;appiajp202020081251. doi: 10.1176/appi.ajp.2020.20081251

Ng J, Rosenblat JD, Lui LMW, *et al.* Efficacy of ketamine and esketamine on functional outcomes in treatment-resistant depression: A systematic review. *Journal of Affective Disorders*. 2021;293:285–94. doi: 10.1016/j.jad.2021.06.032

Walsh Z, Mollaahmetoglu OM, Rootman J, *et al.* Ketamine for the treatment of mental health and substance use disorders: comprehensive systematic review. *BJPsych Open*. 2022;8:e19. doi: 10.1192/bjo.2021.1061

Gastaldon C, Raschi E, Kane JM, *et al.* Post-Marketing Safety Concerns with Esketamine: A Disproportionality Analysis of Spontaneous Reports Submitted to the FDA Adverse Event Reporting System. *Psychother Psychosom*. 2020;1–8. doi: 10.1159/000510703

Chang H, Huang M-C, Chen L-Y. Major Depressive Disorder Induced by Chronic Ketamine Abuse: A Case Report. *Prim Care Companion CNS Disord*. Published Online First: 23 June 2016. doi: 10.4088/PCC.15l01881

Liang HJ, Tang KL, Chan F, *et al.* Ketamine Users Have High Rates of Psychosis and/or Depression. *Journal of Addictions Nursing*. 2015;26:8–13. doi: 10.1097/JAN.0000000000000060

Morgan CJA, Muetzelfeldt L, Curran HV. Ketamine use, cognition and psychological wellbeing: a

comparison of frequent, infrequent and ex-users with polydrug and non-using controls. *Addiction*. 2009;104:77–87. doi: 10.1111/j.1360-0443.2008.02394.x

Alario AA, Niciu MJ. (Es)Ketamine for Suicidal Ideation and Behavior: Clinical Efficacy. *Chronic Stress*. 2022;6:24705470221128016. doi: 10.1177/24705470221128017

Shamabadi A, Ahmadzade A, Hasanzadeh A. Ketamine for suicidality: An umbrella review. *Brit J Clinical Pharma*. 2022;88:3990–4018. doi: 10.1111/bcp.15360

Witt K, Potts J, Hubers A, *et al.* Ketamine for suicidal ideation in adults with psychiatric disorders: A systematic review and meta-analysis of treatment trials. *Aust N Z J Psychiatry*. 2020;54:29–45. doi: 10.1177/0004867419883341

Jollant F, Colle R, Nguyen TML, *et al.* Ketamine and esketamine in suicidal thoughts and behaviors: a systematic review. *Therapeutic Advances in Psychopharmacology*. 2023;13:204512532311513. doi: 10.1177/20451253231151327

Juul S, Faltermeier P, Siddiqui F, *et al.* Challenges in the selection and measurement of outcomes in psychiatric trials. *BMJ EBM*. 2025;bmjebm-2024-113171. doi: 10.1136/bmjebm-2024-113171

Xu C, Furuya-Kanamori L, Islam N, *et al.* Should studies with no events in both arms be excluded in evidence synthesis? *Contemporary Clinical Trials*. 2022;122:106962. doi: 10.1016/j.cct.2022.106962

Liu D, Liu RY, Xie M. Exact Meta-Analysis Approach for Discrete Data and its Application to 2 × 2 Tables With Rare Events. *Journal of the American Statistical Association*. 2014;109:1450–65. doi: 10.1080/01621459.2014.946318

Fountoulakis KN, Saitis A, Schatzberg AF. Esketamine Treatment for Depression in Adults: A PRISMA Systematic Review and Meta-Analysis. *AJP*. 2025;182:259–75. doi: 10.1176/appi.ajp.20240515

Chen C-C, Zhou N, Hu N, *et al.* Acute Effects of Intravenous Sub-Anesthetic Doses of Ketamine and Intranasal Inhaled Esketamine on Suicidal Ideation: A Systematic Review and Meta-Analysis. *NDT*. 2023;Volume 19:587–99. doi: 10.2147/ndt.s401032

Page MJ, McKenzie JE, Bossuyt PM, *et al.* The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;n71. doi: 10.1136/bmj.n71

Higgins JPT, Altman DG, Gotzsche PC, *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928–d5928. doi: 10.1136/bmj.d5928

Xu C, Furuya-Kanamori L, Zorzela L, *et al.* A proposed framework to guide evidence synthesis practice for meta-analysis with zero-events studies. *Journal of Clinical Epidemiology*. 2021;135:70–8. doi: 10.1016/j.jclinepi.2021.02.012

Plöderl M, Hengartner MP, Bschor T, *et al.* Commentary to 'antidepressants and suicidality: A re-analysis of the re-analysis'. *J Affect Disord*. 2020;273:252–3. doi: 10.1016/j.jad.2020.04.025

Jansen K, Holling H. Random-effects meta-analysis models for the odds ratio in the case of rare events under different data-generating models: A simulation study. *Biometrical J*. 2023;65:2200132. doi: 10.1002/bimj.202200132

Higgins JPT, Thomas CJ, Chandler J, *et al.* Cochrane Handbook for Systematic Reviews of Interventions version 6.5 (updated August 2024). Cochrane 2024.

Wan X, Wang W, Liu J, *et al.* Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014;14:135. doi: 10.1186/1471-2288-14-135

Burger J, Capobianco M, Lovern R, *et al.* A Double-Blinded, Randomized, Placebo-Controlled Sub-Dissociative Dose Ketamine Pilot Study in the Treatment of Acute Depression and Suicidality in a Military Emergency Department Setting. *Mil Med*. 2016;181:1195–9.

R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing 2025.

Sumner R.L., McMillan R., Spriggs M.J., *et al.* Ketamine Enhances Visual Sensory Evoked Potential Long-term Potentiation in Patients With Major Depressive Disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2020;5:45–55. doi: 10.1016/j.bpsc.2019.07.002

Fava M, Freeman MP, Flynn M, *et al.* Double-blind, placebo-controlled, dose-ranging trial of intravenous ketamine as adjunctive therapy in treatment-resistant depression (TRD). *Molecular Psychiatry*. 2020;25:1592–603. doi: 10.1038/s41380-018-0256-5

Anderson IM, Blamire A, Branton T, *et al.* Ketamine augmentation of electroconvulsive therapy to improve neuropsychological and clinical outcomes in depression (Ketamine-ECT): a multicentre, double-blind, randomised, parallel-group, superiority trial. *The lancet Psychiatry*. 2017;4:365. doi: 10.1016/S2215-0366(17)30077-9

Lii TR, Smith AE, Flohr JR, *et al.* Randomized trial of ketamine masked by surgical anesthesia in patients with depression. *Nat Mental Health*. 2023;1:876–86. doi: 10.1038/s44220-023-00140-x

Dwyer J.B., Landeros-Weisenberger A., Johnson J.A., *et al.* Efficacy of intravenous ketamine in adolescent treatment-resistant depression: A randomized midazolam-controlled trial. *Am J Psychiatry*. 2021;178:352–62. doi: 10.1176/appi.ajp.2020.20010018

Fineberg SK, Choi EY, Shapiro-Thompson R, *et al.* A pilot randomized controlled trial of ketamine in Borderline Personality Disorder. *Neuropsychopharmacology*. 2023;48:991–9. doi: 10.1038/s41386-023-01540-4

Glue P, Neehoff S, Beaglehole B, *et al.* Ketamine for treatment-resistant major depressive disorder: Double-blind active-controlled crossover study. *J Psychopharmacol*. 2024;38:162–7. doi: 10.1177/02698811241227026

Sharma T, Guski LS, Freund N, *et al.* Suicidality and aggression during antidepressant treatment: systematic review and meta-analyses based on clinical study reports. *BMJ*. Published Online First: 27 January 2016. doi: 10.1136/bmj.i65

Le Noury J, Nardo JM, Healy D, *et al.* Restoring Study 329: efficacy and harms of paroxetine and imipramine in treatment of major depression in adolescence. *BMJ*. 2015;h4320. doi: 10.1136/bmj.h4320

Witt K, Townsend E, Arensman E, *et al.* Psychosocial Interventions for People Who Self-Harm: Methodological Issues Involved in Trials to Evaluate Effectiveness. *Archives of Suicide Research*. 2020;24:S32–93. doi: 10.1080/13811118.2019.1592043

Abbar M, Demattei C, El-Hage W, *et al.* Ketamine for the acute treatment of severe suicidal ideation: Double blind, randomised placebo controlled trial. *BMJ: British Medical Journal*. 2022;376.

Nikolin S, Rodgers A, Schwaab A, *et al.* Ketamine for the treatment of major depression: a systematic review and meta-analysis. *eClinicalMedicine*. 2023;62:102127. doi: 10.1016/j.eclinm.2023.102127

Feng W, Chen C, Zeng Y, *et al.* Efficacy of single and repeated ketamine administration for suicidal ideation in adults with psychiatric disorders: A meta-analysis. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2025;136:111152. doi: 10.1016/j.pnpbp.2024.111152

Glue P, Loo C, Fam J, *et al.* Extended-release ketamine tablets for treatment-resistant depression: a randomized placebo-controlled phase 2 trial. *Nat Med*. 2024;30:2004–9. doi: 10.1038/s41591-024-03063-x

Daly E.J., Trivedi M.H., Janik A., *et al.* Efficacy of Esketamine Nasal Spray Plus Oral Antidepressant Treatment for Relapse Prevention in Patients with Treatment-Resistant Depression: A Randomized Clinical Trial. *JAMA Psychiatry*. 2019;76:893–903. doi: 10.1001/jamapsychiatry.2019.1189

Morgan CJA, Curran HV, the Independent Scientific Committee on Drugs (ISCD). Ketamine use: a review. *Addiction*. 2012;107:27–38. doi: 10.1111/j.1360-0443.2011.03576.x

Liu Y, Lin D, Wu B, *et al.* Ketamine abuse potential and use disorder. *Brain Research Bulletin*. 2016;126:68–73. doi: 10.1016/j.brainresbull.2016.05.016

Abdallah C.G., Roache J.D., Gueorguieva R., *et al.* Dose-related effects of ketamine for antidepressant-resistant symptoms of posttraumatic stress disorder in veterans and active duty military: a double-blind, randomized, placebo-controlled multi-center clinical trial. *Neuropsychopharmacology*. 2022;47:1574–81. doi: 10.1038/s41386-022-01266-9

Ahmed GK, Elserogy YM, Elfadl GMA, *et al.* Antidepressant and anti-suicidal effects of ketamine in treatment-resistant depression associated with psychiatric and personality comorbidities: A double-blind randomized trial. *J Affect Disord*. 2023;325:127–34. doi: 10.1016/j.jad.2023.01.005

Anderson IM, Blamire A, Branton T, *et al.* Randomised controlled trial of ketamine augmentation of electroconvulsive therapy to improve neuropsychological and clinical outcomes in depression (Ketamine-ECT study). *Efficacy Mech Eval*. 2017;4:1–112. doi: 10.3310/eme04020

Canuso C.M., Singh J.B., Fedgchin M., *et al.* Efficacy and safety of intranasal esketamine for the rapid reduction of symptoms of depression and suicidality in patients at imminent risk for suicide: Results of a double-blind, randomized, placebo-controlled study. *Am J Psychiatry*. 2018;175:620–30. doi: 10.1176/appi.ajp.2018.17060720

Chen X, Hou X, Bai D, *et al.* Efficacy and Safety of Flexibly Dosed Esketamine Nasal Spray Plus a Newly Initiated Oral Antidepressant in Adult Patients with Treatment-Resistant Depression: A Randomized, Double-Blind, Multicenter, Active-Controlled Study Conducted in China and USA. *NDT*. 2023;Volume 19:693–707. doi: 10.2147/NDT.S391096

Colla M, Offenhammer B, Scheerer H, *et al.* Oral prolonged-release ketamine in treatment-resistant depression - A double-blind randomized placebo-controlled multicentre trial of KET01, a novel ketamine formulation - Clinical and safety results. *J Psychiatr Res*. 2024;173:124–30. doi: 10.1016/j.jpsychires.2024.03.002

Dakwar E., Nunes E.V., Hart C.L., *et al.* A single ketamine infusion combined with mindfulness-based behavioral modification to treat cocaine dependence: A randomized clinical trial. *Am J Psychiatry*. 2019;176:923–30. doi: 10.1176/appi.ajp.2019.18101123

Dakwar E., Levin F., Hart C.L., *et al.* A Single Ketamine Infusion Combined With Motivational Enhancement Therapy for Alcohol Use Disorder: A Randomized Midazolam-Controlled Pilot Trial. *Am J Psychiatry*. 2020;177:125–33. doi: 10.1176/appi.ajp.2019.19070684

Daly E.J., Singh J.B., Fedgchin M., *et al.* Efficacy and safety of intranasal esketamine adjunctive to oral antidepressant therapy in treatment-resistant depression: A randomized clinical trial. *JAMA Psychiatry*. 2018;75:139–48. doi: 10.1001/jamapsychiatry.2017.3739

Domany Y., Bleich-Cohen M., Tarrasch R., *et al.* Repeated oral ketamine for out-patient treatment of resistant depression: Randomised, double-blind, placebo-controlled, proof-of-concept study. *Br J Psychiatry*. 2019;214:20–6. doi: 10.1192/bjp.2018.196

Domany Y, Shelton RC, McCullumsmith CB. Ketamine for acute suicidal ideation. An emergency department intervention: a randomized, double-blind, placebo-controlled, proof-of-concept trial. *Depression and anxiety*. 2020;37:224. doi: 10.1002/da.22975

Domany Y, McCullumsmith CB. Single, Fixed-Dose Intranasal Ketamine for Alleviation of Acute Suicidal Ideation. An Emergency Department, Trans-Diagnostic Approach: a Randomized, Double-Blind, Placebo-Controlled, Proof-of-Concept Trial. *Archives of suicide research*. 2022;26:1250. doi: 10.1080/13811118.2021.1878078

Duek O, Korem N, Li Y, *et al.* Long term structural and functional neural changes following a single infusion of Ketamine in PTSD. *Neuropsychopharmacology*. 2023;48:1648–58. doi: 10.1038/s41386-023-01606-3

Feder A, Parides MK, Murrough JW, *et al.* Efficacy of intravenous ketamine for treatment of chronic posttraumatic stress disorder: a randomized clinical trial. *JAMA psychiatry*. 2014;71:681. doi: 10.1001/jamapsychiatry.2014.62

Feder A, Costi S, Rutter SB, *et al.* A Randomized Controlled Trial of Repeated Ketamine Administration for Chronic Posttraumatic Stress Disorder. *American journal of psychiatry*.

2021;178:193. doi: 10.1176/appi.ajp.2020.20050596

Fedgchin M, Trivedi M, Daly EJ, *et al.* Efficacy and Safety of Fixed-Dose Esketamine Nasal Spray Combined With a New Oral Antidepressant in Treatment-Resistant Depression: Results of a Randomized, Double-Blind, Active-Controlled Study (TRANSFORM-1). *Int J Neuropsychopharmacol*. 2019;22:616–30. doi: 10.1093/ijnp/pyz039

George D, Gálvez V, Martin D, *et al.* Pilot Randomized Controlled Trial of Titrated Subcutaneous Ketamine in Older Patients with Treatment-Resistant Depression. *American journal of geriatric psychiatry*. 2017;25:1199. doi: 10.1016/j.jagp.2017.06.007

Grabski M, McAndrew A, Lawn W, *et al.* Adjunctive Ketamine With Relapse Prevention-Based Psychological Therapy in the Treatment of Alcohol Use Disorder. *Am J Psychiatry*. 2022;179:152–62. doi: 10.1176/appi.ajp.2021.21030277

Grunebaum MF, Ellis SP, Keilp JG, *et al.* Ketamine versus midazolam in bipolar depression with suicidal thoughts: A pilot midazolam-controlled randomized clinical trial. *Bipolar Disord*. 2017;19:176–83. doi: 10.1111/bdi.12487

Grunebaum MF, Galfalvy HC, Choo T-H, *et al.* Ketamine for Rapid Reduction of Suicidal Thoughts in Major Depression: A Midazolam-Controlled Randomized Clinical Trial. *Am J Psychiatry*. 2018;175:327–35. doi: 10.1176/appi.ajp.2017.17060647

Hu Y.-D., Xiang Y.-T., Fang J.-X., *et al.* Single i.v. ketamine augmentation of newly initiated escitalopram for major depression: results from a randomized, placebo-controlled 4-week study. *Psychol Med*. 2016;46:623–35. doi: 10.1017/S0033291715002159

Ionescu D.F., Bentley K.H., Eikermann M., *et al.* Repeat-dose ketamine augmentation for treatment-resistant depression with chronic suicidal ideation: A randomized, double blind, placebo controlled trial. *J Affective Disord*. 2019;243:516–24. doi: 10.1016/j.jad.2018.09.037

Ionescu DF, Fu DJ, Qiu X, *et al.* Esketamine Nasal Spray for Rapid Reduction of Depressive Symptoms in Patients With Major Depressive Disorder Who Have Active Suicide Ideation With Intent: results of a Phase 3, Double-Blind, Randomized Study (ASPIRE II). *The international journal of neuropsychopharmacology*. 2021;24:22. doi: 10.1093/ijnp/pyaa068

Jones GH, Vecera CM, Ruiz AC, *et al.* A Randomized, Double-Blind, Placebo-Controlled Pilot Trial of the Acute Antisuicidal and Antidepressant Effects of Intranasal (R,S)-Ketamine in Severe Unipolar and Bipolar Depression With and Without Comorbid Alcohol Use Disorder. *J Clin Psychiatry*. 2024;85:23m14974. doi: 10.4088/JCP.23m14974

Lapidus KAB, Levitch CF, Perez AM, *et al.* A randomized controlled trial of intranasal ketamine in major depressive disorder. *Biol Psychiatry*. 2014;76:970–6. doi: 10.1016/j.biopsych.2014.03.026

Leal GC, Souza-Marques B, Mello RP, *et al.* Arketamine as adjunctive therapy for treatment-resistant depression: A placebo-controlled pilot study. *J Affect Disord*. 2023;330:7–15. doi: 10.1016/j.jad.2023.02.151

Lijffijt M, Murphy N, Iqbal S, *et al.* Identification of an optimal dose of intravenous ketamine for late-life treatment-resistant depression: a Bayesian adaptive randomization trial.

Neuropsychopharmacology. 2022;47:1088–95. doi: 10.1038/s41386-021-01242-9

Loo CK, Gálvez V, O'Keefe E, *et al.* Placebo-controlled pilot trial testing dose titration and intravenous, intramuscular and subcutaneous routes for ketamine in depression. *Acta psychiatrica Scandinavica*. 2016;134:48. doi: 10.1111/acps.12572

Loo C, Glozier N, Barton D, *et al.* Efficacy and safety of a 4-week course of repeated subcutaneous ketamine injections for treatment-resistant depression (KADS study): randomised double-blind active-controlled trial. *Br J Psychiatry*. 2023;223:533–41. doi: 10.1192/bjp.2023.79

Milak M.S., Rashid R., Dong Z., *et al.* Assessment of Relationship of Ketamine Dose with Magnetic Resonance Spectroscopy of Glx and GABA Responses in Adults with Major Depression: A Randomized Clinical Trial. *JAMA Netw Open*. 2020;3:e2013211. doi: 10.1001/jamanetworkopen.2020.13211

Murrough JW, Iosifescu DV, Chang LC, *et al.* Antidepressant Efficacy of Ketamine in Treatment-Resistant Major Depression: A Two-Site Randomized Controlled Trial. *Am J Psychiatry*. 2013;170:1134–42. doi: 10.1176/appi.ajp.2013.13030392

Murrough JW, Soleimani L, DeWilde KE, *et al.* Ketamine for rapid reduction of suicidal ideation: a randomized controlled trial. *Psychol Med*. 2015;45:3571–80. doi: 10.1017/S0033291715001506

Ochs-Ross R, Daly EJ, Zhang Y, *et al.* Efficacy and Safety of Esketamine Nasal Spray Plus an Oral Antidepressant in Elderly Patients With Treatment-Resistant Depression—TRANSFORM-3. *Am J Geriatr Psychiatry*. 2020;28:121–41. doi: 10.1016/j.jagp.2019.10.008

Ohtani Y, Tani H, Nomoto-Takahashi K, *et al.* Efficacy and safety of intravenous ketamine treatment in Japanese patients with treatment-resistant depression: A double-blind, randomized, placebo-controlled trial. *Psychiatry Clin Neurosci*. 2024;78:765–75. doi: 10.1111/pcn.13734

Pattanaseri K, Lortrakul J, Jaisin K, *et al.* A randomized controlled pilot study of daily intravenous ketamine over three days for treatment-resistant depression. *BMC Psychiatry*. 2024;24:512. doi: 10.1186/s12888-024-05951-5

Phillips J.L., Norris S., Talbot J., *et al.* Single, repeated, and maintenance ketamine infusions for treatment-resistant depression: A randomized controlled trial. *Am J Psychiatry*. 2019;176:401–9. doi: 10.1176/appi.ajp.2018.18070834

Popova V, Daly EJ, Trivedi M, *et al.* Efficacy and Safety of Flexibly Dosed Esketamine Nasal Spray Combined With a Newly Initiated Oral Antidepressant in Treatment-Resistant Depression: A Randomized Double-Blind Active-Controlled Study. *Am J Psychiatry*. 2019;176:428–38. doi: 10.1176/appi.ajp.2019.19020172

Price RB, Spotts C, Panny B, *et al.* A Novel, Brief, Fully Automated Intervention to Extend the Antidepressant Effect of a Single Ketamine Infusion: A Randomized Clinical Trial. *Am J Psychiatry*. 2022;179:959–68. doi: 10.1176/appi.ajp.20220216

Rodriguez CI, Kegeles LS, Levinson A, *et al.* Randomized controlled crossover trial of ketamine in obsessive-compulsive disorder: proof-of-concept. *Neuropsychopharmacology*. 2013;38:2475. doi: 10.1038/npp.2013.150

Schiweck C., Lutin E., De Raedt W., *et al.* Twenty-Four-Hour Heart Rate Is a Trait but Not State Marker for Depression in a Pilot Randomized Controlled Trial With a Single Infusion of Ketamine. *Front Psychiatry*. 2021;12:696170. doi: 10.3389/fpsyt.2021.696170

Shiroma P.R., Thuras P., Wels J., *et al.* A randomized, double-blind, active placebo-controlled study of efficacy, safety, and durability of repeated vs single subanesthetic ketamine for treatment-resistant depression. *Transl Psychiatry*. 2020;10:206. doi: 10.1038/s41398-020-00897-0

Singh J.B., Fedgchin M., Daly E.J., *et al.* A double-blind, randomized, placebo-controlled, dose-frequency study of intravenous ketamine in patients with treatment-resistant depression. *Am J Psychiatry*. 2016;173:816–26. doi: 10.1176/appi.ajp.2016.16010037

Singh JB, Fedgchin M, Daly E, *et al.* Intravenous Esketamine in Adult Treatment-Resistant Depression: A Double-Blind, Double-Randomization, Placebo-Controlled Study. *Biol Psychiatry*. 2016;80:424–31. doi: 10.1016/j.biopsych.2015.10.018

Smith-Apeldoorn SY, Veraart JKE, Kamphuis J, *et al.* Oral esketamine in patients with treatment-resistant depression: a double-blind, randomized, placebo-controlled trial with open-label extension. *Mol Psychiatry*. 2024;29:2657–65. doi: 10.1038/s41380-024-02478-9

Sos P, Klirova M, Novak T, *et al.* Relationship of ketamine's antidepressant and psychotomimetic effects in unipolar depression. *Neuroendocrinol Lett*. 2013;34:287–93.

Su T-P, Chen M-H, Li C-T, *et al.* Dose-Related Effects of Adjunctive Ketamine in Taiwanese Patients with Treatment-Resistant Depression. *Neuropsychopharmacol*. 2017;42:2482–92. doi: 10.1038/npp.2017.94

Su T-P, Li C-T, Lin W-C, *et al.* A Randomized, Double-Blind, Midazolam-Controlled Trial of Low-Dose Ketamine Infusion in Patients With Treatment-Resistant Depression and Prominent Suicidal Ideation. *Int J Neuropsychopharmacol*. 2023;26:331–9. doi: 10.1093/ijnp/pyad014

Takahashi N, Yamada A, Shiraishi A, *et al.* Efficacy and safety of fixed doses of intranasal Esketamine as an add-on therapy to Oral antidepressants in Japanese patients with treatment-resistant depression: a phase 2b randomized clinical study. *BMC Psychiatry*. 2021;21:526. doi: 10.1186/s12888-021-03538-y

Taylor JH, Landeros-Weisenberger A, Coughlin C, *et al.* Ketamine for social anxiety disorder: A randomized, placebo-controlled crossover trial. *Neuropsychopharmacology*. 2018;43:325–33. doi: 10.1038/npp.2017.194

Tiger M, Veldman ER, Ekman CJ, *et al.* A randomized placebo-controlled PET study of ketamine´s effect on serotonin1B receptor binding in patients with SSRI-resistant depression. *Translational psychiatry*. 2020;10:159. doi: 10.1038/s41398-020-0844-4

Wang S, Deng C-M, Zeng Y, *et al.* Efficacy of a single low dose of esketamine after childbirth for

mothers with symptoms of prenatal depression: randomised clinical trial. *BMJ*. 2024;385:e078218. doi: 10.1136/bmj-2023-078218

102 Xiao C, Zhou J, Li A, *et al.* Esketamine vs Midazolam in Boosting the Efficacy of Oral Antidepressants for Major Depressive Disorder: A Pilot Randomized Clinical Trial. *JAMA Netw Open*. 2023;6:e2328817. doi: 10.1001/jamanetworkopen.2023.28817

103 Yang SQ, Zhou YY, Yang ST, *et al.* Effects of different doses of esketamine intervention on postpartum depressive symptoms in cesarean section women: A randomized, double-blind, controlled clinical study. *J Affect Disord*. 2023;339:333–41. doi: 10.1016/j.jad.2023.07.007

104 Zarate CA, Singh JB, Carlson PJ, *et al.* A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Archives of general psychiatry*. 2006;63:856. doi: 10.1001/archpsyc.63.8.856

105 Zarate CA, Brutsche NE, Ibrahim L, *et al.* Replication of ketamine's antidepressant efficacy in bipolar depression: a randomized controlled add-on trial. *Biological psychiatry*. 2012;71:939. doi: 10.1016/j.biopsych.2011.12.010

106 Zhou Y, Lan X, Wang C, *et al.* Effect of Repeated Intravenous Esketamine on Adolescents With Major Depressive Disorder and Suicidal Ideation: A Randomized Active-Placebo-Controlled Trial. *J Am Acad Child Adolesc Psychiatry*. 2024;63:507–18. doi: 10.1016/j.jaac.2023.05.031

Reduction of Suicide Ideation by Drug over Time