Genetic and Cellular Architecture of Breast Cancer Risk in Multi-Ancestry Studies of 159,297 Cases and 212,102 Controls

James L. Li[1,*], Maria Zanti[2,*], Jacob Williams[3,*], Om Jahagirdar[3], Guochong Jia[4], Alistair Turcan[5], Qiang Hu[6], Jean-Tristan Brandenburg[7,8], Li Yan[6], Weang-Kee Ho[9,10], Jingmei Li[11], José Patricio Miranda[12,13,14], Devika Godbole[3], Julie-Alexia Dias[15], Xiaomeng Zhang[16], Leila Dorling[16], Wenlong Carl Chen[7,8,40], Nicholas Boddicker[17], Ying Wang[18], Alicia Martin[18,19,20], Yan Dora Zhang[21], Joe Dennis[16], Esther M. John[22,23], Gabriela Torres-Mejia[24], Larry Kushi[25,26], Jeffrey Weitzel[27], Susan L. Neuhausen[28], Luis Carvajal-Carmona[29,30], Christopher Haiman[31], Elad Ziv[32], Laura Fejerman[33,34,35], Wei Zheng[4], Dezheng Huo[1], Douglas Easton[16,36], Stephen J. Chanock[3], Nilanjan Chatterjee[37,38], Peter Kraft[3], Montserrat Garcia-Closas[39], Wendy S.W. Wong[3], Kyriaki Michailidou[2], Qianqian Zhu[6], Martin Jinye Zhang[5], Diptavo Dutta[3], Thomas U. Ahearn[3,†], Haoyu Zhang[3,†]

[1] Department of Public Health Sciences, University of Chicago, Chicago, IL, USA

[2] Biostatistics Unit, The Cyprus Institute of Neurology and Genetics, Nicosia, Cyprus

[3] Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

[4] Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, TN, USA

[5] Ray and Stephanie Lane Computational Biology Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA

[6] Department of Biostatistics and Bioinformatics, Roswell Park Comprehensive Cancer Center, Buffalo, NY, USA

[7] Sydney Brenner Institute for Molecular Bioscience, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

[8] Strengthening Oncology Services Research Unit, Faculty of Health Sciences,

University of the Witwatersrand, Johannesburg, South Africa

[9] School of Mathematical Sciences, Faculty of Science and Engineering, University of

Nottingham Malaysia, Semenyih, Selangor, Malaysia

[10] Cancer Research Malaysia, Subang Jaya, Selangor, Malaysia

[11] Genome Institute of Singapore, Agency for Science, Technology and Research

(A*STAR), Singapore

[12] Department of Nutrition, Diabetes and Metabolism, School of Medicine, Pontificia

Universidad Católica de Chile, Santiago, Chile

[13] PhD Program in Epidemiology, Pontificia Universidad Católica de Chile, Santiago,

Chile

[14] Advanced Center for Chronic Diseases, Pontificia Universidad Católica de Chile &

Universidad de Chile, Santiago, Chile

[15] Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA,

USA

[16] Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary

Care, University of Cambridge, Cambridge, UK

[17] Department of Quantitative Health Sciences, Mayo Clinic, Rochester, MN, USA

[18] Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston,

MA, USA

[19] Department of Medicine, Harvard Medical School, Boston, MA, USA

[20] Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard,

Cambridge, MA, USA

[21] Department of Statistics & Actuarial Science, School of Computing and Data Science,

The University of Hong Kong, Hong Kong SAR, China

[22] Departments of Epidemiology & Population Health and of Medicine (Oncology),

Stanford University School of Medicine, Stanford, CA, USA

[23] Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA

[24] Instituto Nacional de Salud Pública, Cuernavaca, Mexico

[25] UC Davis Genome Center, University of California, Davis, Davis, CA, USA

[26] Division of Research, Kaiser Permanente Northern California, Oakland, CA, USA

[27] Division of Precision Prevention, University of Kansas Comprehensive Cancer
Center, Kansas City, KS, USA

[28] Department of Population Sciences, Beckman Research Institute of City of Hope,
Duarte, CA, USA

[29] Department of Biochemistry and Molecular Medicine, School of Medicine, University
of California at Davis, Davis, CA, USA

[30] The Health Equity, Leadership, Science and Community Laboratory, Genome Center,
University of California at Davis, Davis, CA, USA

[31] Department of Preventive Medicine, Norris Comprehensive Cancer Center, Keck
School of Medicine, University of Southern California, Los Angeles, CA, USA

[32] Division of General Internal Medicine, Department of Medicine, University of
California, San Francisco, San Francisco, CA, USA

[33] Department of Public Health Sciences, University of California Davis, Davis, CA, USA

[34] Genome Center, University of California Davis, Davis, CA, USA

[35] UC Davis Comprehensive Cancer Center, University of California Davis, Davis, CA,
USA

[36] Centre for Cancer Genetic Epidemiology, Department of Oncology, University of
Cambridge, Cambridge, UK

[37] Department of Biostatistics, Johns Hopkins University, Baltimore, MD, USA

[38] Department of Oncology, Johns Hopkins School of Medicine, Baltimore, MD, USA

[39] Cancer Epidemiology and Prevention Research Unit, The Institute of Cancer
Research and Imperial College London, London, UK

[40] National Cancer Registry, a Division of the National Institute for Communicable

Diseases, National Health Laboratory Service, Johannesburg, South Africa

[*]These authors equally contributed to the manuscript

[†]Correspondence to: Thomas U. Ahearn (thomas.ahearn@nih.gov), and Haoyu Zhang

(haoyu.zhang2@nih.gov).

SUMMARY

Breast cancer genome-wide association studies (GWAS) have identified over 200

independent genome-wide significant susceptibility markers. However, most studies

have focused on one or two ancestral groups. We examined breast cancer genetic

architecture using GWAS summary statistics from African (AFR), East Asian (EAS),

European (EUR) and Hispanic/Latina (H/L) samples, totaling 159,297 cases and

212,102 controls, comprising the largest multi-ancestry study of breast cancer to date.

The logit-scale heritability of breast cancer ranged from h^2=0.47 (SE = 0.07) in EAS to

AFR h^2=0.61 (SE = 0.10), with no significant differences across ancestries (p=0.63).

The estimated number of susceptibility markers in a sparse normal-mixture effects

model also varied from 4,446 (SE = 3,100) in EAS to 8,308 (SE = 2,751) in AFR, but

differences were not significant across ancestries (p=0.55). Cross-sample genetic

correlations varied, with the strongest correlation between EUR and EAS ($\rho = 0.79$, SE

= 0.08) and weakest between AFR and H/L ($\rho = 0.26$, SE = 0.24). Common variants in

regulatory elements were enriched for genetic association across samples. By

integrating the GWAS summary statistics with the Tabula Sapiens scRNA-seq atlas, we

identified ancestry-shared associations between breast cancer and specific cell types,

including innate immune cells, secretory epithelial cells and stromal cells. Collectively,

these results support a largely shared polygenic architecture of breast cancer across

ancestries, with consistent enrichment of common regulatory variants and convergent

cellular signatures identified through single-cell analyses.

INTRODUCTION

Breast cancer is the most common cancer among women worldwide, affecting over two million individuals annually[1]. Genome-wide association studies (GWAS) have identified more than 200 genome-wide significant susceptibility markers associated with breast cancer risk[2–7]. However, most GWAS have focused on populations of European (EUR) and East Asian (EAS) ancestry samples[2,4–9], while studies in African (AFR) [3,10–13] and Hispanic/Latina (H/L)[14,15] samples are limited, and often with smaller sample sizes. Existing studies have both replicated risk markers identified across ancestry populations, as well as identified population-specific risk markers. Furthermore, polygenic risk score (PRS) models trained in European samples often exhibit lower predictive performance in other genetic ancestry samples[16], emphasizing the need for a broader understanding of breast cancer genetics across different ancestries.

We examined the genetic architecture of breast cancer using GWAS summary statistics from a total of 159,297 cases and 212,102 controls from AFR, EAS, EUR and H/L samples (**Supplementary Figure 1, Supplementary Table 1**). We estimated genome-wide single-nucleotide polymorphism (SNP)-based heritability, the number of susceptibility genetic markers, and assessed genetic correlations across samples. We further investigated enrichment of regulatory elements associated with breast cancer risk, and applied the single-cell disease-relevance score (scDRS+) method to identify cell types implicated in breast cancer across ancestry samples. These findings further refine our understanding of breast cancer genetics and highlight the influence of common variants and regulatory elements across ancestries.

RESULTS

Logit-scale SNP-based heritability

We used linkage-disequilibrium (LD) score regression[17] to estimate the logit-scale SNP-based heritability (h^2), which quantifies the proportion of variance in the log-odds of disease explained by common genetic variants under a logistic regression model (**Methods**). To obtain a robust estimate of the heritability, we conducted sensitivity

analyses by applying various SNP inclusion criteria, including different minor allele

frequency thresholds (MAF > 0.01 or MAF > 0.05), and SNP arrays (HapMap3[18] or

HapMap3 + Multi-Ethnic Genotyping Array (MEGA)[19]). LD scores were computed using

1000 Genomes Project (1000G)[20] Phase 3 for AFR, EAS and EUR samples, and

Admixed American (AMR) as reference for H/L samples. For AFR, we additionally

generated estimates using a subset of 2,731 controls from the African Ancestry Breast

Cancer Genetics (AABCG) Consortium.

Heritability estimates for EUR and EAS samples remained consistent across different

SNP inclusion criteria hat varied by minor allele frequency thresholds (MAF > 0.01 vs.

MAF > 0.05) and SNP sets (HapMap3 vs. HapMap3 + MEGA) (**Supplementary Table**

2). Thus, we adopted the most comprehensive SNP set (MAF > 0.01 and HapMap3 plus

MEGA SNPs) for our final analyses. In contrast, heritability estimates for AFR varied

significantly depending on the LD reference panel (h^2 =0.27, SE=0.11 to h^2 =0.61,

SE=0.10; **Supplementary Table 2)**. Given that the GWAS summary statistics for AFR

were primarily from U.S. samples (85.3%)[12], while AFR in 1000G are mostly from

African continental samples, AABCG likely provides more appropriate reference for LD

estimation in this context. For H/L, heritability estimates were less stable due to the

smaller GWAS sample size, leading to larger standard errors. To enhance robustness,

we applied the more conservative MAF > 0.05 threshold.

Across ancestry samples, SNP-based heritability estimates ranged from EAS h^2=0.466

(SE=0.066) to AFR h^2=0.614 (SE=0.095), with EUR h^2=0.501 (SE=0.050) and H/L

h^2=0.588 (SE=0.360) showing intermediate values (**Table 1**). While point estimates

varied modestly, we found no statistically significant evidence of heterogeneity in SNP-

based heritability across ancestry samples (Cochran's Q = 1.72, p = 0.63,

Supplementary Figure 2). Overall, these findings suggest common susceptibility

variants moderately contribute to breast cancer risk across samples. The slightly higher

point estimate in AFR could reflect the broader range of genetic variation captured by a

larger set of well-imputed SNPs (n=1,628,778) (**Table 1**).

The polygenicity of breast cancer

Breast cancer exhibits a high degree of polygenicity across populations. Using the GENESIS package[21,22], we modeled the distribution of marker effect sizes for AFR, EAS and EUR samples as a sparse mixture of normal. Due to a limited sample size, the model for H/L did not converge, precluding a reliable estimate. Across the three analyzed samples, marker effect sizes showed no evidence of inflation (**Supplementary Figure 3**). The estimated polygenicity (number of markers with non-zero effects) was highest in AFR (8,308, SE=2,751), followed by EUR (5,235, SE=1,191) and EAS (4,446, SE=3,100, **Supplementary Table 3**). We found no statistically significant heterogeneity in polygenicity across samples (Cochran's Q = 1.20, p = 0.55), that observed differences are consistent with sampling variation. These findings support a broadly shared polygenic basis of breast cancer across ancestries, with modest differences that may reflect sample-specific power or LD structure.

We used GENESIS to project the proportion of heritability explained and the accuracy of polygenic risk scores (PRS) at increasing GWAS sample sizes (**Figure 1A, Supplementary Table 4**). Polygenic risk scores evaluated in GENESIS were based on the clumping and thresholding (CT-PRS) approach[23,24], applied independently to each sample using ancestry-specific GWAS summary statistics and corresponding LD reference panels. The proportion of genetic variance explained was defined as the variance captured by the CT-PRS divided by the logit-scale heritability of common variants. At a GWAS sample size of 100,000 cases and 100,000 controls, the proportion of genetic variance explained was 26.2% in AFR, 39.4% in EAS, and 38.6% in EUR (**Supplementary Table 4**). Projected area under the curve (AUC) values for CT-PRS followed a similar trend. At a sample size of 100,000 cases and 100,000 controls, the expected AUC was 61.1% for AFR, 63.4% for EAS and 62.1% for EUR (**Figure 1B, Supplementary Table 4**). Asymptotically (i.e., at infinite sample size), the estimated maximum AUCs for CT-PRS were 71% for AFR, and 71% for EAS, and 69% for EUR (**Supplementary Table 3**).

Cross-sample genetic correlations of breast cancer

We next assessed cross-sample genetic correlations of breast cancer using *Popcorn*[25], which estimates genetic effect correlations accounting for LD differences across ancestries. Using variants from the HapMap3 panel, we identified the strongest genetic correlations between EUR and EAS (ρ: 0.79, SE: 0.08) and EUR and H/L (ρ: 0.68, SE: 0.21), indicating substantial shared genetic architecture between these samples (**Figure 2A, Supplementary Table 5**). In contrast, genetic correlations involving AFR were consistently lower, with estimates ranging from AFR and H/L ($\rho = 0.26$, SE = 0.24) to AFR and EUR ($\rho = 0.42$, SE = 0.14), suggesting greater divergence in effect sizes of variants tagging in AFR. We further evaluated whether expanding the variant set to include both HapMap3 and MEGA array variants altered correlation patterns and the estimates remained broadly consistent (**Figure 2B, Supplementary Table 5**).

Genomic enrichment of breast cancer heritability

To investigate the regulatory landscape underlying breast cancer heritability, we estimated the proportion of SNP-based heritability attributable to 73 functional genomic annotations using stratified LD-score regression[26,27] (**Methods**, **Supplementary Table 6**). These annotations, based on the baseline-LD v2.2 model, include a curated set of coding, regulatory, and evolutionary genomic features previously shown to capture trait heritability[26–28]. Annotations definitions and data sources are described in the **Methods**. In the EUR sample, several annotations showed strong, statistically significant enrichment. The highest enrichment was found in ancient promoter regions (32.42-fold, $p = 1.98 \times 10^{-4}$), followed by transcription factor binding sites (TFBS) (5.79-fold, $p = 9.16 \times 10^{-5}$), and the promoter-associated histone mark H3K4me3 (4.9-fold, $p = 8.56 \times 10^{-5}$). We also observed significant enrichment in super-enhancers (3.01-fold, $p = 3.73 \times 10^{-11}$), and enhancer-associated marks H3K4me1 (2.52-fold, $p = 1.28 \times 10^{-7}$) and H3K27ac (2.16-fold, $p = 2.25 \times 10^{-10}$), highlighting the contribution of regulatory elements to genetic risk. In the EAS sample, the only annotation reaching statistical significance after Bonferroni correction ($p < 6.84 \times 10^{-4}$) was super-enhancers (2.73-fold, $p = 2.6 \times 10^{-7}$). In the AFR sample, significant enrichment was found for H3K27ac (2.15-fold, $p = 2.59 \times 10^{-4}$). No statistically significant enrichment was observed in the H/L sample. Patterns of

enrichment for annotations that were significantly enriched in at least one ancestry

sample were generally consistent across ancestries (**Supplementary Figure 4**). A

heterogeneity test across samples for these five annotations revealed no statistically

significant differences after Bonferroni correction.

Cell-type specific associations with breast cancer risk

To identify cell types associated with breast cancer risk across ancestries, we applied

scDRS+[29][30], to GWAS summary statistics from each ancestry and the Tabula Sapiens

scRNA-seq atlas[31] (133 cell types across 24 tissues). scDRS+ computes disease

relevance score for individual cells based on expression of GWAS-prioritized genes,

then aggregates these scores to test cell-type enrichments. For robustness, we

prioritized cell types showing associations in more than three ancestry samples for

robustness, reporting both fine-mapped (**Figure 3A, Supplementary Table 7**) and

marginal results (**Figure 3B; Supplementary Table 8**).

First, we observed associations in several innate immune cell types, including classical

monocytes (false discovery rate (FDR) <0.05 in EAS and EUR; FDR<0.1 in H/L and

AFR, **Figure 3A**) and neutrophils (FDR<0.05 in EAS; FDR<0.1 in H/L and EUR, **Figure**

3A), consistent with the established role of myeloid cells in breast cancer biology and

the tumor microenvironment[32–37]. Second, we detected associations for lacrimal gland

functional unit cells (FDR<0.05 in EAS and EUR; FDR<0.1 in H/L and AFR, **Figure 3A**)

and skeletal muscle satellite stem cells (FDR<0.05 in EAS and EUR; FDR<0.1 in H/L,

Figure 3A). These associations are unlikely to reflect direct etiological cell types for

breast cancer but may instead arise from shared epithelial or stromal regulatory

programs in mammary tissue. For example, luminal epithelial cells of mammary gland

were non-significant in fine-mapping but showed marginal association (FDR<0.05 in

AFR; FDR≈0.07-0.10 for EAS, EUR, and H/L, **Supplementary Table 8**), while

fibroblasts of breast were likewise non-significant in fine-mapping but displayed

marginal associations in EAS (FDR=0.049), H/L (FDR=0.02), and AFR (FDR=0.008),

with a weaker signal in EUR (FDR=0.11). Third, these findings are supported by

consistent marginal association signals across all samples (**Figure 3B**), which we

included to mitigate the interpretative challenges inherent in fine-mapping correlated cell
populations.

To assess shared cellular signatures across ancestry samples, we examined the
correlation of marginal scDRS+ association scores for the key cell types, neutrophils,
classical monocytes, and skeletal muscle satellite stem cells. The association scores for
these cell types were highly correlated between ancestry pairs (**Supplementary Figure**
5), suggesting conserved cellular mechanisms underlying genetic susceptibility. UMAP
visualizations further demonstrated overlapping disease-relevant cell clusters across
samples (**Figure 3C, D**).

DISCUSSION

We present a compressive cross-ancestry evaluation of the genetic architecture of
breast cancer using large-scale GWAS summary statistics from AFR, EAS, EUR and
H/L samples. With 220,498 cases and 200,182 controls, this is the largest such
evaluation to date. Our results support a largely shared polygenic architecture of breast
cancer across ancestries, with consistent enrichment of common regulatory variants
and convergent cellular signatures identified through single-cell analyses. While point
estimates of heritability and polygenicity varied across samples, we did not detect statistically
significant heterogeneity, suggesting that observed differences likely reflect sampling variation,
LD structure, or reference panel choice rather than fundamental biological divergence.

Breast cancer exhibited moderate logit-scale SNP-based heritability across all groups,
with the highest point estimate observed in AFR. These estimates were robust in EUR
and EAS, but more sensitive to LD reference panel choice in AFR, where using sample-
matched LD from AABCG controls significantly increased heritability compared to using
AFR from 1000G (**Supplementary Table 2**). This highlights the importance of
appropriate ancestry-specific LD modeling in admixed populations[38–40]. While the point
estimates varied, our formal test of heterogeneity found no statistically significant
differences in heritability across samples (p=0.64). These results are broadly consistent
with prior reports based on subsets of our underlying data. For example, Michailidou et
al. (2017)[7], using primarily OncoArray data (a large component of our EUR sample),

estimated that common variants explained ~41% of the familial relative risk,

corresponding to a logit-scale heritability of approximately 0.568, closely matching our

EUR estimate of 0.501. Likewise, our AFR estimate of 0.614 is comparable to the logit-

scale heritability (referred to as frailty-scale) of 0.667 reported in the recent African

ancestry sample breast cancer GWAS[12], which used the same summary statistics but a

different LD reference panel. While heritability has been estimated in prior studies,

these have largely focused on individual ancestry groups and used varying

methodologies; to our knowledge, our study provides the first systematic, cross-

ancestry comparison of logit-scale SNP-based heritability using harmonized models.

We focused on the logit scale, as it provides an interpretable and robust estimate of

genetic variance explained by GWAS variants and connects directly to familial risk[41],

without requiring assumptions about disease prevalence.

We also observed a high degree of polygenicity across ancestry samples. The

estimated number of susceptibility markers with non-zero effects in a sparse mixture

model ranged from 4,446 in EAS to 8,308 in AFR. While these differences may reflect

true biological variation, such as lower LD in African genomes allowing more markers to

be distinguished statistically, our heterogeneity test found no significant differences in

polygenicity across samples (p=0.55). Interestingly, the AFR sample also had the

highest SNP-based heritability estimate, suggesting a potentially larger contribution of

common variants to risk. However, PRS performance in AFR remained lower compared

to EUR and EAS at comparable sample sizes[42]. This finding is consistent with our

projections (**Figure 1B**), which demonstrate that PRS accuracy is not only a function of

total heritability, but also of the effect-size distribution, i.e., the degree of polygenicity,

and the LD structure of the population. Thus, the higher heritability in AFR may be offset

by greater polygenicity and LD complexity, both of which reduce the efficiency of risk

prediction models based on common variants.

Genetic correlations were strongest between EUR and EAS ($\rho = 0.79$), and weakest for

comparison involving AFR, reflecting ancestral differences in LD structure and variant

tagging. These findings may reflect known patterns of human migration and population

divergence indicated from populations genetics, including an initial divergence in line

with the "Out-of-Africa" migration model, with subsequent and more recent separation

between European, East Asian, and American populations[43]. The moderately strong

correlation between EUR and H/L ($\rho = 0.68$) likely reflects the substantial European

ancestry present in many H/L individuals. However, it is important to note that this

average may obscure heterogeneity within H/L populations: individuals with higher

proportions of Indigenous American ancestry would be expected to show lower genetic

correlation with EUR populations. The H/L summary statistics analyzed here were

derived from both U.S.-based cohorts (e.g., NC-BCFR/SFBCS, RPGEH, MEC) and

Latin American studies (COLUMBUS and CAMA), encompassing a mix of Mexican,

Central American, and South American individuals with variable proportions of

European, Indigenous American, and African ancestry. Similarly, the AFR GWAS data

primarily reflect African-American individuals with admixed West African and European

ancestry. As such, the estimated genetic correlations represent average effect-size

sharing across ancestrally diverse groups and may mask finer-scale heterogeneity.

Future analyses incorporating local ancestry or ancestry-stratified GWAS may help to

disentangle these layered genetic relationships more precisely.

Stratified heritability enrichment revealed consistent contributions from regulatory

elements, including promoters, enhancers, and histone marks (e.g., H3K27ac,

H3K4me1), across ancestry samples. While some differences in the strength of

enrichment were observed, heterogeneity tests did not identify statistically significant

variation in enrichment patterns across samples. In the H/L sample, no annotations

reached statistical significance, likely due to limited GWAS sample size and reduced

power to detect enrichment.

Integrating GWAS with single-cell transcriptomic data via scDRS+[29], we identified

consistent enrichments in classical monocytes, neutrophils, lacrimal gland functional

unit cell, and skeletal muscle satellite stem cells for breast cancer-associated gene

expression across ancestries. These findings suggest that genetic risk variants may

exert their effects through immune, epithelial, and stromal cells, such as inflammation,

innate immune responses, or tissue repair, rather than solely within breast epithelial

cells. Monocyte and neutrophil associations align with prior evidence of myeloid roles in

tumor inflammation and progression [32–37]. The lacrimal gland functional unit cells and

skeletal muscle satellite stem cell associations are unlikely to reflect direct etiological

cell types for breast cancer but may instead arise from shared epithelial or stromal

regulatory programs in mammary tissue , such as those observed for luminal epithelial

cells of mammary gland (FDR<0.05 in EAS; FDR<0.2 in EUR and AFR) and fibroblasts

of breast (non-significant in fine-mapping but showing marginal association at

FDR<0.05 in EAS,H/L and AFR). Alternatively, they could support emerging roles for

muscle or stromal niches in metastatic reactivation [44]. While scDRS+ does not directly

identify the tissue of origin, it provides a hypothesis-generating framework that links

genetic risk to specific cellular contexts based on expression profiles. The cross-

ancestry consistency reinforces shared biological pathways underlying breast cancer

risk in diverse populations.

This is the largest cross-ancestry investigation of breast cancer architecture; however,

our study was limited by relatively small sample sizes particularly in non-European

samples. Our findings highlight the value of large samples from diverse populations in

breast cancer GWAS, which enables more powerful modeling of the underlying

polygenic risk by leveraging similar and distinct linkage disequilibrium patterns across

populations. Notably, important efforts are being made to increase samples size of

breast cancer GWAS among women of Hispanic/Latina descent by the Latin American

Genomics of Breast Cancer Consortium (LAGENO-BC) and among women of African

descent by AABCG[12]. Moreover, the Confluence Project is building a large, diverse

research resource of approximately 300,000 breast cancer cases and 300,000 controls

by coordinating collaborations between LAGENO-BC, AABCG[12], BCAC, the

Consortium of Investigators of Modifiers of BRCA1/2 (CIMBA), the Male Breast Cancer

Genetics Consortium (MERGE), and other breast cancer studies and clinical trials, that

will help support and lead to an improved understanding of breast cancer etiology, and

improvements in risk stratification and prevention across diverse populations. In

summary, our study refines the understanding of breast cancer genetics across

populations and highlights the value of multi-ancestry studies of breast cancer for

gaining better insights in breast cancer etiology and polygenic risk prediction and

stratification.

Table 1. Sample-specific estimates of the logit-scale SNP-based heritability

Sample	Cases	Controls	MAF cutoff	LD Score	No. of SNPs	Heritability[e]
AFR[c]	18,034	22,104	MAF > 0.01	AABCG	1,628,778	0.614 (0.095)
EAS[b]	20,393	86,329	MAF > 0.01	1000G	1,189,628	0.466 (0.066)
EUR[a]	118,474	96,201	MAF > 0.01	1000G	1,415,774	0.501 (0.050)
H/L[d]	2,396	7,468	MAF > 0.05	1000G	1,138,603	0.588 (0.360)

Summary statistics from [a]the Breast Cancer Association Consortium (BCAC) (iCOGs and OncoArray) [b] BCAC (OncoArray) and Biobank Japan [c]Summary statistics from the African Ancestry Breast Cancer Genetic (AABCG) consortium [d] NC-BCFR/SFBCS (Northern California Breast Cancer Family Registry/ San Francisco Bay Area Breast Cancer Study), RPGEH (Research project on genes environment and health), MEC (Multiethnic cohort), COLUMBUS and CAMA (Cancer de mama) studies/consortia. Variants included in the analyses were on HapMap3 + Multi-Ethnic Genotyping Array (MEGA). [e] logit-scale heritability (standard error). The logit-scale heritability (also known as frality-scale heritability) is defined as $\sigma^2_{GWAS} = Var(\sum_{m=1}^{M} \beta_m G_m)$, where G_m is the standardized genotype for the mth SNP, β_m is the true log odds ratio for the mth SNP and M is the total number of causal SNPs among the GWAS variants. AFR, African; H/L, Hispanic and Latina; EAS, East Asian; EUR, European; MAF, minor allele frequency; LD, linkage disequilibrium; SNPs, single nucleotide polymorphisms.

Figure 1. Projected genetic variance explained and predictive performance of clumping and thresholding (CT) polygenic risk scores (PRSs) across samples at varying GWAS sample sizes. A) Estimated proportion of GWAS heritability explained by CT PRS, as a function of total GWAS sample sizes, using the GENESIS framework. The proportion was calculated as the variance explained by the PRS divided by the logit-scale heritability of common variants. B) Projected area under the curve (AUC) for CT PRS models at different sample sizes. Detailed projection of explained genetic variance and AUC at different sample size are provided in **Supplementary Table 4**. The red dashed line indicates a total sample size of 200,000 individuals (100,000 cases and 100,000 controls). Abbreviations: AFR, African; EAS, East Asian; EUR, European.

Figure 2. Heatmaps for cross-sample genetic correlations of breast cancer estimated by POPCORN. for A) HapMap3 variants and B) HapMap3 and Multi-Ethnic Genotyping Array variants combined. <u>Note</u>: African, AFR; Hispanic/Latina, H/L; East Asian, EAS; European, EUR.

Figure 3: scDRS+ results on Tabula Sapiens. One asterisk indicates FDR<0.1, two

asterisks indicates FDR<0.05. A) Fine-mapped associations from scDRS+, showing

FDR values for cell types with at least one ancestry having FDR < 0.05; all shown

associations are also marginally significant. B) Marginal associations from scDRS+,

showing FDR values for cell types with associations across all traits. C) MAP of the cell

types shown in panel A. D) UMAPs of scDRS+ scores across ancestries (EAS, EUR,

AFR, AMR) for the same cell types as in C. Red indicates higher disease relevance.

METHODS

Study samples and data collection

We conducted analyses using breast cancer GWAS summary statistics from large

genetic consortia focused on AFR, EAS, EUR and H/L samples, with a total of 159,297

cases and 212,102 controls across large genetic consortia. Sample labels (AFR, EAS,

EUR, H/L) were based on published descriptions of contributing GWAS. For AFR, data

were obtained from AABCG, with 18,034 cases and 22,104 controls[12]. For EAS, we

conducted a fixed-effect meta-analysis to combine the GWAS summary statistics

generated from the BCAC[7] and Biobank Japan[45], comprising a total of 20,393 cases

and 86,329 controls. For EUR, we used summary statistics from BCAC (iCOGs and

OncoArray) with a combined sample size of 118,474 cases and 96,201 controls[4]. H/L

ancestry summary statistics were derived from a meta-analysis of five studies[15]: NC-

BCFR/SFBCS (Northern California Breast Cancer Family Registry/ San Francisco Bay

Area Breast Cancer Study), RPGEH (Research Program on Genes, Environment and

Health), MEC (Multiethnic cohort), COLUMBUS and CAMA (Cancer de mama) including

2,396 cases and 7,468 controls. **Supplementary Table 1** summarizes the studies

included in the analyses. Processing of meta-analyzed GWAS summary statistics for

subsequent analysis is detailed in **Supplementary Figure 1**.

Sources of genetic data for LD score construction

We constructed ancestry-specific LD score reference panels using unrelated individuals

from Phase 3 of the 1000G[20], excluding first- and second-degree relatives as well as

individuals with first-cousin-level relatedness due to inbreeding[46]. For the EUR, EAS,

and H/L samples, LD scores were computed using 484 EUR, 480 EAS, and 317 AMR

individuals in the 1000G reference panel, respectively. For the AFR sample, we

estimated LD scores using a set of 2,730 AFR women controls from the AABCG

consortium. As a sensitivity analysis, we also generated LD scores using 575 AFR

individuals from 1000G for comparison. The LD scores were conducted using LDSC

with a LD window of 1MB using the flag "--l2 --ld-window-kb 1000".

Estimating ancestry-specific heritability of breast cancer

We estimated the logit-scale sample-specific SNP-based heritability for overall breast cancer risk using LD score regression[17] and using the ancestry-sample-specific GWAS summary statistics. On the logit scale, heritability is defined as: $\sigma_{GWAS}^2 = Var(\sum_{m=1}^{M} \beta_m G_m)$, where G_m is the standardized genotype for the mth SNP, β_m is the underlying log odds ratio, and M is the total number of causal SNPs among the GWAS variants. This quantity, also referred to as frailty-scale heritability, represents the variance in log-odds of disease explained by common genetic variants. To estimate the logit-scale heritability using LD-score regression, we computed the effective sample size as follows:

$$N_j = \left(2s_j^2 p_j (1 - p_j)\right)^{-1}$$

where s_j is the standard error of the log odds ratio, and p_j is the effect allele frequency for variant j. For imputed variants, we adjusted the effective sample size to account for imputation uncertainty:

$$N_j = r_j^2 \left(2s_j^2 p_j (1 - p_j)\right)^{-1}$$

where r_j^2 is the imputation quality score for variant j. To reduce the influence of poorly powered variants, we excluded those with effective sample sizes in the lowest 10% quantile and retained only the top 90% for heritability analysis. To assess the robustness of our heritability estimates, we conducted sensitivity analyses using varying SNP inclusion criteria, including: 1) using variants present in HapMap3 alone versus variants in HapMap3 plus the MEGA array, and 2) varying the MAF thresholds of between MAF > 0.01 or MAF > 0.05.

Assessing polygenicity of breast cancer across samples

To assess the polygenicity of genetic effects of breast cancer risk within each sample, we utilized the GENESIS framework[21,22], which models GWAS summary statistics under a three-component mixture model. Analyses were constrained to variants with MAF > 0.05 and in HapMap3 panel, following the guidance from GENESIS. For EUR and EAS

samples, we used the precomputed LD provided by GENESIS. For the AFR and H/L

samples, we constructed ancestry-specific LD-scores using individuals from the 1000G

AMR samples, and the 2730 AA controls from AABCG, as described in the heritability

analysis section. Following GENESIS's guidance, LD scores were computed by defining

tagging variants as those within a 1Mb window size and with an estimated LD

coefficient r^2 above 0.1. GENESIS models the distribution of genetic marker effect sizes

using a mixture of three components, including two normal distributions for non-null

genetic markers with distinct variance components. This allows the model to capture

both highly polygenic background signals and a subset of genetic markers with larger

effect sizes. From the fitted model, we estimated the total number of susceptibility

genetic markers with non-zero effect sizes. We also calculated the proportion of logit-

scale heritability explained by a CT-PRS. The CT-PRS[23,24] is a commonly used method

that selects approximately independent variants through LD clumping and optimizes p-

value thresholds to maximize predictive performance. Finally, we projected the

expected AUC for CT-PRS at varying GWAS sample sizes based on the model

estimates.

Examining genetic correlations of breast cancer risk between samples

To estimate cross-sample genetic correlations in breast cancer risk, we utilized the

Popcorn package[25]. Cross-ancestry LD reference panels were built from the same

ancestry-specific sources used in heritability analyses. We applied a MAF filter >0.01 for

EUR, EAS, and AFR samples, and >0.05 for the H/L sample to ensure robustness in LD

estimation. We then utilized the default settings in Popcorn to compute pairwise, cross-

sample genetic correlations for 1) variants in the HapMap3 reference panel and 2) the

combined set of HapMap3 and MEGA array variants.

Genomic enrichment analysis using stratified LD-score regression

We used S-LDSC to estimate the contribution of functional genomic annotations to

SNP-based heritability of breast cancer[26,27]. S-LDSC models GWAS test statistics as a

function of LD to variants in annotated regions, allowing partitioning of heritability while

accounting for LD structure and overlapping annotations. We used 73 binary

annotations from the baseline-LD v2.2 model[26–28], which includes regulatory, coding,

and evolutionary features compiled from ENCODE, Roadmap Epigenomics, and

comparative genomics data. Key annotations discussed in the Results include:

1. Ancient promoters: Promoter elements whose sequence aligns to species that

diverged before the placental–marsupial split, representing deeply conserved regulatory

regions[28].

2. TFBS: Genomic regions identified from aggregated ENCODE ChIP-seq[47] and DNase

I footprinting data[48], representing sites of transcription factor occupancy.

3. H3K4me3: A histone modification marking active promoter regions, aggregated

across tissues from ENCODE[47] and Roadmap Epigenomics[49].

4. H3K4me1: A mark associated with primed or poised enhancers[27].

5. H3K27ac: A mark of active enhancers and promoters[27].

6. Super-enhancers: Large regulatory domains composed of clustered enhancers with

high H3K27ac signal intensity[50].

We restricted analyses to SNPs from the HapMap3 panel with minor allele

frequency >5%. For EUR and EAS, we used precomputed LD scores

(https://zenodo.org/records/10515792). For AFR and H/L, we computed ancestry-

specific LD scores using reference samples from the AABCG consortium (AFR) and

1000 Genomes AMR (H/L), combining annotations from EUR and EAS sources.

Enrichment was defined as the proportion of heritability explained by SNPs in each

annotation divided by the proportion of SNPs overlapping that annotation.

Cell-type specific associations analysis using scDRS+

To identify cell types associated with breast cancer risk, we applied scDRS+[29,30], an

improved version of scDRS[30] with enhanced accuracy through single-cell imputation

and modeling of associations among related cell populations, to GWAS summary

statistics from each ancestry alongside the Tabula Sapiens scRNA-seq atlas[31] (27,051

cells from 133 cell types across 24 tissues; 50 cell types with >100 cells). This included

580 cells from the mammary tissue (cell types with >100 cells: luminal epithelial cells of

mammary gland, T cells, fibroblast of breast), and immune cell types (cell types

with >300 cells across all tissues: macrophage, CD8-positive, alpha-beta T cell, CD4-

positive, alpha-beta T cell, memory B cell, monocyte, plasma cell, B cell, T cell, classical

monocyte, NK cell, innate lymphoid cell, immune cell, CD8-positive, alpha-beta memory

T cell, neutrophil, CD4-positive, alpha-beta memory T cell, naive thymus-derived CD4-

positive, alpha-beta T cell).

To enhance data quality, we restricted analyses to fluorescence-activated cell-sorting

cells. For each ancestry, we incorporated the top 1,000 MAGMA-prioritized genes[51] into

the scDRS+ framework to calculate cell-level disease relevance scores. We first

identified marginally associated cells using an FDR threshold of 0.1 (analogous to

scDRS), followed by joint fine-mapping to identify cells associated with breast cancer

across ancestries. Cell-type-disease enrichments were tested by aggregating cell-level

scores (marginal and joint) following the scDRS pipeline[30].

To evaluate cross-sample concordance, we calculated the correlation of scDRS+

marginal scores between each pair of ancestries across all cells in a given cell type. We

compared these to an empirical null distribution derived from control scores to obtain p-

values for significant nonzero correlation. For all analyses, variant sets were restricted

to HapMap3 + MEGA SNPs, using MAF > 0.01 for AFR, EAS, and EUR, and MAF >

0.05 for H/L.

REFERENCES

1. Sung, H., Ferlay, J., Siegel, R.L., Laversanne, M., Soerjomataram, I., Jemal, A., and Bray, F. (2021). Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J. Clin. *71*, 209–249. https://doi.org/10.3322/CAAC.21660.

2. Jia, G., Ping, J., Shu, X., Yang, Y., Cai, Q., Kweon, S.S., Choi, J.Y., Kubo, M., Park, S.K., Bolla, M.K., et al. (2022). Genome- and transcriptome-wide association studies of 386,000 Asian and European-ancestry women provide new insights into breast cancer genetics. Am. J. Hum. Genet. *109*, 2185–2195. https://doi.org/10.1016/J.AJHG.2022.10.011.

3. Adedokun, B., Du, Z., Gao, G., Ahearn, T.U., Lunetta, K.L., Zirpoli, G., Figueroa, J., John, E.M., Bernstein, L., Zheng, W., et al. (2021). Cross-ancestry GWAS meta-analysis identifies six breast cancer loci in African and European ancestry women. Nat. Commun. *12*. https://doi.org/10.1038/S41467-021-24327-X.

4. Zhang, H., Ahearn, T.U., Lecarpentier, J., Barnes, D., Beesley, J., Qi, G., Jiang, X., O'Mara, T.A., Zhao, N., Bolla, M.K., et al. (2020). Genome-wide association study identifies 32 novel breast cancer susceptibility loci from overall and subtype-specific analyses. Nat. Genet. *52*, 572–581. https://doi.org/10.1038/S41588-020-0609-2.

5. Michailidou, K., Beesley, J., Lindstrom, S., Canisius, S., Dennis, J., Lush, M.J., Maranian, M.J., Bolla, M.K., Wang, Q., Shah, M., et al. (2015). Genome-wide association analysis of more than 120,000 individuals identifies 15 new susceptibility loci for breast cancer. Nat. Genet. *47*, 373–380. https://doi.org/10.1038/ng.3242.

6. Michailidou, K., Hall, P., Gonzalez-Neira, A., Ghoussaini, M., Dennis, J., Milne, R.L., Schmidt, M.K., Chang-Claude, J., Bojesen, S.E., Bolla, M.K., et al. (2013). Large-scale genotyping identifies 41 new loci associated with breast cancer risk. Nat. Genet. *45*, 353–361. https://doi.org/10.1038/ng.2563.

7. Michailidou, K., Lindström, S., Dennis, J., Beesley, J., Hui, S., Kar, S., Lemaçon, A., Soucy, P., Glubb, D., Rostamianfar, A., et al. (2017). Association analysis identifies 65 new breast cancer risk loci. Nature *551*, 92–94. https://doi.org/10.1038/nature24284.

8. Shu, X., Long, J., Cai, Q., Kweon, S.S., Choi, J.Y., Kubo, M., Park, S.K., Bolla, M.K., Dennis, J., Wang, Q., et al. (2020). Identification of novel breast cancer susceptibility loci in meta-analyses conducted among Asian and European descendants. Nat. Commun. *11*, 1–9. https://doi.org/10.1038/s41467-020-15046-w.

9. Han, M.R., Long, J., Choi, J.Y., Low, S.K., Kweon, S.S., Zheng, Y., Cai, Q., Shi, J., Guo, X., Matsuo, K., et al. (2016). Genome-wide association study in East Asians identifies two novel breast cancer susceptibility loci. Hum. Mol. Genet. *25*, 3361. https://doi.org/10.1093/HMG/DDW164.

10. Feng, Y., Stram, D.O., Rhie, S.K. yong, Millikan, R.C., Ambrosone, C.B., John, E.M., Bernstein, L., Zheng, W., Olshan, A.F., Hu, J.J., et al. (2014). A comprehensive examination of breast cancer risk loci in African American women. Hum. Mol. Genet. *23*, 5518–5526. https://doi.org/10.1093/HMG/DDU252,.

586   11.   Huo, D., Feng, Y., Haddad, S., Zheng, Y., Yao, S., Han, Y.J., Ogundiran, T.O.,
587   Adebamowo, C., Ojengbede, O., Falusi, A.G., et al. (2016). Genome-wide association
588   studies in women of African ancestry identified 3q26.21 as a novel susceptibility locus for
589   oestrogen receptor negative breast cancer. Hum Mol Genet. *25*, 4835–4846.
590   https://doi.org/10.1093/HMG/DDW305.

591   12.   Jia, G., Ping, J., Guo, X., Yang, Y., Tao, R., Li, B., Ambs, S., Barnard, M.E., Chen, Y.,
592   Garcia-Closas, M., et al. (2024). Genome-wide association analyses of breast cancer in
593   women of African ancestry identify new susceptibility loci and improve risk prediction.
594   Nat. Genet. *56*, 819–826. https://doi.org/10.1038/s41588-024-01736-4.

595   13.   Hayat, M., Chen, W.C., Babb de Villiers, C., Hyuck Lee, S., Curtis, C., Newton, R.,
596   Waterboer, T., Sitas, F., Bradshaw, D., Muchengeti, M., et al. (2025). Genome-wide
597   association study identifies common variants associated with breast cancer in South
598   African Black women. Nat. Commun. *16*, 3542. https://doi.org/10.1038/S41467-025-
599   58789-0,.

600   14.   Hoffman, J., Fejerman, L., Hu, D., Huntsman, S., Li, M., John, E.M., Torres-Mejia, G.,
601   Kushi, L., Ding, Y.C., Weitzel, J., et al. (2019). Identification of novel common breast
602   cancer risk variants at the 6q25 locus among Latinas 06 Biological Sciences. Breast
603   Cancer Res. *21*. https://doi.org/10.1186/S13058-018-1085-9,.

604   15.   Fejerman, L., Ahmadiyeh, N., Hu, D., Huntsman, S., Beckman, K.B., Caswell, J.L.,
605   Tsung, K., John, E.M., Torres-Mejia, G., Carvajal-Carmona, L., et al. (2014). Genome-
606   wide association study of breast cancer in Latinas identifies novel protective variants on
607   6q25. Nat. Commun. *5*, 1–8. https://doi.org/10.1038/ncomms6260.

608   16.   Roberts, E., Howell, S., and Evans, D.G. (2023). Polygenic risk scores and breast cancer
609   risk prediction. Breast *67*, 71–77. https://doi.org/10.1016/j.breast.2023.01.003.

610   17.   Bulik-Sullivan, B.K., Loh, P.-R., Finucane, H.K., Ripke, S., Yang, J., Patterson, N., Daly,
611   M.J., Price, A.L., and Neale, B.M. (2015). LD Score regression distinguishes confounding
612   from polygenicity in genome-wide association studies. Nat. Genet. *47*, 291–295.
613   https://doi.org/10.1038/ng.3211.

614   18.   Altshuler, D.M., Gibbs, R.A., Peltonen, L., Schaffner, S.F., Yu, F., Dermitzakis, E.,
615   Bonnen, P.E., De Bakker, P.I.W., Deloukas, P., Gabriel, S.B., et al. (2010). Integrating
616   common and rare genetic variation in diverse human populations. Nature *467*, 52–58.
617   https://doi.org/10.1038/NATURE09298.

618   19.   Bien, S.A., Wojcik, G.L., Zubair, N., Gignoux, C.R., Martin, A.R., Kocarnik, J.M., Martin,
619   L.W., Buyske, S., Haessler, J., Walker, R.W., et al. (2016). Strategies for Enriching
620   Variant Coverage in Candidate Disease Loci on a Multiethnic Genotyping Array. PLoS
621   One *11*, 167758. https://doi.org/10.1371/JOURNAL.PONE.0167758.

622   20.   Auton, A., Abecasis, G.R., Altshuler, D.M., Durbin, R.M., Bentley, D.R., Chakravarti, A.,
623   Clark, A.G., Donnelly, P., Eichler, E.E., Flicek, P., et al. (2015). A global reference for
624   human genetic variation. Nature *526*, 68–74. https://doi.org/10.1038/nature15393.

625   21.   Zhang, Y., Qi, G., Park, J.H., and Chatterjee, N. (2018). Estimation of complex effect-size
626   distributions using summary-level statistics from genome-wide association studies across
627   32 complex traits. Nat. Genet. *50*, 1318–1326. https://doi.org/10.1038/S41588-018-0193-
628   X.

629   22.   Zhang, Y.D., Hurson, A.N., Zhang, H., Choudhury, P.P., Easton, D.F., Milne, R.L.,
630   Simard, J., Hall, P., Michailidou, K., Dennis, J., et al. (2020). Assessment of polygenic
631   architecture and risk prediction based on common variants across fourteen cancers. Nat.
632   Commun. *11*, 1–13. https://doi.org/10.1038/s41467-020-16483-3.

633   23.   Purcell, S.M., Wray, N.R., Stone, J.L., Visscher, P.M., O'Donovan, M.C., Sullivan, P.F.,
634   Ruderfer, D.M., McQuillin, A., Morris, D.W., Oĝdushlaine, C.T., et al. (2009). Common
635   polygenic variation contributes to risk of schizophrenia and bipolar disorder. Nature *460*,
636   748–752. https://doi.org/10.1038/NATURE08185.

637   24.   Wray, N.R., Goddard, M.E., and Visscher, P.M. (2007). Prediction of individual genetic
638   risk to disease from genome-wide association studies. Genome Res. *17*, 1520–1528.
639   https://doi.org/10.1101/GR.6665407.

640   25.   Brown, B.C., Ye, C.J., Price, A.L., and Zaitlen, N. (2016). Transethnic Genetic-Correlation
641   Estimates from Summary Statistics. Am. J. Hum. Genet. *99*, 76–88.
642   https://doi.org/10.1016/J.AJHG.2016.05.001.

643   26.   Gazal, S., Finucane, H.K., Furlotte, N.A., Loh, P.R., Palamara, P.F., Liu, X., Schoech, A.,
644   Bulik-Sullivan, B., Neale, B.M., Gusev, A., et al. (2017). Linkage disequilibrium-dependent
645   architecture of human complex traits shows action of negative selection. Nat. Genet. *49*,
646   1421–1427. https://doi.org/10.1038/NG.3954,.

647   27.   Finucane, H.K., Bulik-Sullivan, B., Gusev, A., Trynka, G., Reshef, Y., Loh, P.R., Anttila,
648   V., Xu, H., Zang, C., Farh, K., et al. (2015). Partitioning heritability by functional
649   annotation using genome-wide association summary statistics. Nat. Genet.
650   https://doi.org/10.1038/ng.3404.

651   28.   Hujoel, M.L.A., Gazal, S., Hormozdiari, F., van de Geijn, B., and Price, A.L. (2019).
652   Disease Heritability Enrichment of Regulatory Elements Is Concentrated in Elements with
653   Ancient Sequence Age and Conserved Function across Species. Am. J. Hum. Genet.
654   *104*, 611–624. https://doi.org/10.1016/J.AJHG.2019.02.008.

655   29.   Turcan, A., Zhang, M., and Hou, K. (2024). Accurate polygenic association of disease to
656   individual cells in single-cell RNA-seq data. In Presented at: American Society of Human
657   Genetics (ASHG) Annual Meeting.

658   30.   Zhang, M.J., Hou, K., Dey, K.K., Sakaue, S., Jagadeesh, K.A., Weinand, K.,
659   Taychameekiatchai, A., Rao, P., Pisco, A.O., Zou, J., et al. (2022). Polygenic enrichment
660   distinguishes disease associations of individual cells in single-cell RNA-seq data. Nat.
661   Genet. *54*, 1572–1580. https://doi.org/10.1038/S41588-022-01167-Z,.

662   31.   Jones, R.C., Karkanias, J., Krasnow, M.A., Pisco, A.O., Quake, S.R., Salzman, J., Yosef,
663   N., Bulthaup, B., Brown, P., Harper, W., et al. (2022). The Tabula Sapiens: A multiple-
664   organ, single-cell transcriptomic atlas of humans. Science (1979) *376*.
665   https://doi.org/10.1126/SCIENCE.ABL4896,.

666   32.   Zheng, C., Xu, X., Wu, M., Xue, L., Zhu, J., Xia, H., Ding, S., Fu, S., Wang, X., Wang, Y.,
667   et al. (2023). Neutrophils in triple-negative breast cancer: an underestimated player with
668   increasingly recognized importance. Breast Cancer Res. *25*.
669   https://doi.org/10.1186/S13058-023-01676-7,.

670   33.   Obeagu, E.I., and Obeagu, G.U. (2024). Exploring neutrophil functionality in breast
671   cancer progression: A review. Medicine (United States) *103*, E37654.
672   https://doi.org/10.1097/MD.0000000000037654,.

34. Camargo, S., Moskowitz, O., Giladi, A., Levinson, M., Balaban, R., Gola, S., Raizman, A.,
Lipczyc, K., Richter, A., Keren-Khadmy, N., et al. (2025). Neutrophils physically interact
with tumor cells to form a signaling niche promoting breast cancer aggressiveness. Nat.
Cancer *6*. https://doi.org/10.1038/S43018-025-00924-3,.
35. Zhao, Y., Liu, Z., Liu, G., Zhang, Y., Liu, S., Gan, D., Chang, W., Peng, X., Sung, E.S.,
Gilbert, K., et al. (2023). Neutrophils resist ferroptosis and promote breast cancer
metastasis through aconitate decarboxylase 1. Cell Metab. *35*, 1688-1703.e10.
https://doi.org/10.1016/j.cmet.2023.09.004.
36. Wang, Y.H., Shen, C.Y., Lin, S.C., Kuo, W.H., Kuo, Y.T., Hsu, Y.L., Wang, W.C., Lin,
K.T., and Wang, L.H. (2021). Monocytes secrete CXCL7 to promote breast cancer
progression. Cell Death Dis. *12*. https://doi.org/10.1038/S41419-021-04231-4,.
37. Kresovich, J.K., O'Brien, K.M., Xu, Z., Weinberg, C.R., Sandler, D.P., and Taylor, J.A.
(2020). Prediagnostic Immune Cell Profiles and Breast Cancer. JAMA Netw. Open *3*.
https://doi.org/10.1001/JAMANETWORKOPEN.2019.19536,.
38. Hou, K., Gogarten, S., Kim, J., Hua, X., Dias, J.A., Sun, Q., Wang, Y., Tan, T., Atkinson,
E.G., Martin, A., et al. (2024). Admix-kit: an integrated toolkit and pipeline for genetic
analyses of admixed populations. Bioinformatics *40*.
https://doi.org/10.1093/BIOINFORMATICS/BTAE148.
39. Zhong, Y., Perera, M.A., and Gamazon, E.R. (2019). On Using Local Ancestry to
Characterize the Genetic Architecture of Human Traits: Genetic Regulation of Gene
Expression in Multiethnic or Admixed Populations. Am. J. Hum. Genet. *104*, 1097–1115.
https://doi.org/10.1016/J.AJHG.2019.04.009,.
40. Yang, Z., Wang, C., Posadas-Garcia, Y.S., Añorve-Garibay, V., Vardarajan, B., Estrada,
A.M., Sohail, M., Mayeux, R., and Ionita-Laza, I. (2025). Fine-mapping in admixed
populations using CARMA-X, with applications to Latin American studies. Am. J. Hum.
Genet. https://doi.org/10.1016/J.AJHG.2025.02.020,.
41. Pharoah, P.D.P., Antoniou, A., Bobrow, M., Zimmern, R.L., Easton, D.F., and Ponder,
B.A.J. (2002). Polygenic susceptibility to breast cancer and implications for prevention.
Nat. Genet. *31*, 33–36. https://doi.org/10.1038/ng853.
42. Gao, G., Zhao, F., Ahearn, T.U., Lunetta, K.L., Troester, M.A., Du, Z., Ogundiran, T.O.,
Ojengbede, O., Blot, W., Nathanson, K.L., et al. (2022). Polygenic risk scores for
prediction of breast cancer risk in women of African ancestry: a cross-ancestry approach.
Hum. Mol. Genet. *31*, 3133–3143. https://doi.org/10.1093/HMG/DDAC102.
43. McEvoy, B.P., Powell, J.E., Goddard, M.E., and Visscher, P.M. (2011). Human population
dispersal "Out of Africa" estimated from linkage disequilibrium and allele frequencies of
SNPs. Genome Res. *21*, 821–829. https://doi.org/10.1101/GR.119636.110,.
44. Hagerling, C., Gonzalez, H., Salari, K., Wang, C.Y., Lin, C., Robles, I., van Gogh, M.,
Dejmek, A., Jirström, K., and Werb, Z. (2019). Immune effector monocyte–neutrophil
cooperation induced by the primary tumor prevents metastatic progression of breast
cancer. Proc. Natl. Acad. Sci. *116*, 21704–21714.
https://doi.org/10.1073/PNAS.1907660116,.
45. Sakaue, S., Kanai, M., Tanigawa, Y., Karjalainen, J., Kurki, M., Koshiba, S., Narita, A.,
Konuma, T., Yamamoto, K., Akiyama, M., et al. (2021). A cross-population atlas of

genetic associations for 220 human phenotypes. Nat. Genet. *53*, 1415–1424.
https://doi.org/10.1038/S41588-021-00931-X,.

46. Gazal, S., Sahbatou, M., Babron, M.C., Genin, E., and Leutenegger, A.L. (2015). High
level of inbreeding in final phase of 1000 Genomes Project. Sci. Rep. *5*.
https://doi.org/10.1038/SREP17453,.

47. Dunham, I., Kundaje, A., Aldred, S.F., Collins, P.J., Davis, C.A., Doyle, F., Epstein, C.B.,
Frietze, S., Harrow, J., Kaul, R., et al. (2012). An integrated encyclopedia of DNA
elements in the human genome. Nature *489*, 57–74. https://doi.org/10.1038/nature11247.

48. Trynka, G., Sandor, C., Han, B., Xu, H., Stranger, B.E., Liu, X.S., and Raychaudhuri, S.
(2012). Chromatin marks identify critical cell types for fine mapping complex trait variants.
Nat. Genet. *45*, 124–130. https://doi.org/10.1038/ng.2504.

49. Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M.,
Yen, A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., et al. (2015).
Integrative analysis of 111 reference human epigenomes. Nature *518*, 317–329.
https://doi.org/10.1038/NATURE14248,.

50. Hnisz, D., Abraham, B.J., Lee, T.I., Lau, A., Saint-André, V., Sigova, A.A., Hoke, H.A.,
and Young, R.A. (2013). Super-enhancers in the control of cell identity and disease. Cell
155, 934. https://doi.org/10.1016/j.cell.2013.09.053.

51. de Leeuw, C.A., Mooij, J.M., Heskes, T., and Posthuma, D. (2015). MAGMA: Generalized
Gene-Set Analysis of GWAS Data. PLoS COMPUT. Biol. *11*.
https://doi.org/10.1371/JOURNAL.PCBI.1004219,.

ACKNOWLEDGEMENTS

The analysis utilized DNANexus and the high-performance computation Biowulf cluster at the National Institutes of Health (NIH), USA. L.C-C. is grateful for financial support from NIH (grants R56CA280636, D43CA260689, U54CA233306, U54CA283766 and U54CA280811) and The Heart, BrEast, and BrAin HeaLth Equity Research (HEAL HER) program, a program made possible by residual class settlement funds in the matter of April Krueger v. Wyeth, Inc., Case No. 03-cv-2496 (US District Court, SD of Calif.). J.Weitzel is supported by the Breast Cancer Research Foundation (grant 24-210). This research was supported in part by the Intramural Research Program of the NIH. The contributions of the NIH authors were made as part of their official duties as NIH federal employees, are in compliance with agency policy requirements, and are considered Works of the United States Government. However, the findings and conclusions presented in this paper are those of the authors and do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human Services.

AUTHOR CONTRIBUTIONS

T.U.A and H.Z. conceived the project. J.L.L. and J.Williams conducted the heritability analyses. J.W., Q.H. and L.Y. conducted the polygenicity analyses. O.J. conducted the cross-sample genetic correlation analyses. J.Williams and M.Z. conducted the enrichment analyses. A.T. conducted scDRS+ analyses. J.W. computed the LD-scores for EAS, EUR and H/L. G.J. computed the LD-scores and MAGMA analyses on AABCG controls. Statistical analyses were conducted under the guidance of K.M., Q.Z., M.J.Z., D.D., T.U.A., and H.Z.; J.L.L., M.J.Z., J. Williams, O.J., A.T., T.U.A. and H.Z. wrote the first draft. J.T.B., L.F., W.Z., S.J.C., N.C., P.K., M.G.C., M.J.Z., T.U.A and H.Z. reviewed and edited the draft. All co-authors reviewed and approved the final version of the manuscript.

DATA AVAILABILITY

AFR GWAS summary statistics are available at GWAS Catalog (GCST90296719)

BCAC EAS GWAS summary statistics are available at:
https://www.ccge.medschl.cam.ac.uk/breast-cancer-association-consortium-bcac/data-data-access/summary-results/gwas-summary-results

Biobank Japan GWAS summary statistics are available at: https://pheweb.jp/pheno/BrC

BCAC EUR GWAS summary statistics are available at:
https://www.ccge.medschl.cam.ac.uk/breast-cancer-association-consortium-bcac/data-data-access/summary-results/gwas-summary-associations

The H/L GWAS summary statistics are available by contacting Dr. Elad Ziv (elad.ziv@ucsf.edu) to gain access.

CODE AVAILABILITY

All data analysis code is available at: https://github.com/confluence-breast-cancer-consortia/pre-confluence-analysis/tree/main

LDSC and stratified LDSC are available at: https://github.com/bulik/ldsc

GENESIS is available at: https://github.com/yandorazhang/GENESIS

779   Popcorn is available at: https://github.com/brielin/Popcorn

780   scDRS is available at: https://github.com/martinjzhang/scDRS