β_1- and β_2-adrenergic Receptor Haplotypes Regulate Therapeutic Responses to Placebo and the Biased Ligand β-blocker Bucindolol

Ian A. Carroll,[1,2,3] Dobromir Slavov,[1,2] Eric R. Jonas,[1] Wayne A. Minobe,[1] Lori A. Walker,[1]

Amrut V. Ambardekar,[1,2] Timothy A. McKinsey,[1,2] Ammon D. Knaupp,[4] Donghwa Kim,[5]

Stephen B. Liggett,[3,5,6] Michael R. Bristow[1,2,3,*]

[1]Department of Medicine (Cardiology) University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA

[2]Section of Pharmacogenomics, University of Colorado Cardiovascular Institute, Boulder (80309) and Aurora (80045) CO, USA

[3]Genvara Biopharma, Westminster, CO 80021 USA

[4]BYU Genomic and Bioinformatic Center, Provo UT 84602, USA

[5]Department of Medicine and Center for Personalized Medicine, University of South Florida Morsani College of Medicine, Tampa, FL 33612, USA

[6]Department of Molecular Pharmacology and Physiology, University of South Florida Morsani College of Medicine, Tampa, FL 33612, USA

*Corresponding Author

Word Counts: Abstract 350, Text 7871, Figure Legends 683

ABSTRACT

Background: *ADRB1* and *ADRB2,* encoding cardiac myocyte β_1- and β_2-adrenergic receptors (ARs)

that mediate pathologic myocardial remodeling in response to chronically increased signaling, contain

N-terminus haplotype variants capable of influencing agonist- or biased ligand-induced receptor

internalization that uncouples canonical signaling and initiates EGFR/ERK1/2 cardioprotection.

Methods: In two heart failure (HF) clinical trial genetic substudies we investigated effects of

internalizing vs. internalization-resistant *ADRB1*/*ADRB2* haplotypes on clinical or biomarker responses

to the biased ligand β-blocker bucindolol vs. placebo or vs. the nonbiased β_1-antagonist metoprolol, and

in haplotyped isolated human heart preparations we measured ERK1/2 activation in response to these

same interventions.

Results: In subjects with $\geq$3 internalizing *ADRB1*+*ADRB2* haplotypes (6.7% subcohort) placebo

treatment was associated with fewer clinical events compared to subjects with internalization-resistant

haplotypes (Odds Ratio (OR) 0.28, 95% CI (0.10, 0.82)). In contrast, placebo treatment in subjects with

$\geq$3 internalization-resistant haplotypes (70% subcohort) was associated with more clinical events in

comparison to subjects with internalizing haplotype counterparts (OR 1.64 (1.46, 1.84)). Bucindolol

treatment was equal to placebo in the $\geq$3 internalizing subcohort, but was superior to placebo in the

internalization-resistant subcohort (bucindolol vs. placebo OR 0.49 (0.41, 0.58)). In subjects with all 4

haplotypes internalization-resistant (25% subcohort), bucindolol vs. placebo reduced time to first event

rates by 62.3$\pm$17.5% (P <0.01, 1.68$\pm$0.34 fold > the all-haplotypes parent population and additive to

1.92$\pm$0.58 fold when the *ADRB1* haplotype contained Arg389 rather than Gly389). The same bucindolol

vs. placebo pattern was observed for NT-proBNP or norepinephrine reduction vs. metoprolol. In these

comparisons *ADRB2* and *ADRB1* haplotypes behaved similarly, and although the haplotypes differed in

frequency between Black and non-Black subjects, within haplotypes there were no by-race differences

in therapeutic effects. Bucindolol but not metoprolol activated ERK1/2 signaling in isolated ventricular

preparations with ≥3 internalization-resistant haplotypes.

Conclusions: 1) Both β_1- and β_2-AR haplotypes regulate therapeutic responses in HF; internalizing

species confer protection against clinical events in placebo-treated subjects, while in internalization-

resistant haplotypes the biased ligand β-blocker bucindolol but not the non-biased ligand metoprolol is

associated with favorable effects. 2) The biased ligand cardioprotective effect may be related to

internalization-dependent or -independent ERK1/2 activation.

BACKGROUND

Human cardiac myocytes contain 3 types of β-adrenergic receptors (β-ARs), with β_1-ARs dominating but β_2-ARs also potentially involved in the regulation of cardiac function and capable of mediating pathologic effects of chronic adrenergic stimulation in the failing heart [1, 2]. In heart failure (HF) increased norepinephrine (NE) signaling contributes to myocardial pathologic remodeling, the inhibition of which results in the beneficial effects of β-blockers in subjects with reduced ejection fraction (HFrEF, LVEF <0.40) [1, 3, 4]. Although chronically increased cardiac adrenergic drive has a net myopathic effect on the myocardium [1, 4] including predisposing to arrhythmias [5], certain cellular mechanisms can counter these adverse effects. The most obvious of these are β-AR desensitization and/or downregulation that diminish adrenergic signaling through canonical cAMP/PKA and CaMKII pathways [1]. In addition, for both β_1- [6−8] and β_2-ARs [9−11] desensitization also includes signaling through an alternative, EGFR/ERK1/2 cardioprotective pathway consisting of receptor phosphorylation, binding to β-arrestins, endocytic internalization and endosomal trafficking to fates that include lysosomal degradation [7, 8]. In this series of events receptor internalization is a critical step, as inhibitors of endocytic vesicle formation block EGFR transactivation and ERK1/2 phosphorylation [10, 12]. β-adrenergic agonists classically activate this pathway [6−8, 10−12], but with some differences [13] so can β-blocker "biased ligands"[14−17].

On short-term (≤1 hour) exposure to agonists some β-AR polymorphic variants differ from their counterparts in their abilities to desensitize, internalize, and/or traffic intracellularly. For *ADRB1* the more prominently desensitizing and internalizing β_1-AR variant is Gly49 [18], which compared to Ser49 also exhibits differences in subsequent intracellular trafficking [19, 20] that are present in both the Arg389 and Gly389 haplotypes [20]. In addition, when in a laboratory-constructed Gly389 haplotype, Gly49 is associated with greater agonist-induced ERK1/2 phosphorylation than the Ser49Gly389 variant [21]. For β_2-ARs, the Arg16Gln27 haplotype exhibits rapid agonist desensitization *in vivo* compared to

other haplotypes [22], and rapidly and substantially desensitizes [23] and internalizes [23–25] on agonist

exposure in cells transfected with the original human β_2-AR cDNA containing the Arg16Gln27

haplotype [26]. Thus both *ADRB1* and *ADRB2* N-terminus nonsynonymous SNPs can affect

internalization.

The *ADRB1* position 49 and the *ADRB2* position 16 or 27 polymorphisms do not influence signal

transduction efficiency [18–20, 24]. However, the *ADRB1* Ser49Gly polymorphism is in strong linkage

disequilibrium (LD) with the nonsynonymous C-terminus SNP Arg389Gly [27] whose encoded variants

have major effects on signal transduction capacity and agonist affinity [28, 29]. The *ADRB2* position 16

and 27 variants are also in tight linkage disequilibrium [27]. *ADRB1* 49/389 haplotypes can therefore

encode receptors with greater (Arg389) or lesser (Gly389) signaling capacity or agonist affinity that

internalize to a greater (Gly49) or lesser (Ser49) extent in the presence of agonists or potentially to

biased ligands, while the *ADRB2* Arg16Gln27 vs. other encoded haplotypes have the potential for more

vs. less internalization.

In the MAP-MI study [27] we recently reported that the *ADRB1* Gly49Arg389 haplotype is

associated with a reduction in incident ventricular fibrillation (VF) in acute myocardial infarction. There

was also evidence of a VF protective signal in subjects with the *ADRB2* Arg16Gln27 haplotype [27].

These findings were interpreted as potentially due to EGFR/ERK1/2 cardioprotective pathway signaling

[6, 12] that internalization of both β_1- [8, 12, 30] and β_2-ARs [10, 31, 32] are known to activate when

occupied by an agonist, in this case NE. An earlier study had found that the *ADRB1* Gly49

polymorphism, in genotyping studies typically present only as the Gly49Arg389 haplotype, is associated

with improved survival in dilated cardiomyopathy patients, also suggesting a myocardial protective

mechanism [33]. Since HFrEF human ventricles contain 35-40% β_2-ARs that are coupled to inotropic

and pro-arrhythmic pathways and when overexpressed in model systems to a cardiomyopathy [1,2],

there is also a rationale for β_2-AR internalization affecting outcomes in HFrEF.

Here we test the hypothesis that internalization or internalization-related effects of *ADRB1* and

ADRB2 common haplotype variants influence clinical outcomes in HFrEF patients treated with placebo,

the biased ligand β-blocker bucindolol, or the nonbiased β-blocker metoprolol succinate. The 2 clinical

trial cohorts investigated [29, 34] were haplotyped for *ADRB1* and *ADRB2* variants, and one of them

(BEST Trial Adrenergic Receptor Polymorphism Substudy [29, 35]) included populations with

substantial racial heterogeneity that affects haplotype frequency and thus provides an opportunity for

internal confirmation by investigating separate genetic subpopulations. The presented data support the

utility of pharmacogenetic targeting of protein haplotypes [36], something that has long held promise

[37, 38] but has been difficult to implement.

METHODS

Clinical Trials with Genetic Substudies

BEST Adrenergic Receptor Polymorphism Substudy

This 1040 patient pharmacogenomics Substudy [29] of the Beta-blocker Evaluation of Survival Trial

(BEST, NCT00000560) [35], entitled "Pharmacogenomics of Beta-adrenergic Receptor Polymorphisms

and the Response to Beta-blockers in Heart Failure," had the primary aim of "delineating the

relationship between β_1-AR and β_2-AR polymorphisms and the clinical response to β-blockers."

Secondary aims were to "determine the relationship between β_1-AR and β_2-AR polymorphisms and the

natural history of moderate to severe heart failure in patients using the placebo arm of BEST," and added

after the trial was completed, to "see whether these polymorphisms can account for the racial differences

in the response to bucindolol that have been observed in BEST." Two polymorphisms in *ADRB1*

(positions 49 and 389) and 3 in *ADRB2* (positions 16, 27, and 164), all originally discovered and

characterized in the laboratory of the substudy's Principal Investigator (Stephen Liggett), were

prospectively identified for investigation, and the *ADRB2* Thr164Ile polymorphism was later removed

from consideration because of a minor allele frequency (0.012) insufficient for analysis.

The BEST pharmacogenomics substudy patients had advanced HFrEF (NYHA Class III-IV HF,

LVEFs ≤ 0.35) with elevated plasma NE levels (**Table S1, Supplemental Material**) who were

randomized 1:1 to treatment with placebo or the nonselective β_1-, β_2-AR blocker/biased ligand

bucindolol. Eligible subjects gave written informed consent for the parent trial [35] and as an option for

peripheral blood DNA extraction and genotyping in the Substudy [29]. The primary endpoint of the

pharmacogenomics substudy was time to all-cause mortality or cardiac transplantation.

GENETIC-AF

The GENETIC-AF trial (NCT01970501*)*, a Phase 2 trial that randomized HFrEF and HFmrEF (HF

with LVEF 0.40-0.49) patients to bucindolol or metoprolol succinate who were *ADRB1* Arg389Arg

genotype (N=267) and at risk for developing recurrent AF [34], included a voluntary DNA Bank

substudy for which 138 patients signed written informed consent. This substudy utilized banked DNA

from extracted whole blood samples used for the qualifying *ADRB1* Arg389Arg testing, which was

subsequently genotyped for *ADRB1* Ser49Gly and *ADRB2* position 27 and 16 polymorphisms. The

Primary Aim of the Substudy was association of adrenergic receptor polymorphisms with NT-proBNP

and NE biomarker measurements, with changes from baseline in these biomarkers being the primary

endpoints. The primary endpoint of the parent trial was time to AF, atrial flutter or ACM

(AF/AFL/ACM).

Study Approvals

The BEST trial and its pharmacogenomics substudy were approved by ethical committees at the

NHLBI, Department of Veterans affairs (Palo Alto VA, location of the statistical analysis center), the

DNA Oversight Committee and each investigational site. The GENETIC-AF study and its DNA

substudy were approved at each investigational site. Participants in each clinical trial signed written

informed consent for both the parent trial and separately for voluntary DNA sampling that could be used

for genetic or other analyses with de-identification of individual subjects.

Race as Modifier of Polymorphism and Haplotype Frequencies.

The race/ethnicity case report form classifications in both clinical trials were "White, Black, Hispanic, Asian/Pacific Islander, American Indian/Alaskan, Other" determined by self-identification. In the current as well as previous reports [35], Blacks were assigned one racial category, and all other races were classified as non-Blacks.

Genotyping and Haplotyping

BEST Pharmacogenomics Substudy genotyping for *ADRB1* Arg389Gly, *ADRB1* Ser49Gly, *ADRB2* Gln27Glu, and *ADRB2* Gly16Arg was conducted by Restriction Fragment Length Polymorphism (RFLP) conducted in DNA extracted from whole blood as previously described [29, 39] with results subsequently confirmed by DNA sequencing. For GENETIC-AF subjects the *ADRB1* Arg389Arg genotype necessary for trial enrollment was determined by TaqMan® SNP genotyping as previously described [34], while the *ADRB1* Ser49Gly genotype and *ADRB2* genotypes were measured by sequencing SNP flanking-primer PCR amplicons using DNA extracted for the *ADRB1* Arg389Gly genotyping. Haplotyping in both substudies was via an expectation-maximization (EM) algorithm for maximum likelihood estimates, using the "Haplo.stats" package available in R [27, 40]. Linkage disequilibrium was tested using the R based "genetics" package (https://CRAN.R-project.org/package=genetics). Haplotypes for pairwise SNPs were only ambiguous in the double heterozygote case; thus, for both *ADRB1* and *ADRB2* 20 double genotype heterozygote DNA samples were subjected to long-read sequencing [41] that confirmed the likelihood reads of missingness of 1 haplotype from each gene (Gly49Gly389 and Arg16Glu27, the double minor allele variants of each haplotype).

Quantitation and Statistical Analysis

General Strategy for Clinical Trial Substudy Data

The overall aim of the statistical analysis was to compare clinical event and biomarker outcomes in

HF subjects with more vs. fewer internalizing β-adrenergic receptor (β-AR) haplotypes, across 2 clinical

trials where haplotyping was performed and which had extensive outcomes databases. We thus designed

a statistical approach to be consistent with each trial's original statistical analysis plan, but would also

address the current primary hypothesis that β-AR haplotypes with different internalization potential

affect HF outcomes. Because of variations in cohort sample sizes internalization subdivision definitions

varied slightly between analyses (described in detail in the second paragraph of **Supplemental**

Material, **Conventions and Presentation Formats** and summarized in **Table S2**) but always included a

comparison that encompassed internalizing vs. internalization-resistant (*ADRB1+ADRB2*) haplotypes.

The greater sample size of the BEST Pharmacogenomics Substudy allowed more gradation in

estimation of dose-response effects of haplotype internalization (i.e., sufficient sample size for analyses

using three mutually-exclusive tiers and two separate treatment arms), whereas the GENETIC-AF DNA

substudy, limited in sample size and restricted in *ADRB1* genotype to Arg389Arg, maintained the

directionality of this classification by confining the analysis to two mutually-exclusive tiers that were

also analyzed in the BEST cohort.

Genotype and Haplotype Analyses

Genotype-level analyses are inherently confounded by effects related to other loci in linkage

disequilibrium with the polymorphism of interest. This can be reconciled by analysis at the haplotype

level, where 3 haplotypes exist for both *ADRB1* (Gly49Arg389, Ser49Arg389, Ser49Gly389) and

ADRB2 (Arg16Gln27, Gly16Gln27, Gly16Glu27), which manifest independently at the genome

(different chromosomes) or statistical (no detectable linkage disequilibrium and satisfying Hardy-

Weinberg equilibrium) level. A standard Bonferroni adjustment for internalizing vs. internalization-

resistant haplotypes for both *ADRB1* and *ADRB2* would therefore yield $\alpha = .025$ for an endpoint

analysis, which was considered the adjusted α-level.

Clinical Endpoints

The clinical endpoints prospectively chosen for analysis included the primary cardiovascular

endpoints for both the BEST Pharmacogenomic and GENETIC-AF DNA substudies (ACM/Tx [29] and

AF/AFL/ACM [34], respectively), the primary endpoint of the BEST parent trial [35] (ACM) that was a

secondary endpoint in its substudy, 2 secondary endpoints from the BEST parent trial (CVM, HFH)

[35], a composite endpoints from these secondaries (ACM/HFH), a previously reported primary

endpoint used for BEST trial data (CVM/HFH) [42] that has become a favored clinical outcome

assessment in HF trials, and 1 post hoc endpoint (CVH) requested by the FDA [43]. Clinical endpoints

were analyzed by forest plots, structural equation modeling (SEM), and time to first event (Kaplan-

Meier) curves. The methodology for these analyses is further described in **Supplemental Material**,

Methods.

Isolated Human Heart Studies

Tissue Bath Preparations

Explanted failing hearts from end stage heart failure patients with nonischemic or ischemic

cardiomyopathies were obtained at the time of cardiac transplantation and immediately transported to

the laboratory in iced Tyrode's solution. Right ventricular trabeculae (N=10-14) of uniform size (1-2 x

6-8 mm) were mounted in tissue baths in Tyrode's solution at 37 and field stimulated to contract at 60

bpm, as previously described [29]. Following a 1 hour equilibration period 1-2 trabeculae/group were

incubated with vehicle, isoproterenol (1umol/L), bucindolol HCL(1umol/L) or metoprolol tartrate (10

umol/L) for either 5, 10 or 60 minutes. At the end of its incubation period each trabeculum was removed

from the bath, flash frozen in liquid nitrogen and stored at -80°C.

pERK1/2 Immunoblotting

These details are in **Supplemental Materials**, **Supplemental Methods**. **Statistical Testing**

Statistical Testing

Details of the SEM analyses, including a listing of covariates used for adjustment, can be found in

Supplemental Methods, Structural Equation Modeling. Time to first event Hazard Ratios (HzR)

were generated by Cox proportional hazards models (**Supplemental Methods, Time to First Event**

Analyses), with first clinical event Odds Ratios (ORs) obtained from logistic regression models.

Confidence intervals for linkage disequilibrium coefficients were calculated using 10,000 nonparametric

bootstrapped estimates. Treatment effects between bucindolol vs. control (placebo or metoprolol

succinate) were calculated from HzRs as: % treatment effect = ((1-HzR) x 100). Normality was

determined by the Anderson Darling test. For normally distributed data assessing one clinical endpoint t-

tests were used, and for >1 endpoint one-way ANOVA with a Holm-Sidak's test was employed. Analysis

of ERK1/2 phosphorylation was by a one-way ANOVA mixed model with repeat measures and a Holm-

Sidak multiple comparison test. For non-normally distributed data Wilcoxon or Friedman's/Kruskal-

Wallis tests with a Dunn's multiplicity adjustment were employed.

RESULTS

BEST Adrenergic Receptor Polymorphism Substudy

Allele, Genotype, and Haplotype Frequencies; Linkage Disequilibrium

A description of the formatting and terminology used for genotypes, haplotypes and diplotypes is in

Table S2 and the first paragraph of **Supplemental Material**. Allele frequencies of *ADRB1* and *ADRB2*

polymorphisms by Black and non-Black self-identified races are in **Table S3**, and are similar to those

previously reported for European- vs. African-Ancestry in the BEST Substudy [44]. Blacks have higher

minor allele frequencies for *ADRB1* Gly389 and Gly49, and for *ADRB2* Arg16. *ADRB2* Glu27 is lower

frequency in Blacks.

Genotypes and theoretical vs. identified diplotypes are in **Table S4.** For both *ADRB1* and *ADRB2*, of

10 theoretical diplotypes only 6 were identified; diplotypes with *ADRB1* Gly49Gly389 and *ADRB2*

Arg16Glu27 haplotypes are missing. In each gene identified diplotypes consist of 3 homozygotes, 2

single heterozygotes, and 1 double heterozygote. There is strong LD between *ADRB1* positions 49 and

389 polymorphisms, as well as among *ADRB2* positions 16, 27 and 164 (**Table S5**); the latter

polymorphism was very low frequency (minor allele 0.012, **Table S3**) and was not further investigated.

The absence of *ADRB1* Gly49Gly389and *ADRB2* Arg16Glu27 haplotypes was confirmed by long-read

DNA sequencing conducted in 26 *ADRB1* 49/389 and 50 *ADRB2* 16/27 double heterozygotes. **Table S6**

also gives haplotypes by sex within race categories, where there is no difference between sexes within

either race category.

Diplotype and haplotype frequencies by race classification are in **Tables 1** and **S6**, respectively.

Gly49Arg389 and Ser49Gly389 haplotypes are more prevalent in Blacks, while the major allele

haplotype Ser49Arg389 is more prevalent in non-Blacks (**Table S6**). For *ADRB2* Gly16Gln27 as well as

Arg16Gln27 are more prevalent in Blacks. The internalizing haplotypes Gly49Arg389 (*ADRB1*) and

Arg16Gln27 (*ADRB2*) are respectively 1.71 (P<0.0001) and 1.21 (P=0.004) fold more frequent in

Blacks vs. non-Blacks. Internalization-resistant haplotypes are more prevalent in non-Blacks compared

to Blacks, by 1.14 (P <0.0001) and 1.15 (P=0.004) fold in *ADRB1* and *ADRB2*, respectively (**Table S6**).

For *ADRB1* diplotypes (**Table 1**), any that contain at least 2 total glycines in positions 49 plus 389

are more prevalent in Blacks, while the Ser49Arg389 homozygous diplotype is more common in non-

Blacks. The Gly49Arg389 homozygous diplotype is 2.34 fold more common in Blacks (P =0.012), as is

the homozygous Ser49Gly389 diplotype (by 2.43 fold, P <0.0001).

Baseline Characteristics

Baseline characteristics by diplotype, race and treatment group are in **Table S1**. For diplotypes the

main differentiating characteristic is race (**Tables S1** and **S6)**. In comparison to non-Blacks, Blacks are 5

years younger and have higher resting heart rates and systolic blood pressures. There are no differences

in baseline characteristics between treatment groups.

β-adrenergic Receptor Haplotype Internalization Potential Affects Clinical Outcomes

Clinical Events in Internalizing vs. Internalization-resistant Haplotypes

As an initial test of the hypothesis that *ADRB1* and *ADRB2* haplotypes with different internalization

potential affect clinical outcomes, in placebo- and bucindolol-treated subjects in the BEST substudy

unadjusted odds ratios (ORs (95% C.I.s)) for first clinical event rates were compared between defined

{*ADRB1+ADRB2*} haplotype internalizing vs. internalization-resistant groups. The index internalizing

haplotypes were *ADRB1* Gly49Arg389 and *ADRB2* Arg16Gln27 as tabulated in {*ADRB1+ADRB2*}

diplotype combinations, which were compared to all other diplotype combinations (see classification

nomenclature, **Table S2**). In the classification descriptors "number" is the sum of the *ADRB1+ADRB2*

haplotypes in the diplotype combination, "D" (double) indicates that at least 1 defined haplotype is

present in both genes, "I" is internalizing, and "IR" is internalization-resistant.

Figures 1-3 (all-LVEF cohort) and **S1-S3** (cohort restricted to baseline LVEFs of ≥0.20) are forest

plots of the between-haplotypes ORs for 7 clinical endpoints that include the primary endpoints for the

BEST and GENETIC-AF substudies, respectively time to all-cause mortality or cardiac transplantation

(ACM/Tx)[28] and time to first atrial fibrillation or flutter (AF/AFL/ACM) [34]. To provide a robust

statistical analysis as well as to correct for possible confounding variables, a covariate adjusted

structural equation model (SEM) endpoint analysis is provided in **Table S7** where clinical events are

modeled against increasing numbers of internalization-resistant haplotypes from 0 to 4. All analyses are

constructed from the first clinical events in subjects who had complete *ADRB1* and *ADRB2* genotyping.

In **Figures 1-3** and **S1-S3** {*ADRB1+ADRB2*} diplotype combinations are organized by 3 sections or tiers

based on the total number of internalizing {*ADRB1+ADRB2*} index haplotypes. **Tier 1** is 3-4 copies of

ADRB1 AND *ADRB2* internalizing haplotypes ("Double-Internalizing", ≥**3DI**); **Tier 2** has 2 copies of

internalizing haplotypes ("Incompletely Internalizing", **2I**); and **Tier 3** has 0-1 copies of internalizing (3-

4 internalization-resistant) haplotypes ("Double Internalization-resistant", **≥3DIR**). The sequence of

presentation is placebo treated patients (**Figures 1 and S1**), bucindolol patients (**Figures 2 and S2**), and

bucindolol vs. placebo treatment effects (**Figures 3 and S3**). ORs in **Figures 1/S1** and **2/S2** are

generated by comparing the haplotype pairs in Column 1 to all other haplotype pairs not defined in

Column 1, and treatment effect ORs (**Figures 3/S3**) are based on bucindolol vs. placebo event ORs.

Placebo Treatment in Subjects with {ADRB1+ADRB2} Internalizing vs. Internalization-resistant

Haplotypes is Associated with Lower Clinical Event Rates

In the All-LVEF cohort the endpoint analysis for placebo treatment (**Figure 1**) indicates that subjects

within **Tier 1 (≥3DI)** have OR point estimates to the left of the line of identity, ranging from 0.29 (0.12,

0.68) for cardiovascular mortality or heart failure hospitalization (CVM/HFH) to 0.48 (0.23, 1.00) for

cardiovascular hospitalization (CVH). The mean for all 7 endpoints is 0.39 (0.27, 0.55). In contrast, for

the **Tier 3 (≥3DIR)** subjects ORs are to the right of the identity line, between 1.45 (0.88, 2.39) for CVM

and 1.78 (1.22, 2.58) for CVH, mean 1.64 (1.46, 1.84). The **Tier 2 (2I),** subjects position between **Tiers**

1 and **3**, slightly to the left of identity, with a mean OR of 0.76 (0.66, 0.88). By an aggregate analysis of

all 7 endpoints, **Tier 1** event rates and their ORs are significantly different from **Tier 3** (P=0.005). The

less severe LV dysfunction cohort (LVEF ≥0.20) placebo forest plot (**Figure S1**) is very similar to the

All-LVEF cohort, with a Tier 1 vs Tier 3 aggregate OR difference also P=0.005. The **Tier 1** mean OR

for the 7 endpoints was 0.28 (0.10, 0.82), with a **Tier 3** of 2.16 (1.81, 2.58).

The SEM analysis in the All-LVEF cohort, adjusted for both dependencies between endpoints and

direct effects of 13 covariates, confirms an increase in the placebo event rate slopes progressing from 0

to 4 internalization-resistant haplotypes in **Tier 1** through **Tier 3 (Table S7)**. All 7 endpoints in both

cohorts have positive individual slopes, indicating increases in event likelihood as the number of

internalization-resistant haplotypes is increased. Five of the 7 endpoints in the All-LVEF cohort satisfy P

<0.05, with 4 of 7 in LVEF ≥0.20. By Likelihood Ratio Testing the slopes show no evidence of

heterogeneity (χ^2 P =0.58, 0.78 for All-LVEF, LVEF $\geq$0.20 respectively), indicating that the effect of

internalization does not differ across endpoints. A follow-up SEM with equality constraints for this

component of the structural model yields a significant positive effect of internalization tier on event

likelihood across all 7 endpoints (P = 0.002 and 0.022 for the All-LVEF and $\geq$0.20 cohorts, respectively)

(**Table S7**).

Bucindolol Treatment is Unaffected by Haplotype Internalizing Potential

In contrast to placebo treatment, the bucindolol forest plot OR point estimates for the All-LVEF

(**Figure 2**) and LVEF $\geq$0.20 cohorts (**Figure S2**) exhibit minimal tendency for departure from the line of

identity, with a nonsignificant SEM analysis (**Table S7**). In contrast to placebo, no estimated effect of

increasing internalization-resistant haplotypes is statistically significant (**Table S7**).

Bucindolol vs. Placebo Treatment Effects are Inversely Related to Haplotype Internalizing

Potential

Figures 3 and **S3** contain between-treatment group ORs within internalization haplotype tiers, and

demonstrate a pattern commensurate with the within-placebo and -bucindolol subjects in **Figures 2** and

S2. In both the All-LVEF (**Figure 3**) and LVEF $\geq$0.20 (**Figure S3**) cohorts the **$\geq$3DI Tier 1** ORs,

although statistically nonsignificant, are to the right of the identity line, while in **Tier 3** the **$\geq$3DIR** ORs

are <1.0 with the majority (5 of 7 in **Figure 3**, all 7 in **Figure S3**) statistically significant. The mean OR

of the 7 endpoints for the All-LVEF cohort in **Tier 3** is 0.66 (0.58,0.76), while the LVEF $\geq$0.20 cohort is

0.49 (0.41, 0.58).

The SEM analysis (**Table S7**) difference between the placebo and bucindolol group slopes indicates

consistent treatment effects across all endpoints in both LVEF cohorts. Bucindolol minus placebo slope

values vary from -0.11 (HFH) to -0.27 (ACM/Tx) in the All LVEF Cohort, and -0.22 (AF/AFL/ACM) to

-0.38 (ACM/Tx) in LVEF $\geq$0.20, indicting the bucindolol treatment effect is consistently against the

presence of internalizing and in favor of internalization-resistant haplotypes. The All 7 Endpoints

interaction P values are statistically significant in both cohorts, P=0.048 (All LVEF) and P=0.003 (LVEF

≥0.20), respectively indicating the slopes of the placebo and bucindolol groups are different (**Table S7**),

and that bucindolol treatment negates the deleterious effects of increasing numbers of internalization-

resistant haplotypes observed in placebo-treated subjects.

ADRB1 and ADRB2 Haplotypes Are Associated with Similar Internalization to Non-internalization

Dose-Responses

Placebo Treatment

Figures S4-S6 contain All-LVEF cohort data for the internalizing *ADRB1* Gly49Arg389 haplotype

held constant at 1 or 2 copies and paired with decreasing copies (from 2 to 0) of the *ADRB2*

internalizing, Gln27/Arg16 haplotype. ORs are calculated and plotted as for **Figures 1-3**, representing

an internalizing to internalization-resistant *ADRB2* haplotype dose-response for the 7 clinical endpoints.

Within each clinical endpoint there is a progressive increase in ORs for all 7 endpoints as the copy

number of *ADRB2* Arg16Gln27 decreases, from an OR range of 0.15 − 0.47 (mean 0.28±0.12) for 2

copy to 1.03 − 1.59 (mean 1.34±0.19) for 0 copies (P=0.016). For the 7 clinical endpoints, 8 of the 14

ORs in the Arg16Gln27 2 or 1-2 copy groups are statistically significantly reduced (upper bound <1.00),

with none changed in the remaining 3 categories with fewer copy numbers. The converse, haplotype

downtitration of *ADRB1* Gly49Arg389 in the presence of constant, 1-2 copies of *ADRB2* Arg16Gln27

(**Figure S7**) yields similar data.

Bucindolol Treatment

In contrast, the internalizing to internalization-resistant haplotype dose-response is not present in

patients treated with bucindolol (**Figures S5, S8**). In **Figure S5,** with downtitration of the internalizing

ADRB2 haplotype Arg16Gln27, ORs are on either side and close to the 1.0 identity line with none

statistically significant. The downtitration of the *ADRB1* internalizing haplotype (**Figure S8**) reveals a

slightly different pattern, with 2 copies of Gly49Arg389 demonstrating a lower OR (ranging from 0 to

0.73, mean 0.32±0.29) than all other copies in all 7 endpoints. Also, for mortality dominant endpoints

(CVM, ACM/Tx, ACM), haplotypes with 0 copies of Gly49Arg389 have ORs consistently >1.0 (range

1.25-1.68, mean 1.52±0.16). These 2 deviations from Arg16Gln27 downtitration are likely due to the

respective favorable and relatively unfavorable impacts of bucindolol in subjects with *ADRB1*

Arg389Arg and Gly389 genotypes [29].

Bucindolol vs. Placebo Treatment Effects

Bucindolol vs. placebo treatment effects with downtitration of internalizing haplotypes (**Figures S6**

and **S9**) generally indicate an overall progressive decrease in ORs in all endpoints, or the opposite

pattern to that for placebo in **Figures S4** and **S7**. The one exception is consistently low ORs for 2 copy

Gly49Arg389 in all 7 endpoints, again likely due to the favorable impact of the *ADRB1* Arg389Arg

genotype on bucindolol treatment effects [30].

Hazard Ratios for Time to Clinical Events, Internalizing vs. Internalization-resistant Haplotypes

Placebo or Bucindolol Treatment in Internalizing vs. Internalization-resistant Haplotypes

Clinical event hazard ratios (HzRs) differ from odds ratios by incorporating the element of time,

which can increase the sensitivity for detection of treatment effect differences. For placebo treated

patients in the BEST substudy All-LVEF cohort, **Figure S10** contains time to first event curves and

HzRs comparing ≥2 Double Internalizing (≥**2DI**, 1-2 copies of *ADRB1* Arg389Gly16 AND 1-2 copies of

ADRB2 Arg16Gln27) to subjects with all other (≥2 Not Double Internalizing, ≥**2IR**) haplotypes, for 4 of

the clinical endpoint ORs plotted in **Figures 1-3** and **S1-S3**. In all 4 endpoints placebo-treated subjects

with ≥**2DI** haplotypes exhibit progressive, favorable separation in the Kaplan-Meier (K-M) curves

compared to ≥**2IR** haplotypes, with HzRs ranging from 0.62 (0.40, 0.97) to 0.69 (0.48, 1.00). **Figure**

S11 contains the same comparisons for bucindolol-treated patients in the All-LVEF cohort, where curve

separation does not occur for ≥**2DI** vs. ≥**2IR**subjects; HzRs are near 1.0 for all 4 endpoints, and event

rates are similar to the ≥**2DI** placebo curves in **Figure S10**.

Bucindolol vs. Placebo Treatment Effects in Maximum Internalization-resistant Haplotypes

As an extreme evaluation of the consequences of internalization-resistant *ADRB1* and *ADRB2*

haplotypes on treatment effects, in the LVEF ≥0.20 cohort **Figure 4** shows bucindolol vs. placebo time

to first event curves for "Maximum Double Internalization-resistant" (**4DIR**) *ADRB1* and *ADRB2*

combination diplotypes (2 copies of an *ADRB1* Ser49 containing haplotype AND 2 copies of an *ADRB2*

non-Arg16Gln27 diplotype). HzRs (95% C.I.s) range from 0.17 (0.05, 0.56) for ACM/Tx (substudy

primary endpoint) to 0.53 (0.33, 0.84) for CVH.

ADRB1 and ADRB2 Haplotype Effects Are Equivalent in Black and Non-Black Race

As shown in **Tables 1** and **S1**, Black vs. other races exhibits differences in common genetic variation

within β-ARs, as in other neurohumoral [44] and non-neurohumoral systems [45]. **Figure 5** consists of

bucindolol vs. placebo time to event curves in the All-LVEF cohort by Black or non-Black racial

categories for the high prevalence composite endpoints of time to first CVM/HFH and CVH, in subjects

with Double Internalizing (**≥2DI)**, and Not Double Internalizing (**≥2IR)** haplotypes. For **≥2DI** subjects

due to favorable effects in placebo-treated subjects there is no evidence of a bucindolol treatment effect

in either Blacks or non-Blacks. In contrast, in **≥2IR** subjects for both endpoints there is evidence of a

treatment effect, with little or no difference in HzRs between Black and non-Black subjects (HzRs

ranging from 0.64 (0.39, 1.06) for CVH in Blacks to 0.76 (0.61, 0.96) for CVH in non-Blacks, nearly

identical HzRs in CVM/HFH). These data indicate that within the **≥2IR** haplotype phenotype there are

no differences between Blacks and non-Blacks for enhancement of bucindolol treatment effects.

The haplotype dose response progressing from ≥2 Double Internalizing to ≥2 Internalization-

resistant in placebo treated patients (**Figures 1, S1**) is observed in both Blacks and non-Blacks (**Figure**

S12). For both groups the 7 endpoint ORs in **Tier 1**, **≥3DI** subjects are consistently to the left of the line

of identity, **Tier 3**, ≥3DIR point estimates are to the right of 1.0, and **2I** Incompletely Internalizing

estimates are intermediate, positioned around 1.0.

GENETIC-AF DNA Substudy

Internalizing/Internalization-resistant Haplotypes

GENETIC-AF Substudy subjects baseline characteristics by *ADRB1* or *ADRB2* diplotype are in

Table S8. The small sample size precluded assessment of clinical events, but the trial included

peripheral venous plasma NT-proBNP and NE biomarker measurements at baseline (pre-randomization

to bucindolol or metoprolol succinate) and at 4, 12 and 24 weeks (trial end) of efficacy follow-up [34].

The genotypes, haplotypes and diplotypes are in **Table S9.** Subject diplotypes were divided by

{*ADRB1+ADRB2*} diplotype combination into ≥**2DI** and ≥**2IR** groups as for the BEST substudy, and

biomarker changes were assessed in the 2 treatment groups (**Figure 6**, **Tables S10-S13**). Biomarkers

were also analyzed in **4DIR** haplotype subjects (**Tables S10-S13**). Neither biomarker was normally

distributed, and both were analyzed by non-parametric tests.

Baseline characteristics within *ADRB1* or *ADRB2* diplotype cohorts (**Table S8**) or in the treatment

arms do not differ meaningfully. NT-proBNP is a biomarker for both AF and HF [34], and thus is a

suitable surrogate for GENETIC-AF's primary endpoint of time to symptomatic AF/AFL/ACM in a HF

population. In bucindolol-treated subjects with maximum internalization-resistant, **4DIR** haplotypes

(**Table S10**), at 4, 12 and 24 weeks NT-proBNP is decreased by respective amounts of 737±312 pg/ml

(mean±sem, P= 0.009), 493±329 pg/ml (P=0.15) and 778±266 pg/ml (P= 0.005). In contrast, metoprolol

treated **4DIR** patients (**Table S11**) exhibit no statistically significant changes at any timepoint.For

ADRB1+ADRB2 ≥**2DI** haplotypes, neither bucindolol nor metoprolol treated subjects exhibit changes in

NT-proBNP at any timepoint (**Figure 6).** In contrast, in ≥**2IR** haplotype subjects treated with bucindolol

NT-proBNP is reduced from a baseline of 1364±1494 pg/ml by 252±242 pg/ml (mean±sem, P=0.004 vs.

baseline), 361±158 pg/ml (P=0.043), and 380±143 pg/ml (P=0.003) at 4, 12 and 24 weeks respectively

(**Figure 6**). Metoprolol-treated patients with ≥**2IR** haplotypes have no significant changes from baseline

(**Figure 6).**

NE measurements are in **Figure 6** and in **Tables S12** and **S13**. For bucindolol-treated subjects with

4DIR haplotypes the pattern is similar to NT-proBNP; there are statistically significant reductions at 4,

12 and 24 weeks (**Table S12**). Also similar to NT-proBNP, in **4DIR** haplotypes there are no reductions

in NE at any timepoint in the Metoprolol Arm (**Table S13**). Results for ≥**2DI**, Double Internalizing

haplotypes are also similar to NT-proBNP, with no significant changes at any timepoint in either the

bucindolol or metoprolol arms (**Figure 6**)). The ≥**2IR,** bucindolol group (**Figure 6**) results are also

similar to those for NT-proBNP, with statistically significant reductions at 4, 12 and 24 weeks. In the

metoprolol arm ≥**2IR** subjects there are no changes in NE at any timepoint. (**Figure 6**).

Expansion of the Bucindolol Pharmacogenetic Target Population and Efficacy by the Inclusion of

ADRB1, ADRB2 **Internalization-resistant Haplotypes**

For bucindolol vs. placebo treatment effects, efficacy enhancement by internalization-resistant β-AR

haplotypes in the BEST substudy creates the potential for enlarging bucindolol's pharmacogenetically

targeted population beyond *ADRB1* Arg389Arg [29, 34]. Also, within the expanded target population

there is the potential for subgroups yielding treatment effects superior to *ADRB1* Arg389Arg [29, 34].

For the LVEF ≥0.20 cohort, treatment effects for all listed pharmacogenetic groups (**Table S15, columns**

2-8) exceed that for the All-Genotypes parent index cohort (Column 1), at a significance level of at least

P <0.05. In addition, in 4 of the 6 internalization-resistant diplotype-containing subgroups (Columns 3-

8) treatment effects statistically exceed those of the Column 2 *ADRB1* Arg389Arg genotype index

pharmacogenetic enhancement subgroup, and all 6 are quantitatively greater. Moreover, the combined

ADRB1 AND *ADRB2* Maximum Double Internalization-resistant (**4DIR**) diplotype subgroup (Column

6) treatment effect exceeds the Column 2 *ADRB1* Arg389Arg group by 1.34 fold (P=0.0017) and the

ADRB1 Arg389Arg group by 1.68 fold (P <0.01) (**Table S15**, **Figure S13**). The combination of the

4DIR subgroup with non-overlapping *ADRB1* Arg389Arg genotype subjects (Column 7) creates a

subgroup that is 33% larger than the Column 2 index pharmacogenetic group (N=441 vs. 331), with a

1.03 fold marginally higher (P=0.064) treatment effect on the paired endpoints analysis. Finally, the

requirement that both the *ADRB1* Ser49Arg389 homozygous diplotype AND internalization-resistant

ADRB2 diplotypes be present (Column 8, 10% of the total cohort, the *ADRB1* Arg389 Ser49

homozygous diplotype subset of 4DIR) creates an increase in treatment effect of 1.52 fold >Column 2

(P=0.004), and 1.92 fold >Column1 (P=0.01) (**Figure S13**). Analogous data for the All LVEF cohort are

similar (**Table S16**).

ERK1/2 Phosphorylation in Human Isolated Right Ventricular Trabeculae

In order to confirm that bucindolol acts as a biased ligand in the human heart and is capable of

activating a putative cardioprotective pathway, ERK1/2 phosphorylation (pERK1/2) was assessed in

preparations of right ventricular (RV) trabeculae isolated from 6 human explanted failing hearts, 5 of

which had ≥3 copies of {*ADRB1+ADRB2*} (**Table S17**). After a 5- (N=1) or 10-minute (N=5) tissue

bath incubation, bucindolol (P=0.0027 vs. Vehicle) but not metoprolol (P=0.38) is associated with an

increase in pERK1/2 compared to vehicle (**Figure 7, Table S17**). Isoproterenol's pERK1/2/total ERK is

similar to bucindolol's (respectively 1.36±0.36 vs. 1.40±0.27, P=0.73) (**Table S17**) but due to greater

variability (**Figure 7B**) it is not statistically significant (P=0.104 vs. Vehicle). At 60 minutes bucindolol

uniquely is decreased compared to values at 5/10 minutes, 0.87±0.55 vs. 1.40±0.27, P=0.037 for

interaction with time.

DISCUSSION

For same-allele nonsynonymous SNPs it is the encoded protein haplotype encompassing each

polymorphic locus that has the potential to modify biologic effects [36−38, 46]. This has been

previously shown for *ADRB2* haplotypes expressed in COS-7 cells, where mRNA, protein and

pharmacologic phenotype are haplotype encoded [47]. The *ADRB1* 49/389 haplotype would be expected

to encode a higher function β_1-AR that easily internalizes on agonist exposure (Gly49Arg389), a higher

function receptor relatively resistant to internalization (Ser49Arg389), and a lower function receptor

resistant to internalization (Ser49Gly389) [27]. The 4th possibility, Gly49Gly389, would theoretically

have lower function and would easily internalize, but was not found in this and most other genotyping

studies [27, 48], presumably due to strong linkage disequilibrium. The β_2-AR has 2 loci that in tandem

regulate internalization, with the Arg16Gln27 haplotype exhibiting a high level of agonist mediated

internalization [9, 23−25] and/or rapid desensitization [22, 23]. Similar to *ADRB1*, only 3 *ADRB2*

haplotypes are usually found in genetic studies including in the current report, with Arg16Glu27 absent.

The absence of 1 of 4 possible haplotypes in each β-AR gene allows near certain maximum likelihood

haplotyping from genotype data, and long-read DNA sequencing confirmed that for both *ADRB1* and

ADRB2 double heterozygote genotypes only 1 expected haplotype was present (respectively

Gly49Arg389/Ser49Gly389 and Arg16Gln27/Gly16Glu27).

***ADRB1*, *ADRB2* Haplotypes in the BEST Adrenergic Receptor Polymorphism Substudy.**

The data from the BEST Adrenergic Receptor Polymorphism substudy indicated that placebo

treatment results in reduced clinical event rates in patients who have *ADRB1* and *ADRB2* haplotypes

encoding internalizing receptor variants. In a HF patient treated with placebo internalization would

presumably occur on exposure to increased NE levels, present in both clinical trial populations. In the

BEST substudy, across 7 clinical endpoints placebo-treated subjects with ≥3DI internalizing haplotypes

had lower clinical event rates with ORs <1.0 compared to counterpart haplotypes, whereas patients with

internalization-resistant (≥3DIR) haplotypes exhibited higher event rates with ORs >1.0. These results

are consistent with data from the MAP-MI study, with those previous observations extended to the

combination of {*ADRB1*+*ADRB2*} internalizing haplotypes.

In contrast, subjects in the bucindolol group exhibited more uniform effects across internalization

tiers, with event rates similar to placebo in internalizing haplotypes and no statistically significant event

rate differences in ORs between haplotype tiers. As a result, on unadjusted analyses bucindolol vs.

placebo treatment effect event rate ORs were lower in subjects with internalization-resistant (≥3DIR)

haplotypes and trended lower in Incompletely Internalizing (2I) subjects. Structural equation modeling

of 7 clinical endpoint events progressing from 0 to 4 internalization-resistant haplotypes confirmed the

effects suggested by the unadjusted analyses, as placebo-treated subjects exhibited an increase in event

rate slopes and bucindolol treatment was associated with no changes. These differences led to bucindolol

vs. placebo treatment effects consistently favoring internalization-resistant haplotypes. The event rate

OR data were further corroborated by time to clinical event analyses, where placebo treatment resulted

in a reduction in HzRs in subjects with internalizing (≥2DI) compared to internalization-resistant (≥2IR)

haplotypes. In contrast, for bucindolol treatment there was no difference between subjects with ≥2DI vs.

≥2IR haplotypes, with HzRs similar to placebo in the ≥2DI haplotype group. That is, when there is at

least 1 internalizing haplotype for *ADRB1* AND *ADRB2*, placebo in the presence of a β-agonist (NE) has

a favorable effect approximately equivalent to bucindolol. With bucindolol treatment the effects of NE

on receptor internalization in subjects with ≥2DI, internalizing haplotypes would presumably be

blocked, with biased ligand effects leading to internalization/and or blockade of any adverse effects

mediated through canonical cell surface signaling. In contrast, in subjects with ≥2IR, internalization-

resistant haplotypes, bucindolol exhibited efficacy in excess of placebo with statistically significant

treatment effects ranging from 24% to 36% for time to effect analyses regardless of race. The

therapeutic preference of bucindolol for internalization-resistant haplotypes was underscored by results

in 4DIR subjects, who exhibited treatment effects from 51% to 83%. Therefore, compared to placebo,

the biased ligand β-blocker bucindolol was associated with a more favorable treatment effect in subjects

with internalization-resistant haplotypes, which constituted approximately 70% of the BEST substudy

All LVEF cohort.

ADRB1, ADRB2 **Haplotypes in the GENETIC-AF Trial**

The limited sample size of the GENETIC-AF DNA substudy was nevertheless sufficient to detect β-

AR haplotype effects on NT-proBNP and NE. Both NT-proBNP and NE plasma levels were reduced by

bucindolol compared to metoprolol, similar to results in the entire cohort [34, 49]. In the substudy the

reductions in both biomarkers occurred only in the Bucindolol Arm in subjects with 4DIR or ≥2IR

internalization-resistant haplotypes. In contrast, metoprolol succinate was not associated with a

reduction in NT-proBNP or NE in subjects with internalization-resistant haplotypes. Thus relative to

internalization category haplotypes the effect of the nonbiased antagonist metoprolol was different from

bucindolol, and from placebo in the BEST substudy where clinical events were reduced in ≥2DI

subjects. This suggests that in subjects with ≥2DI, internalizing haplotypes metoprolol blocked the

beneficial, receptor internalizing effects of NE, but did not produce any favorable effects on its own.

This could be because metoprolol is a nonbiased ligand, or because as β_1-AR selective antagonist β_2-AR

adverse canonical signaling was unaffected.

Haplotype Data Indicate that Both *ADRB1* and *ADRB2* Internalizing Variants Affect Heart

Failure Clinical Events and Bucindolol Treatment Effects.

Human cardiac myocytes contain β_2-ARs that are coupled to positive inotropic effects via the

canonical cAMP/PKA pathway [1] and to ERK1/2 signaling following internalization [10]. When

overexpressed in mouse hearts human β_2-ARs mediate histopathological effects [2, 51], and in failing

human ventricles the proportion of β_2-ARs/Total β-ARs is increased, from 20-25% to 35-40% [1].

However, the numerically dominant β_1-AR signaling pathway is more biologically adverse, and in

clinical trials mortality reduction outcomes are similar for β_1-AR selective blocking agents and the

nonselective, biased ligand β-blocker carvedilol [1]. Despite these observations, in the current study

with placebo treatment *ADRB2* internalizing haplotypes were associated with improved clinical

outcomes, and internalization-resistant haplotypes conferred improved bucindolol treatment effects.

Furthermore, haplotype copy number dose-response indicated that *ADRB2* effects were equivalent to

those for *ADRB1* haplotypes. These data support that in the human heart myocardial β_2-ARs are

operational in the natural history of adrenergically-mediated pathologic ventricular remodeling, and that

their pharmacologic manipulation can have therapeutic consequences. Moreover, for clinical endpoints

the combination of *ADRB1* and *ADRB2* internalization-resistant haplotypes was additive, as might be

expected based on the known differences in their encoded receptor signaling pathways [1].

Effects of Race

As it does for the *ADRB1* Arg389Gly polymorphism [44], race also affected β-AR internalization

variants, and was the only baseline characteristic in the BEST adrenergic receptor polymorphism

substudy that consistently varied by diplotype. Blacks compared to non-Blacks had higher frequencies

of internalizing and lower frequencies of internalization-resistant haplotypes, which suggests that in the

absence of other confounding variables placebo treatment (or presumably no treatment) would favor

Blacks based on the beneficial impact of a greater frequency of internalizing *ADRB1* and *ADRB2*

haplotypes. In contrast, some of bucindolol's efficacy enhancement is dependent on the presence of

internalization-resistant haplotype encoded receptors, for which non-Blacks had a slight advantage.

However, as shown in the current and a previous genotype study [44], within the same haplotypes

Blacks have therapeutic responses to bucindolol that are equivalent to those in non-Blacks. These

observations underscore the value of pharmacogenetic targeting in predicting drug response, which can

override any contribution by baseline characteristics such as race or other potentially confounding

variables.

Biased Ligand Properties of Bucindolol

A biased β-adrenergic receptor ligand can be defined as a compound that can signal through both G-

protein dependent and independent pathways [52, 53], favoring one or the other in certain contexts [47].

Cardiac myocyte surface membrane β_1- or β_2-ARs typically signal through the G-protein-cAMP/PKA or

CaMKII pathway [1], and internalized receptors [8, 10, 12, 30−32] through EGFR/ERK1/2. As it applies

to bucindolol or the structurally similar compound carvedilol [53], in systems that are hypersensitive for

the detection of G protein mediated adenylate cyclase stimulation/cAMP generation both compounds

exhibit weak β-agonist activity, but also can activate MAP kinase ERK1/2 independent of G-proteins

[14] qualifying them as biased ligands. For both compounds the β-agonist activity is insufficient for

activation of the β_1- or β_2-AR pathway in the human heart either in vitro [29, 54, 55] or in vivo [56, 57];

data from the current study indicate that in failing human hearts bucindolol is capable of activating

ERK1/2 signaling, further qualifying it as a biased ligand.

A comparison of factors influencing the biased ligand properties of bucindolol and carvedilol

includes: similar ligand binding characteristics that differ from other β-AR antagonists when co-

crystallized with the avian β_1-AR [53]; higher potency of bucindolol for ERK1/2 activation [14]; similar

guanine nucleotide modulatable binding (a characteristic of agonists) in human β_1- and β_2-ARs [54, 55];

and for bucindolol but not carvedilol, following 24 hours of incubation downregulation of β_1- and β_2-

ARs with no reduction in cognate mRNA [58] (evidence of internalization leading to receptor

degradation from trafficking to lysosomes [59, 60]. These latter experiments were performed in both

cultured avian cardiac myocytes (primarily β_1-ARs) and hamster β_2-AR containing smooth muscle cells

that express internalization-resistant β-AR haplotypes [61, 62]. In addition, bucindolol's or carvedilol's

ERK1/2 activation can be blocked by endocytosis/internalization inhibitors [63].

In the current study the increase in ERK1/2 phosphorylation in response to bucindolol occurred

rapidly (after a 10 minute incubation) as noted by others for isoproterenol [12] and nebivolol [17], and

converted to dephosphorylation by 60 minutes. The degree of early activation was quantitatively similar

to the statistically nonsignificant change for isoproterenol, and the nonbiased β-blocker metoprolol did

not activate ERK1/2 signaling. Five of the 6 isolated RV trabeculae preparations were from failing hearts

with a majority of internalization-resistant (*ADRB1+ADRB2*) haplotypes (four ≥3DIR and one 4DIR).

These findings support the possibility that the favorable effects of bucindolol in internalization-resistant

receptors could have been due to "cardioprotective" signaling mediated either by biased ligand-induced

receptor internalization [15−17] or ERK1/2 signaling independent of internalization [64, 65]. That a

biased ligand β-blocker can internalize relatively resistant β-AR haplotypes and activate ERK1/2

signaling has been previously established with carvedilol [16], in β_1-ARs expressed from the original

human β_1-AR cDNA that is Ser49Gly389 [66].

Carvedilol's interaction with β-AR haplotypes in clinical HF settings has not been investigated.

Despite their similarities bucindolol and carvedilol have somewhat different properties, including

moderate sympatholysis with bucindolol but not by carvedilol [1, 50] that likely contributes to the

former's enhanced effectiveness in HF patients homozygous for *ADRB1* Arg389 [1, 29, 67, 68]. Unlike

bucindolol, carvedilol has not exhibited selective enhancement of transplant-free survival in subjects

with *ADRB1* Arg389Arg [69, 70] genotypes. Bucindolol is also a more potent β_2-AR antagonist than

carvedilol, by approximately 5-8 fold [55].

Haplotype Targeting Expands the Pharmacogenetic Heart Failure or Atrial Fibrillation

Prevention Potential of Bucindolol

In BEST substudy subjects with LVEFs ≥0.20 the enhanced treatment effects of bucindolol in

subjects with both *ADRB1* and *ADRB2* Maximum Double Internalization-resistant (4DIR) diplotypes (a

25% subgroup) resulted in a 1.34 fold increase in treatment effects compared to the *ADRB1* Arg389Arg

index population, and 1.68 fold increase compared to the All-Genotypes cohort. If *ADRB1* Ser49Arg389

homozygous diplotypes AND 4DIR diplotypes were considered (a 10% subgroup) the treatment effect

enhancement compared to the index *ADRB1* Arg389Arg group was by 1.52 fold in the LVEF ≥0.20

cohort and by 1.40 fold in the All-LVEFs cohort, with respective increases vs. the All-Genotypes cohort

of 1.92 and 2.06 fold. Thus, efficacy enhancement through targeting internalization-resistant *ADRB1* and

ADRB2 haplotypes exceeds that for *ADRB1* Arg389Arg, and when combined with this genotype using

AND logic yields an approximate doubling in effectiveness.

When the *ADRB1* AND *ADRB2* ≥4DIR subgroup was combined with *ADRB1* Arg389Arg subjects

by OR logic the result was a 60-62% subgroup that was essentially identical to *ADRB1* Arg389Arg for

treatment effects, in both LVEF cohorts. As a result, in the BEST pharmacogenomics substudy the

addition of internalization-resistant *ADRB1* and *ADRB2* haplotypes ("haplotype targeting") to the

current *ADRB1* Arg389Arg targeting resulted in a larger population eligible for benefit, increasing from

45-47% to 60-62% of the all-genotype parent population. Since the BEST pharmacogenetic substudy

was enriched in Blacks, who have a lower frequency of both *ADRB1* Arg389 and internalization-

resistant *ADRB1* and *ADRB2* haplotypes, the race-adjusted eligibility for therapeutic enhancement in the

entire U.S. population would increase from 49% to 65%. Thus, by using the *ADRB1* Arg389Arg OR

(*ADRB1* AND *ADRB2* Internalization-resistant) joint criterion, the population eligible for treatment

enhancement can be increased by 33%.

Heart Failure Natural History Implications

The effects in the placebo-treated group could have implications for HF natural history, which

suggest that patients with ≥ 2 (*ADRB1+ADRB2*) internalizing haplotypes have a more favorable

prognosis, may not respond any better than placebo to a biased ligand β-blocker and could have worse

outcomes when treated with a non-biased β-blocker. Equivalently, in comprising a likely substantial

majority of HF patients (approximately 70%), patients with 3 or more internalization-resistant

haplotypes constitute a major at-risk population.

Potential Limitations

These data were generated in 2 previously conducted clinical trials, one of which [35] was from an

era when HFrEF therapy did not include two (sacubitril-valsartan and SGLT2 inhibitors) of the current

"Four Pillars" [71] of HF drug treatment, and the other [34] was prior to the regulatory approval and

therapeutic implementation of SGLT2is. Although the tested hypothesis was generated from

observations made in a more recent study [27], further clinical outcomes testing of the β-AR haplotype

internalization hypothesis is warranted in contemporary HF populations.

In addition to the biased ligand/receptor internalization/cardioprotective signaling hypothesis there

are other possible explanations for the differential effects of apparent receptor internalization treatment

associated clinical events. These include non-EGFR/ERK1/2 subcellular signal trafficking [72] of

encoded internalized haplotypes, and internalization-based exclusion of cell surface β-ARs from adverse

canonical signaling. In the latter paradigm the benefit of internalized, desensitized β-ARs in the presence

of placebo is simply to attenuate the adverse effects of cell surface canonical signaling [16, 73], while

bucindolol as a high affinity β_1-, β_2-AR antagonist [1, 54, 55] exerts a greater therapeutic effect when

receptors are non-internalized. These possible mechanistic explanations will need to be further

investigated in cardiac myocyte model systems expressing human β-AR haplotypes, using a non-

radioligand receptor internalization assay to accommodate the slow β_1- and β_2-AR offset kinetics of

bucindolol.

Conclusions

In HFrEF and HFmrEF patients, polymorphic loci in the intronless *ADRB1* and *ADRB2* genes

encoding receptor haplotypes that regulate internalization affect HF and AF clinical events as well as

biomarker behavior, in both placebo- and bucindolol-treated subjects. Including *ADRB1* and *ADRB2*

internalization-resistant haplotypes in bucindolol's pharmacogenetic targeting results in efficacy

enhancement similar to that produced by the *ADRB1* Arg389Arg genotype, and restricting the

*ADRB1*contribution to Arg389 in a internalization-resistant diplotype combination produces an additive

effect as would be expected by polymorphisms affecting different mechanisms. Haplotype

pharmacogenetic targeting is feasible in HF, and can identify populations at differential risk for serious

outcomes as well as candidates for increased therapeutic effectiveness.

Acknowledgements

Funding sources: This work was supported by National Heart, Lung, and Blood Institute (NHLBI) Grants HL052318, HL48013, HL071609, and HL077101. Additional support is from Genvara Biopharma, Westminster, CO. DNA samples from the BEST trial were provided by the BEST DNA Bank, co-sponsored by the NHLBI and the Department of Veterans Affairs Cooperative Studies. DNA Samples for the GENETIC-AF trial were provided by ARCA Biopharma (Westminster, CO), the trial's sponsor. We thank Rachel Rosenberg for manuscript and submission handling.

Author contributions

Conceptualization, M.R.B., I.A.C., S.B.L.; Methodology, I.A.C., M.R.B., D.S., W.A.M., E.R.J., L.A.W., A.D.K., A.V.A., S.B.L., D.K.; Software, I.A.C.; Validation, M.R.B., I.A.C., W.A.M., D.S.; Formal Analysis, I.A.C., M.R.B., D.S., W.A.M., L.A.W.; A.D.K. Investigation, S.B.L., M.R.B., I.A.C., D.S., E.R.J., W.A.M., L.A.W., D.K., A.D.K.; Resources, M.R.B., S.B.L., I.A.C., A.V.A., T.A.M; Data Curation, I.A.C, M.R.B., D.S.; Writing– Original Draft, M.R.B., I.A.C., D.S., W.A.M.; Writing – Review & Editing, M.R.B., I.A.C., S.B.L., T.A.M.,L.A.W.; Visualization, I.A.C. E.R.J., M.R.B., W.A.M; Supervision, M.R.B., S.B.L., T.A.M, D.S, A.V.A.; Project Administration, M.R.B., S.B.L.; Funding Acquisition, M.R.B., S.B.L.

Conflict of Interest Statement

M.R.B., I.A.C. and S.B.L. own stock in and are co-founders of Genvara Biopharma (Westminster, CO USA, current sponsor of bucindolol), and are Inventors on a patent covering some of the manuscript's content. The data contained in the manuscript were generated in M.R.B.'s and S.B.L.'s laboratories at the University of Colorado and University of South Florida respectively, and aspects of it contained in patent submissions are licensed to Genvara by these universities. M.R.B. and I.A.C. have partial salary support from Genvara; M.R.B. is Chief Science and Medical Officer of Genvara, and I.A.C. is its VP of Information Systems. M.R.B. and S.B.L. are also Principal Investigators on a submitted NIH grant related to this work.

Availability of Data and Materials

Outputs and code for structural equation modeling and haplotyping data are available on GitHub (https://github.com/CUAMC-SoM-Cardiology/M_Bristow-Lab). Group level data not included in the manuscript are available on request. De-identified subject level data for use in a research context will be available on request.

REFERENCES

1. Bristow MR. Treatment of chronic heart failure with β-adrenergic receptor antagonists: a convergence of receptor pharmacology and clinical cardiology. Circ Res. 2011;109(10):1176-94.

2. Liggett SB, Tepe NM, Lorenz JN, Canning AM, Jantz TD, Mitarai S, et al. Early and delayed consequences of beta(2)-adrenergic receptor overexpression in mouse hearts: critical role for expression level. Circulation. 2000;101(14):1707-14.

3. Tatman PD, Kao DP, Chatfield KC, Carroll IA, Wagner JA, Jonas ER, et al. An extensive β1-adrenergic receptor gene signaling network regulates molecular remodeling in dilated cardiomyopathies. JCI Insight. 2023;8(16):e169720.

4. Eichhorn EJ, Bristow MR. Medical therapy can improve the biological properties of the chronically failing heart: a new era in the treatment of heart failure. Circulation. 1996;94(9):2285-96.

5. Arora R. Recent insights into the role of the autonomic nervous system in the creation of substrate for atrial fibrillation: implications for therapies targeting the atrial autonomic nervous system. Circ Arrhythm Electrophysiol. 2012;5(4):850-59.

6. Noma T, Lemaire A, Naga Prasad SV, Barki-Harrington L, Tilley DG, Chen J, et al. Beta-arrestin-mediated beta1-adrenergic receptor transactivation of the EGFR confers cardioprotection. J Clin Invest. 2007;117(9):2445-58.

7. Rapacciuolo A, Suvarna S, Barki-Harrington L, Luttrell LM, Cong M, Lefkowitz RJ, Rockman HA. Protein kinase A and G protein-coupled receptor kinase phosphorylation mediates beta-1 adrenergic receptor endocytosis through different pathways. J Biol Chem. 2003;278(37):35403-11.

8. Tilley DG, Kim IM, Patel PA, Violin JD, Rockman HA. Beta-Arrestin mediates beta1-adrenergic receptor-epidermal growth factor receptor interaction and downstream signaling. J Biol Chem. 2009;284(30),20375-86.

9. von Zastrow M, Kobilka BK. Ligand-regulated internalization and recycling of human beta 2-adrenergic receptors between the plasma membrane and endosomes containing transferrin receptors. J Biol Chem. 1992;267(5):3530-38.

10. Daaka Y, Luttrell LM, Ahn S, Della Rocca GJ, Ferguson SS, Caron MG, Lefkowitz RJ. Essential role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. J Biol Chem. 1998;273(2):685-88.

11. Chen H, Ma N, Xia J, Liu J, Xu Z. β2-Adrenergic receptor-induced transactivation of epidermal growth factor receptor and platelet-derived growth factor receptor via Src kinase promotes rat cardiomyocyte survival. Cell Biol Int. 2012;36(3):237-44.

12. Grisanti LA, Talarico JA, Carter RL, Yu JE, Repas AA, Radcliffe SW, et al. β-Adrenergic receptor-mediated transactivation of epidermal growth factor receptor decreases cardiomyocyte apoptosis through differential subcellular activation of ERK1/2 and Akt. J Mol Cell Cardiol. 2014;72:39-51.

13. Han SO, Xiao K, Kim J, Wu JH, Wisler JW, Nakamura N, Freedman NJ, Shenoy SK. MARCH2 promotes endocytosis and lysosomal sorting of carvedilol-bound β(2)-adrenergic receptors. J Cell Biol. 2012;199(5):817-30.

14. Galandrin S, Bouvier M. Distinct signaling profiles of beta1 and beta2 adrenergic receptor ligands toward adenylyl cyclase and mitogen-activated protein kinase reveals the pluridimensionality of efficacy. Mol Pharmacol. 2006;70(5):1575-84.

15. Wisler JW, DeWire SM, Whalen EJ, Violin JD, Drake MT, Ahn S, et al. A unique mechanism of beta-blocker action: carvedilol stimulates beta-arrestin signaling. Proc Natl Acad Sci U S A. 2007;104(42):16657-62.

16. Kim IM, Tilley DG, Chen J, Salazar NC, Whalen EJ, Violin JD, Rockman HA. Beta-blockers alprenolol and carvedilol stimulate beta-arrestin-mediated EGFR transactivation. Proc Natl Acad Sci U.S.A. 2008;105(38):14555-60.

17. Erickson CE, Gul R, Blessing CP, Nguyen J, Liu T, Pulakat L, et al. The β-blocker Nebivolol is a GRK/β-arrestin biased agonist. PLoS One. 2013;8(8): e71980.

18. Levin MC, Marullo S, Muntaner O, Andersson B, Magnusson Y. The myocardium-protective Gly-49 variant of the beta 1-adrenergic receptor exhibits constitutive activity and increased desensitization and down-regulation. J Biol Chem. 2002; 277(34): 30429-35.

19. Rathz DA, Brown KM, Kramer LA, Liggett SB. Amino acid 49 polymorphisms of the human beta1-adrenergic receptor affect agonist-promoted trafficking. J Cardiovasc Pharmacol. 2002;39(2):155-60.

20. Sandilands A, Yeo G, Brown MJ. O'Shaughnessy KM. Functional responses of human beta1 adrenoceptors with defined haplotypes for the common 389R>G and 49S>G polymorphisms. Pharmacogenetics. 2004;14(6):343-49.

21. Zhang F, Steinberg SF. S49G and R389G polymorphisms of the β□-adrenergic receptor influence signaling via the cAMP-PKA and ERK pathways. Physiol Genomics. 2013;45(23):1186-92.

22. Dishy V, Sofowora GG, Xie HG, Kim RB, Byrne DW Stein CM, Wood,AJ. The effect of common polymorphisms of the beta2-adrenergic receptor on agonist-mediated vascular desensitization. N Engl J Med. 2001;345(14):1030-35.

23. January B, Seibold A, Whaley B, Hipkin RW, Lin D. Schonbrunn A, et al. Beta2-adrenergic receptor desensitization, internalization, and phosphorylation in response to full and partial agonists. J Biol Chem. 1997;272 (38):23871-79.

24. Green SA, Turki J, Innis M, Liggett SB. Amino-terminal polymorphisms of the human beta 2-adrenergic receptor impart distinct agonist-promoted regulatory properties. Biochemistry. 1994;33(32):9414-19.

25. Suzuki T, Nguyen CT, Nantel F, Bonin H, Valiquette M, Frielle T, Bouvier M. Distinct regulation of beta 1- and beta 2-adrenergic receptors in Chinese hamster fibroblasts. Mol Pharmacol. 1992;41(3):542-48.

26. Kobilka BK, Dixon RA, Frielle T, Dohlman HG, Bolanowski MA, Sigal IS, et al. cDNA for the human beta 2-adrenergic receptor: a protein with multiple membrane-spanning domains and encoded by a gene whose chromosomal location is shared with that of the receptor for platelet-derived growth factor. Proc Natl Acad Sci U S A. 1987;84(1):46-50.

27. Chevalier P, Roy P, Bessiere F, Morel E, Ankou B, Morgan G, et al. Impact of neuroeffector adrenergic receptor polymorphisms on incident ventricular fibrillation during acute myocardial ischemia. J Am Heart Assoc. 2023;12(6): e025368.

28. Mason DA, Moore JD, Green SA, Liggett SB. A gain-of-function polymorphism in a G-protein coupling domain of the human β1-adrenergic receptor. J Biol Chem. 1999;274(18):12670-74.

29. Liggett SB, Mialet-Perez J, Thaneemit-Chen S, Weber SA, Greene SM, Hodne D. A polymorphism within a conserved β1-adrenergic receptor motif alters cardiac function and β-blocker response in human heart failure. Proc Natl Acad Sci U.S.A. 2006;103(30):11288-93.

30. Eichel K, Jullié D, von Zastrow M. β-Arrestin drives MAP kinase signalling from clathrin-coated structures after GPCR dissociation. Nat Cell Biol. 2016;18(3):303-10.

31. Maudsley S, Pierce KL, Zamah AM, Miller WE, Ahn S, Daaka Y, et al. The beta(2)-adrenergic receptor mediates extracellular signal-regulated kinase activation via assembly of a multi-receptor complex with the epidermal growth factor receptor. J Biol Chem. 2000;275(3):9572-80.

32. Kwon Y, Mehta S, Clark M, Walters G, Zhong Y, Lee HN, et al. Non-canonical β-adrenergic activation of ERK at endosomes. *Nature*. 2022;611(7934):173-79.

33. Börjesson M, Magnusson Y, Hjalmarson A, Andersson B. A novel polymorphism in the gene coding for the beta(1)-adrenergic receptor associated with survival in patients with heart failure. Eur Heart J. 2000;21(22):1853-58.

34. Piccini JP, Abraham WT, Dufton C, Carroll IA, Healey JS, van Veldhuisen J, et al. Bucindolol for the maintenance of sinus rhythm in a genotype-defined HF population. JACC Heart Fail. 2019;7(7):586-98.

35. Eichhorn EJ, Domanski MJ, Krause-Steinrauf H, Bristow MR, Lavori PW, BEST Investigators. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. N Engl J Med. 2001;344(22):1659-67.

36. Spooner W, McLaren W, Slidel T, Finch DK, Butler R, Campbell J, et al. Haplosaurus computes protein haplotypes for use in precision drug design. Nat Commun. 2018;9(1):4128.

37. Judson R, Stephens JC, Windemuth AA. The predictive power of haplotypes in clinical response. Pharmacogenomics. 2000;1(1):15-26.

38. Drysdale CM, McGraw D., Stack CB, Stephens JC, Judson RS, Nandabalan K, et al. Complex promoter and coding region beta 2-adrenergic receptor haplotypes alter receptor expression and predict in vivo responsiveness. Proc Natl Acad Sci U S A. 2000;97(19):10483-88.

39. Small KM, Rathz, DA, Liggett, SB. Identification of adrenergic receptor polymorphisms. Methods Enzymol. 2002;343:459-75.

40. Sinnwell JP, Schaid DJ. HAPLO.STATS: Statistical analysis of haplotypes with traits and covariates when linkage phase is ambiguous. R package version 1.8.7. https://CRAN.R-project.org/package=haplo.stats. Accessed 4/16/2024.

41. Porubsky D, Ebert P, Audano PA, Vollger MR, Harve W.T, Marijon P, et al. Fully phased human genome assembly without parental data using single-cell strand sequencing and long reads. Nat Biotechnol. 2021;39(3):302-08.

42. Shen L, Jhund PS, Mogensen UM, Køber L, Claggett B, Rogers JK, McMurray JJ, et al. Re-examination of the BEST trial using composite outcomes, including emergency department visits. JACC Heart Fail. 2017;5(8):591-99.

43. O'Connor CM, Fiuzat M, Carson PE, Anand I, Plehn JF, Gottlieb SS, et al. Combinatorial pharmacogenetic interactions of bucindolol and β1, α2C adrenergic receptor polymorphisms. PLoS One. 2012;7(10): e44324.

44. Taylor MR, Sun AY, Davis G, Fiuzat M, Liggett SB, Bristow MR. Race, common genetic variation, and therapeutic response disparities in heart failure. JACC Heart Fail. 2014 2(6):561-72.

45. Tishkoff SA, Verrelli BC. Patterns of human genetic diversity: implications for human evolutionary history and disease. Annu Rev Genomics Hum Genet. 2003; 4:293-40.

46. Stephens JC, Schneider JA, Tanguay DA, Choi J, Acharya T, Stanley SE, et al. Haplotype variation and linkage disequilibrium in 313 human genes. Science. 2001;293(5529):489-93.

47. Panebra A, Wang WC, Malone MM, Pitter DR, Weiss ST, Hawkins GA, Liggett SB. Common ADRB2 haplotypes derived from 26 polymorphic sites direct beta2-adrenergic receptor expression and regulation phenotypes. PLoS One. 2010;5(7):e11819.

48. Forleo C, Sorrentino S, Guida P, Romito R, De Tommasi E, Iacoviello M, Pitzalis M. Beta1- and beta2-adrenergic receptor polymorphisms affect susceptibility to idiopathic dilated cardiomyopathy. J Cardiovasc Med (Hagerstown). 2007;8(8):589-95.

49. Piccini JP, Dufton C, Carroll IA, Healey JS, Abraham WT, Khaykin Y, et al. Bucindolol decreases atrial fibrillation burden in patients with heart failure and the *ADRB1* Arg389Arg genotype. Circ Arrhythm Electrophysiol. 2021;14(8):e009591.

50. Bristow MR, Abraham WT, Yoshikawa T, White M, Hattler BG, Crisman TS, et al. Second- and third-generation beta-blocking drugs in chronic heart failure. Cardiovasc Drugs Ther. 1997;11(Suppl 1):291-96.

51. Freeman K, Lerman I, Kranias EG, Bohlmeyer T, Bristow MR, Lefkowitz RJ, et al. Alterations in cardiac adrenergic signaling and calcium cycling differentially affect the progression of cardiomyopathy. J Clin Invest. 2001;107(8): 967-74.

52. Drake MT, Violin JD, Whalen EJ, Wisler JW, Shenoy, SK, Lefkowitz RJ. Beta-arrestin-biased agonism at the beta2-adrenergic receptor. J Biol Chem. 2008;283(9):5669-76.

53. Warne T, Edwards PC, Leslie AG, Tate CG. Crystal structures of a stabilized β1-adrenoceptor bound to the biased agonists bucindolol and carvedilol. Structure. 2012;20(5):841-49.

54. Hershberger RE, Wynn JR, Sundberg L, Bristow MR. Mechanism of action of bucindolol in human ventricular myocardium. J Cardiovasc Pharmacol 1990;15(6):959–67.

55. Yoshikawa T, Port JD, Asano K, Chidiak P, Bouvier M, Dutcher D, et al. Cardiac adrenergic receptor effects of carvedilol. Eur Heart J. 1996;17(Suppl B):B8-B16.

56. Bristow MR, Roden RL, Lowes BD, Gilbert EM, Eichhorn EJ. The role of third generation β-blocking agents in chronic heart failure. Clin Cardiol. 1998;21(12 Suppl 1):I3-13.

57. Bristow MR, O'Connell JB, Gilbert EM, French WJ, Leatherman G, Kantrowitz NE, et al. Dose-response of chronic beta-blocker treatment in heart failure from either idiopathic dilated or ischemic cardiomyopathy. Circulation. 1994 89(4):1632-42.

58. Asano K, Zisman LS, Yoshikawa T, Headly V, Bristow MR, Port DJ. Bucindolol, a nonselective B1- and B2- adrenergic receptor antagonist, decreases β-adrenergic receptor density in cultured embryonic chick cardiac myocyte membranes. J Cardiovasc Pharmacol. 2001;37(6):678-91.

59. Moore RH, Tuffaha A, Millman EE, Dai W, Hall HS, Dickey B.F, Knoll BJ. Agonist-induced sorting of human beta2-adrenergic receptors to lysosomes during downregulation. J Cell Sci. 1999;112(Pt 3):329-38.

60. Nakagawa T, Asahi M. β1-adrenergic receptor recycles via a membranous organelle, recycling endosome, by binding with sorting nexin27. J Membr Biol 2013;246(7):571-579.

61. Chicken *ADRB1*, human equivalent haplotype is Ser49Arg389. https://www.ncbi.nlm.nih.gov/nuccore/XM_040703024.2?report=GenBank . Accessed 3/10/2025.

62. Hamster *ADRB2*, human equivalent haplotype is Gly16Glu27. https://www.ncbi.nlm.nih.gov/nucleotide/XM_027402776.2?report=genbank&log$=nuclalign&blast_rank=1&RID=WYT2ZYU3016 . Accessed 3/10/2025.

63. Ajumobi, T. Beta-Blockers Act through clathrin-dependent internalization and EGFR transactivation to promote ERK Phosphorylation. (2014). Graduate Thesis Collection. *267*. https://digitalcommons.butler.edu/grtheses/267. Accessed 4/24/2024.

64. Galandrin S, Oligny-Longpré G, Bonin H, Ogawa K, Galés C, Bouvier M**.** Conformational rearrangements and signaling cascades involved in ligand-biased mitogen-activated protein kinase signaling through the beta1-adrenergic receptor. Mol Pharmacol. 2008;74(1):162-72.

65. Huang J, Sun Y, Huang XY. Distinct roles for Src tyrosine kinase in beta2-adrenergic receptor signaling to MAPK and in receptor internalization. *J Biol Chem*. 2004;279(20):21637-42.

66. Frielle T, Collins S, Daniel KW, Caron MG, Lefkowitz RJ, Kobilka BK. Cloning of the cDNA for the human beta 1-adrenergic receptor. Proc Natl Acad Sci U S A. 1987;84(22):7920-24.

67. Aleong RG, Sauer WH, Robertson AD, Liggett SB, Bristow MR. Prevention of atrial fibrillation by bucindolol is dependent on the beta389 Arg/Gly adrenergic receptor polymorphism. JACC: Heart Fail. 2012;1(4):338-44

68. Aleong RG, Aleong RG, Sauer WH, Robertson AD, Liggett SB, Bristow MR. Adrenergic receptor polymorphisms and prevention of ventricular arrhythmias with bucindolol in patients with chronic heart failure. Circ Arrhythm Electrophysiol. 2013;6(1):137-43.

69. Sehnert AJ, Daniels SE, Elashoff M, Wingrove JA, Burrow CR, Horne B, et al. Lack of association between adrenergic receptor genotypes and survival in heart failure patients treated with carvedilol or metoprolol. J Am Coll Cardiol. 2008;52(8):644-51.

70. White HL, de Boer RA, Maqbool A, Greenwood D, van Veldhuisen DJ, Cuthbert R, et al. An evaluation of the beta-1 adrenergic receptor Arg389Gly polymorphism in individuals with heart failure: a MERIT-HF sub-study. Eur J Heart Fail. 2003*;*5(4):463-68.

71. Straw S, McGinlay M, Witte KK. Four pillars of heart failure: contemporary pharmacological therapy for heart failure with reduced ejection fraction. Open Heart. 2021;8(1):e001585.

72. Wei W, Smrcka, AV. Subcellular beta-adrenergic receptor signaling in cardiac physiology and disease. J Cardiovasc Pharmacol. 2022 80(3):334-41.

73. Tan LB, Benjamin IJ, Clark WA. Beta adrenergic receptor desensitization may serve a cardioprotective role. Cardiovasc Res. 1992;26(6):608-14.

TABLES

Table 1. *ADRB1*, *ADRB2* diplotype frequencies (*f*) by race, BEST Adrenergic Receptor Polymorphism Substudy.

Gene	Diplotype	Black N = 205 *ADRB1*, 207 *ADRB2* (*f*)	non-Black N = 824 *ADRB1*, 833 *ADRB2* (*f*)	All Races N = 1029 *ADRB1*, 1040 *ADRB2* (*f*)	B/non-B Fold Difference	P value, Black vs. non-Black
ADRB1	Ser49Arg389 Ser49Arg389	21 (0.10)	281 (0.34)	302 (0.29)	0.29	<0.0001
	Gly49Arg389 Gly49Arg389	14 (0.068)	24 (0.029)	38 (0.037)	2.34	0.012
	Ser49Gly389 Ser49Gly389	35 (0.17)	58 (0.070)	93 (0.090)	2.43	<0.0001
	Ser49Arg389 Gly49Arg389	30 (0.15)	119 0.14)	149 (0.145)	1.07	0.91
	Gly49Arg389 Ser49Gly389	40 (0.20)	61 (0.074)	101 (0.098)	2.70	<0.0001
	Ser49Arg389 Ser49Gly389	65 (0.32)	281 (0.34)	346 (0.34)	0.94	0.56
ADRB2	Gly16Gln27 Gly16Gln27	19 (0.092)	51 (0.061)	70 (0.067)	1.51	0.12
	Arg16Gln27 Arg16Gln27	41 (0.20)	137 (0.16)	178 (0.17)	1.25	0.26
	Gly16Glu27 Gly16Glu27	7 (0.034)	136 (0.16)	143 (0.14)	0.21	<0.0001
	Arg16Gln27 Gly16Gln27	77 (0.37)	133 (0.16)	210 (0.20)	2.31	<0.0001
	Arg16Gln27 Gly16Glu27	35 (0.17)	244 (0.29)	279 (0.27)	0.59	0.0002
	Gly16Gln27 Gly16Glu27	28 (0.14)	132 (0.16)	160 (0.15)	0.23	0.45
	Not Arg16Gln27 Arg16Gln27	166 0.80	696 0.84	862 0.83	0.95	0.26
	1 or 2 copies Arg16Gln27	153 (0.74)	514 (0.62)	667 (0.64)	1.19	0.0012
	No copies Arg16Gln27	54 (0.26)	319 (0.38)	373 (0.36)	0.68	0.0012

f = Diplotype frequency (Diplotype N/Total N).

Main Effects of Internalization Tier within Placebo Patients
(Unadjusted, All LVEF Subjects; N = 521)

†Intrnlz H'type #	Tier	Endpoint	Effect	p-value	P(Event) by Haplo (Y/N) [N]	Odds Ratio
3-4		CVM		0.129	8.3%[36], 18.8%[485]	0.39 [0.12, 1.31]
3-4		AF/AFL/ACM		0.165	14.3%[28], 26.4%[375]	0.46 [0.16, 1.37]
3-4	1	CVM/HFH		0.004**	19.4%[36], 45.4%[485]	0.29 [0.12, 0.68]
3-4	(≥3DI)	Heart Failure Hosp		0.019*	16.7%[36], 36.9%[485]	0.34 [0.14, 0.84]
3-4		ACM/Tx		0.105	11.1%[36], 23.1%[485]	0.42 [0.14, 1.20]
3-4		Cardiovascular Hosp		0.049*	30.6%[36], 47.8%[485]	0.48 [0.23, 1.00]
3-4		ACM		0.083	8.3%[36], 20.8%[485]	0.35 [0.10, 1.15]
2		CVM		0.51	16.2%[136], 18.7%[385]	0.84 [0.50, 1.42]
2		AF/AFL/ACM		0.26	21.5%[107], 27.0%[296]	0.74 [0.44, 1.25]
2	2	CVM/HFH		0.21	39.0%[136], 45.2%[385]	0.77 [0.52, 1.15]
2	(2I)	Heart Failure Hosp		0.190	30.9%[136], 37.1%[385]	0.76 [0.50, 1.15]
2		ACM/Tx		0.31	19.1%[136], 23.4%[385]	0.77 [0.48, 1.26]
2		Cardiovascular Hosp		0.038*	39.0%[136], 49.4%[385]	0.66 [0.44, 0.98]
2		ACM		0.30	16.9%[136], 21.0%[385]	0.76 [0.46, 1.27]
0-1		CVM		0.145	19.8%[349], 14.5%[172]	1.45 [0.88, 2.39]
0-1		AF/AFL/ACM		0.071	28.4%[268], 20.0%[135]	1.58 [0.96, 2.61]
0-1	3	CVM/HFH		0.005**	47.9%[349], 34.9%[172]	1.71 [1.17, 2.50]
0-1	(≥3DIR)	Heart Failure Hosp		0.011*	39.3%[349], 27.9%[172]	1.67 [1.12, 2.48]
0-1		ACM/Tx		0.064	24.6%[349], 17.4%[172]	1.55 [0.97, 2.46]
0-1		Cardiovascular Hosp		0.003**	51.3%[349], 37.2%[172]	1.78 [1.22, 2.58]
0-1		ACM		0.053	22.3%[349], 15.1%[172]	1.62 [0.99, 2.63]

†Number of internalizing haplotypes

Odds Ratio of Event (axis: 0.1 0.25 0.5 1 2 4 10)

Figure 1. All LVEF cohort patients randomized to placebo in the BEST Adrenergic Receptor Polymorphism substudy, forest plot of *ADRB1+ADRB2* combination diplotypes arranged by the number of internalizing haplotypes (Column 1), displaying Odds Ratios (ORs, Columns 4 and 7) between event rates (Column 6) in the Column 1 diplotypes vs. all other combination diplotypes. The *ADRB1*, *ADRB2* diplotype combinations are displayed by "≥3 Double Internalizing" combinations in **Tier 1** (≥3DI, 1 or 2 copies of *ADRB1* Gly49Arg389 AND *ADRB2* Arg16Gln27, for a total of 3-4 copies of an internalizing haplotype); "Incompletely Internalizing" in **Tier 2** (**2I**), 2 total copies of internalizing haplotypes, either 1 copy of *ADRB1* AND *ADRB2*, or 2 copies on one of them); and "≥3 Double Internalization-resistant" haplotype combinations in **Tier 3** (≥3DIR, 0-1 total copies of *ADRB1* Gly49Arg389 AND *ADRB2* Arg16Gln27, 3-4 copies of internalization-resistant haplotypes). The counterpart, comparison haplotypes used to generate ORs follow OR logic, i.e. if either the *ADRB1* or the *ADRB2* haplotype doesn't match the cognate Column 1 haplotype then the pair becomes a member of the counterpart group. Column 6 lists the event rate and [N] in the paired haplotypes in column 1, followed by the event rate and [N] in the counterpart, comparator groups. CVM=Cardiovascular Mortality; AF/AFL/ACM=Atrial fibrillation, Atrial Flutter or All-Cause Mortality; CVM/HFH=CVM or Heart Failure hospitalization; HFH=Heart Failure Hospitalization; ACM/Tx=ACM or Cardiac Transplantation; CVH=Cardiovascular Hospitalization; ACM=All-Cause Mortality. For the AF/AFL/ACM endpoint, only non-AF patients at the time of randomization are analyzed, resulting in a smaller N than for the 6 HF endpoints. For all endpoints only first events are considered. Odds Ratios (ORs) are calculated by:

[(Index event rate in %) / (100 − Index event rate)] / [(Comparator event rate) / (100 − Comparator event rate)], or in Row 1 (8.3/91.7) / (18.8/81.2) = 0.39.

Figure 2. All LVEF cohort BEST substudy patients randomized to bucindolol in the BEST Adrenergic Receptor Polymorphism substudy. The forest plot is constructed as for **Figure 1**.

Main Effects of Treatment within Internalization Tier
(Unadjusted, All LVEF Subjects; N = 1029)

†Intrnlz H'type #	Tier	Endpoint	Effect	p-value	P(Event) by Rx (B/P) [N]	Odds Ratio
3-4		CVM		0.97	8.6%[35], 8.3%[36]	1.03 [0.19, 5.49]
3-4		AF/AFL/ACM		0.41	23.1%[26], 14.3%[28]	1.80 [0.45, 7.28]
3-4	1	CVM/HFH		0.53	25.7%[35], 19.4%[36]	1.43 [0.47, 4.40]
3-4	(≥3DI)	Heart Failure Hosp		0.51	22.9%[35], 16.7%[36]	1.48 [0.46, 4.82]
3-4		ACM/Tx		0.69	14.3%[35], 11.1%[36]	1.33 [0.33, 5.44]
3-4		Cardiovascular Hosp		0.74	34.3%[35], 30.6%[36]	1.19 [0.44, 3.21]
3-4		ACM		0.43	14.3%[35], 8.3%[36]	1.83 [0.40, 8.34]
2		CVM		0.30	11.7%[120], 16.2%[136]	0.68 [0.33, 1.41]
2		AF/AFL/ACM		0.38	16.7%[96], 21.5%[107]	0.73 [0.36, 1.48]
2	2	CVM/HFH		0.35	33.3%[120], 39.0%[136]	0.78 [0.47, 1.31]
2	(2I)	Heart Failure Hosp		0.66	28.3%[120], 30.9%[136]	0.88 [0.52, 1.52]
2		ACM/Tx		0.38	15.0%[120], 19.1%[136]	0.75 [0.39, 1.44]
2		Cardiovascular Hosp		0.60	35.8%[120], 39.0%[136]	0.87 [0.53, 1.45]
2		ACM		0.55	14.2%[120], 16.9%[136]	0.81 [0.41, 1.60]
0-1		CVM		0.022*	13.3%[353], 19.8%[349]	0.62 [0.42, 0.93]
0-1		AF/AFL/ACM		0.113	22.4%[268], 28.4%[268]	0.73 [0.49, 1.08]
0-1	3	CVM/HFH		0.001**	35.7%[353], 47.9%[349]	0.60 [0.45, 0.82]
0-1	(≥3DIR)	Heart Failure Hosp		0.005**	29.2%[353], 39.3%[349]	0.64 [0.47, 0.87]
0-1		ACM/Tx		0.045*	18.4%[353], 24.6%[349]	0.69 [0.48, 0.99]
0-1		Cardiovascular Hosp		0.005**	40.8%[353], 51.3%[349]	0.65 [0.49, 0.88]
0-1		ACM		0.093	17.3%[353], 22.3%[349]	0.73 [0.50, 1.05]

†Number of internalizing haplotypes

Odds Ratio: 0.1 0.25 0.5 1 2 4 10

Figure 3. All LVEF cohort subjects, forest plot of bucindolol vs. placebo treatment effect (bucindolol event rate vs. placebo event rate) Odds Ratios (ORs) by internalizing haplotype number in the Column 1 *ADRB1+ADRB2* combination diplotypes. [Ns] are the number of subjects in the bucindolol and placebo groups respectively; otherwise set-up is as in Figures 1 and 2.

Figure 4. Time to first event curves in bucindolol vs. placebo treated patients with Maximum Double Internalization-resistant (**4DIR**), total of 4 internalization-resistant haplotypes), LVEF ≥0.20 cohort N=184. **A.** All-cause Mortality or Cardiac Transplantation (ACM/Tx); **B.** Heart Failure Hospitalization (HFH); **C.** Cardiovascular Mortality or Heart Failure Hospitalization (CVM/HFH); **D.** Cardiovascular Hospitalization (CVH).

Fig 5. Time to first event curves curves in Blacks or non-Blacks in the All LVEF cohort for the CVM/HFH (**A**) and CVH (**B**) time to first event endpoints, with haplotype categories of: Double Internalizing (≥2**DI**) = ≥1 *ADRB1* AND *ADRB2* internalizing haplotypes (*ADRB1* Gly49Arg389, *ADRB2* Arg16Gln27); Double internalization-resistant (≥2**IR**) = all other haplotypes. The ≥2DI subpopulation is 18.4% of subjects treated with placebo.

Fig 6. GENETIC-AF study, peripheral venous plasma levels of NT-proBNP (top panel) or norepinephrine (bottom panel) for subjects with ≥2 double internalizing *ADRB1* AND *ADRB2* haplotypes (**≥2DI**) or ≥2 Internalization-resistant haplotypes (**≥2IR**) in the Bucindolol or Metoprolol Groups. Central tendencies of data are in mean±SD, with non-parametric tests used for determining statistical significance.

Figure 7. A. ERK1/2 phosphorylation (pERK) and total ERK in preparations of right ventricular trabeculae incubated for 10 minutes with vehicle (Veh); bucindolol (Buc, 1e-6M); metoprolol (Met, 1e-5M); or isoproterenol (Iso, 1e-6M) (patient 5 in Table S17). **B**. Box plots, median and 25, 75 percentile plus individual values of pERK1/2/total ERK1/2. L-panel, Data at 5 (N=1) and 10 minutes incubation (N=5); R-panel, 60 minutes incubation. Drug concentrations are bucindolol 1e-6M, metoprolol 1e-5M and isoproterenol 1e-6M. Vehicle is H_2O. Data are normally distributed by the Anderson-Darling test; statistical analysis (Table S17) is by ANOVA/mixed model with repeat measures and a Holm-Sidak's multiple comparison test compared to Vehicle control. *P=0.0027; †P=0.037 for lower value compared to 5/10 minutes (min) and interaction with time; ‡5 min incubation, all other points are 10 min.

Main Effects of Internalization Tier within Placebo Patients
(Unadjusted, All LVEF Subjects; N = 521)

†Intrnlz H'type #	Tier	Endpoint	Effect	p-value	P(Event) by Haplo (Y/N) [N]	Odds Ratio
3-4		CVM		0.129	8.3%[36], 18.8%[485]	0.39 [0.12, 1.31]
3-4		AF/AFL/ACM		0.165	14.3%[28], 26.4%[375]	0.46 [0.16, 1.37]
3-4	**1**	CVM/HFH		0.004**	19.4%[36], 45.4%[485]	0.29 [0.12, 0.68]
3-4	**(≥3DI)**	Heart Failure Hosp		0.019*	16.7%[36], 36.9%[485]	0.34 [0.14, 0.84]
3-4		ACM/Tx		0.105	11.1%[36], 23.1%[485]	0.42 [0.14, 1.20]
3-4		Cardiovascular Hosp		0.049*	30.6%[36], 47.8%[485]	0.48 [0.23, 1.00]
3-4		ACM		0.083	8.3%[36], 20.8%[485]	0.35 [0.10, 1.15]
2		CVM		0.51	16.2%[136], 18.7%[385]	0.84 [0.50, 1.42]
2		AF/AFL/ACM		0.26	21.5%[107], 27.0%[296]	0.74 [0.44, 1.25]
2	**2**	CVM/HFH		0.21	39.0%[136], 45.2%[385]	0.77 [0.52, 1.15]
2	**(2I)**	Heart Failure Hosp		0.190	30.9%[136], 37.1%[385]	0.76 [0.50, 1.15]
2		ACM/Tx		0.31	19.1%[136], 23.4%[385]	0.77 [0.48, 1.26]
2		Cardiovascular Hosp		0.038*	39.0%[136], 49.4%[385]	0.66 [0.44, 0.98]
2		ACM		0.30	16.9%[136], 21.0%[385]	0.76 [0.46, 1.27]
0-1		CVM		0.145	19.8%[349], 14.5%[172]	1.45 [0.88, 2.39]
0-1		AF/AFL/ACM		0.071	28.4%[268], 20.0%[135]	1.58 [0.96, 2.61]
0-1	**3**	CVM/HFH		0.005**	47.9%[349], 34.9%[172]	1.71 [1.17, 2.50]
0-1	**(≥3DIR)**	Heart Failure Hosp		0.011*	39.3%[349], 27.9%[172]	1.67 [1.12, 2.48]
0-1		ACM/Tx		0.064	24.6%[349], 17.4%[172]	1.55 [0.97, 2.46]
0-1		Cardiovascular Hosp		0.003**	51.3%[349], 37.2%[172]	1.78 [1.22, 2.58]
0-1		ACM		0.053	22.3%[349], 15.1%[172]	1.62 [0.99, 2.63]

†Number of internalizing haplotypes

0.1   0.25 0.5   1   2   4   10

Odds Ratio of Event

Main Effects of Internalization Tier within Bucindolol Patients
(Unadjusted, All LVEF Subjects; N = 508)

†Intrnlz H'type #	Tier	Endpoint	Effect	p-value	P(Event) by Haplo (Y/N) [N]	Odds Ratio
3-4		CVM		0.46	8.6%[35], 12.9%[473]	0.63 [0.19, 2.13]
3-4		AF/AFL/ACM		0.79	23.1%[26], 20.9%[364]	1.14 [0.44, 2.93]
3-4	**1**	CVM/HFH		0.26	25.7%[35], 35.1%[473]	0.64 [0.29, 1.40]
3-4	**(≥3DI)**	Heart Failure Hosp		0.44	22.9%[35], 29.0%[473]	0.73 [0.32, 1.64]
3-4		ACM/Tx		0.62	14.3%[35], 17.5%[473]	0.78 [0.30, 2.08]
3-4		Cardiovascular Hosp		0.54	34.3%[35], 39.5%[473]	0.80 [0.39, 1.64]
3-4		ACM		0.73	14.3%[35], 16.5%[473]	0.84 [0.32, 2.24]
2		CVM		0.72	11.7%[120], 12.9%[388]	0.89 [0.47, 1.68]
2		AF/AFL/ACM		0.23	16.7%[96], 22.4%[294]	0.69 [0.38, 1.26]
2	**2**	CVM/HFH		0.77	33.3%[120], 34.8%[388]	0.94 [0.61, 1.45]
2	**(2I)**	Heart Failure Hosp		0.95	28.3%[120], 28.6%[388]	0.99 [0.63, 1.55]
2		ACM/Tx		0.44	15.0%[120], 18.0%[388]	0.80 [0.46, 1.41]
2		Cardiovascular Hosp		0.39	35.8%[120], 40.2%[388]	0.83 [0.54, 1.27]
2		ACM		0.46	14.2%[120], 17.0%[388]	0.81 [0.45, 1.43]
0-1		CVM		0.46	13.3%[353], 11.0%[155]	1.25 [0.69, 2.25]
0-1		AF/AFL/ACM		0.33	22.4%[268], 18.0%[122]	1.31 [0.76, 2.26]
0-1	**3**	CVM/HFH		0.37	35.7%[353], 31.6%[155]	1.20 [0.80, 1.80]
0-1	**(≥3DIR)**	Heart Failure Hosp		0.63	29.2%[353], 27.1%[155]	1.11 [0.73, 1.69]
0-1		ACM/Tx		0.33	18.4%[353], 14.8%[155]	1.30 [0.77, 2.17]
0-1		Cardiovascular Hosp		0.26	40.8%[353], 35.5%[155]	1.25 [0.85, 1.85]
0-1		ACM		0.39	17.3%[353], 14.2%[155]	1.26 [0.74, 2.14]

†Number of internalizing haplotypes

Odds Ratio of Event (axis: 0.1 0.25 0.5 1 2 4 10)

Main Effects of Treatment within Internalization Tier
(Unadjusted, All LVEF Subjects; N = 1029)

†Intrnlz H'type #	Tier	Endpoint	Effect	p-value	P(Event) by Rx (B/P) [N]	Odds Ratio
3-4		CVM		0.97	8.6%[35], 8.3%[36]	1.03 [0.19, 5.49]
3-4		AF/AFL/ACM		0.41	23.1%[26], 14.3%[28]	1.80 [0.45, 7.28]
3-4	1	CVM/HFH		0.53	25.7%[35], 19.4%[36]	1.43 [0.47, 4.40]
3-4	(≥3DI)	Heart Failure Hosp		0.51	22.9%[35], 16.7%[36]	1.48 [0.46, 4.82]
3-4		ACM/Tx		0.69	14.3%[35], 11.1%[36]	1.33 [0.33, 5.44]
3-4		Cardiovascular Hosp		0.74	34.3%[35], 30.6%[36]	1.19 [0.44, 3.21]
3-4		ACM		0.43	14.3%[35], 8.3%[36]	1.83 [0.40, 8.34]
2		CVM		0.30	11.7%[120], 16.2%[136]	0.68 [0.33, 1.41]
2		AF/AFL/ACM		0.38	16.7%[96], 21.5%[107]	0.73 [0.36, 1.48]
2	2	CVM/HFH		0.35	33.3%[120], 39.0%[136]	0.78 [0.47, 1.31]
2	(2I)	Heart Failure Hosp		0.66	28.3%[120], 30.9%[136]	0.88 [0.52, 1.52]
2		ACM/Tx		0.38	15.0%[120], 19.1%[136]	0.75 [0.39, 1.44]
2		Cardiovascular Hosp		0.60	35.8%[120], 39.0%[136]	0.87 [0.53, 1.45]
2		ACM		0.55	14.2%[120], 16.9%[136]	0.81 [0.41, 1.60]
0-1		CVM		0.022*	13.3%[353], 19.8%[349]	0.62 [0.42, 0.93]
0-1		AF/AFL/ACM		0.113	22.4%[268], 28.4%[268]	0.73 [0.49, 1.08]
0-1	3	CVM/HFH		0.001**	35.7%[353], 47.9%[349]	0.60 [0.45, 0.82]
0-1	(≥3DIR)	Heart Failure Hosp		0.005**	29.2%[353], 39.3%[349]	0.64 [0.47, 0.87]
0-1		ACM/Tx		0.045*	18.4%[353], 24.6%[349]	0.69 [0.48, 0.99]
0-1		Cardiovascular Hosp		0.005**	40.8%[353], 51.3%[349]	0.65 [0.49, 0.88]
0-1		ACM		0.093	17.3%[353], 22.3%[349]	0.73 [0.50, 1.05]

†Number of internalizing haplotypes

Odds Ratio
0.1   0.25   0.5   1   2   4   10

A. ACM/Tx, BEST: Maximum Double Internalization-resistant *ADRB1* AND *ADRB2* (≥4DIR) LVEF ≥ 20

Event Rates:
Bucindolol Group: 3/86 (3.5%)
Placebo Group: 20/98 (20.4%)
Hazard Ratio = 0.17 (0.05,0.56)
P-value = 0.004

B. AF/AFL/ACM, BEST: Maximum Double Internalization-resistant *ADRB1* AND *ADRB2* (≥4DIR) LVEF ≥ 20

Event Rates:
Bucindolol Group: 7/67 (10.4%)
Placebo Group: 19/76 (25%)
Hazard Ratio = 0.41 (0.17,0.97)
P-value = 0.042

C. CVM/HFH, BEST: Maximum Double Internalization-resistant *ADRB1* AND *ADRB2* (≥4DIR) LVEF ≥ 20

Event Rates:
Bucindolol Group: 19/86 (22.1%)
Placebo Group: 41/98 (41.8%)
Hazard Ratio = 0.49 (0.28,0.85)
P-value = 0.011

D. CVH, BEST: Maximum Double Internalization-resistant *ADRB1* AND *ADRB2* (≥4DIR) LVEF ≥ 20

Event Rates:
Bucindolol Group: 28/86 (32.6%)
Placebo Group: 53/98 (54.1%)
Hazard Ratio = 0.53 (0.33,0.84)
P-value = 0.007

A.

Anti-phosphoERK1,2

kDa

50 –

37 –

Veh Veh Buc Buc Met Met Iso Iso

Total-ERK1,2

50 –

37 –

Veh Veh Buc Buc Met Met Iso Iso

B.