Refining the genetic landscape of anophthalmia and microphthalmia: a comprehensive framework with deep learning and updated gene panels

Mara I. Maftei, MSc[1]; Lucas G.N. Spink, BSc[1]; Oriol Gracia Carmona, PhD[2,3]; Simona Mikula Mrstakova, PhD[1]; Lean Abahreh, BSc[1]; Robin Hayes, BSc[1]; Aitor Bañon, PhD[1]; María Elisa Cuevas, BSc[4]; Katherine Cid, BSc[4]; Raul Araya-Secchi, PhD[5]; Franca Fraternali, PhD[2,7,8]; Jing Yu, PhD[9]; Gavin Arno, PhD[1,10*]; Rodrigo M. Young, PhD[1,4*].

1. Institute of Ophthalmology, UCL, London, UK

2. Research Department of Structural and Molecular Biology, Division of Biosciences, UCL, UK

3. Randall Centre for Cell & Molecular Biophysics, King's College London, London,UK

4. Center for Integrative Biology, Universidad Mayor, Santiago, Chile

5. Facultad de Ingenieria, Universidad San Sebastian, Santiago, Chile

6. Centro Cientifico y Tecnologico de Excelencia, Fundacion Ciencia & Vida, Santiago, Chile.

7. Institute of Structural and Molecular Biology, UCL, London, UK

8. Department of Biological Sciences, Birkbeck, UCL, London UK

9. Novo Nordisk Research Centre Oxford, Oxford, UK

10. Greenwood Genetic Center, Greenwood, South Carolina, USA.

* Authors for correspondence.

Abstract

Importance

Anophthalmia and microphthalmia (A/M) are rare congenital eye disorders with a low molecular diagnosis rate, which limits clinical management and genetic counselling. Improved detection and interpretation of pathogenic variants is essential for advancing diagnosis and care in affected individuals.

Objective

To improve the molecular diagnostic yield in A/M patients by refining the methodology of variant investigation and association using an updated rigorously curated gene panel, and a refined bioinformatic pipeline incorporating structural variant detection, *in silico* Artificial Intelligence assisted predictive tools, and molecular dynamics simulations.

Methodology

We curated an updated A/M gene panel through a systematic literature review and screened for rare variants in these genes using data from the UK's 100,000 Genomes Project, a national whole-genome sequencing initiative conducted by Genomics England. The cohort comprised 306 individuals recruited to the Rare Disease programme with a clinical diagnosis of anophthalmia or microphthalmia, recorded either as the primary phenotype or within HPO, SNOMED, or ICD-10 terms. Variants, including loss-of-function, missense, RNA splicing, and structural variants, were annotated with deep learning tools (AlphaMissense, SpliceAI), and missense variants were further assessed using REVEL, Missense3D, and molecular dynamics simulations.

Results

We identified pathogenic or likely pathogenic variants in 37 (12.1%) individuals, with an additional 23 (7.5%) harbouring strong candidate variants of uncertain significance. Our literature review identified the biggest contributors to A/M phenotypes to be *MFRP*, *OTX2*, *PRSS56* and *SOX2*, each with over 100 patients reported in the literature, with a total number of 124 genes found be associated to A/M. Variants from our screen were most often found in genes with high A/M association, but also included novel findings within genes with a weaker association to A/M such as *ACTG1*, *HDAC6*, *RERE* and *SIX3;* adding support to their disease relevance.

Conclusions and Relevance

This study increased the diagnostic yield in A/M patients recruited to the 100KGP, and provides further evidence of genotype-phenotype associations within the aetiology of A/M. We also provide an updated framework for enhancing clinical genetic diagnosis in A/M that may inform broader strategies for other complex congenital disorders. However, as molecular diagnosis of A/M remains low, further research in understanding the genetic aetiology of A/M is necessary.

Introduction

Cost-effective, high-throughput whole genome sequencing has significantly improved our understanding of the genetic basis of rare diseases[1,2]. However, many cases remain unsolved, limiting genetic counselling and patient care[3,4]. In particular, the molecular diagnosis rate in anophthalmia (A; absence of the eye) and microphthalmia (M; small eye) is only 13–33%[5–14]. Although A/M is untreatable, identifying a genetic cause remains crucial to enable appropriate clinical management, monitoring for associated complications, and enabling cascade testing and family reproductive planning[12,15].

A/M constitutes a spectrum of disease that can present unilaterally or bilaterally; occur in isolation or alongside ocular anomalies such as coloboma or anterior segment dysgenesis[11,12]; and often form part of syndromic presentations with other systemic abnormalities[16]. More than 100 genes have been associated to A/M, with the literature reporting *SOX2* as the most frequently disrupted gene (15–40% of diagnosed cases), followed by *OTX2* and *RAX*[10,11,14]. Methodological limitations and knowledge gaps may contribute to the low molecular diagnosis rate of A/M. Clinical and genetic heterogeneity, such as incomplete penetrance, variable expressivity, and mosaicism can complicate variant interpretation for molecular diagnosis[11,12]. Furthermore, underlying environmental factors, such as foetal alcohol syndrome, maternal vitamin A deficiency, exposure to toxins and certain infections during gestation may also impact the outcome of A/M without a monogenic cause[17].

The 100,000 Genomes Project (100KGP)[18] has enabled the systematic aggregation of genomic and clinical data of rare disease patients in the UK, providing an opportunity to investigate genetically heterogeneous conditions such as A/M in a phenotypically profiled cohort[1]. To address the persistently low molecular diagnosis rate of A/M, we set out to implement a new genomic variant interrogation pipeline to enable comprehensive evaluation of all A/M participants within the 100KGP cohort. This approach involved curating an A/M gene panel based on a systematic literature review, incorporating SV discovery, and utilising Alphamissense and SpliceAI, deep learning-based *in silico predictors,* to interpret variant pathogenicity. Our strategy led to the identification of 60 candidate variants in 56 previously unsolved participants, thereby strengthening the genotype–phenotype correlations for *ACTG1*, *HDAC6*, *RERE* and *SIX3* which are less commonly associated with A/M.

Methods

<u>A/M Cohort Selection</u>

Participants with a diagnosis of A/M were identified in the 100KGP dataset using relevant Human Phenotype Ontology (HPO) terms, ICD-10 codes, and/or SNOMED terms. This approach yielded a cohort of 306 individuals from 283 unrelated families. Notably, 118 participants from 108 families were recruited with A/M as the disease term, while the remaining 188 participants had A/M included in their HPO terms or clinical (SNOMED or ICD-10) terms, but were recruited under a different clinical condition. Of note, in compliance with Genomics England Airlock policy, detailed clinical terms are not included herein[18].

<u>A/M Gene Panel Curation</u>

A 124 A/M-associated gene panel (eTable1) was generated based on the Genomics England PanelApp 'Green' and 'Amber' genes (2016)[19], and literature reviews: Patel et al. (2018)[10], Harding and Moosajee (2019)[11], and Plaisancié et al. (2019)[12]. Additional genes identified in recent studies were included: *EPHA2*[13,20,21], *GJA8*[1,32-38], *NR6A1*[22], and *PRR12*[23,24]. We systematically reviewed published reports of variants in genes included in the panel, using PubMed keyword searches "gene", "human", "microphthalmia" or "anophthalmia". Only individuals harbouring a variant and explicitly described to have anophthalmia, microphthalmia, or documented axial length measurements consistent with an A/M diagnosis were included. Genes having pathogenic variants reported in at least five individuals across two or more families were defined as those with the strongest evidence for an association.

<u>Genomic Variant Calling and Annotation</u>

Structural variants were identified using the SVRare pipeline[25], with genomic coordinates aligned to the GRCh38 (hg38) assembly. Small variants were filtered for rarity, with an allele frequency (AF) of ≤0.001 in gnomAD v3.25[26] and the Genomics England research cohort, with retention of missense and splice-altering variants meeting in silico deleteriousness thresholds (AlphaMissense >0.340; SpliceAI >0.5). (see Supplementary Methods). To avoid artifacts or miscalls, all candidate variants were

confirmed by inspection using the Integrative Genomics Viewer (IGV[27]), with a minimum read depth of 15x and mapping quality ≥ 10. Subsequently, all true missense calls were annotated manually with REVEL[28] and Missense3D[29]. Furthermore, all missense variants located within functional domains were subjected to molecular dynamics (MD) simulations through a Root Mean Square Fluctuation analysis (see Supplementary Methods).

Results

Candidate variant identification

Sixty variants (in 56 individuals, eTable2, Fig.S1) within the A/M gene panel passed our filtering criteria: 16 pLOF (predicted loss-of-function), 7 structural variants (SV), 6 candidate splice-disrupting variants and 31 missense variants (summarised in eTable2). 37 participants (eTable2; 12.1% of the A/M cohort) harboured pathogenic or likely pathogenic variants and 23 (7.5%) carried candidate rare variants (AF ≤0.001) of unknown significance (VUS) with high pathogenicity scores. We consider 8 of these VUS (*ABCB6*:p.Met421Ile and p.Glu446Gly; *ALDH1A3*:p.Phe35Ser; *CHD7*:p.His1682Arg; *CRYBB2*:p.Arg145Pro; *NHS*:c.565+103674G>T; *OTX2*:p.Lys100Glu; and *TFAP2A*:p.Ala299Asp) to be likely pathogenic based on their location within critical functional domains, consistently high in silico pathogenicity predictions, and/or their strong specificity to the observed phenotypes. Participants 15, 43 and 46 carried candidate compound heterozygous variants in *ERCC2, PRSS56* and *RAB3GAP1*, respectively (eTable2).

Predicted loss-of-function alleles

All participants harbouring pLOF variants had phenotypes, inheritance and segregation patterns consistent with the clinical presentations associated with these gene defects (eTables 2 and 3). The variants *KMT2D:*p.Met1379ValfsTer52 (participant 23) and *SOX2:*p.Asn24ArgfsTer65 (participant 53) had both been reported in unrelated cases of A/M[30–36].

A recurrent *PAX2* frameshift variant (p.Val26GlyfsTer28) was identified in participant 35. This variant was also present in six other 100KGP participants, five with renal-coloboma syndrome features

without microphthalmia, and one unaffected relative. This variant has been well documented as pathogenic and has variable expressivity including kidney hypoplasia and ocular involvement[37–39].

Possible mosaicism in A/M genes

Three pLoF variants were observed affecting 10-20% of WGS reads, suggesting possible mosaicism: *BCOR*:p.Ser286AlafsTer92 (21.4% affected reads, participant 6, Fig.S1A), *BCOR*:p.Val1286CysfsTer8 (11.5% affected reads, participant 8, Fig.S1B), and *MYRF*:p.Pro190ArgfsTer19 (18.5% affected reads, participant 26, Fig.S13). Participant 26 expressed phenotypes milder than what is expected for a *MYRF* LOF variant, possibly due to mosaicism[40–42] (eTable2), while the participants 6 and 8 have clinical phenotypes (eTable2) in keeping with the known phenotypic variability in mosaic *BCOR* cases[43,44].

Structural variant analysis

Six SVs impacting the coding sequence of *ATOH7*, *HCCS*, *NDP*, *NHS*, *OTX2* and *PTCH1* were identified and classified as pathogenic or likely pathogenic according to ACMG/ClinGen guidelines[45] (eTable2).

Large deletions encompassing entire genes within our panel were identified in participants 5, 18 and 53 (eTable2). Participant 5 carries a 14.6 Mb (10:57,171,179-71,785,816) heterozygous deletion (Fig.S2A) spanning 72 coding genes including *ATOH7*, *ANK3*, *NEUROG3*, and *NODAL*, all involved in early embryonic development. Participant 18 and her affected mother (18_M) carry a 471.5 kb (X:10,945,466-11,417,014) deletion encompassing *HCCS* (Fig.S2B), and three other coding genes: *ARHGAP6, AMELX, MIR548AX*. Participant 53 harbours a *de novo* 707.7 kb (14:56,783,923-57,491,668; Fig.S2C) deletion deleting the entire *OTX2*, and *EXOC5, AP5M1, NAA30, CCDC198*.

SVs partially deleting genes were identified in participants 28, 29 and 41 (Fig.S2D-F, eTable2). Participant 28 inherited a hemizygous 4.4 kb (X:43,954,691-43,959,091) deletion from his heterozygous, unaffected mother. This variant resulted in the deletion of exon 2 in *NDP*, leading to start codon loss (Fig.2A, Fig.S2D). Participant 29 inherited a hemizygous 44.5 kb (X:17,153,721-17,599,035) deletion of *NHS* exon 1 from his mother (29_M) who expresses similar, but milder

phenotypes (Fig.2B, Fig.S2E, eTable 3). Finally, a *de novo* 1.8 kb (9:95,475,579-95,477,447) deletion of *PTCH1* exons 11–12 was identified in participant 44 (Fig.1Y, Fig.S2F).

Identification of candidate RNA splice-altering variants

Six variants that potentially disrupt RNA splicing were identified: *NHS*:c.565+103674G>T (Fig.2B, participant 30), *PAX6*:c.-128-2del (Fig.2C, participant 36), *PORCN*:c.1087+5G>A (Fig.2D, participant 4)*, PRR12*:c.4773+1G>C (Fig.2E, participant 41), *RARB*:c.609+5G>C (Fig.2F, participant 47), *RERE:c.2197A>G* (Fig.2G, participant 48). The variants in *PAX6* and *PRR12* carried by participants 36 and 41 were predicted to be pathogenic, while the variants in *NHS*, *PORCN*, *RARB* and *RERE* are classified as VUS.

The *NHS*:c.565+103674G>T and *PORCN*:c.1087+5G>A variants were both identified hemizygously in male participants (30 and 4, respectively), inherited from their unaffected heterozygous mothers (eTable 5). For the *NHS*:c.565+103674G>T variant, SpliceAI predicts an acceptor site gain (Δscore=0.74) 13bp downstream of the variant position (Fig.2B). This could lead to pseudo-exon inclusion in the transcript. Notably, this variant was also identified within another 100KGP participant with HPO terms in keeping with Nance Horan Syndrome, but not affected with microphthalmia. Functional investigation of the splice impact would clarify its pathogenicity in these cases.

The *PORCN* variant, located at the conserved +5 position of intron 12, showed a SpliceAI donor loss score of 0.47 and a high-confidence alternative donor gain site 31bp downstream (Δscore=0.92, Fig.2D). Use of this alternate donor gain would result in a 36bp intronic inclusion and an in-frame insertion (p.His362_Val363ins12) resulting in a disruption of the MBOAT domain, potentially disrupting Wnt signalling[46].

Heterozygous variants in *RARB*:c.609+5G>C and *RERE*:c.2197A>G (p.Met733Val) were identified in participants 47 and 48, respectively. The first, in intron 4 is predicted to abolish the canonical donor site (Δscore=0.75) and activate a cryptic donor site 19bp upstream (Δscore=0.64), leading to deletion of 14bp of exon 4 from the transcript, and resulting in a predicted frameshift p.Gly199GlufsTer3 (Fig.2F). The *RERE* missense variant (p.Met733Val), while scoring low on missense pathogenicity

prediction (AlphaMissense score 0.069), is predicted to create a strong cryptic splice donor site within exon 18 (Δscore=0.97). This would lead to the excision of 1,199 nucleotides, causing a large deletion and frameshift (p.Met733ValfsTer35 Fig.2G).

<u>Missense variant analysis</u>

We identified thirteen heterozygous missense variants in genes associated with autosomal dominant A/M, two variants in genes associated with X-linked A/M, two homozygous variants in recessive A/M genes and two compound heterozygous genotypes (eTable2). Of these 31 missense variants, 15 were classified as pathogenic or likely pathogenic with the remaining as VUS.

All VUS were predicted pathogenic by AlphaMissense, with corresponding high REVEL scores (eTable2). Noteworthy, dominant *CHD7*, *GDF6*, *KMT2D* and *SMCHD1* variants were inherited from apparently unaffected parents (eTable2, eTable3) either suggesting these are not pathogenic or exhibit incomplete/non penetrance[10–12,47–49]. The VUSs in *ABCB6*, *COL4A1*, *CRYBB2*, *GDF6*, *KERA*, *KMT2D* (p.Leu241Gln) *MYRF, OTX2*, *PRSS56* (p.Gly107Glu), *SMCHD1* and *TFAP2A* impacted functional domains: ABC transporter transmembrane domain (*ABCB6*, Fig.1A), a collagenous domain (*COL4A1*, Fig.1G), a 'Greek key' domain (*CRYBB2*, Fig.1H), a TGF-β domain (*GDF6*, Fig.1J), a Leucine-rich repeat (*KERA*, Fig.1M), a PHD finger (*KMT2D*, Fig.1N), DNA binding domains (*MYRF*, Fig.1Q; *OTX2*, Fig.1S; *TFAP2A*, Fig.1AD) and a serine protease inhibitor domain (*PRSS56*, Fig1.W). While not affecting functional domains, the VUS variants in *ALDH1A3*, *CHD7*, *KMT2D* and *SMCHD1* (p.Glu721Gln), were carried by participants expressing phenotypes in keeping with with those reported for similar variants in these genes[48,50,51]. Notably, the homozygous p.Phe35Ser variant in *ALDH1A*3, was found in the same participant who carried the heterozygous *PORCN*:c.1087+5G>A variant. As both are VUS, further investigation is necessary, including RNA analysis for a splice defect in *PORCN,* to help determine which is more likely to be causative[52–54].

Molecular dynamic simulations of protein domains can be used to study the effect of amino acid changes by analysing the Root Mean Square Fluctuation (RMSF) of atoms[55,56]. To add evidence to the pathogenicity of the identified missense VUS, we performed RMSF analyses on the missense variants within functional domains in *GDF6, MYRF, OTX2, PAX6, PORCN, SIX3, TFAP2A*, focusing on the Cα

atoms to assess potential impacts on structural stability (Fig.3A). Comparisons centred on the top 10% of residues exhibiting the greatest fluctuation differences. The *PAX6* variants (p.Arg26Gly and p.Asn138His) displayed a clear destabilising effect on the whole modelled protein region (Fig.3A-C). *TFAP2A*:Arg239Gln also displayed a significant destabilising effect (Fig.3A, D). In contrast, the *SIX3*:p.Arg192His variant demonstrated a stabilising effect, suggesting increased conformational rigidity in the protein fold (Fig.2A, E).

Compound heterozygous variants

Participant 15 harboured two missense variants in *ERCC2* (Fig.1I, eTable2). The maternally inherited allele harboured c.1802G>A, p.Arg601Gln, with c.1939C>T, p.Arg647Cys presumed in trans (paternal DNA unavailable; eTable3). Participant 46 carried a maternally inherited frameshift variant in *RAB3GAP1* (p.Glu796ValfsTer12) and a large paternally inherited 346.5kb deletion (2:135,165,337-135,511,837; Fig.S2G) encompassing part of *RAB3GAP1* and *ZRANB3,* resulting in the loss of the final exon of *RAB3GAP1* (Fig.1Y, Fig.S2G, eTable2). Participant 43 carried a likely pathogenic frameshift p.Val525CysfsTer55 in *PRSS56* on the paternal allele, and a VUS missense p.Arg480Pro variant on the maternal allele (Fig.1W, eTable2).

Discussion

This study describes the genetic findings in 60/306 individuals recruited to the UK 100KGP for a curated 124 gene panel implicated in A/M. Our analysis using updated variant interpretation tools yielded an overall candidate variant identification rate of 19.6%. However, the overall diagnostic rate for A/M is likely higher due to prescreening for variants within *OTX2* and *SOX2* prior to recruitment to the 100KGP, these genes together account for approximately 20% of A/M cases[11,12,56–59]. The estimated diagnostic rate is projected to approximately 35.7%.

Importantly, our analysis included all participants recruited with A/M regardless of solved status; 12 variants (from 12 participants) identified here, had been previously reported and published. One additional previously reported variant, *PDGFRA*:p.Thr432Met was not retained in our final candidate

list, as *PDGFRA* currently lacks sufficient evidence for association with A/M and was excluded from our curated gene panel.

From our literature search to generate a curated A/M gene panel, we better characterised the genetic aetiology of A/M. Historically, *SOX2*, *OTX2*, and *RAX* were considered the top three contributors to A/M[10,11]; however, we identified that *SOX2*, *PRSS56*, *MFRP*, and *OTX2* were the most frequent contributors to A/M, each with over 100 patients reported. It may also be important to note that variants in both *MFRP* and *PRSS56* cause microphthalmia strictly affecting the posterior segment. As expected, the majority of pathogenic variants identified in the screening occurred in genes with strong associations to A/M such as *BCOR*, *OTX2*, *PAX6* and *TFAP2A*. Our findings in *ERCC2* and *NDP* further reinforce their association to A/M, as both previously had five reported cases[61–69]. We also identified a likely pathogenic variant in *HDAC6* and a strong VUS candidate in *ACTG1*, both genes with limited evidence of association with A/M; where the association is supported by either a single previously reported affected individual, or where the association has only been inferred[70–73]. We further identified a likely pathogenic variant in *SIX3* (previously three A/M cases reported, eTable1) and *RERE* (two cases reported, eTable1), adding further support for the association of these genes to A/M.

This study identified seven structural variants representing 16% of all the variants underscoring the impact of SVs in A/M and the importance of including SV interrogation tools such as SVRare[25] into clinical pipelines. The application of deep learning-based predictors, AlphaMissense and SpliceAI, to large-scale variant datasets provides an effective approach for variant prioritisation. Despite not being explicitly trained on ACMG/AMP criteria, 66.7% (40/60) of the variants flagged by these algorithms, and included in this study, met the pathogenic or likely pathogenic classification. The structural and stability effects of missense variants were further evaluated using Missense3D and molecular dynamic simulations. RMSF analysis revealed local flexibility changes indicative of altered fold stability, as observed for PAX6 variants (Fig.3C, D). While Missense3D predicted no structural damage for *PAX6*:p.Asn139His and *SIX3*, MD simulations indicated destabilisation; conversely, *PORCN*:p.Phe250Ser was predicted as damaging by Missense3D but showed no instability in RMSF, though its previous association with microphthalmia supports pathogenicity[52,55,74]. Such discrepancies

highlight that many pathogenic variants act through mechanisms beyond fold stability, such as protein–protein interactions, impaired complex formation, or dysregulated localization, which conventional MD cannot capture[56]. This may also explain the limited correlation between AlphaMissense pathogenicity scores and structural instability predictions, as AlphaMissense integrates broader evolutionary and functional features[75].

Overall, applying our variant interpretation pipeline could increase the molecular diagnosis rate for the A/M cohort by 12.1% with ACMG guidelines. Despite this effort, 80.4% of the cohort remains unsolved, and compared to other ophthalmic disease cohorts in the 100K GP, the A/M diagnostic yield remains low. Factors contributing to this may include yet unidentified A/M-associated genes, regulatory variants, epigenetic regulation, non-Mendelian inheritance and non-genetic phenocopies, none of which were investigated here.

Altogether, our study advances both the precision and scope of A/M genetic diagnosis by offering a rigorously curated gene panel and a refined analytical framework that integrates structural variant detection and deep learning-based prediction algorithms. In addition, we present additional affected individuals harbouring variants in genes with uncertain associations to A/M, consolidating their affiliation in causing A/M.

Acknowledgements

This research was made possible through access to data in the National Genomic Research Library, which is managed by Genomics England Limited (a company owned by the Department of Health and Social Care). The National Genomic Research Library holds data provided by patients and collected by the NHS as part of their care and data collected as part of their participation in research. The National Genomic Research Library is funded by the National Institute for Health Research and NHS England. The Wellcome Trust Cancer Research UK, and the Medical Research Council have also funded research infrastructure.

This research was supported by the NIHR Moorfields Biomedical Research Centre (GA), the Moorfields Eye Charity Career Development award (GR001155 to RMY), Springboard Awards (GR001210 and GR001662 to RMY), PhD Studentship (GR001661 to MIM), Medical Research

Council (MR/X001067/1 to RMY), FONDECYT (1221843 to RMY), Biotechnology and Biological Sciences Research Council (BB/T002212/1 to FF as principal investigator) and the British Heart Foundation grant (RG/F/22/110079 to FF and OGC).

Competing interests

The authors declare no competing interests.

Data and resource availability

All relevant data and resources can be found within the article and its supplementary information.

Author contribution

MM: methodology; formal analysis; investigation; writing – original draft; writing – review and editing; visualisation

LS: formal analysis; investigation; writing

OGC: methodology; formal analysis; investigation; writing

SM, LA, RH, AB, MEC, KC, RAS, FF, JY: investigation

GA: methodology; validation; formal analysis; investigation; review and editing; supervision.

RY: conceptualisation; methodology; validation; formal analysis; investigation; resources; writing – original draft; writing – review and editing; visualisation; supervision; project administration; funding acquisition.

References

1. Smedley D, Smith KR, Martin A, et al. 100,000 Genomes Pilot on Rare-Disease Diagnosis in Health Care — Preliminary Report. *N Engl J Med*. 2021;385(20):1868-1880. doi:10.1056/nejmoa2035790

2. Shieh JTC. Genomic technologies to improve variation identification in undiagnosed diseases. *Pediatr Neonatol*. 2023;64:S18-S21. doi:10.1016/j.pedneo.2022.10.002

3. Turro E, Astle WJ, Megy K, et al. Whole-genome sequencing of patients with rare diseases in a national health system. *Nature*. 2020;583(7814):96-102. doi:10.1038/s41586-020-2434-2

4. Marwaha S, Knowles JW, Ashley EA. A guide for the diagnosis of rare and undiagnosed disease: beyond the exome. *Genome Med*. 2022;14(1). doi:10.1186/s13073-022-01026-w

5. Schneider A, Bardakjian T, Reis LM, Tyler RC, Semina EV. Novel SOX2 mutations and genotype-phenotype correlation in anophthalmia and microphthalmia. *Am J Med Genet A*. 2009;149A(12):2706-2715. doi:10.1002/ajmg.a.33098

6. Shah SP, Taylor AE, Sowden JC, et al. Anophthalmos, Microphthalmos, and Coloboma in the United Kingdom: Clinical Features, Results of Investigations, and Early Management. *Ophthalmology*. 2012;119(2):362-368. doi:10.1016/j.ophtha.2011.07.039

7. Chassaing N, Causse A, Vigouroux A, et al. Molecular findings and clinical data in a cohort of 150 patients with anophthalmia/microphthalmia. *Clin Genet*. 2013;86(4):326-334. doi:10.1111/cge.12275

8. Gerth-Kahlert C, Williamson KA, Ansari M, et al. Clinical and mutation analysis of 51 probands with anophthalmia and/or severe microphthalmia from a single center. *Mol Genet Genomic Med*. 2013;1(1):15-31. doi:10.1002/mgg3.2

9. Slavotinek AM, Garcia ST, Chandratillake G, et al. Exome sequencing in 32 patients with anophthalmia/microphthalmia and developmental eye defects. *Clin Genet*. 2015;88(5):468-473. doi:10.1111/cge.12543

10. Patel A, Hayward JD, Tailor V, et al. The Oculome Panel Test. *Ophthalmology*. 2019;126(6):888-907. doi:10.1016/j.ophtha.2018.12.050

11. Harding P, Moosajee M. The Molecular Basis of Human Anophthalmia and Microphthalmia. *J Dev Biol*. 2019;7(3). doi:10.3390/jdb7030016

12. Plaisancié J, Ceroni F, Holt R, et al. Genetics of anophthalmia and microphthalmia. Part 1: Non-syndromic anophthalmia/microphthalmia. *Hum Genet*. 2019;138(8-9):799-830. doi:10.1007/s00439-019-01977-y

13. Harding P, Gore S, Malka S, Rajkumar J, Oluonye N, Moosajee M. Real-world clinical and molecular management of 50 prospective patients with microphthalmia, anophthalmia and/or ocular coloboma. *Br J Ophthalmol*. 2022;107(12):bjo-2022. doi:10.1136/bjo-2022-321991

14. Reis LM, Semina EV. Conserved genetic pathways associated with microphthalmia, anophthalmia, and coloboma. *Birth Defects Res Part C Embryo Today Rev*. 2015;105(2):96-113. doi:10.1002/bdrc.21097

15. Richardson R, Sowden J, Gerth-Kahlert C, Moore AT, Mariya Moosajee. Clinical utility gene card for: Non-Syndromic Microphthalmia Including Next-Generation Sequencing-Based Approaches. *Eur J Hum Genet*. 2017;25(4):512-512. doi:10.1038/ejhg.2016.201

16. Slavotinek A. Genetics of anophthalmia and microphthalmia. Part 2: Syndromes associated with anophthalmia–microphthalmia. *Hum Genet*. 2018;138(8-9):831-846. doi:10.1007/s00439-018-1949-1

17. Goyal S, Tibrewal S, Ratna R, Vanita V. Genetic and environmental factors contributing to anophthalmia and microphthalmia: Current understanding and future directions. *World J Clin Pediatr*. 2025;14(2). doi:10.5409/wjcp.v14.i2.101982

18. The National Genomic Research Library v5.1, Genomics England. 2020. doi:doi:10.6084/m9.figshare.4530893/7

19. Anophthalmia or microphthalmia (Version 1.51). Genomicsengland.co.uk. 2016. https://panelapp.genomicsengland.co.uk/panels/34/

20. Harding P, Toms M, Schiff E, et al. EPHA2 Segregates with Microphthalmia and Congenital Cataracts in Two Unrelated Families. *Int J Mol Sci*. 2021;22(4):2190. doi:10.3390/ijms22042190

21. Courdier C, Gemahling A, Guindolet D, et al. EPHA2 biallelic disruption causes syndromic complex microphthalmia with iris hypoplasia. *Eur J Med Genet*. 2022;65(10):104574. doi:10.1016/j.ejmg.2022.104574

22. Neelathi UM, Ullah E, George A, et al. Variants in NR6A1 cause a novel oculo-vertebral-renal (OVR) syndrome. *PubMed*. Published online 2024. doi:10.1101/2024.11.09.24316578

23. Reis LM, Costakos D, Wheeler PG, et al. Dominant variants in PRR12 result in unilateral or bilateral complex microphthalmia. *Clin Genet*. 2021;99(3):437-442. doi:10.1111/cge.13897

24. Chowdhury F, Wang L, Al-Raqad M, et al. Haploinsufficiency of PRR12 causes a spectrum of neurodevelopmental, eye, and multisystem abnormalities. *Genet Med*. 2021;23(7):1234-1245. doi:10.1038/s41436-021-01129-6

25. Yu J, Szabo A, Pagnamenta AT, et al. SVRare: discovering disease-causing structural variants in the 100K Genomes Project. *MedRxiv Cold Spring Harb Lab*. Published online 2021. doi:10.1101/2021.10.15.21265069

26. Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*. 2020;581(7809):434-443. doi:10.1038/s41586-020-2308-7

27. Robinson JT, Thorvaldsdóttir H, Winckler W, et al. Integrative genomics viewer. *Nat Biotechnol*. 2011;29(1):24-26. doi:10.1038/nbt.1754

28. Ioannidis NM, Rothstein JH, Pejaver V, et al. REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am J Hum Genet*. 2016;99(4):877-885. doi:10.1016/j.ajhg.2016.08.016

29. Ittisoponpisan S, Islam SA, Khanna T, Alhuzimi E, David A, Sternberg MJE. Can Predicted Protein 3D Structures Provide Reliable Insights into whether Missense Variants Are Disease Associated? *J Mol Biol*. 2019;431(11):2197-2212. doi:10.1016/j.jmb.2019.04.009

30. Zenteno JC, Perez-Cano HJ, Aguinaga M. Anophthalmia-esophageal atresia syndrome caused by an SOX2 gene deletion in monozygotic twin brothers with markedly discordant phenotypes. *Am J Med Genet A*. 2006;140A(18):1899-1903. doi:10.1002/ajmg.a.31384

31. Schneider A, Bardakjian TM, Zhou J, et al. Familial recurrence ofSOX2anophthalmia syndrome: Phenotypically normal mother with two affected daughters. *Am J Med Genet A*. 2008;146A(21):2794-2798. doi:10.1002/ajmg.a.32384

32. Suzuki J, Azuma N, Dateki S, et al. Mutation spectrum and phenotypic variation in nine patients with SOX2 abnormalities. *J Hum Genet*. 2014;59(6):353-356. doi:10.1038/jhg.2014.34

33. Cheon CK, Ko JM. Kabuki syndrome: clinical and molecular characteristics. *Korean J Pediatr*. 2015;58(9):317. doi:10.3345/kjp.2015.58.9.317

34. Bögershausen N, Gatinois V, Riehmer V, et al. Mutation Update for Kabuki Syndrome GenesKMT2DandKDM6Aand Further Delineation of X-Linked Kabuki Syndrome Subtype 2. *Hum Mutat*. 2016;37(9):847-864. doi:10.1002/humu.23026

35. Errichiello E, Gorgone C, Giuliano L, et al. SOX2: Not always eye malformations. Severe genital but no major ocular anomalies in a female patient with the recurrent c.70del20 variant. *Eur J Med Genet*. 2018;61(6):335-340. doi:10.1016/j.ejmg.2018.01.011

36. Zhang Y, Zhang X, Long R, Yu L. A novel deletion mutation of the SOX2 gene in a child of Chinese origin with congenital bilateral anophthalmia and sensorineural hearing loss. *J Genet*. 2018;97(4):1007-1011. doi:10.1007/s12041-018-0970-4

37. Xiong HY, Shi YQ, Zhong C, et al. Detection of De Novo PAX2 Variants and Phenotypes in Chinese Population: A Single-Center Study. *Front Genet*. 2022;13. doi:10.3389/fgene.2022.799562

38. Sanyanusin P, McNoe LA, Sullivan MJ, Weaver RG, Eccles MR. Mutation of PAX2 in two siblings with renal-coloboma syndrome. *Hum Mol Genet*. 1995;4(11):2183-2184. doi:10.1093/hmg/4.11.2183

39. Ford B, Rupps R, Lirenman D, et al. Renal-coloboma syndrome: Prenatal detection and clinical spectrum in a large family. *Am J Med Genet*. 2001;99(2). doi:10.1002/1096-8628(2000)9999:999%3C00::AID-AJMG1143%3E3.0.CO;2-F

40. Garnai SJ, Brinkmeier ML, Emery B, et al. Variants in myelin regulatory factor (MYRF) cause autosomal dominant and syndromic nanophthalmos in humans and retinal degeneration in mice. Anderson MG, ed. *PLOS Genet*. 2019;15(5):e1008130. doi:10.1371/journal.pgen.1008130

41. Rossetti LZ, Glinton K, Yuan B, et al. Review of the phenotypic spectrum associated with haploinsufficiency of MYRF. *Am J Med Genet A*. 2019;179(7):1376-1382. doi:10.1002/ajmg.a.61182

42. Siggs OM, Awadalla MS, Souzeau E, et al. The genetic and clinical landscape of nanophthalmos and posterior microphthalmos in an Australian cohort. *Clin Genet*. 2020;97(5):764-769. doi:10.1111/cge.13722

43. Hilton E, Johnston JJ, Whalen S, et al. BCOR analysis in patients with OFCD and Lenz microphthalmia syndromes, mental retardation with ocular anomalies, and cardiac laterality defects. *Eur J Hum Genet*. 2009;17(10):1325-1335. doi:10.1038/ejhg.2009.52

44. Ragge N, Isidor B, Bitoun P, et al. Expanding the phenotype of the X-linked BCOR microphthalmia syndromes. *Hum Genet*. 2018;138(8-9):1051-1069. doi:10.1007/s00439-018-1896-x

45. Riggs ER, Andersen EF, Cherry AM, et al. Technical standards for the interpretation and reporting of constitutional copy-number variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics (ACMG) and the Clinical Genome Resource (ClinGen). *Genet Med*. 2019;22. doi:10.1038/s41436-019-0686-8

46. Yu J, Liao PJ, Xu W, et al. Structural model of human PORCN illuminates disease-associated variants and drug-binding sites. *J Cell Sci*. 2021;134(24):jcs259383. doi:10.1242/jcs.259383

47. Asai-Coakwell M, French CR, Ye M, et al. Incomplete penetrance and phenotypic variability characterize Gdf6-attributable oculo-skeletal phenotypes. *Hum Mol Genet*. 2009;18(6):1110-1121. doi:10.1093/hmg/ddp008

48. Cuvertino S, Hartill V, Colyer A, et al. A restricted spectrum of missense KMT2D variants cause a multiple malformations disorder distinct from Kabuki syndrome. *Genet Med*. 2020;22(5):867-877. doi:10.1038/s41436-019-0743-3

49. Gregory LC, Gergics P, Nakaguma M, et al. The phenotypic spectrum associated with OTX2 mutations in humans. *Eur J Endocrinol*. 2021;185(1):121-135. doi:10.1530/EJE-20-1453

50. Shaw ND, Brand H, Kupchinsky ZA, et al. SMCHD1 mutations associated with a rare muscular dystrophy can also cause isolated arhinia and Bosma arhinia microphthalmia syndrome. 2017;49(2):238-248. doi:10.1038/ng.3743

51. Zentner GE, Layman WS, Martin DM, Scacheri PC. Molecular and phenotypic aspects of CHD7 mutation in CHARGE syndrome. *Am J Med Genet A*. 2010;152A(3):674-686. doi:10.1002/ajmg.a.33323

52. Madan S, Liu W, Lu JT, et al. A non-mosaic PORCN mutation in a male with severe congenital anomalies overlapping focal dermal hypoplasia. *Mol Genet Metab Rep*. 2017;12:57-61. doi:10.1016/j.ymgmr.2017.06.002

53. Wawrocka A, Walczak-Sztulpa J, Pawlak M, Gotz-Wieckowska A, Krawczynski MR. Non-syndromic anophthalmia/microphthalmia can be caused by a PORCN variant inherited in X-linked recessive manner. *Am J Med Genet - Part A*. 2020;185(1):250-255. doi:10.1002/ajmg.a.61938

54. Kesim Y, Ceroni F, Damián A, et al. Clinical and genetic analysis further delineates the phenotypic spectrum of ALDH1A3-related anophthalmia and microphthalmia. *Eur J Hum Genet*. 2023;31(10):1175-1180. doi:10.1038/s41431-023-01342-8

55. Buric F, Viknander S, Fu X, et al. Amino acid sequence encodes protein abundance shaped by protein stability at reduced synthesis cost. *Protein Sci*. 2024;34(1). doi:10.1002/pro.5239

56. Beltran A, Jiang X, Shen Y, Lehner B. Site-saturation mutagenesis of 500 human protein domains. *Nature*. 2025;637. doi:10.1038/s41586-024-08370-4

57. Ragge NK, Brown AG, Poloschek CM, et al. Heterozygous Mutations of OTX2 Cause Severe Ocular Malformations. *Am J Hum Genet*. 2005;76(6):1008-1022. doi:10.1086/430721

58. Chassaing N, Sorrentino S, Davis EE, et al. OTX2mutations contribute to the otocephaly-dysgnathia complex. *J Med Genet*. 2012;49(6):373-379. doi:10.1136/jmedgenet-2012-100892

59. Williamson KA, FitzPatrick DR. The genetic architecture of microphthalmia, anophthalmia and coloboma. *Eur J Med Genet*. 2014;57(8):369-380. doi:10.1016/j.ejmg.2014.05.002

60. Li X, Xiao H, Su Y, et al. Clinical features of patients with mutations in genes for nanophthalmos. *Br J Ophthalmol*. 2024;108(12):1679-1687. doi:10.1136/bjo-2023-324931

61. Kondo D, Noguchi A, Tamura H, et al. Elevated Urinary Levels of 8-Hydroxy-2′-deoxyguanosine in a Japanese Child of Xeroderma Pigmentosum/Cockayne Syndrome Complex with Infantile Onset of Nephrotic Syndrome. *Tohoku J Exp Med*. 2016;239(3):231-235. doi:10.1620/tjem.239.231

62. Graham JM, Anyane-Yeboa K, Raams A, et al. Cerebro-Oculo-Facio-Skeletal Syndrome with a Nucleotide Excision–Repair Defect and a Mutated XPD Gene, with Prenatal Diagnosis in a Triplet Pregnancy. *Am J Hum Genet*. 2001;69(2):291-300. doi:10.1086/321295

63. Farmaki E, Nedelkopoulou N, Delli F, Sarafidis K, Zafeiriou DI. Brittle Hair, Photosensitivity, Brain Hypomyelination and Immunodeficiency: Clues to Trichothiodystrophy. *Indian J Pediatr*. 2017;84(1):89-90. doi:10.1007/s12098-016-2209-9

64. Janani R, Chermakani P, Sundaresan P, Shetty S, Rai K. A rare case in a child with mild trichothiodystrophy associated with ERCC2 gene. *Indian J Ophthalmol - Case Rep*. 2022;2(4):962-964. doi:10.4103/ijo.IJO_1217_22

65. Horibata K, Kono S, Ishigami C, et al. Constructive rescue of TFIIH instability by an alternative isoform of XPD derived from a mutated XPD allele in mild but not severe XP-D/CS. *J Hum Genet*. 2015;60(5):259-265. doi:10.1038/jhg.2015.18

66. Berger W, Meindl A, Van De Pol TJR, et al. Isolation of a candidate gene for Norrie disease by positional cloning. *Nat Genet*. 1992;1(3):199-203. doi:10.1038/ng0692-199

67. Payabvash S, Anderson JS, Nascene DR. Bilateral persistent fetal vasculature due to a mutation in the Norrie disease protein gene. *Neuroradiol J*. 2015;28(6):623-627. doi:10.1177/1971400915609350

68. Deml B, Reis LM, Lemyre E, Clark RD, Kariminejad A, Semina EV. Novel mutations in PAX6, OTX2 and NDP in anophthalmia, microphthalmia and coloboma. *Eur J Hum Genet*. 2015;24(4):535-541. doi:10.1038/ejhg.2015.155

69. Lin M, Lu Y, Sui Y, et al. A novel c.287G>T *NDP* missense mutation in a Chinese family with Norrie disease. *Ophthalmic Genet*. 2020;41(4):338-340. doi:10.1080/13816810.2020.1759106

70. Khalifa O, Al-Sahlawi Z, Imtiaz F, et al. Variable expression pattern in Donnai-Barrow syndrome: Report of two novel LRP2 mutations and review of the literature. *Eur J Med Genet*. 2015;58(5):293-299. doi:10.1016/j.ejmg.2014.12.008

71. Chassaing N, Siani V, Carles D, et al. X-linked dominant chondrodysplasia with platyspondyly, distinctive brachydactyly, hydrocephaly, and microphthalmia. *Am J Med Genet A*. 2005;136A(4):307-312. doi:10.1002/ajmg.a.30570

72. Simon D, Laloo B, Barillot M, et al. A mutation in the 3′-UTR of the *HDAC6* gene abolishing the post-transcriptional regulation mediated by hsa-miR-433 is linked to a new form of dominant X-linked chondrodysplasia. *Hum Mol Genet*. 2010;19(10):2015-2027. doi:10.1093/hmg/ddq083

73. Kantarci S, Casavant D, Prada C, et al. Findings from aCGH in patients with congenital diaphragmatic hernia (CDH): A possible locus for Fryns syndrome. *Am J Med Genet A*. 2006;140A(1):17-23. doi:10.1002/ajmg.a.31025

74. Frisk S, Grandpeix-Guyodo C, Silwerfeldt KP, et al. Goltz syndrome in males: A clinical report of a male patient carrying a novel PORCN variant and a review of the literature. *Clin Case Rep*. 2018;6(11):2103-2110. doi:10.1002/ccr3.1783

75. Cheng J, Novati G, Pan J, et al. Accurate proteome-wide missense variant effect prediction with AlphaMissense. *Science*. 2023;381(6664). doi:10.1126/science.adg7492

Figure 1. Schematic of coding variants found within A/M participants of the 100K GP

A-AD) Schematic of the coding variants identified by our screening within anophthalmic and microphthalmic patients recruited to the 100KGP. Predicted-loss-of-function variants are represented in red and missense in yellow. The participants who harbour each variant is presented in parentheses.

Figure 2. Schematic of variants affecting non-coding regions found within A/M participants of the 100KGP

A-H) Schematic of the variants affecting non-coding regions that were identified by our screening within anophthalmic and microphthalmic patients recruited to the 100KGP. Splice variants are represented in green, with the participants who harbour each variant presented in parentheses. AG: acceptor gain, AL: acceptor loss, DG: donor gain, DL: donor loss.

Figure 3. Root Mean Square Fluctuation for missense variants

A) Box plot showing Root Mean Square Fluctuation (RMSF) differences (in Å) for backbone Cα atoms, relative to the wild-type (wt) mean RMSF for each system. Differences were calculated using the top 10% of residues with the highest fluctuation differences in each system. Asterisks indicate statistical significance: $p < 0.05$(*). B-E) Protein structural representation of RMSF for those which achieved statistical significance in (A). Regions in red represent mutant flexibility compared to wt, while blue represents rigidity. Yellow represents the position of the amino acid substitution.

Gene	[GEL PanelApp]	[Harding and Moosajee, 2019]	[Plaisancie et al., 2019]	[Patel et al., 2019]	Occurance in the four reviews	Number of patients in literature	Number of familes in literature
SOX2	1	1	1	1	4	161	147
PRSS56	1	1	1	1	4	161	100
MFRP	1	1	1	1	4	129	103
OTX2	1	1	1	1	4	104	64
BCOR	1	1	1	1	4	79	66
FOXE3	1	1	1	0	3	75	48
VSX2	1	1	1	1	4	74	34
TMEM98	1	1	1	1	4	66	13
RAB3GAP1	1	1	1	1	4	59	47
ALDH1A3	1	1	1	1	4	59	39
STRA6	1	1	1	1	4	48	32
PORCN	0	1	1	1	3	38	32
PAX6	1	1	1	1	4	35	28
MYRF	1	0	0	0	1	35	14
GJA1	0	1	0	1	2	30	17
HCCS	1	1	1	1	4	29	27
BMP4	1	1	1	1	4	27	26
SIX6	1	1	1	1	4	27	20
SMCHD1	0	1	1	0	2	26	25
RAB3GAP2	1	1	1	1	4	25	18
GJA8	0	0	1	0	1	25	20
MAB21L2	1	1	1	1	4	23	20
CHD7	0	1	1	0	2	23	23
RARB	1	1	1	1	4	22	17
TFAP2A	0	1	1	1	3	19	17
RAX	1	1	1	1	4	18	17
PXDN	0	1	1	0	2	17	12
SMOC1	1	1	1	1	4	15	10
GDF6	1	1	1	1	3	15	15
C12ORF57	0	1	1	1	3	14	13
NHS	0	1	0	0	1	14	14
YAP1	0	1	1	0	2	13	6
TBC1D20	1	0	1	1	3	12	7
RAB18	1	1	1	0	3	12	7
CRYAA	0	1	0	1	2	11	6
EPHA2	0	0	0	0	0	11	4
PAX2	1	1	1	0	3	9	9
GDF3	1	1	1	1	3	9	8
ERCC6	1	1	0	0	1	9	8
PRR12	0	0	0	0	0	9	8
TMX3	0	1	1	1	3	8	8
NAA10	0	1	1	1	3	8	3
COL4A1	1	1	1	0	3	8	7
LRP5	0	1	0	0	1	8	6
CRPPA	0	1	0	0	1	8	8
PTCH1	0	1	1	1	3	7	7
KMT2D	1	1	0	0	2	7	7
PITX3	1	1	1	0	2	7	6
ZEB2	0	1	0	0	1	7	7
FAT1	1	0	0	0	1	7	5

Gene							
TENM3	1	1	1	1	4	6	5
ATOH7	0	1	1	1	3	6	2
SHH	1	1	0	1	3	6	6
FAM111A	0	1	0	1	2	6	5
SLC24A5	0	1	0	0	1	6	6
FKTN	0	1	0	0	1	6	6
ALX1	0	1	0	0	1	6	3
ABCB6	1	1	1	1	3	5	5
TBC1D32	0	1	0	1	2	5	5
CRYBA4	0	1	0	1	2	5	4
HMX1	0	1	0	1	2	5	2
MITF	0	1	1	0	2	5	5
POMT1	0	1	0	0	1	5	5
NDP	0	1	0	0	1	5	4
MAPRE2	0	1	0	0	1	5	5
DOCK6	0	1	0	0	1	5	5
ERCC2	1	0	0	0	0	5	5
TUBGCP4	1	0	0	0	0	5	4
KERA	0	0	0	0	0	5	2
FREM1	1	1	1	1	4	4	4
RBP4	0	1	1	1	3	4	4
FRAS1	1	1	1	0	3	4	3
SALL4	0	1	1	1	3	4	3
FREM2	1	1	0	0	2	4	4
PQBP1	0	1	0	1	2	4	2
FOXC1	1	1	0	0	2	4	4
TUBB	0	1	0	0	1	4	3
SRD5A3	0	1	0	0	1	4	4
CAPN15	1	0	0	0	1	4	3
PLK4	1	0	0	0	0	4	2
B3GLCT	1	0	0	0	0	4	4
Six3	0	1	1	1	3	3	3
BMP7	0	1	1	1	3	3	3
B3GALNT2	0	1	0	0	1	3	3
SALL1	0	1	0	0	1	3	2
RPGRIP1L	0	1	0	0	1	3	2
POMGNT1	0	1	0	0	1	3	3
KIF11	0	1	0	0	1	3	3
CRYBB2	0	0	0	1	1	3	3
FKRP*	0	1	0	0	1	3	3
POMT2	0	1	0	0	1	3	3
C16orf62	1	0	0	0	0	3	3
SMO	1	1	1	0	3	2	2
ERCC5*	1	1	0	1	2	2	2
FOXL2*	0	1	0	1	2	2	2
PDE6D	0	1	0	1	2	2	1
GLI2*	0	1	0	0	1	2	2
CEP290*	0	1	0	0	1	2	2
RERE*	0	1	0	0	1	2	2
OLFM2	0	0	1	0	1	2	2
MAF*	0	0	1	0	1	2	2

DPYD*	0	0	0	1	1	2	2
SMG9	0	1	0	0	1	2	2
ACTB*	0	0	1	0	1	2	2
SNX3	0	1	0	0	1	2	2
FANCL*	0	1	0	0	1	2	2
NR6A1	0	0	0	0	0	2	2
CYP1B1*	1	0	0	0	0	2	2
FNBP4	0	1	1	1	3	1	1
VAX1	1	1	1	1	3	1	1
ERCC1	1	1	0	1	2	1	1
PITX2*	1	0	0	0	1	1	1
IPO13	0	1	0	0	1	1	1
SCLT1*	0	0	0	1	1	1	1
TCTN2*	0	1	0	0	1	1	1
CSPP1*	0	1	0	0	1	1	1
TMEM216*	0	1	0	0	1	1	1
ACTG1*	0	0	1	0	1	1	1
GTF2H5*	1	0	0	0	0	1	1
BMPR1B*	0	0	0	0	0	1	1
HMGB3	0	1	1	1	3	0	0
HDAC6	1	1	0	0	1	0	0
LRP2*	0	0	1	0	1	0	0
DAG1*	0	1	0	0	1	0	0
COX7B*	0	0	1	0	1	0	0
GRIP1*	1	0	0	0	1	0	0
PAX3*	0	0	1	0	1	0	0
SOX10*	0	0	1	0	1	0	0

Table 1. Genes reported to be associated with A/M in literature, and corresponding publications

This panel was generated by aggregating the genes associated by A/M according to the Genomics England PanelApp and the three reviews stated. A 1 within the PanelApp and three review columns, indicates that they have reported that gene as associated with A/M. Every gene above the black threshold has pathogenic variants reported in at least five individuals across two or more families and are deemed to have a confident association with A/M. Genes between the black and red thresholds are deemed to have a weak association with A/M, due to pathogenic variants reported in only one to four individuals, or with family data heavily suggesting an association. Genes with an asterisk (*) appear only in syndromic presentations with sporadic A/M features, indicating these may be incidental and may not represent true associations.

Participant	Participant phenotype	Sex	Gene	Chromosomal position	Variant	Variant type	Zygosity	Inheritance	ACMG classification	AlphaMissense	REVEL	Missense3D	SpliceAI	Gnomad AF (v3.1.2)
1	Microphthalmia and coloboma	F	ABCB6	2:219214974 C>T	c.1263G>A (p.Met421Ile)	Missense	Het	AD (M)	VUS: PM2, PP2	0.952	0.68	No structural damage detected	/	0
2	Microphthalmia and nystagmus	F	ABCB6	2:219214438 T>C	c.1337A>G (p.Glu446Gly)	Missense	Het	AD (P)	VUS: PM2, PP3	0.577	0.78	No structural damage detected	/	0
3	Microphthalmia, hypoplasia of the optic nerve	M	ACTG1	17:81511304 G>A	c.686C>T (p.Thr229Ile)	Missense	Het	AD (M)	Likely pathogenic: PM1, PM2, PP2, PP3	0.433	0.77	No structural damage detected	/	0
4			ALDH1A3	15:100885271T>C	c.104T>C (p.Phe35Ser)	Missense	Hom	AR	VUS: PP3, PP2	0.966	0.71	No structural damage detected	/	0
	Anophthalmia	F	PORCN	X:48515958 G>A	c.1087+5G>A	Splice	Het	XLD (M)	VUS: PM2, PP3	/	/	/	DG: 0.92	0
5	Microphthalmia, multiple systemic abnormalities	M	ATOH7 (HERC4, NRBF2, TACR2, CCAR1, CTNNA3, JMJD1C, NPFFR1, SLC16A9, DNAJC12, IPMK, SLC29A3, EGR2, DDX50, PBLD, REEP3, SIRT1, KIF1BP, PCBD1, VSIR, C10orf105, TMEM26, TFAM, PALD1, ANK3, LRRC20, CCDC6, CISD1, HK1, PRF1, TSPAN15, TBATA, BICC1, DNA2, EIF4EBP2, UBE2D1, SLC25A16, AC067752.1, RUFY2, STOX1, AIFM2,A DAMTS14, LRRTM3, COL13A1, HNRNPH3, MYPN, ADO, TET1, SRGN, VPS26A, NEUROG3, NODAL, HKDC1, FAM241B, TYSND1, RHOBTB1, DDX21, CABCOCO1, UNC5B, FAM13C, ARID5B, MRLN, CDK1, RTKN2, ZNF365, PHYHIPL, CDH23, SAR1A, SUPV3L1, PPA1, SGPL1, H2AFY2,)	10:57171179-71785816		Deletion	Het	AR (M)	Pathogenic: 1A (0), 2A (1), 2B (0), 2C (0), 2G (0), 2H (0), 3C (0.9), 4L (0.15), 5D (0.3)	/	/		/	/
6	Microphthalmia, glaucoma, cataracts	F	BCOR	X:40074489 CT>C	c.856del (p.Ser286AlafsTer92)	Frameshift	Mosaic	N/A	Likely pathogenic: PVS1, PM2	/	/		/	0.0000008295
7	Microphthalmia, glaucoma, cataracts	F	BCOR	X:40064590 CTATA>C	c.3244_3247del (p.Tyr1082 ValfsTer30)	Frameshift	Het	N/A	Likely pathogenic: PVS1, PM2	/	/		/	0
8	Microphthalmia, intellectual disability, cleft palate	F	BCOR	X:40063063 CA>C	c.3855del (p.Val1286CysfsTer8)	Frameshift	Mosaic	De novo	Likely pathogenic: PVS1, PM2	/	/		/	0
9	Microphthalmia, autism	F	BCOR	X:40054315 TG>T	c.4759del (p.Gln1587ArgfsTer31)	Frameshift	Het	N/A	Pathogenic: PVS1, PS4, PM6	/	/		/	0
10	Microphthalmia and cataracts	M	BMPR1B	4:95148798G>A	c.1127 G>A (p.Arg376His)	Missense	Het	N/A	Likely pathogenic: PS2, PM2, PP3	0.989	0.62	**Buried H-bond breakage**: This substitution disrupts all side-chain / side-chain H-bond(s) and/or side-chain / main-chain bond(s) H-bonds formed by a buried Arg residue (RSA 4.8%)- Arg376 NH1 to Pro432 O (2.86), Arg376 NH1 to Ser433 O (3.0), Arg376 NH2 to Pro426 O (2.52). **Buried/exposed switch**: This substitution results in a change between buried and exposed state of the target variant residue. Arg is buried (RSA 4.8%) and His is exposed (RSA 14.6%).	/	0.000006196
11	Microphthalmia, coloboma and global developmental delay.		CHD7	8:60741663 AC>A	c.232del (p.Gln78LysfsTer5)	Frameshift	Het	De novo	Likely pathogenic: PVS1, PS2, PM2	/	/		/	0
12	Microphthalmia, hypoplasia of the optic nerve	M	CHD7	8:60845058 A>G	c.5045A>G (His1682Arg)	Missense	Het	AD (M)	VUS: PM2, PP2, PP3	0.990	0.74	No structural damage detected	/	0.000002478
13	Microphthalmia and retinal dystrophy	M	COL4A1	13:110178176 C>A	c.2514G>T (p.Met838Ile)	Missense	Het	N/A	VUS: PM1, PM2, PM5, PP2	0.533	0.51	No structural damage detected	/	0
14	Microphthalmia, cataracts, global developmental delay	M	CRYBB2	22:25229563 G>C	c.434G>C (p.Arg145Pro)	Missense	Het	AD (P)	VUS: PM2, PP2, PP3	0.991	0.86	**Disallowed Phi/Psi angle**: The phi/psi angles are in the favored region for the wild-type residue but are in the outlier region for the variant residue.	/	0
15	Microphthalmia	F	ERCC2	19:45353112 C>T	c.1802G>A (p.Arg601Gln)	Missense	Het	AR (M)	Likely pathogenic: PM1, PM2, PM5, PP3	0.996	0.67	**Cavity altered**: The substitution leads to the contraction of cavity v	/	0.0003155
				19:45352613 G>A	c.1939C>T (p.Arg647Cys)	Missense	Het	N/A	Likely pathogenic: PM1, PM2, PM3, PP3	0.866	0.77	No structural damage detected	/	0.00003966
16	Microphthalmia	M	GDF6	8:96144663 G>A	c.1268C>T (p.Thr423Ile)	Missense	Het	AD (P)	VUS: PM2. PP3	0.623	0.78	No structural damage detected	/	0.0000006841
17	Microphthalmia, glaucoma and cataracts.	F	GJA8	1:147908245T>G	c.290T>G (p.Val97Gly)	Missense	Het	N/A	Likely pathogenic: PM1, PM2, PM6, PP2, PP3, PP5	0.594	0.85	No structural damage detected	/	0
18	Microphthalmia	F							Pathogenic: 1A (0), 2A (1), 2H (0), 3A (0), 5D (0)	/	/		/	/
18_M	Microphthalmia	F	HCCS (ARHGAP6, AMELX, MIR548AX)	X:10945466-11417014	/	Deletion	Het	XLD (M)		/	/		/	/
19	Anophthalmia	F	HDAC6	X:48823934 G>C	c.3316G>C (p.Ala1106Pro)	Missense	Het	N/A	Likely pathogenic: PM2, PM6, PP2, PP3	0.734	0.60	**Cavity altered**: The substitution leads to the expansion of cavity volume by 139.536 Å^3.	/	0
20	Microphthalmia, microcornea and anterior segment dysgenesis	M												
20_S	Microphthalmia, microcornea and anterior segment dysgenesis	M	KERA	12:91055473G>A	c.809C>T (p.Ser270Leu)	Missense	Hom	N/A	VUS: PM2	0.929	0.5	No structural damage detected	/	0.00002647
21	Microphthalmia, coloboma, intellectual disability	M	KMT2D	12:49053593 A>T	c.722T>A (p.Leu241Gln)	Missense	Het	AD (M)	VUS: PM2, PP2	0.539	0.54	No structural damage detected	/	0
22	Microphthalmia and coloboma	M	KMT2D	12:49049698 C>T	c.3890G>A (p.Arg1297His)	Missense	Het	AD (M)	VUS: PM2, PP2	0.805	0.46	No structural damage detected	/	0.000006569
23	Microphthalmia, coloboma, global developmental delay	M	KMT2D	12:49048064 CAT>C	c.4135_4136del (p.Met1379ValfsTer52)	Frameshift	Het	De novo	Pathogenic: PVS1, PS2, PS4, PM2	/	/		/	0
24	Microphthalmia	M	MAB21L2	4:150583408A>T	c.379A>T (p.Lys127Ter)	Stop Gain	Het	De novo	Likely pathogenic: PVS1, PM2	/	/		/	0
25	Microphthalmia	F	MFRP	11:119342977 TG>T	c.1150delC (p.His384ThrfsTer94)	Frameshift	Hom	AR (P/-)	Pathogenic: PVS1, PM3, PM2	/	/		/	0.000003731
26	Microphthalmia, global developmental delay.		MYRF	11:61770345 G>GCC	c.565_566dup (p.Pro190ArgfsTer19)	Frameshift	Mosaic	De novo	Pathogenic: PVS1, PS2, PM2	/	/		/	0.0000006645
27	Microphthalmia, coloboma, global developmental delay	M	MYRF	11:61776323 G>T	c.1390G>T (p.Val464Phe)	Missense	Het	AD (P)	VUS: PM2, PP2, PP3	0.925	0.23	No structural damage detected	/	0.00001055
28	Microphthalmia, hearing impairment and autism.	M	NDP	X:43954691-43959091		Deletion	Hemi	XLR (M)	Pathogenic: 1A (0), 2B (0), 2C (0.9), 3A (0),	/	/		/	/
29	Microphthalmia, supernumerary teeth	M							Likely pathogenic: 1A (0), 2B (0), 2C (0.90), 3A (0), 5D (0)					
29_M	Microphthalmia, cataracts	F	NHS	X:17153721-17599035		Deletion	Hemi	XLsD (M)		/	/		/	/
30	Microphthalmia, glaucoma	M	NHS	X:17479996 G>T	c.565+103674G>T	Splice	Hemi	XLsD (M)	VUS: PM2, PP3, PP4, BS2	/	/		AG: 0.74	0.0001889
31	Microphthalmia, intellectual disability and delayed gross motor development	M	NR6A1	9:124536055 C>T	c.902G>A (p.Trp301Ter)	Stop Gain	Het	N/A	Likely pathogenic: PVS1, PM2	/	/		/	0
32	Microphthalmia, microtia	F	OTX2	14:56804190 G>A	c.271C>T (p.Gln91Ter)	Stop gain	Het	N/A	Pathogenic: PVS1, PS4	/	/		DG: 0.88	0
33	Microphthalmia, coloboma, intellectual disability	F	OTX2	14:56802331 T>C	c.298A>G (p.Lys100Glu)	Missense	Het	AD (P)	VUS: PM1, PM2, PP3	1.000	0.77	No structural damage detected	/	0
34	Microphthalmia	M	OTX2 (EXOC5, AP5M1, NAA30, CCDC198)	14:56783923-57491668		Deletion	Het	De novo	Pathogenic: 1A (0), 2A (1), 2H (0), 3A (0), 4L (0.1), 5A (0.3)	/	/		/	/
35	Microphthalmia and nystagmus.	M	PAX2	10:100749771 C>CG	c.76dup (p. Val26GlyfsTer28)	Frameshift	Het	De novo	Pathogenic: PVS1, PS4, PM2	/	/		/	0
36	Anophthalmia or microphthalmia, and aniridia	F	PAX6	11:31806926 CT>C	c.-128-2del	Splice	Het	N/A	Pathogenic: PVS1, PS2, PM2, PP1, PP3	/	/		AL:0.65	0

No.	Phenotype	Sex	Gene	Genomic position	cDNA (protein) change	Type	Zygosity	Inheritance	Classification	Score 1	Score 2	Structural damage	Splice	AF
37	Microphthalmia, coloboma, cataracts	M										**Buried charge replaced:** This substitution replaces a buried charged residue (Arg, RSA 6.8%) with an uncharged residue (GLY). **Buried salt bridge breakage:** This substitution disrupts a salt bridge formed by OE1 atom of Glu226 and NH1 atom of ARG 261 (distance: 3.114 Å). The wild-type residue has RSA of 6.8%. **Cavity altered:** The substitution leads to the expansion of cavity volume by 120.528 Å^3. **Buried H-bond breakage:** This substitution disrupts all side-chain / side-chain H-bond(s) and/or side-chain / main-chain bond(s) H-bonds formed by a buried Arg residue (RSA 6.8%)- Arg26A NE to Leu21 O (2.82), Arg26 NH2 Ile66 O (2.90).		0
37_M	Microphthalmia, cataracts, nystagmus	F	PAX6	11:31802769 G>C	c.76C>G (p.Arg26Gly)	Missense	Het	AD	Pathogenic: PS4, PM1, PP2, PM2, PM5, PP3	1.000	0.93			0
38	Microphthalmia, nystagmus and coloboma	M	PAX6	11:31800843T>C	c.413A>G (p.Asn138Ser)	Missense	Het	De novo	Likely pathogenic: PM1, PP2, PM2, PM5, PP3	0.982	0.8	No structural damage detected		0
39	Microphthalmia, **global developmental delay**, **intellectual disability** and **delayed speech**	F	PAX6	11:31800844 T>G	c.412A>C (p.Asn138His)	Missense	Het	De novo	Pathogenic: PS2, PM1, PM2, PM5, PP2, PP3	0.998	0.94	No structural damage detected		0
40	Microphthalmia, coloboma, **intellectual disability**	M	PORCN	X:48514269 C>T	c.749C>T (p.Ser250Phe)	Missense	Hemi	XLR (M)	Likely pathogenic: PM2, PP3, PP4, PP5	0.876	0.91	**Buried/exposed switch:** This substitution results in a change between buried and exposed state of the target variant residue. Ser is buried (RSA 4.6%) and Phe is exposed (RSA 17.7%).		8.25E-07
41	Microphthalmia, coloboma, developmental delay	M	PRR12	19:49601919 G>C	c.4773+1G>C	Splice	Het	De novo	Pathogenic: PVS1, PM2, PM3	/	/		DL: 1.00	0
42	Microphthalmia	M	PRSS56	2:232522034 G>A	c.320G>A (p.Gly107Glu)	Missense	Hom	N/A	VUS: PM2, PP3	0.831	N/A	No structural damage detected		0.000006569
				2:232524762 G>C	c.1439G>C (p.Arg480Pro)	Missense	Het	AR (M)	VUS: PM2, PM3	0.395	N/A	No structural damage detected		0
43	Anophthalmia or microphthalmia	F	PRSS56	2:232525264 TG>T	c.1573delG (p.Val525CysfsTer55)	Frameshift	Het	AR (P)	Likely pathogenic: PVS1, PM2, PP5	/	/			0.00006574
44	Microphthalmia and disorder of bone.	M	PTCH1	9:95475579-95477447		Deletion	Het	De novo	Likely pathogenic: 1A (0), 2B (0), 2E (0.45), 3A (0), 5A (0.45)	/	/			/
45	Microphthalmia	F	PTCH1	9:95453509 G>GC	c.3417dup (p.Leu1140AlafsTer5)	Frameshift	Het	De novo	Pathogenic: PVS1, PS2, PM2	/	/			0
46	Microphthalmia, cataracts, global developmental delay	M	RAB3GAP1	2:135162745 TAGAAA>T	c.2387_2390del (p.Glu796ValfsTer12)	Frameshift	Het	AR (M)	Pathogenic: PVS1, PM2, PM3	/	/		AL: 0.98	0
			RAB3GAP1 (ZRANB3)	2:135165337-135511837	/	Deletion	Het	AR (P)	Likely pathogenic: 1A (0), 2B (0), 2D (0.9), 3A (0), 4L (0.15), 4N (-0.10)	/	/		/	/
47	Microphthalmia, coloboma	F	RARB	3:25569923 G>C	c.609+5G>C	Splice	Het	N/A	VUS: PM2, PP3	/	/		DL: 0.75	0
48	Microphthalmia, coloboma, **hearing impairment**	M	RERE	1:8361310 T>C	c.2197A>G (p.Met733Val)	Splice	Het	N/A	VUS: PM2, PP3	0.069			DG: 0.97	0.000005588
49	Microphthalmia, coloboma	F	SIX3	2:44942679 G>A	c.575G>A (p.Arg192His)	Missense	Het	De novo	Pathogenic: PS2, PM1, PM2, PP2, PP3	1.000	0.94	No structural damage detected		0
50	Microphthalmia, coloboma	M	SLC24A5	1:108157481 C>T	c.650G>A (p.Arg217His)	Missense	Het	N/A	Pathogenic: PS2, PS3, PM2, PM5, PP3	0.946	0.89	No structural damage detected		0
51	Microphthalmia, aplasia/agenesis of the nose	M	SMCHD1	18:2697034 A>G	c.1043A>G (p.His348Arg)	Missense	Het	De novo	Pathogenic: PS2, PS4, PM2, PP2, PP5	0.925	0.45	Cavity altered: The substitution leads to the contraction of cavity vo		0
52	Microphthalmia, **hearing impairment**	M	SMCHD1	18:2707821 G>C	c.2161G>C (p.Glu721Gln)	Missense	Het	AD (P)	VUS: PM2, PM2	0.738	0.28	No structural damage detected		0.000006574
53	Microphthalmia and glaucoma	M	SOX2	3:181712418 GGCGGCGGCGG	c.70_89del (p.Asn24ArgfsTer65)	Frameshift	Het	De novo	Pathogenic: PVS1, PS2, PS3, PM2, PP5	/	/			0
54	Microphthalmia, coloboma, **hearing impairment**	M	TFAP2A	6:10404562 C>T	c.716G>A (p.Arg239Gln)	Missense	Het	De novo	Pathogenic: PS2, PM1, PM2, PP2, PP3	1.000	0.92	**Buried charge replaced:** This substitution replaces a buried charged residue (ARG, RSA 1.6%) with an uncharged residue (GLN). **Buried H-bond breakage:** This substitution disrupts all side-chain / side-chain H-bond(s) and/or side-chain / main-chain bond(s) H-bonds formed by a buried ARG residue (RSA 1.6%): Arg239 NE to Glu244 OE2 (3.22), Arg239 NH1 to Glu244 OE1 (3.35), Arg239 NH1 to Thr292 O (2.47), Arg239 NH1 to Leu294 O (3.83), Arg239 NH2 to Thr292 O (3.60), Arg239 NH2 Leu294 O (3.23). **Buried salt bridge breakage:** This substitution disrupts a salt bridge formed by NE atom of Arg239 and OE2 atom of Glu244 (distance: 3.224 Å). The wild-type residue has RSA of 1.6%.		6.20E-07
55	Microphthalmia, coloboma, **hearing impairment**	M	TFAP2A	6:10404509 T>C	c.769A>G (p.Arg257Gly)	Missense	Het	De novo	Pathogenic: PS3, PS4, PM1, PM2, PM5, PP2, PP3	1.000	0.97	No structural damage detected		0
56	Microphthalmia, coloboma and cataracts	F	TFAP2A	6:10400571G>T	c.890C>A (p.Ala299Asp)	Missense	Het	N/A	VUS: PM2, PP3, PP2	1	0.999	**Buried hydrophilic introduced:** This substitution replaces a buried hydrophobic residue (Ala, RSA 0.0%) with a hydrophilic residue (Asp, RSA 0.0%). **Buried charge introduced:** This substitution replaces a buried uncharged residue (Ala, RSA 0.0%) with a charged residue (Asp).		0

Table 2. Candidate variants identified within the 100KGP anophthalmia and microphthalmia cohort

Participant numbers in bold are ones who have been previously molecular diagnosed. Phenotypes in bold are ones which are associated with the respective gene. Inheritance of the variant is stated whether it was inherited maternally (M), or paternally (P), de novo, or whether there is insufficient parental data (N/A). AD: Autosomal dominant, AF: Allele frequency, AG: Acceptor gain, AL: Acceptor loss, AR: Autosomal recessive, DG: Donor gain, DL: Donor loss, Het: Heterozygous, Hom: Homozygous, VUS: variant of uncertain significance, XLD: X-linked dominant, XLR: X-linked recessive, XLsD: X-linked semi-dominant

Participant	Recruited rare disease	Gene	Chromosomal position	HGVS c	HGVS p	Zygosity	Inheritance	Pedigree	ACMG classification
1	Ocular malformation - Anophthalmia or microphthalmia	ABCB6	2:219214974 C>T	c.1263G>A	p.Met421Ile	Het	AD (M)	Trio, unaffected parents	VUS: PM2, PP2
2	Ocular malformation - Anophthalmia or microphthalmia	ABCB6	2:219214438 T>C	c.1337A>G	p.Glu446Gly	Het	AD (P)	Trio, unaffected parents	VUS: PM2, PP3
3	Neurodevelopmental disorder - Intellectual disability	ACTG1	17:81511304 G>A	c.686C>T	p.Thr229Ile	Het	AD (M)	Trio, unaffected parents	Likely pathogenic: PM1, PM2, PP2, PP3
4	Ocular malformation - Anophthalmia or microphthalmia	ALDH1A3	15:100885271T>C	c.104T>C	p.Phe35Ser	Hom	AR	Trio, unaffected parents	VUS: PM2, PP3, PP2
		PORCN	X:48514269 C>T	c.749C>T	p.Ser250Phe	Hemi	XLR (M)	Duo, unaffected mother	Likely pathogenic: PM2, PP3, PP4, PP5
5	Neurodevelopmental disorder - Intellectual disability	ATOH7 (HERC4, NRBF2, TACR2, CCAR1, CTNNA3, JMJD1C, NPFFR1, SLC16A9, DNAJC12, IPMK, SLC29A3, EGR2, DDX50, PBLD, REEP3, SIRT1, KIF1BP, PCBD1, VSIR, C10orf105, TMEM26, TFAM, PALD1, ANK3, LRRC20, CCDC6, CISD1, HK1, PRF1, TSPAN15, TBATA, BICC1, DNA2, EIF4EBP2, UBE2D1, SLC25A16, AC067752.1, RUFY2, STOX1, AIFM2,A DAMTS14, LRRTM3, COL13A1, HNRNPH3, MYPN, ADO, TET1, SRGN, VPS26A, NEUROG3, NODAL, HKDCI, FAM241B, TYSND1, RHOBTB1, DDX21, CABCOCO1, UNC5B, FAM13C, ARID5B, MRLN, CDK1, RTKN2, ZNF365, PHYHIPL, CDH23, SAR1A, SUPV3L1, PPA1, SGPL1, H2AFY2.)	10:57171179-71785816	/	/	Het	AR (M)	Trio, affected mother, affected sibling	Pathogenic: 1A (0), 2A (1), 2B (0), 2C (0), 2G (0), 2H (0), 3C (0.9), 4L (0.15), 5D (0.3)
6	Anterior segment anomalies - Cataracts	BCOR	X:40074489 CT>C	c.856del	p.Ser286AlafsTer92	Mosaic	N/A	Duo, unaffected mother	Likely pathogenic: PVS1, PM2
7	Anterior segment anomalies - Cataracts	BCOR	X:40064590 CTATA>C	c.3244_3247del	p.Tyr1082 ValfsTer30	Het	N/A	Duo, unaffected father	Likely pathogenic: PVS1, PM2
8	Neurodevelopmental disorder - Intellectual disability	BCOR	X:40063063 CA>C	c.3855del	p.Val1286CysfsTer8	Mosaic	De novo	Trio, unaffected parents	Likely pathogenic: PVS1, PM2
9	Ocular malformation - Anophthalmia or microphthalmia	BCOR	X:40054315 TG>T	c.4759del	p.Gln1587ArgfsTer31	Het	N/A	Duo, unaffected mother	Pathogenic: PVS1, PS4, PM6
10	Ocular malformation - Anophthalmia or microphthalmia	BMPR1B	4:95148798G>A	c.1127 G>A	p.Arg376His	Het	N/A	Duo, unaffected mother	Likely pathogenic: PS2, PM2, PP3
11	Ocular malformation - Anophthalmia or microphthalmia	CHD7	8:60741663 AC>A	c.232del	p.Gln78LysfsTer5	Het	De novo	Trio, unaffected parents	Pathogenic: PVS1, PS2, PM2
12	Ocular malformation - Anophthalmia or microphthalmia	CHD7	8:60845058 A>G	c.5045A>G	p.His1682Arg	Het	AD (M)	Trio, unaffected parents	VUS: PM2, PP2, PP3
13	Posterior segment anomalies - Familial exudative vitreoretinopathy	COL4A1	13:110178176 C>A	c.2514G>T	p.Met838Ile	Het	N/A	Duo, unaffected father	VUS: PM1, PM2, PM5, PP2,
14	Neurodevelopmental disorder - Intellectual disability	CRYBB2	22:25229563 G>C	c.434G>C	p.Arg145Pro	Het	AD (P)	Trio, unaffected parents	VUS: PM2, PP2, PP3
15	Ocular malformation - Anophthalmia or microphthalmia	ERCC2	19:45353112 C>T	c.1802G>A	p.Arg601Gln	Het	AR (M)		Likely pathogenic: PM1, PM2, PM5, PP3
			19:45352613 G>A	c.1939C>T	p.Arg647Cys	Het	N/A	Duo, unaffected mother	Likely pathogenic: PM1, PM2, PM3, PP3
16	Ocular malformation - Anophthalmia or microphthalmia	GJA8	1:147908245T>G	c.290T>G	p.Val97Gly	Het	N/A	Duo, unaffected mother	Likely pathogenic: PM1, PM2, PM6, PP2, PP3, PP5
17	Ocular malformation - Anophthalmia or microphthalmia	GDF6	8:96144663 G>A	c.1268C>T	p.Thr423Ile	Het	AD (P)	Trio, unaffected parents	VUS: PM2. PP3
18 18_M	Ocular malformation - Anophthalmia or microphthalmia	HCCS (ARHGAP6, AMELX, MIR548AX)	X:10945466-11417014	/	/	Het	XLD (M)	Duo, affected mother	Pathogenic: 1A (0), 2A (1), 2H (0), 3A (0), 5D (0)
19	Ocular malformation - Anophthalmia or microphthalmia	HDAC6	X:48823934 G>C	c.3316G>C	p.Ala1106Pro	Het	N/A	Duo, unaffected mother	Likely pathogenic: PM2, PM6, PP2, PP3
20 20_S	Ocular malformation - Anophthalmia or microphthalmia	KERA	12:91055473G>A	c.809C>T	p.Ser270Leu	Hom	N/A	Trio, unaffected father, affected	VUS: PM2
21	microphthalmia	KMT2D	12:49053593 A>T	c.722T>A	p.Leu241Gln	Het	AD (M)	Trio, unaffected parents	VUS: PM2, PP2
22	Ocular malformation - Anophthalmia or microphthalmia	KMT2D	12:49049698 C>T	c.3890G>A	p.Arg1297His	Het	AD (M)	Duo, unaffected mother	VUS: PM2, PP2
23	Ocular malformation - Anophthalmia or microphthalmia	KMT2D	12:49048064 CAT>C	c.4135_4136del	p.Met1379ValfsTer52	Het	De novo	Trio, unaffected parents	Pathogenic: PVS1, PS2, PS4, PM2
24	Neurodevelopmental disorder - Holoprosencephaly	MYRF	11:61770345 G>GCC	c.565_566dup	p.Pro190ArgfsTer19	Mosaic	De novo	Trio, unaffected parents	Pathogenic: PVS1, PS2, PM2

#	Recruited disease	Gene	Genomic position	c.	p.	Zygosity	Inheritance	Pedigree	Classification
25	Neurodevelopmental disorder - Intellectual disability	MYRF	11:61776323 G>T	c.1390G>T	p.Val464Phe	Het	AD (P)	Duo, unaffected father	VUS: PM2, PP2
26	Ocular malformation - Anophthalmia or microphthalmia	MAB21L2	4:150583408A>T	c.379A>T	p.Lys127Ter	Het	De novo	Trio, unaffected parents	Likely pathogenic: PVS1, PM2
27	Ocular malformation - Anophthalmia or microphthalmia	MFRP	11:119342977TG>T	c.1150delC	p.His384ThrfsTer94	Hom	AR (P/-)	Duo, unaffected father	Pathogenic: PVS1, PM3, PM2
28	Hypothalamic and pituitary disorders - Idiopathic hypogonadotropic hypogonadism	NDP	X:43954691-43959091	/	/	Hemi	XLR (M)	Duo, unaffected mother	Pathogenic: 1A (0), 2B (0), 2C (0.9), 3A (0),
29 29_M	Anterior segment anomalies - Cataracts	NHS	X:17153721-17599035	/	/	Hemi	XLsD (M)	Duo, affected mother	Likely pathogenic: 1A (0), 2B (0), 2C (0.90), 3A (0), 5D (0)
30	Ocular malformation - Anophthalmia or microphthalmia	NHS	X:17479996 G>T	c.565+103674G>T	/	Hemi	XLsD (M)	Trio, unaffected parents	Likely Pathogenic: PS4, PM2, PP3, PP4
31	Ocular malformation - Anophthalmia or microphthalmia	NR6A1	9:124536055 C>T	c.902G>A	p.Trp301Ter	Het	N/A	Duo, unaffected mother	Likely pathogenic: PVS1, PM2
32	Neurodevelopmental disorder - Intellectual disability	OTX2	14:56,804,190 G>A	c.271C>T	p.Gln91Ter	Het	N/A	Singleton	Pathogenic: PVS1, PS4
33	Multisystem groups - Undiagnosed monogenic disorder	OTX2	14:56802331 T>C	c.298A>G	p.Lys100Glu	Het	AD (P)	Trio, unaffected parents	VUS: PM1, PM2, PP3
34	Ocular malformation - Anophthalmia or microphthalmia	OTX2 (EXOC5, AP5M1, NAA30, CCDC	14:56783923-57491668	/	/	Het	De novo	Trio, unaffected parents	Pathogenic: 1A (0), 2A (1), 2H (0), 3A (0), 4L (0.1), 5A (0.3)
35	Ocular malformation - Anophthalmia or microphthalmia	PAX2	10:100749771 C>CG	c.76dup	p. Val26GlyfsTer28	Het	De novo	Trio, unaffected parents	Pathogenic: PVS1, PS2, PS4, PM2
36	Ocular malformation - Anophthalmia or microphthalmia	PAX6	11:31806926 CT>C	c.-128-2del	/	Het	N/A	Duo, unaffected mother	Pathogenic: PVS1, PS2, PM2, PP1, PP3
37 37_M	Neurodevelopmental disorder - Intellectual disability	PAX6	11:31802769 G>C	c.76C>G	p.Arg26Gly	Het	AD	Trio, affected mother, unaffected father	Pathogenic: PS4, PM1, PP2, PM2, PM5, PP3
38	Ocular malformation - Anophthalmia or microphthalmia	PAX6	11:31800843T>C	c.413A>G	p.Asn138Ser	Het	De novo	Trio, unaffected parents	Likely pathogenic: PM1, PP2, PM2, PM5, PP3
39	Neurodevelopmental disorder - Intellectual disability	PAX6	11:31800844 T>G	c.412A>C	p.Asn138His	Het	De novo	Trio, unaffected parents	Pathogenic: PS2, PM1, PM2, PM5, PP2, PP3
40	Ocular malformation - Anophthalmia or microphthalmia	PORCN	X:48514269 C>T	c.749C>T	p.Ser250Phe	Hemi	XLR (M)	Duo, unaffected mother	Likely pathogenic: PM2, PP3, PP4, PP5
41	Ocular malformation - Anophthalmia or microphthalmia	PRR12	19:49601919 G>C	c.4773+1G>C	/	Het	De novo	Trio, unaffected parents	Pathogenic: PVS1, PM2, PP3
42	Ocular malformation - Anophthalmia or microphthalmia	PRSS56	2:232522034G>A	c.320G>A	p.Gly107Glu	Hom	N/A	Singleton	VUS: PM2, PP3
43	Ocular malformation - Anophthalmia or microphthalmia	PRSS56	2:232524762G>C	c.1439G>C	p.Arg480Pro	Het	AR (M)		VUS: PM2, PM3
	Ocular malformation - Anophthalmia or microphthalmia		2:232525264TG>T	c.1573delG	p.Val525Cysfs*55	Het	AR (P)	Trio, unaffected parents	Likely pathogenic:PVS1, PM2, PP5
44	Ocular malformation - Anophthalmia or microphthalmia	PTCH1	9:95453509 G>GC	c.3417dup	p.Leu1140AlafsTer5	Het	De novo	Trio, unaffected parents	Pathogenic: PVS1, PS2, PM2
45	Ocular malformation - Anophthalmia or microphthalmia	PTCH1	9:95475579-95477447	/	/	Het	De novo	Trio, unaffected parents	Likely pathogenic: 1A (0), 2B (0), 2E (0.45), 3A (0), 5A (0.45)
46	Multisystem groups - Undiagnosed monogenic disorder	RAB3GAP1	2:135162745 TAGAAT>T	c.2387_2390del	p.Glu796ValfsTer12	Het	AR (M)		Pathogenic: PVS1, PM2, PM3
		RAB3GAP1 (ZRANB3)	2:135165337-135511837	/	/	Het	AR (P)	Trio, unaffected parents	Likely pathogenic: 1A (0), 2B (0), 2D (0.9), 3A (0), 4L (0.15), 4N (-0.10)
47	Multisystem groups - Undiagnosed monogenic disorder	RARB	3:25569923 G>C	c.609+5G>C	/	Het	N/A	Singleton	VUS: PM2, PP3
48	Non-syndromic hearing loss - congenital hearing impairment	RERE	1:8361310 T>C	c.2197A>G	p.Met733Val	Het	N/A	Duo, unaffected father	VUS: PM2, PP3
49	Ocular malformation - Anophthalmia or microphthalmia	SIX3	2:44942679 G>A	c.575G>A	p. Arg192His	Het	De novo	Trio, unaffected parents	Pathogenic: PS2, PM1, PM2, PP2, PP3
50	Dysmorphic disorders - Coarse facial features including Coffin-Siris-like disorders	SLC24A5	1:108157481 C>T	c.650G>A	p.Arg217His	Het	N/A	Trio, unaffected father and unaffected family member	Pathogenic: PS2, PS3, PM2, PM5, PP3
51	Structural renal and and urinary tract disease - Cystic kidney disease	SMCHD1	18:2697034 A>G	c.1043A>G	p.His348Arg	Het	De novo	Quintet, unaffected parents and two unaffected siblings	Pathogenic: PS2, PS4, PM2, PP2, PP5
52	Ocular malformation - Anophthalmia or microphthalmia	SMCHD1	18:2707821 G>C	c.2161G>C	p.Glu721Gln	Het	AD (P)	Trio, unaffected parents	VUS: PM2, PP2
53	Structural renal and and urinary tract disease - Congenital anomaly of the kidneys	SOX2	3:181712418 GGCGGCGGCGGCA	c.70_89del	p.Asn24ArgfsTer65	Het	De novo	Trio, unaffected parents	Pathogenic: PVS1, PS2, PS3, PM2, PP5
54	Ocular malformation - Coloboma	TFAP2A	6:10404562 C>T	c.716G>A	p.Arg239Gln	Het	De novo	Trio, unaffected parents	Pathogenic: PS2, PM1, PM2, PP2, PP3
55	Ultra-rare undiagnosed monogenic disorder	TFAP2A	6:10404509 T>C	c.769A>G	p.Arg257Gly	Het	De novo	Trio, unaffected parents	Pathogenic: PS3, PS4, PM1, PM2, PM5, PP2, PP3
56	Ocular malformation - Anophthalmia or microphthalmia	TFAP2A	6:10400571G>T	c.890C>A	p.Ala299Asp	Het	N/A	Duo, unaffected mother	VUS: PM2, PP3, PP2

Table 3. Recruitment and pedigree information of participants in the 100KGP anophthalmia and microphthalmia cohort

Phenotypes in bold are ones which are associated with the respective gene. Recruited disease refers to the disorder for which the family was recruited to the 100K GP. Inheritance of the variant is stated whether it was inherited maternally (M), or paternally (P), de novo, or whether there is insufficient parental data (N/A). AD: Autosomal dominant, AF: Allele frequency, AG: Acceptor gain, AL: Acceptor loss, AR: Autosomal recessive, DG: Donor gain, DL: Donor loss, Het: Heterozygous, Hom: Homozygous, VUS: variant of uncertain significance, XLD: X-linked dominant, XLR: X-linked recessive, XLsD: X-linked semi-dominant.