A walk in the park – identifying healthy greenspaces using scents

1. William T. Kay[a,*] https://orcid.org/0000-0002-2561-4856

2. Anya Lindström Battle[a] https://orcid.org/0000-0001-6356-4688

3. Mahal Humberstone[a]

4. Molly Tucker[a] https://orcid.org/0009-0002-5463-4239

5. Kieran E. Storer[a] https://orcid.org/0000-0001-9844-7174

6. Geoffrey Kite[b] https://orcid.org/0000-0003-4482-0409

7. Katherine Willis[a] https://orcid.org/0000-0002-6763-2489

[a] Department of Biology, University of Oxford, South Parks Road, Oxford OX1 3RB, UK

[b] Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK.

* Corresponding author: william.kay@biology.ox.ac.uk

Societal Impact Statement

As urbanisation accelerates globally, access to nature is increasingly recognised as vital for public health and wellbeing. We captured and analysed plant-emitted airborne "scent signatures" across Oxford's urban greenspaces to assess their potential health relevance. We found that sites differ in levels of health-promoting volatiles and harmful pollutants, and that these compounds vary with environmental conditions such as temperature and humidity. Our findings provide a novel framework for evaluating urban greenspaces, informing planning, air-quality management, and public health strategies to create greener, healthier cities.

1. Abstract

1.   Biogenic volatile organic compounds (bVOCs) emitted by plants can promote stress reduction and cognitive benefits, whereas anthropogenic VOCs (aVOCs) common in urban air, such as BTEX compounds, pose health risks. We aimed to characterise and compare these airborne chemicals across urban greenspaces and develop a novel, health-oriented metric for site evaluation.

2.   Air samples were collected across six Oxford greenspaces on a single date using Tenax™ filters and analysed by GC–MS. Data were processed using metabolomics-style workflows, identifying 245 biologically relevant compounds. Multivariate analyses compared site profiles, and one site was monitored over 12 months to assess environmental influences.

3.   Greenspaces exhibited significantly different volatile signatures, including variation in both health-promoting bVOCs and harmful aVOCs. Temperature, humidity, wind speed, and rainfall significantly influenced the presence and abundance of beneficial compounds.

4.   These results highlight the importance of considering airborne chemistry in urban planning and public health decision-making. Integrating VOC profiling into planning and public health strategies could support delivery of higher-quality, health-promoting urban environments and inform guidance on optimal timing for outdoor activity.

2. Keywords

Greenspace chemical ecology, Gas Chromatography-Mass Spectrometry, volatile analysis, Volatile Organic Compounds, Biogenic Volatile Organic Compounds, terpenes, health

3.   Introduction

Biogenic volatile organic compounds (bVOCs) are a chemically diverse group of carbon-based secondary metabolites produced across all kingdoms of life (Wang et al. 2024). They function in stress protection, signalling, and communication within and between species (Weisskopf, Schulz, and Garbeva 2021) and many are released into the air via passive diffusion from different tissues. In plants, bVOCs such as isoprene, monoterpenes, and sesquiterpenes account for almost all biogenic emissions (Guenther et al. 2012). Among these, monoterpenes such as alpha-pinene, beta-pinene, and limonene are key contributors to both atmospheric chemistry and human well-being. Numerous studies link exposure to plant-emitted terpenes with lowered cortisol, blood pressure, and heart rate, as well as improved mood, cognition, and immune function (Zeng et al. 2022; Park et al. 2010; Li 2010; Bratman et al. 2015). By contrast, anthropogenic VOCs (aVOCs) derived from transport, plastics, and industrial sources - particularly benzene, toluene, ethylbenzene, and xylene (collectively known as the BTEX compounds) - are well-known urban pollutants associated with respiratory and cardiovascular diseases and cancer (Gao et al. 2023).

Because of their location the 'scentscapes' (the total volatile profile in a specific time and place) of urban greenspaces are likely to comprise both bVOCs and aVOCs. Understanding the balance between these "healthy" and "unhealthy" volatiles is therefore essential for assessing how urban greenspaces influence well-being, and for driving policy for positive changes in urban development.

Here, we present a new analytical pipeline for characterising ambient air VOCs *in situ*, integrating field sampling and thermal desorption gas chromatography mass spectrometry (TD-GC-MS) analysis. Although similar methods have been developed for headspace sampling (Yin et al. 2022; de Carvalho Couto et al. 2024; Reyrolle et al. 2024), none have yet been published for the analysis of ambient air. We apply this approach to six Oxford greenspaces to test whether site location determines the distribution of biogenic versus anthropogenic volatiles - and to identify those environments most likely to promote human health.

Additionally, it is well known that the release of bVOCs varies in time. For instance, plants coordinate

their volatile production with the availability of substrates from photosynthesis, which varies

considerably over the course of a day. Indeed, the production of most plant volatiles is under diurnal

and/or circadian control; on a longer timescale, the effects of factors such as seasonality, phenology,

and other environmental effects including temperature and humidity on the emission of volatiles

have been well established (Joo et al. 2018; Yang et al. 2024; Mu et al. 2022; Li et al. 2023).

Firsly, we analysed volatiles on a single date as a snapshot across six outdoor greenspace sites.

Secondly, we sampled a single site – the Botanical Garden outdoors – over a full year, linking

differences in volatile fingerprints to seasonal environmental changes and transient weather events.

Our work aimed to assess three main hypotheses: (i) that a site's location is a significant driver of

volatile difference enabling the separation of greenspaces by their scentscape, (ii) that urban

greenspace volatile profiles change over time dependent on environmental variables, and iii) there is

sufficient distinction between the presence and abundance of both bVOCs and aVOCs between

urban green spaces that it is possible to determine which may be more healthy/unhealthy for human

recreation.

4.   Materials and Methods

4.1   Site locations and sampling strategy

Six urban greenspace sites around Oxford were chosen as sampling sites due to their (i) public

accessibility, (ii) different ecologies, and (iii) contrasting distance from the city centre. We selected

five outdoor greenspaces (Botanic Garden Outdoors, University Parks Dense, University Parks Open,

Warneford Meadow, and Wytham Woods) and one indoor site (the Rainforest Glasshouse at the

Oxford Botanical Gardens) (Figure 1A).

To investigate the characteristic VOC profiles of the selected sites, we employed a collection protocol

adapted from Walker et al (2023) with the addition of housing the pump and apparatus within a

bespoke metal box (VOKSBOX, B). Ambient air was captured using a GilAir® pump (Sensidyne, LP, St.

Petersburg, FL, USA) connected simultaneously to 4 replicate Markes (Markes international Ltd.

Bridgend, UK) Tenax® TA tubes using a flow rate of 400ml/min for 2 hours (12 L of air per tube).

Samples were collected at a height of 1.5m – based on the approximate level of the adult human

nose while walking - between the hours of 11:00 - 13:00 (+/- 1 hour) GMT. Tubes were capped

immediately after collection and stored at - 20 °C for up to 14 days before GC-MS analysis.

A pilot dataset was collected on 27 November 2024 and used to assess the hypothesis that sampling

sites have distinct volatile profiles (Figure 2). As well as looking at a snapshot of the scentscapes

across the six urban greenspaces, we also wanted to explore how these scentscapes change across

time and in the face of environmental variation (Figure 3). Because of the known impacts of

seasonality on bVOC emissions we focused only on biogenic volatiles. Due to the higher numbers of

biogenic volatiles present in the Botanical Outdoors location, and as this was likely to be the site with

the highest footfall, we also sampled this site on 20 January 2025, 04 March 2025, 30 April 2025, 30

June 2025, 24 July 2025, and 04 September 2025. This allowed us to capture a range of temperature,

humidity, precipitation, and wind conditions (Figure 3A). All Botanical Outdoors samples during the

year were taken from 11:00 to 13:00 in exactly the same GPS location and height.

Site-specific temperature and relative humidity at the time of sampling were recorded using a Kestrel

Drop D2 data logger (Nielsen-Kellerman Company, Boothwyn, PA, USA). Historical meteorological

data were obtained from multiple sources: air temperature data were retrieved from the Oxford

Radcliffe Weather Station, and mean values recorded between 11:00 and 13:00 were used to

coincide with the sampling window. Precipitation, humidity, wind speed, and wind direction data

were obtained from the Kidlington weather station via Weather Underground

(www.wunderground.com).

Figure 1: A: Map of the sampling locations used in this study. (Botanic Garden Outdoors; Botanic Garden Rainforest Glasshouse; University Parks Dense; University Parks Open; Warneford Meadow; Wytham Woods). Numbers on roads indicate Annual Average Daily Traffic (AADF) for 2023 as representative data for each site (GOV.UK 2023) B: Sampling box (VOKSBOX) developed for this work on a tripod at 1.5m height. Air from the GilAir® sample pump is split up to 6 ways using brass splitters with silicone tubing. C-E: A flow chart of the methods used in this paper for acquisition, identification, and processing of compounds. Red text shows number of compounds identified after each processing step. F: Example chromatogram resulting from sample collection in Botanical Gardens Outdoors sampling site and GCMS protocol - annotated with positions of alkanes used to build Kovats retention index.

4.2 Identification and analysis of VOC profiles

GC-MS Protocol: For identification of VOC profiles (Figure 1C-E), we employed gas chromatography

mass spectrometry (GC-MS) analysis at the Jodrell laboratory, Royal Botanic Gardens, Kew. Samples

were stored at room temperature for at least 2 hours before desorption using a TD100-xr automated

thermal desorption (TD) unit from Markes (Markes international Ltd. Bridgend, UK). Samples were

desorbed on to a Tenax® TA packed cold trap (set to 4 °C) by heating the sampling tubes for 10 mins

at 230 °C using a flow rate of 50 ml/min helium. Samples were purged from the cold trap onto a J&W

DB-5ms GC column (30 m × 0.25 mm × 0.25 μm) within an Agilent 8890 GC System (Agilent

Technologies, Santa Clara, CA, USA) by heating from 4 °C to 300 °C over 5 mins at a purge flow rate of

1ml/min Helium. Chromatography was achieved by heating the column at an initial temperature of

40 °C at a rate of 3 °C/min to 220 °C at a flow rate of 1 ml/min Helium. Data were collected in

electron ionization (EI) mode using an Agilent 5977C GC/ MSD over a mass range of 50–400 amu.

Library Development: The GC-MS library used for identification of compounds was a modified

version of the NIST20 Wiley Registry 12th edition mass spectral library minimised to compounds

existing in the Chemical Entities of Biological Interest (ChEBI) database (Degtyarenko et al. 2007).

Only compounds with entries for retention indexes (RI) on semi-standard non-polar columns (to

match the J&W DB-5ms column used) were selected. To avoid mismatches, isomers with the same RI

were combined (e.g. (−)-alpha-pinene / (+)-alpha-pinene, became alpha-pinene). The final library

used (W12N20MAIN_chebi_WK_edit.MSP) contained 3,027 unique compounds.

Pre-Analysis: To aid in the correct identification of compounds, a non-isothermal Kovats index

(Kovats 1958) was built using retention times from C8 – C20 alkane standards (Merck KGaA,

Darmstadt, Germany,, 04070-1ML, Figure 1F). This index uses known compounds to produce

accurate estimations of where all compounds will elute from the GC column (retention times) thus

allowing precise identification. We validated this index using 60 pure terpene standards (Restek

Corporation, Bellefonte, PA, USA, 34142 and 34143).

Compound Identification: Sample analysis was performed using the Automated Mass Spectral

Deconvolution and Identification System (AMDIS) version 2.73. Compound identification relied on

spectral and RI matching, with a minimum match factor threshold of 70 (Net), a RI window of +/- 10,

and a very strong match factor penalty. Deconvolution settings were component width: 25; adjacent

peak subtraction: two; shape requirements: medium; sensitivity: high; and resolution: medium. The

analysis was restricted to a single identification per library compound.

Post Processing: The relative amount (%) of each compound within a sample was used for all analysis

in this paper. This was calculated as based on the Total Ion Count (TIC) of the sample / peak area of

each compound. No absolute quantification was carried out in this work due to not having

concentration curves for all compounds identified. Best-match compounds for each retention time

and replicate tube were then chosen based on Net match score. Peak areas of any compounds

identified in blank control tubes were removed from the peak areas of all samples within a GC-MS

run. Missing values for compounds (i.e. those compounds found only in some technical replicate

tubes) were zero-filled.

4.3   Statistics and data visualisation for volatile analysis

For all VOC data (Figure 2, Figure 3), TIC (total ion count) data for all compounds was converted to

relative amount (%) using the formula ion count/total ion count of sample.

For the single (November 2024) snapshot, multivariate analysis were performed using all 245

retained compounds following post-processing and filtering. Compound abundance data were

square-root transformed to reduce the influence of highly abundant compounds and to mitigate zero

inflation, following established ecological recommendations (Legendre and Legendre 2012; 2013;

Chang and Lin 2011).

For Principal Component Analysis (PCA, Figure 2A) transformed data were mean-centred and scaled

to unit variance to ensure comparability among compounds with differing absolute abundances. PCA

was conducted using the FactoMineR package with visualisation provided by factoextra in R

(Kassambara and Mundt 2017; Lê, Josse, and Husson 2008). To assess compositional differences in

VOC profiles among sampling locations, a permutational multivariate analysis of variance

(PERMANOVA) was performed using the adonis2 function from the vegan R package, based on

Euclidean distances and 9,999 permutations (Oksanen 2015).

Hierarchical Cluster Analysis (HCA, Figure 2B) was conducted using Ward's minimum variance linkage

method with Euclidean distance, a combination that favours the formation of compact and internally

coherent clusters. Cluster structure identified by HCA was subsequently used to define groups for

downstream exploratory analyses. Partial Least Squares Discriminant Analysis (PLS-DA) was applied

as a supervised exploratory method to characterise compounds contributing most strongly to

differences among the three major clusters identified by HCA. PLS-DA was implemented using the pls

and mixOmics R packages (Rohart et al. 2017). To minimise over-interpretation, PLS-DA results were

not used for hypothesis testing but to identify patterns in variable contributions. The top 10

compounds with the highest Variable Importance in Projection (VIP) scores for each cluster are

reported in Table 1.

To examine differences in VOC composition using ecologically interpretable groupings, compounds

were classified into six functional classes: aliphatics, aromatics, monoterpenes, sesquiterpenes,

terpenoids, and others (including esters and nitrogen- and sulphur-containing compounds) (Figure 2C

and D). For each class, compound richness (number of compounds) and summed relative abundance

were calculated per location. One-way ANOVAs were used to test for differences among locations,

with post-hoc Tukey's HSD tests applied to identify pairwise differences in both summed abundance

(Table S 2 & Table S 3) and compound richness (Table S 4 & Table S 5).

Spatial variation in terpene abundance was visualised using a hierarchical heatmap (Figure 2E) in

which colour intensity represents compound abundance scaled within individual locations.

To contextualise VOC profiles in relation to anthropogenic influence, the relative abundance of all

terpenes and terpenoids was expressed as a ratio to the summed relative abundance of

anthropogenic BTEX compounds (Figure 2F, benzene, toluene, ethylbenzene, and p-xylene;

∑terpenes/∑BTEX), following established approaches for assessing biogenic versus anthropogenic

dominance (Malik et al. 2025).

For analysis of VOC profiles over time in the single sampling site (Figure 3), relative amount data was

used. The average relative amounts of all identified biogenic volatiles were summed within six

compound classes and compared across the 7 time-points.

All statistics were run using R version 4.4.1 (2024-06-14 ucrt) within RStudio 2024.09.0+375

"Cranberry Hibiscus" release for windows. All graphs were generated using the ggplot2 (Wickham

2011) package.

4.4 Statistical modelling of meteorological drivers of bVOC emissions

To investigate environmental drivers of bVOC emissions within a single greenspace (Botanical

Gardens Outdoors) across the calendar year, linear mixed-effects models were fitted to compound

class–level emission data from the Oxford Botanic Garden Outdoor site (BO) using R (v.4.x). Peak

areas were averaged across technical replicates for each sampling date prior to analysis. Compounds

were grouped into five functional classes: monoterpenes, sesquiterpenes, terpenoids, aromatics,

aliphatics, and others (esters, nitrogen-, sulphur- and halogen-containing compounds), and summed

abundances were calculated for each class per sampling date. Response variables were log-

transformed using log(1 + x) to stabilise variance and reduce right skew. Predictor variables included

mean daily temperature (°C), relative humidity (%), daily precipitation (mm), and wind speed (m s⁻¹).

All predictors were centred and scaled (z-score transformation) prior to modelling to facilitate

comparison of effect sizes. For each compound class, a linear mixed-effects model was fitted using

the lme4 package, with sampling date included as a random intercept to account for repeated

measurements over time:

$$log(1 + y_i) = \beta 0 + k \sum \beta k X k, i + b_i + \varepsilon_i$$

where y_i is the summed abundance of a given compound class on sampling date i, X_k are scaled

meteorological predictors, b_i is the random effect of sampling date, and ε_i is the residual error.

Statistical significance of fixed effects was assessed using Wald tests implemented via the lmerTest

package. Model coefficients and standard errors were extracted for visualisation as coefficient plots

and heatmaps, with colour intensity representing the direction and magnitude of standardised

effects. Model structure was defined a priori based on hypothesised meteorological drivers, and

model fit was assessed to ensure consistency across compound classes.

5. Results

5.1 A snapshot in time: distinctive fingerprints of six urban greenspaces

Our analysis pipeline resulted in 245 unique compounds being identified within our 6 Oxford

sampling sites (Table S 1). A PCA (Figure 2A) using all identified compounds as predictor variables

revealed significant site-specific clustering (PERMANOVA, $F_{5,16}$ = 8.889, p = 0.001) with pair-wise

separation of all sites (p < 0.05) except Wytham Woods (WY) in the Warneford Meadow (WA,

PERMANOVA, p = 0.846). Further HCA analysis further revealed site-level structuring of most VOC

profiles (Figure 2B). Primarily, University Parks Open (UO) and Botanical Gardens Outdoors (BO)

separate from all other sites (Figure 2B Cluster i). PLS-DA VIP analysis (Table 1 and Table S7), due to

higher amounts of many anthropogenic emission-related compounds such as 2,3,3-

trimethylpentane, p-xylene, 2-ethylhexan-1-ol, indane, and ethylbenzene. Plant related compounds

(beta-ocimene, dihydromyrcenol, sabinene, diacetone alcohol, beta-pinene, pinocarvone) were

chiefly responsible for the secondary clustering (Figure 2B Cluster ii) due to presence (or higher

proportions) in the Rainforest Glasshouse (RH) compared with University Parks Dense (UD), Wytham

Woods (WY), and Warneford Meadow (WA). The third most significant cluster (Figure 2B Cluster iii)

splits University Parks Open (UO) and Botanical Gardens Outdoors (BO), again due to the higher

proportion/presence of terpenes/terpenoids such as 2-thujene, farnesane, and bornyl acetate found

in Botanical Gardens Outdoors. Full results of PLS-DA analysis can be found in Table S7.

Table 1: Top 10 compounds from PLS-DA VIP analysis (partial least squares discriminant analysis using variable importance in projection) of the first 3 clusters of the HCA (hierarchical cluster analysis) analysis in Figure 2B: Bold compounds highlight terpenes/terpenoids.

Cluster i
P-xylene (Aromatics), Ethylbenzene (Aromatics), Isooctane (Aliphatics), 2,3,4-trimethylpentane (Aliphatics), Toluene (Aromatics), 3-methylhexane (Aliphatics), Methylcyclohexane (Aliphatics), 2-dodecene (Aliphatics), 1-ethyl-2-methylbenzene (Aromatics), 3-methylheptane (Aliphatics)
Cluster ii
Beta-ocimene (Monoterpenes), Dihydromyrcenol (Terpenoids), Sabinene (Monoterpenes), Beta-pinene (Monoterpenes), Pinocarvone (Terpenoids), Hydrindane (Aliphatics), **Alpha-pinene (Monoterpenes), Copaene (Sesquiterpenes), 4-terpineol (Terpenoids), Beta-phellandrene (Monoterpenes**)
Cluster iii
2-thujene (Monoterpenes), Prehnitene (Aromatics), **Farnesane (Sesquiterpenes),** P-isopropenylmethylbenzene (Aromatics), Bromodichloromethane (Others (Esters, N, S, Halo)), 1,4-

diethylbenzene (Aromatics), **Tricyclene (Monoterpenes),** 2,2,4,6,6-pentamethylheptane (Aliphatics), **Bornyl acetate (Terpenoids),** Propylbenzene (Aromatics)

5.1.1 Urban greenspaces show distinctive mix of compound classes

The number of identified terpenoids, monoterpenes and sesquiterpenes was found to be highest in

the Rainforest Glasshouse (13, 15, 10 respectively). These compound numbers were significantly

higher than the other 5 sites (one-way ANOVAs followed by Tukey's HSD post-hoc test p < 0.05,

Figure 2D - full statistics in Table S 4 & Table S 5).Numerically, the Botanical Gardens Outdoors was

the outdoor location with the highest counts of terpenoids, monoterpenes and sesquiterpenes (8,

13, 2 respectively), closely followed by Wytham (9, 12, 1) In terms of relative abundance (Figure 2C -

full statistics in Table S 2 & Table S 3), Botanic Garden Outdoors contained a significantly higher

proportion of aromatic compounds compared to other sites suggesting a strong influence from

anthropogenic sources. The Rainforest Glasshouse showed significantly higher proportion of

monoterpenes, whereas sesquiterpenes were shared by the Rainforest Glasshouse and Botanical

Gardens Outdoors (One-way ANOVAs followed by Tukey's HSD post-hoc test p < 0.05).

5.1.2 Presence/absence of plant-based terpenes and terpenoids

A total of 54 terpenes/terpenoids were identified in this work (Figure 2G). Ten compounds - 1-8

cineole (eucalyptol), 3-carene, alpha-calacorene, alpha-pinene, beta-pinene, camphene, camphor,

limonene, menthol, and p-cymene - were detected at every site, with a further two – menthone and

tricyclene – detected in five of the six sampling sites. Again, the Rainforest Glasshouse exhibited the

largest number of identified terpenes/terpenoid compounds overall (38 compounds) of which 16

were not detected at other sites. Of the outdoor sites, Botanical Gardens Outdoors contained the

highest number of identified terpenes/terpenoid (23), of which 6 were uniquely present at this site.

University Parks Dense (17), University Parks Open (14), Warneford Meadow (14) and Wytham Forest

(22) all showed an overlapping but distinctive terpene mix. A full breakdown of compounds unique to

individual sites is shown in Table S 6.

5.1.3 BTEX compounds as predictors of pollution

The ratio of biogenic terpenes/terpenoids to anthropogenic BTEX compounds was significantly higher

in the Rainforest Glasshouse compared to the other sites (Figure 2E, one-way ANOVA followed by

Tukey's HSD post-hoc test p < 0.001). This is in part due to the increased prevalence of

terpenes/terpenoids and in part to the lower proportion of BTEX compounds (Figure 2F) in this

indoor space. The outdoor sites all had a significantly lower ratios but were not significantly different

from each other.

Figure 2: A: PCA which used all 245 identified compounds as predictor variables revealed significant variation between sites, each showing distinctive site-specific clustering (PERMANOVA, $F_{5,16}$ = 6.557, p = 0.001). B: HCA conducted using Ward's linkage and Euclidean distances using all technical replicate samples over 6 urban greenspaces. C: Relative abundance (means) of detected volatile compounds split by classes across sites, expressed as a percentage of the total sample. D: Total number of detected volatile compounds per site, categorized by compound class. Results show that the number of all compounds groups differed between sites. E: Terpene: BTEX ratio of sampling sites (∑terpenes/∑BTEX). Relative amounts of all terpenes/terpenoids as a ratio to the anthropogenic BTEX compound. F: ∑BTEX as % of all identified compounds between sampling sites. G: Terpenes/terpenoid compounds identified across sampling locations. Heatmap showing the scaled relative amount (light green > dark green) or absence (grey) of any terpenes/terpenoid VOCs detected across six sampling locations. Scaling was carried out within single locations. C, D, F: Bars represent the mean (n = 4 technical replicate sampling tubes per site). Letters indicate statistically distinct sampling locations within chemical classes based on one-way ANOVA followed by Tukey's HSD post-hoc test (p < 0.05).

5.2   How do urban greenspace bVOC profiles vary through time under real-world conditions?

Using the same classification system for our identified compounds we counted both compound numbers and summed abundances to visualise each class over time. In terms of the relative abundances, all compound classes peaked during late spring to summer (May to August 2025) with two notable outliers - monoterpenes had an early peak on November 27th, 2024, and terpenoids had a late peak on October 29th, 2025. Interestingly, in terms of the absolute number of compounds identified, it was early spring (04 March 2025) which showed peaks for aliphatics (42), aromatics (14), terpenoids (10) and others (9), with monoterpene counts being more stable over the year.

Mixed-effects modelling revealed that different components of the biogenic scentscape responded in contrasting ways to short-term environmental variation (Figure 3C). Temperature emerged as the most consistent positive driver of terpene emissions, with higher temperatures associated with increased relative abundances of both monoterpenes and sesquiterpenes. Humidity also showed compound-class-specific effects with terpenes exhibiting a significant positive association with increased levels of relative humidity, whereas aromatic and aliphatic compounds were weaker or neutral. Increased wind speed contributed to increases in both aromatic and aliphatic fractions.

Figure 3: Seasonal dynamics and environmental drivers of biogenic volatile organic compounds (bVOCs) at the Botanical Garden outdoor site. (A) Daily temperature (orange) and relative humidity (green), and daily precipitation (bars) with wind speed (line), derived from local and regional meteorological datasets. Vertical dashed lines indicate sampling dates; values above denote site-specific temperature and relative humidity measured during sampling. (B-G) shows monthly mean relative abundances (± SE) of compound classes (aliphatics, aromatics, monoterpenes, sesquiterpenes, terpenoids, and "others" including esters and N-, S- and halogen-containing compounds) from November 2024 to September 2025. Numbers beside points indicate the number of detected compounds contributing to each class per sampling event. (H) Mixed-effects model coefficients linking standardised environmental predictors (temperature, precipitation, humidity and wind speed) to log-transformed relative abundances of each compound class. Tile colour represents effect size (positive = green, negative = brown), and symbols denote statistical significance (* p < 0.05; ** p < 0.01; *** p < 0.001). Models were fitted with sampling date as a random effect to account for temporal structure.

6. Discussion

6.1   bVOCs and aVOCs shape urban greenspace scentscapes

An initial PCA of our six sampling sites on a single date (Nov 27th, 2024) revealed significant site-specific clustering. Further HCA revealed site structuring was due to both bVOCs and aVOCs (Table 1, Figure 2B). We chose to interrogate the first 3 clusters which appeared. The first cluster separated two outdoor sites - Botanical Gardens Outdoors and University Parks Open – from all other sampling sites. PLS-DA VIP analysis showed that this was due to many emission-related aVOCs (Table 1) including the traffic-related BTEX compounds (Figure 2F). These two sites are very close to a busy city centre roads (Figure 1A) suggesting that traffic is the likely main contributor. This is supported by BTEX presence measurements where over 40% of identified volatiles were BTEX compounds in the Botanical Gardens Outdoors and University Parks Open, but just 13% in Wytham Woods and Warneford Meadow which are more isolated from traffic. The relationship between proximity to traffic and BTEX prevalence has not been assessed in this paper but has been previously reported (Hoque et al. 2008).

Interestingly, despite its proximity to University Parks Open (~80 m), University Parks Dense showed markedly lower BTEX levels (43% to 29%). Localised differences in BTEX presence within urban greenspaces has been demonstrated before; for example, Upmanis et al. (2001) reported that concentrations of benzene and toluene were up to one-third lower just 40m into urban parkland within two large Scandinavian cities. A second study also showed decreases in particulate matter (PM_{10} and $PM_{2.5}$) of up to 50% at 200m (Gómez-Moreno et al. 2019). Additionally, the presence of exposed soil and leaf litter layer around trees in University Parks Dense compared to the grassy University Parks Open site may contribute to increased degradation of BTEX compounds through microbial metabolism. A 2012 study, for example, showed that by adding a mixed bacterial consortium to soil, BTEX compounds were more rapidly degraded (Mukherjee and Bordoloi 2012).

Biogenic volatiles, particularly terpenes (Table 1, Figure 2G), were chiefly responsible for the secondary clustering of samples from Rainforest House, and those of Wytham Woods and Warneford Meadow. In total across all sampling sites, we identified 54 unique terpenes/terpenoids. The

Rainforest Glasshouse contained the highest number of terpenes (38) with considerable spread

across the outdoor sampling sites (14 to 23). The presence of these compounds is highly suggestive

of multiple and specific health benefits within each location. For example, 10 terpenes/terpenoids

were ubiquitous across all sampling sites - compounds which have also been found in air samples

elsewhere using similar methods (Gu et al. 2024; Bryant et al. 2022; Walker et al. 2023; Hellén et al.

2024; Jones, Hopkins, and Lewis 2011; Hakola et al. 2003; Detournay et al. 2011). These compounds

alone are cited to have anti-inflammatory, anxiolytic, antimicrobial, immune-boosting,

antidepressant, anaesthetic, gastroprotective, antioxidant, anti-cancer (*in vitro*), immune-

modulatory, sedative, hypolipidemic, analgesic, decongestant, bronchodilatory, mucolytic, and

cognitive enhancing effects (Lee et al. 2022; Vallianou and Hadzopoulou-Cladaras 2016; Salehi et al.

2019; Oliveira et al. 2024; Sun 2007; Woo et al. 2019; Jo et al. 2021). It could be argued, therefore,

that spending time in any of these sites could impart significant health gains. However, the

relationships between these compounds and their benefits are likely concentration dependent,

therefore further methodological work is needed to quantify and make precise recommendations

such as when, and for how long, to visit specific greenspaces. Previously it was found that just 20

minutes is adequate to derive physiological and psychological benefits from time in woodland

(Haluza et al. 2025; Park et al. 2010).

Another useful metric to separate our sites is a ratio of biogenic to anthropogenic compounds. An

example of this can be found in study which assessed air over Covid lockdown in 2020 at an urban

site in India (Malik et al. 2025) in which a sharp increase in the $\sum\alpha$+beta--pinene/$\sum$BTEX ratios during

the strict lockdown phase was hypothesised to be due to a reduction in traffic related activity. We

mirrored this study and found that, on a particular day, the Rainforest House had significantly higher

terpene/BTEX ratio than all the other sites (Figure 2F). This result is likely due to the interplay of two

different factors known to affect the volatile makeup of indoor spaces. On the one hand, this

glasshouse is plant rich and enclosed and as such likely acts as an enclosure for those bVOCs

produced within, and conversely, aVOC ingression may be limited. These results suggest that plant

rich greenhouses may be the most beneficial place to spend time, however, in a 2006-2007 study,

aVOCs were found to be significantly higher in indoor sites than outdoor sites in the vicinity of an

industrial park (Chang et al. 2019), showing that simply being inside is little protection from external

sources of pollution.

6.2   The impacts of seasonality on bVOC emissions in the Botanical Gardens

As well as looking at a snapshot of the scentscapes across the six urban greenspaces, we also

explored how bVOC change across time and in the face of environmental variation in the Botanical

Outdoors location. Using the same classification system for our identified compounds we counted

both compound numbers and summed abundances to visualise each class over time. In terms of the

relative abundances, all compound classes peaked during late spring to summer (May to August

2025) with two notable outliers - monoterpenes had an early peak on November 27th, 2024, and

terpenoids had a late peak on October 29th, 2025. Interestingly, in terms of the absolute number of

compounds identified, it was early spring (March 4th) which showed peaks for aliphatics (42),

aromatics (14), terpenoids (10) and others (9), with monoterpene counts being more stable over the

year.

Higher temperature and relative humidity were the most consistent positive predictors of biogenic

emissions, particularly for terpenes. For temperature this is likely linked to the well-established

positive relationship between emission rates and leaf temperatures via both biosynthesis through

increased enzymatic activity and volatilisation vapour pressure (Guenther et al. 1993; Luo et al.

2025). Aliphatics also showed a positive temperature response, likely driven by enhanced

volatilisation at higher temperatures but also resistance to biodegradation (Abbasian et al. 2015). In

contrast, terpenoids, which are oxygenated terpenes and much more reactive, declined with

increasing temperature. This is consistent with their greater susceptibility to photochemical

degradation at higher temperatures (Atkinson and Arey 2003; Hallquist et al. 2009). The role of

humidity in terpene/terpenoid presence may be linked to an increase in stomatal opening and

subsequent environmental release when relative humidity is higher (Arve and Torre 2015).

Precipitation effects also differed among VOC classes. Terpene abundances showed a positive

association with precipitation. Wetting is known to give a short-term pulse release through

mechanical displacement from soil air spaces after rain (Pugliese et al. 2023) and this was likely the

reason for the high amount recorded on November 27[th], 2024, where 8.10 mm of rain was recorded.

In contrast, aromatics and aliphatics showed a negative association with precipitation on the day of

sampling, consistent with atmospheric dilution, wet deposition, or reduced resuspension of

background hydrocarbons rather than changes in emission strength (Casas et al. 2021). There was a

second rainfall event on September 4[th], 2025 (9.4 mm), however, this followed 16.3 mm of rain on

the previous day which is likely to have negatively affected soil-based terpene release on the day of

sampling. Notably, only two sampling events coincided with measurable rainfall so these patterns

need further exploration in urban greenspaces.

Wind speed effects showed significant positive associations for aromatic and aliphatic compounds.

Given their relatively long atmospheric lifetimes compared to terpenes and terpenoids (Luo et al.

2025; Hellén et al. 2018), these increases are more likely to reflect regional background

concentrations than increases in local emissions. In contrast, terpene emissions showed little

sensitivity to wind speed, consistent with their predominantly local, vegetation-derived sources and

rapid atmospheric reactivity leading to degradation if produced distantly.

Overall, these results indicate that temperature, precipitation, wind speed and humidity all impact on

the detection of different bVOC chemical classes. The absence of a single dominant predictor across

all VOC classes highlights the importance of considering scentscapes as multidimensional chemical

systems, shaped by interacting biological and meteorological processes. However, this class-specific

sensitivity to environmental drivers adds understanding of when and where urban greenspaces may

be most chemically beneficial for human exposure. These results are suggestive that warm, humid,

and (unfortunately) rainy conditions increase the levels of health-promoting terpene/terpenoid

compounds in the air and as such are likely the most beneficial conditions for humans to spend time

in urban greenspaces.

6.3   Methodological advancements, limitations, and future work

As well as providing an understanding of urban greenspace scentscapes over time, this study aimed

to develop a robust and repeatable protocol for measuring the presence and relative amounts of a

wide range of volatiles of interest. Although similar methods have been developed for headspace

sampling (Yin et al. 2022; de Carvalho Couto et al. 2024; Reyrolle et al. 2024), air sampling studies

have historically employed targeted approaches that significantly limit the number of compounds

identified. For instance, previous studies have used between 15 and 66 VOCs across various forest

and urban environments (Walker et al. 2023; Pripdeevech et al. 2025; Mula et al. 2024). In contrast,

our study adopted an untargeted metabolomics-style approach commonly used in biological matrices

to enhance the depth and resolution of VOC profiling (Fernie et al. 2011; Van den Berg et al. 2006;

Rosenthal et al. 2024) to identify 245 unique compounds, of which 54 were unique

terpenes/terpenoids. This is significantly higher than any studies on ambient air so far published

(Walker et al. 2023; Sanaei et al. 2023; Mula et al. 2024). This improved identification capacity is

likely attributable to several key enhancements in our analytical pipeline: greater technical

replication, the use of a Kováts retention index, a more focused compound library, a slow GC

temperature ramp, and the use of deconvolution for resolving co-eluting peaks. Furthermore, the

tight clustering of technical replicates observed in the PCA (Figure 2A) confirms that our collection

methods and downstream processing are sufficiently controlled to ensure analytical reproducibility

and minimise technical variability.

Despite these improvements, our analyses capture only a fraction of the chemical complexity

present. In most samples we detect far more chromatographic features than we can confidently

identify, and the ~245 compounds reported here represent ~10% (or less) of the total compounds

present. This limitation is expected in untargeted ambient-air GC–MS, where a large proportion of

peaks correspond to (i) compounds absent from reference libraries, (ii) compounds lacking retention

index (RI) values on comparable stationary phases, (iii) low-abundance or highly reactive species near

the detection limit, and (iv) co-eluting mixtures that remain partially unresolved even after

deconvolution. As a result, our interpretation focuses on the "identifiable fraction" of the scentscape

rather than the full atmospheric mixture. One major benefit of TD work is the ability for the

'recollection' of the split portion of the GC-MS analysis. Future work intends to use this feature to

analyse the same samples on multiple GC column types (polar/non-polar) – this will help to resolve

significantly more atmospheric compounds.

Identification confidence is a further constraint. We used a minimum Net match threshold of 70,

which is permissive and may increase the risk of imperfect annotations, particularly for isomeric

compounds and low-intensity peaks. Although 99 compounds had a match factor > 90, the mean

match factor for retained identifications was 86. Therefore, a subset of compounds should be

regarded as putative with future work concentrating on targeted confirmation of key marker

compounds using standards. Our identifications are strengthened by combining spectral matching

with RI matching, however, expanding the RI coverage of our library on DB-5–type phases and

continuing to develop an in-house "Oxford greenspace" standard set would improve annotation

rates. Additionally, we did not employ chiral separation in this work, meaning stereoisomers of

compounds such as Alpha-pinene (+/-) were not resolved. These may be important for understanding

biological source and ecological function and as such could be included in future. Finally, to truly

overcome co-elution, two-dimensional gas chromatography (GC x GC) could be used to separate

overlapping peaks without the need for software-based deconvolution.

Our sampling design also imposes important constraints. First, sampling was conducted within a

narrow daytime window (11:00–13:00). As our models show, many bVOCs are affected by

environmental changes, therefore, abundance and composition likely differ through the day. Second,

we used a fixed sampling duration (2 h) and flow rate (100 ml/min). This equates to just 2 minutes of

human breathing. Longer sampling times or higher sampled volumes would likely increase detection

of low-abundance compounds, but at the risk of sorbent saturation, breakthrough of other volatile

species, and increased influence meteorology during the sampling window. Third, while technical

replication was employed, biological/environmental replication was limited: the "snapshot"

comparison is based on a single calendar date, and the year-long analysis focuses on one site. In both

cases, sampling was carried out at a precise location within each site. As such, our site-level

inferences should not be interpreted as definitive rankings of whole sites across seasons and years.

Quantification is another limitation. Relative measures are useful for compositional comparisons, but

absolute quantification will be essential to translate scentscape metrics into health-relevant

exposure estimates. The quantification of compounds such as terpenes will help to support policy-

facing recommendations.

Finally, the environmental predictors included here (temperature, humidity, precipitation, wind

speed) are only a subset of the drivers that shape ambient VOC fingerprints. A more holistic

predictive framework will need to incorporate additional covariates such as (i) local vegetation

composition and phenology (biodiversity and functional traits), (ii) microclimate at the sampling

point (solar radiation, canopy cover, boundary-layer stability, VPD), (iii) soil moisture and microbial

activity, (iv) episodic disturbances (mowing, leaf litter pulses, pest/pathogen outbreaks), and (v)

urban pollution context (traffic intensity, background NO_x/O_3, and atmospheric oxidation capacity).

Incorporating these variables—alongside expanded temporal coverage, multi-year replication, and

partial targeted quantification—will be necessary to move from descriptive classification of

scentscapes to robust prediction of when and where greenspaces maximise health-relevant

exposures.

7.   Author Contributions

William Kay was responsible for methods development, data collection and all data analyses. William

Kay and Anya Lindstrom Battle were both involved in the writing and editing of the manuscript. Kathy

Willis conceived of the initial ideas and methodologies and was also involved in the writing and

editing of the manuscript. Mahal Humberstone, Molly Tucker and Kieran Storer all contributed to in-

field sample collection. Geoffrey Kite was involved in the GC-MS data collection, as well as providing

useful guidance. All authors contributed critically to the drafts and gave final approval for

publication.

8.   Funding

507   This work was funded by the Leverhulme Centre for Nature Recovery, made possible thanks to the

508   generous support of the Leverhulme Trust - Grant number RC-2021-76.

9.   Data Availability Statement

510   Raw AGILENT files are available here: http://doi.org/10.5281/zenodo.18493789

10. Acknowledgments

We would firstly like to thank the University of Oxford Botanic Garden and Wytham Woods for

permission to sample on their sites. We would also like to thank Dr Ben Martin who provided helpful

advice for sampling and analysis as well as the development of early sampling methodologies, and Dr

Luciana Carvalho who always had time to share thoughts and ideas relating to the project. In

addition, we thank Dr Pedro Bota for his generosity of time, availability and equipment. Finally, we

thank all members of the Oxford long-term ecology lab for their ongoing feedback.

11. Conflicts of Interest

The authors declare no conflicts of interest.

12. References

Abbasian, F., Lockington, R., Mallavarapu, M., and Naidu, R. (2015). A Comprehensive Review of Aliphatic Hydrocarbon Biodegradation by Bacteria, *Applied Biochemistry and Biotechnology*, 176: 670–99. https://doi.org/10.1007/s12010-015-1603-5

Arve, L. E., and Torre, S. (2015). Ethylene is involved in high air humidity promoted stomatal opening of tomato (Lycopersicon esculentum) leaves, *Functional Plant Biology*, 42: 376–86. https://doi.org/10.1071/fp14247

Atkinson, R., and Arey, J. (2003). Atmospheric Degradation of Volatile Organic Compounds, *Chemical Reviews*, 103: 4605–38. https://doi.org/10.1021/cr0206420

Bratman, G. N., Daily, G. C., Levy, B. J., and Gross, J. J. (2015). The benefits of nature experience: Improved affect and cognition, *Landscape and urban planning*, 138: 41–50. https://doi.org/10.1016/j.landurbplan.2015.02.005

Bryant, D. J., Nelson, B. S., Swift, S. J., Budisulistiorini, S. H., Drysdale, W. S., Vaughan, A. R., Newland, M. J., Hopkins, J. R., Cash, J. M., and Langford, B. (2022). Biogenic and anthropogenic sources of isoprene and monoterpenes and their secondary organic aerosol in Delhi, India, *Atmospheric Chemistry and Physics Discussions*, 2022: 1–35. https://doi.org/10.1016/j.apr.2019.07.004

Casas, G., Martinez-Varela, A., Vila-Costa, M., Jiménez, B., and Dachs, J. (2021). Rain Amplification of Persistent Organic Pollutants, *Environmental Science & Technology*, 55: 12961–72. https://doi.org/10.1021/acs.est.1c03295

Chang, C.-C., and Lin, C.-J. (2011). LIBSVM: a library for support vector machines, *ACM transactions on intelligent systems and technology (TIST)*, 2: 1–27. https://doi.org/10.1145/1961189.1961199

Chang, T.-Y., Liu, C.-L., Huang, K.-H., and Kuo, H.-W. (2019). Indoor and outdoor exposure to volatile organic compounds and health risk assessment in residents living near an optoelectronics industrial park, *Atmosphere*, 10: 380. https://doi.org/10.3390/atmos10070380

de Carvalho Couto, C., Chávez, D. W. H., Oliveira, E. M. M., Freitas-Silva, O., and Casal, S. (2024). SPME-GC-MS untargeted metabolomics approach to identify potential volatile compounds as markers for fraud detection in roasted and ground coffee, *Food Chemistry*, 446: 138862. https://doi.org/10.1016/j.foodchem.2024.138862

Degtyarenko, K., De Matos, P., Ennis, M., Hastings, J., Zbinden, M., McNaught, A., Alcántara, R.,
Darsow, M., Guedj, M., and Ashburner, M. (2007). ChEBI: a database and ontology for
chemical entities of biological interest, *Nucleic acids research*, 36: D344–D50.
https://doi.org/10.1093/nar/gkm791
Detournay, A., Sauvage, S., Locoge, N., Gaudion, V., Leonardis, T., Fronval, I., Kaluzny, P., and
Galloo, J.-C. (2011). Development of a sampling method for the simultaneous monitoring of
straight-chain alkanes, straight-chain saturated carbonyl compounds and monoterpenes in
remote areas, *Journal of Environmental Monitoring*, 13: 983–90.
https://doi.org/10.1039/c0em00354a
Fernie, A. R., Aharoni, A., Willmitzer, L., Stitt, M., Tohge, T., Kopka, J., Carroll, A. J., Saito, K., Fraser,
P. D., and DeLuca, V. (2011). Recommendations for reporting metabolite data, *The Plant Cell*,
23: 2477–82. https://doi.org/10.1105/tpc.111.086272
Gao, X., Wang, Y., Wu, L., Zheng, F., Sun, N., Liu, G., Liu, Y., Meng, P., Sun, L., and Jing, B. (2023).
The Impact of Anthropogenic VOC Emissions on Atmospheric Pollution: A Case Study of a
Typical Industrialized Area in China, *Atmosphere*, 14: 1586.
https://doi.org/10.3390/atmos14101586
Gómez-Moreno, F. J., Artíñano, B., Ramiro, E. D., Barreiro, M., Núñez, L., Coz, E., Dimitroulopoulou,
C., Vardoulakis, S., Yagüe, C., Maqueda, G., Sastre, M., Román-Cascón, C., Santamaría, J.
M., and Borge, R. (2019). Urban vegetation and particle air pollution: Experimental campaigns
in a traffic hotspot, *Environmental Pollution*, 247: 195–205.
https://doi.org/https://doi.org/10.1016/j.envpol.2019.01.016
GOV.UK. 2023. "Road traffic statistics - Local authority Oxfordshire." In, edited by Department for
Transport. roadtraffic.dft.gov.uk: GOV.UK
Gu, S., Luo, W., Charmchi, A., McWhirter, K. J., Rosenstiel, T., Pankow, J., and Faiola, C. L. (2024).
Limonene Enantiomeric Ratios from Anthropogenic and Biogenic Emission Sources, *Environ*
Sci Technol Lett, 11: 130–35. https://doi.org/10.1021/acs.estlett.3c00794
Guenther, A., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T. a., Emmons, L., and Wang, X.
(2012). The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.
1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model*
Development, 5: 1471–92. https://doi.org/10.5194/gmd-5-1471-2012
Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R. (1993). Isoprene and
monoterpene emission rate variability: model evaluations and sensitivity analyses, *Journal of*
Geophysical Research: Atmospheres, 98: 12609–17. https://doi.org/10.1029/93JD00527
Hakola, H., Tarvainen, V., Laurila, T., Hiltunen, V., Hellén, H., and Keronen, P. (2003). Seasonal
variation of VOC concentrations above a boreal coniferous forest, *Atmospheric Environment*,
37: 1623–34. https://doi.org/10.1016/S1352-2310(03)00014-1
Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J.,
Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T.,
Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W.,
McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D.,
Szmigielski, R., and Wildt, J. (2009). The formation, properties and impact of secondary
organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9: 5155–236.
https://doi.org/10.5194/acp-9-5155-2009
Haluza, D., Kersten, P., Lazic, T., Steinparzer, M., and Godbold, D. (2025). Unlocking the Power of
Nature: Insights from a 20-Minute Forest Visit on Well-Being, *Forests*, 16: 792
Hellén, H., Praplan, A. P., Tykkä, T., Ylivinkka, I., Vakkari, V., Bäck, J., Petäjä, T., Kulmala, M., and
Hakola, H. (2018). Long-term measurements of volatile organic compounds highlight the
importance of sesquiterpenes for the atmospheric chemistry of a boreal forest, *Atmospheric*
Chemistry and Physics, 18: 13839–63
Hellén, H., Tykkä, T., Schallhart, S., Stratigou, E., Salameh, T., and Iturrate-Garcia, M. (2024).
Measurements of atmospheric C 10–C 15 biogenic volatile organic compounds (BVOCs) with
sorbent tubes, *Atmospheric Measurement Techniques*, 17: 315–33.
https://doi.org/10.5194/amt-17-315-2024
Hoque, R. R., Khillare, P., Agarwal, T., Shridhar, V., and Balachandran, S. (2008). Spatial and
temporal variation of BTEX in the urban atmosphere of Delhi, India, *Science of the total*
environment, 392: 30–40. https://doi.org/10.1016/j.scitotenv.2007.08.036
Jo, H., Cha, B., Kim, H., Brito, S., Kwak, B. M., Kim, S. T., Bin, B.-H., and Lee, M.-G. (2021). α-Pinene
enhances the anticancer activity of natural killer cells via ERK/AKT pathway, *International*
Journal of Molecular Sciences, 22: 656. https://doi.org/10.3390/ijms22020656
Jones, C. E., Hopkins, J., and Lewis, A. (2011). In situ measurements of isoprene and monoterpenes
within a south-east Asian tropical rainforest, *Atmospheric Chemistry and Physics*, 11: 6971–
84. https://doi.org/10.5194/acp-11-6971-2011
Joo, Y., Schuman, M. C., Goldberg, J. K., Kim, S. G., Yon, F., Brütting, C., and Baldwin, I. T. (2018).
Herbivore-induced volatile blends with both "fast" and "slow" components provide robust

indirect defence in nature, *Functional Ecology*, 32: 136–49. https://doi.org/10.1111/1365-2435.12947

Kassambara, A., and Mundt, F. 2017. "Package 'factoextra'." In *Extract and visualize the results of multivariate data analyses*, 10.18637.

Kovats, v. E. (1958). Gas-chromatographische charakterisierung organischer verbindungen. Teil 1: retentionsindices aliphatischer halogenide, alkohole, aldehyde und ketone, *Helvetica Chimica Acta*, 41: 1915–32. https://doi.org/10.1002/hlca.19580410703

Lê, S., Josse, J., and Husson, F. (2008). FactoMineR: an R package for multivariate analysis, *Journal of statistical software*, 25: 1–18. https://doi.org/10.18637/jss.v025.i01

Lee, S. H., Kim, D. S., Park, S. H., and Park, H. (2022). Phytochemistry and Applications of Cinnamomum camphora Essential Oils, *Molecules*, 27. https://doi.org/10.3390/molecules27092695

Legendre, P., and Legendre, L. 2012. *Numerical ecology* (Elsevier).

Li, Q. (2010). Effect of forest bathing trips on human immune function, *Environmental health and preventive medicine*, 15: 9–17. https://doi.org/10.1007/s12199-008-0068-3

Li, T., Baggesen, N., Seco, R., and Rinnan, R. (2023). Seasonal and diel patterns of biogenic volatile organic compound fluxes in a subarctic tundra, *Atmospheric Environment*, 292: 119430. https://doi.org/10.1016/j.atmosenv.2022.119430

Luo, R., Lun, X., Gao, R., Wang, L., Yang, Y., Su, X., Habibullah-Al-Mamun, M., Xu, X., Li, H., and Li, J. (2025). A Review of Biogenic Volatile Organic Compounds from Plants: Research Progress and Future Prospects, *Toxics*, 13: 364

Malik, T. G., Gupta, M., Tripathi, N., and Sahu, L. K. (2025). Change in monoterpene concentrations during winter-to-summer transition period and impact of COVID-19 lockdown at an urban site in India, *Atmospheric Environment*, 350: 121141. https://doi.org/10.1016/j.atmosenv.2025.121141

Mu, Z., Asensio, D., Llusia, J., Filella, I., Ogaya, R., Yi, Z., and Penuelas, J. (2022). Annual and seasonal variations in soil volatile organic compound concentrations in a Mediterranean shrubland and holm oak forest, *Geoderma*, 405: 115401. https://doi.org/10.1016/j.geoderma.2021.115401

Mukherjee, A. K., and Bordoloi, N. K. (2012). Biodegradation of benzene, toluene, and xylene (BTX) in liquid culture and in soil by Bacillus subtilis and Pseudomonas aeruginosa strains and a formulated bacterial consortium, *Environmental Science and Pollution Research*, 19: 3380–88. https://doi.org/10.1007/s00449-015-1364-0

Mula, V., Bogdanov, J., Stanoeva, J. P., Zeneli, L., and Zdravkovski, Z. (2024). Monitoring Volatile Organic Compounds in Air Using Passive Sampling: Regional Cross-Border Study between N. Macedonia and Kosovo, *Aerosol and Air Quality Research*, 24: 230170. https://doi.org/10.4209/aaqr.230170

Müller, A., Faubert, P., Hagen, M., Zu Castell, W., Polle, A., Schnitzler, J.-P., and Rosenkranz, M. (2013). Volatile profiles of fungi–chemotyping of species and ecological functions, *Fungal genetics and biology*, 54: 25–33. https://doi.org/10.1016/j.fgb.2013.02.005

Oksanen, J. 2015. "Vegan: community ecology package." In *R package version*, 3.

Oliveira, C. D., Cardoso, M. d. G., Batista, L. R., Alves, E., Rosa, M. B. P., Ferreira, V. R. F., de Souza, L., Pineda, M., Fernandes, A. I., and Nelson, D. L. (2024). The antibacterial, antioxidant, and insecticidal activities of essential oils from Thymus vulgaris L., Salvia officinalis L., and Ocimum basilicum L, *Journal of Food Safety*, 44: e13145. https://doi.org/10.1111/jfs.13145

Park, B. J., Tsunetsugu, Y., Kasetani, T., Kagawa, T., and Miyazaki, Y. (2010). The physiological effects of Shinrin-yoku (taking in the forest atmosphere or forest bathing): evidence from field experiments in 24 forests across Japan, *Environmental health and preventive medicine*, 15: 18–26. https://doi.org/10.1007/s12199-009-0086-9

Pripdeevech, P., Janta, R., Sripahco, T., Meesang, W., Aiyathiti, C., Prabamroong, T., Mahatheeranont, S., Poshyachinda, S., Pongpiachan, S., and Khruengsai, S. (2025). Seasonal volatile organic compound dynamics in urban and forest environments in Thailand: Implications for air quality and secondary pollutants, *Environmental Pollution*, 367: 125565. https://doi.org/10.1016/j.envpol.2024.125565

Pugliese, G., Ingrisch, J., Meredith, L. K., Pfannerstill, E. Y., Klüpfel, T., Meeran, K., Byron, J., Purser, G., Gil-Loaiza, J., and Van Haren, J. (2023). Effects of drought and recovery on soil volatile organic compound fluxes in an experimental rainforest, *nature communications*, 14: 5064. https://doi.org/10.1038/s41467-023-40661-8

Reyrolle, M., Desauziers, V., Pigot, T., Gautier, L., and Le Bechec, M. (2024). Comparison of Untargeted and Markers Analysis of Volatile Organic Compounds with SIFT-MS and SPME-GC-MS to Assess Tea Traceability, *Foods*, 13: 3996. https://doi.org/10.3390/foods13243996

Rohart, F., Gautier, B., Singh, A., and Lê Cao, K.-A. (2017). mixOmics: An R package for 'omics feature selection and multiple data integration, *PLoS computational biology*, 13: e1005752. https://doi.org/10.1371/journal.pcbi.1005752

Rosenthal, K., Lindley, M., Turner, M., Ratcliffe, E., and Hunsicker, E. (2024). Current data processing methods and reporting standards for untargeted analysis of volatile organic compounds using direct mass spectrometry: a systematic review, *Metabolomics*, 20: 42. https://doi.org/10.1007/s11306-024-02104-3

Salehi, B., Upadhyay, S., Erdogan Orhan, I., Kumar Jugran, A., LD Jayaweera, S., A. Dias, D., Sharopov, F., Taheri, Y., Martins, N., and Baghalpour, N. (2019). Therapeutic potential of α- and β-pinene: A miracle gift of nature, *Biomolecules*, 9: 738. https://doi.org/10.3390/biom9110738

Sanaei, A., Herrmann, H., Alshaabi, L., Beck, J., Ferlian, O., Fomba, K. W., Haferkorn, S., van Pinxteren, M., Quaas, J., and Quosh, J. (2023). Changes in biodiversity impact atmospheric chemistry and climate through plant volatiles and particles, *Communications Earth & Environment*, 4: 445. https://doi.org/10.1038/s43247-023-01113-9

Sun, J. 2007. "D-Limonene: safety and clinical applications." In *Alternative medicine review*, 259.

Upmanis, H., Eliasson, I., and Andersson-Sköld, Y. (2001). Case studies of the spatial variation of benzene and toluene concentrations in parks and adjacent built-up areas, *Water, air, and soil pollution*, 129: 61–81. https://doi.org/10.1023/A:1010357914047

Vallianou, I., and Hadzopoulou-Cladaras, M. (2016). Camphene, a Plant Derived Monoterpene, Exerts Its Hypolipidemic Action by Affecting SREBP-1 and MTP Expression, *PloS one*, 11: e0147117. https://doi.org/10.1371/journal.pone.0147117

Van den Berg, R. A., Hoefsloot, H. C., Westerhuis, J. A., Smilde, A. K., and Van der Werf, M. J. (2006). Centering, scaling, and transformations: improving the biological information content of metabolomics data, *BMC genomics*, 7: 1–15. https://doi.org/10.1186/1471-2164-7-142

Walker, H., Jena, A., McEwan, K., Evans, G., and Campbell, S. (2023). Natural volatile organic compounds (NVOCs) are greater and more diverse in UK forests compared with a public garden, *Forests*, 14: 92. https://doi.org/10.3390/f14010092

Wang, L., Lun, X., Wang, Q., and Wu, J. (2024). Biogenic volatile organic compounds emissions, atmospheric chemistry, and environmental implications: a review, *Environmental Chemistry Letters*, 22: 3033–58. https://doi.org/10.1007/s10311-024-01785-5

Weisskopf, L., Schulz, S., and Garbeva, P. (2021). Microbial volatile organic compounds in intra-kingdom and inter-kingdom interactions, *Nature Reviews Microbiology*, 19: 391–404. https://doi.org/10.1038/s41579-020-00508-1

Wickham, H. (2011). ggplot2, *Wiley interdisciplinary reviews: computational statistics*, 3: 180–85. https://doi.org/10.1002/wics.147

Woo, J., Yang, H., Yoon, M., Gadhe, C. G., Pae, A. N., Cho, S., and Lee, C. J. (2019). 3-Carene, a Phytoncide from Pine Tree Has a Sleep-enhancing Effect by Targeting the GABA_A-benzodiazepine Receptors, *Experimental neurobiology*, 28: 593–601. https://doi.org/10.5607/en.2019.28.5.593

Yang, K., Llusià, J., Preece, C., Ogaya, R., Tur, L. M., Mu, Z., You, C., Xu, Z., Tan, Y., and Peñuelas, J. (2024). Impacts of seasonality, drought, nitrogen fertilization, and litter on soil fluxes of biogenic volatile organic compounds in a Mediterranean forest, *Science of the total environment*, 906: 167354. https://doi.org/10.1016/j.scitotenv.2023.167354

Yin, X.-L., Fu, W.-J., Chen, Y., Zhou, R.-F., Sun, W., Ding, B., Peng, X.-T., and Gu, H.-W. (2022). GC-MS-based untargeted metabolomics reveals the key volatile organic compounds for discriminating grades of Yichang big-leaf green tea, *LWT*, 171: 114148. https://doi.org/10.1016/j.lwt.2022.114148

Zeng, W., Liu, H., Hou, S., Qiu, X., Chen, X., Liu, M., Wu, D., and Liu, L. (2022). Explore the Benefits of Natural Air: New Insights from Field and Chamber Tests on Cognitive Performance, *Atmosphere*, 13: 1006. https://doi.org/10.3390/atmos13071006

13.   Supplementary Information

Table S 1:

A full list of the compounds detected on 27ᵗʰ November 2024. List is sorted by retention time (RT). RI is retention index values corresponding to Kovats index. MFᵗᵒᵖ is the highest match factor found over all samples. Green rows show compounds for which we have confirmed RT via standard solutions.

CAS	Name	RT	RI	MFᵗᵒᵖ
107-06-2	1,2-dichloroethane	4.61	649.6	92
590-86-3	3-methylbutanal	4.63	651	79
563-80-4	3-methylbutan-2-one	4.65	652.5	87
71-43-2	Benzene	4.74	658.9	98
56-23-5	Tetrachloromethane	4.77	661.5	96
71-36-3	Butan-1-ol	4.79	663.2	77
110-82-7	Cyclohexane	4.80	663.6	96
565-59-3	2,3-dimethylpentane	4.80	663.7	70
589-34-4	3-methylhexane	4.87	669.1	96
110-83-8	Cyclohexene	5.03	680.8	78
123-75-1	Pyrrolidine	5.07	683.4	81
617-78-7	3-ethylpentane	5.07	683.6	91
107-87-9	Pentan-2-one	5.08	684.4	91
540-84-1	Isooctane	5.10	685.9	97
96-22-0	Pentan-3-one	5.23	695.9	78
142-82-5	Heptane	5.29	699.8	99
10152-76-8	Sulfide, allyl methyl	5.31	700.8	78
79-01-6	Trichloroethene	5.38	703.4	76
110-62-3	Pentanal	5.48	707.5	77
80-62-6	Methyl methacrylate	5.51	708.4	88
109-60-4	Propyl acetate	5.56	710.6	89
625-86-5	2,5-dimethylfuran	5.59	711.5	80
3208-16-0	2-ethylfuran	5.60	712.2	82
75-27-4	Bromodichloromethane	5.63	713.3	87
18936-17-9	2-methylbutanenitrile	5.79	719.4	85
108-87-2	Methylcyclohexane	5.92	724.3	96
589-43-5	2,4-dimethylhexane	5.98	726.7	93
123-51-3	Isoamylol	6.06	730	90
137-32-6	2-methylbutan-1-ol	6.07	730.1	84
108-10-1	Methyl isobutyl ketone	6.15	733.5	96
624-92-0	Dimethyl disulfide	6.36	741.5	86
565-69-5	3-pentanone, 2-methyl -	6.43	744.2	84
565-75-3	2,3,4-trimethylpentane	6.45	744.9	99
556-52-5	Glycidol	6.58	749.9	71
563-16-6	3,3-dimethylhexane	6.61	751.3	96
560-21-4	2,3,3-trimethylpentane	6.63	752	98
123-15-9	2-methyl valeraldehyde	6.71	755	70
589-53-7	4-methylheptane	6.87	761.1	92
108-88-3	Toluene	6.93	763.6	100
544-25-2	Cyclohepta-1,3,5-triene	6.94	764.1	89
589-81-1	3-methylheptane	7.05	768.1	97
71-41-0	Pentan-1-ol	7.20	774.2	70
123-54-6	Acetylacetone	7.24	775.7	72
591-78-6	2-oxohexane	7.50	785.7	78
1632-16-2	3-methyleneheptane	7.52	786.5	77
111-66-0	1-octene	7.60	789.7	90
565-80-0	2,4-dimethyl-3-pentanone	7.64	791	94
3848-24-6	Hexane-2,3-dione	7.64	791.1	94
120-92-3	Cyclopentanone	7.64	791.2	71
141-79-7	Mesityl oxide	7.82	798	70
111-65-9	Octane	7.88	800.2	97
66-25-1	Hexanal	7.91	800.9	92
124-48-1	Chlorodibromomethane	8.02	803.9	82
127-18-4	Tetrachloroethene	8.12	806.4	98
106-36-5	Propyl propionate	8.18	808	89
589-90-2	1,4-dimethylcyclohexane	8.19	808.3	93
1069-53-0	2,3,5-trimethylhexane	8.36	812.6	85
123-86-4	Butyl acetate	8.38	813.2	80
2213-23-2	2,4-dimethylheptane	8.61	818.9	96
123-42-2	Diacetone alcohol	9.29	836.3	90
19549-87-2	2,4-dimethyl-1-heptene	9.31	837	79
108-90-7	Chlorobenzene	9.58	843.9	85
7452-79-1	Ethyl 2-methylbutyrate	9.67	846.1	75
100-41-4	Ethylbenzene	10.15	858.5	100
106-42-3	P-xylene	10.51	867.8	100
2918-13-0	1-hepten-3-one	10.75	873.8	74
123-92-2	Isoamyl acetate	10.75	874	79
106-35-4	Heptan-3-one	11.13	883.7	94
124-11-8	1-nonene	11.20	885.4	86
110-43-0	Heptan-2-one	11.29	887.7	91
100-42-5	Styrene	11.43	891.2	92
108-94-1	Cyclohexanone	11.55	894.3	93
4466-24-4	2-butylfuran	11.59	895.5	73
111-84-2	Nonane	11.77	900.1	97
111-71-7	Heptanal	11.86	901.8	84
142-83-6	2,4-hexadienal	11.86	901.9	74
111-76-2	2-butoxyethanol	12.01	905.2	93
590-01-2	Butyl propionate	12.05	905.9	72
815-17-8	3,3-dimethyl-2-oxobutanoic acid	12.18	908.7	71
67-71-0	Sulfonyldimethane	12.35	912.2	90
15869-93-9	3,5-dimethyloctane	12.74	920.4	82
542-18-7	Monochlorocyclohexane	12.78	921.3	77
508-32-7	Tricyclene	12.83	922.4	93
98-82-8	Cumene	12.84	922.7	83
930-68-7	Cyclohex-2-enone	13.17	929.6	76
4110-44-5	3,3-dimethyloctane	13.29	932.2	80
7785-26-4	Alpha-pinene	13.33	933	100
591-11-7	Beta-angelica lactone	13.72	941.2	71
79-92-5	Camphene	14.12	949.6	96
103-65-1	Propylbenzene	14.23	952.1	89
123-05-7	2-ethylhexanal	14.24	952.3	88
620-14-4	3-ethyltoluene	14.58	959.5	95
100-52-7	Benzaldehyde	14.75	963	97
108-67-8	1,3,5-trimethylbenzene	14.99	968	98
3387-41-5	Sabinene	15.19	972.3	100
28634-89-1	2-thujene	15.19	972.3	97
99-82-1	P-menthane	15.21	972.8	81
464-15-3	Bornane	15.31	974.9	76
611-14-3	1-ethyl-2-methylbenzene	15.42	977.1	98
18172-67-3	Beta-pinene	15.44	977.6	98
108-95-2	Phenol	15.52	979.2	74
110-93-0	Sulcatone	15.70	983.2	91

CAS	Name	RT	RI	Match
872-05-9	1-decene	15.73	983.8	80
100-47-0	Benzonitrile	15.78	984.9	82
123-35-3	Beta-myrcene	15.96	988.6	75
111-13-7	2-octanone	15.97	988.7	79
13475-82-6	2,2,4,6,6-pentamethylheptane	15.98	989.1	95
3777-69-3	2-pentylfuran	16.03	990	77
4551-51-3	Hydrindane	16.10	991.5	87
95-63-6	1,2,4-trimethylbenzene	16.18	993.2	98
288-88-0	1h-1,2,4-triazole	16.30	995.8	73
124-18-5	Decane	16.48	999.7	98
124-13-0	Octanal	16.67	1003.3	95
538-93-2	Isobutylbenzene	16.85	1007	86
99-83-2	Alpha-phellandrene	16.92	1008.5	74
498-15-7	3-carene	16.93	1008.6	97
111-90-0	Diethylene glycol monoethyl ether	17.04	1010.9	76
99-86-5	Alpha-terpinene	17.36	1017.4	92
526-73-8	1,2,3-trimethylbenzene	17.50	1020	96
99-87-6	P-cymene	17.72	1024.4	97
527-84-4	O-cymene	17.73	1024.6	98
535-77-3	M-cymene	17.74	1024.9	94
104-76-7	2-ethylhexan-1-ol	17.86	1027.3	98
5989-27-5	Limonene	17.94	1029	99
555-10-2	Beta-phellandrene	18.03	1030.8	87
470-82-6	1,8-cineole	18.12	1032.5	97
496-11-7	Indane	18.18	1033.7	94
872-50-4	N-methylpyrrolidin-2-one	18.23	1034.7	92
13877-91-3	Beta-ocimene	18.24	1034.9	86
95-13-6	1h-indene	18.64	1043	76
122-78-1	Phenylacetaldehyde	18.68	1043.8	97
90-02-8	Salicylaldehyde	18.73	1044.8	86
105-05-5	1,4-diethylbenzene	18.81	1046.3	93
99-85-4	Gamma-terpinene	19.39	1058	92
493-02-7	Decahydronaphthalene	19.46	1059.4	90
98-86-2	Acetophenone	19.79	1066.1	98
111-87-5	Octan-1-ol	20.03	1070.9	75
18479-58-8	Dihydromyrcenol	20.05	1071.3	96
1195-32-0	P-isopropenylmethylbenzene	20.49	1080.1	83
586-63-0	Isoterpinolene	20.75	1085.4	89
617-94-7	2-phenylpropan-2-ol	20.77	1085.7	73
1195-79-5	Fenchone	20.94	1089.1	80
821-55-6	Nonan-2-one	20.95	1089.4	77
78-70-6	Linalool	21.43	1099	83
1120-21-4	Undecane	21.45	1099.4	96
62108-26-3	2,6,8-trimethyldecane	21.49	1100.1	82
493-01-6	Decalin	21.64	1103.4	76
124-19-6	Nonanal	21.68	1104.2	100
546-80-5	Alpha-thujone	21.80	1106.5	95
3878-55-5	Monomethyl succinate	22.05	1111.6	74
95-93-2	Durene	22.41	1119.2	94
78-59-1	Isophorone	22.49	1120.8	74
4501-58-0	Alpha-campholenaldehyde	22.80	1127.1	79
464-48-2	Camphor	23.83	1148.3	98
488-23-3	Prehnitene	23.92	1150.1	84
1196-31-2	Menthone	24.23	1156.6	91
598-42-5	2-hydroxyacetamide	24.34	1158.8	72
119-64-2	Tetralin	24.46	1161.3	83
140-39-6	Tolylacetate	24.54	1163	78
30460-92-5	Pinocarvone	24.56	1163.4	83
65-85-0	Benzoic acid	24.74	1167.2	85
143-08-8	Nonan-1-ol	24.96	1171.5	77
464-45-9	Borneol	25.06	1173.7	82
150-78-7	1,4-dimethoxybenzene	25.30	1178.6	70
2216-51-5	Menthol	25.31	1178.7	98
562-74-3	4-terpineol	25.44	1181.5	87
91-20-3	Naphthalene	25.68	1186.5	92
693-54-9	Decan-2-one	25.90	1190.9	72
119-36-8	Methyl salicylate	25.99	1192.9	88
112-40-3	Dodecane	26.36	1200.4	99
112-31-2	Decanal	26.61	1205.9	99
1196-01-6	Verbenone	26.72	1208.1	74
7206-26-0	2-dodecene	26.90	1211.9	72
17301-22-3	2,5-dimethylundecane	26.93	1212.7	90
95-16-9	Benzothiazole	27.62	1227.5	75
6485-40-1	Carvone	28.44	1245	94
115-95-7	Linalyl acetate	28.62	1248.8	84
105-60-2	Epsilon-caprolactam	28.87	1254.2	78
112-05-0	Nonanoic acid	29.43	1266.1	95
112-30-1	Decan-1-ol	29.76	1273.3	73
31295-56-4	2,6,11-trimethyldodecane	29.83	1274.7	75
5655-61-8	Bornyl acetate	30.44	1287.8	82
125-12-2	Isobornyl acetate	30.45	1288.1	79
112-12-9	Undecan-2-one	30.64	1292.2	72
90-12-0	1-methylnaphthalene	30.88	1297.3	93
91-57-6	2-methylnaphthalene	30.90	1297.5	94
89-83-8	Thymol	30.93	1298.3	94
98-54-4	4-tert-butylphenol	30.95	1298.7	78
499-75-2	Carvacrol	30.97	1299.1	97
629-50-5	Tridecane	31.02	1300.2	93
112-44-7	Undecanal	31.37	1308.1	96
10235-63-9	Gamma-terpinyl acetate	33.13	1347.6	70
334-48-5	Decanoic acid	33.79	1362.2	93
26730-12-1	4-methyltridecane	33.90	1364.7	78
3891-98-3	Farnesane	34.40	1376	86
3856-25-5	Copaene	34.53	1378.8	89
92-52-4	Biphenyl	34.67	1382.1	88
1120-36-1	1-tetradecene	35.13	1392.4	74
80-46-6	P-t-amylphenol	35.18	1393.5	84
629-59-4	Tetradecane	35.48	1400.3	100
581-42-0	2,6-dimethylnaphthalene	35.74	1406.5	70
112-54-9	Dodecanal	35.87	1409.4	96
475-20-7	Longifolene	36.03	1413.4	75
11028-42-5	Cedrene	36.32	1420.2	73
575-43-9	1,6-dimethylnaphthalene	36.50	1424.5	90
575-37-1	1,7-dimethylnaphthalene	36.54	1425.3	87
575-41-7	1,3-dimethylnaphthalene	36.54	1425.4	82
101-81-5	Diphenylmethane	36.83	1432.1	74
131-11-3	Dimethyl phthalate	37.45	1446.8	92
112-53-8	Dodecan-1-ol	38.63	1475	98
473-13-2	Alpha-selinene	39.33	1491.6	89
17066-67-0	Beta-selinene	39.35	1492	88
13360-61-7	1-pentadecene	39.36	1492.2	72

4630-07-3	Valencene	39.67	1499.6	72
629-62-9	Pentadecane	39.70	1500.2	95
96-76-4	2,4-di-tert-butylphenol	39.88	1504.7	88
5875-45-6	2,5-di-tert-butylphenol	39.92	1505.8	72
10486-19-8	Tridecanal	40.13	1511	87
39029-41-9	Gamma-cadinene	40.32	1515.7	77
132-64-9	Dibenzofuran	40.45	1518.9	83
483-76-1	Delta-cadinene	40.46	1519.3	81
21391-99-1	Alpha-calacorene	41.43	1543.6	86
4621-36-7	3-phenyldecane	42.17	1561.9	84
134-62-3	N,n-diethyl-m-toluamide	42.59	1572.5	76
2050-08-0	Amyl salicylate	42.65	1574.1	73
84-66-2	Diethyl phthalate	43.12	1585.6	94
544-76-3	Hexadecane	43.70	1600.3	97
629-73-2	1-hexadecene	43.73	1600.9	76
10233-13-3	Isopropyl laurate	44.65	1625.3	89
4537-14-8	6-phenylundecane	44.72	1627.1	82
4537-15-9	5-phenylundecane	44.86	1630.8	72
473-16-5	Alpha-eudesmol	45.46	1646.8	84
2156-97-0	Lauryl acrylate	47.20	1692.7	79
6765-39-5	1-heptadecene	47.47	1699.7	86
629-78-7	Heptadecane	47.50	1700.5	96
24157-81-1	2,6-diisopropylnaphthalene	48.23	1720.8	76
2719-62-2	6-phenyldodecane	48.41	1725.6	75
593-45-3	Octadecane	51.12	1800.7	96
118-60-5	Octyl salicylate	51.18	1802.3	85
638-36-8	Phytane	51.34	1806.8	79
110-27-0	Isopropyl myristate	51.92	1823.8	83
504-96-1	Neophytadiene	52.38	1837.2	72
1222-05-5	Musk gx	52.58	1842.9	84
629-92-5	Nonadecane	54.64	1903.1	71
107-06-2	1,2-dichloroethane	4.61	649.6	92

Table S 2: ANOVA results for differences in the summed relative amount of each compound class identified between sampling sites on 27th November, 2024.

term	df	sumsq	meansq	statistic	p.value	Type_7
Location Name	5	0.02	0.00	1.413450768	0.272	Aliphatics
Location Name	5	0.14	0.03	42.28108348	<0.001	Aromatics
Location Name	5	0.10	0.02	56.94394467	<0.001	Monoterpenes
Location Name	5	0.00	0.00	29.87459907	<0.001	Others (Esters, N, S, Halo)
Location Name	5	0.00	0.00	6.986857086	0.001	Sesquiterpenes
Location Name	5	0.00	0.00	4.111920384	0.014	Terpenoids

Table S 3: Summed relative amount of compounds identified within each compound class between sampling sites on 27th November, 2024. Letters indicate groups differences based on Tukey post-hoc tests (p<0.05).

	Aliphatics		Aromatics		Monoterpenes		Others		Sesquiterpenes		Terpenoids	
Location	n	T	n	T	n	T	n	T	n	T	n	T
OxBG Outside	3.89	a	10.57	a	1.07	a	0.26	a	0.02	a	0.11	a
OxBG Rainforest House	2.90	a	1.14	b	3.19	b	0.13	bc	0.06	ab	0.44	a
University Parks Dense	2.75	a	2.34	b	0.27	c	0.12	ab	0.00	b	0.06	a
University Parks Open	2.27	a	3.60	c	0.45	ac	0.27	d	0.00	b	0.04	a
Warneford	2.96	a	1.38	b	0.22	c	0.11	ac	0.00	b	0.06	a
Wytham	6.48	a	2.57	bc	0.40	c	0.06	c	0.00	b	0.33	a

Table S 4: ANOVA results for differences in the number of compounds identified within each compound class between sampling sites on 27th November, 2024.

term	df	sumsq	meansq	statistic	p.value	Type_7
Location Name	5	364.54	72.91	2.435749877	0.080	Aliphatics
Location Name	5	331.04	66.21	4.607413753	0.009	Aromatics
Location Name	5	109.92	21.98	6.503543914	0.002	Monoterpenes
Location Name	5	69.17	13.83	3.124705882	0.037	Others (Esters, N, S, Halo)
Location Name	5	19.50	6.50	7.703703704	0.010	Sesquiterpenes

Location Name	5	167.61	33.52	20.89634002	0.000	Terpenoids

Table S 5: Summed number of compounds identified within each compound class between sampling sites on 27[th] November, 2024. Letters indicate groups differences based on Tukey post-hoc tests ($p<0.05$).

	Aliphatics		Aromatics		Monoterpenes		Others		Sesquiterpenes		Terpenoids	
Location	n	T	n	T	n	T	n	T	n	T	n	T
OxBG Outside	n	Tuk	n	Tuk	n	Tuk	n	Tuk	n	Tuk	n	Tuk
OxBG Rainforest House	65	a	41	b	13	ab	16	b	2	a	8	b
University Parks Dense	67	a	34	ab	15	b	15	ab	10	b	13	c
University Parks Open	60	a	29	a	9	a	9	ab	2	a	6	ab
Warneford	51	a	25	a	9	a	10	ab	1	a	4	a
Wytham	59	a	36	ab	8	a	11	ab	1	a	5	ab

Table S 6: All compounds identified only at a single site during the 27 November 2024 sampling.

Name	Loc	Type_7
2,4-hexadienal	BO	Aliphatics
2,5-dimethylundecane	BO	Aliphatics
2-methyl valeraldehyde	BO	Aliphatics
3,5-dimethyloctane	BO	Aliphatics
4-methyltridecane	BO	Aliphatics
Acetylacetone	BO	Aliphatics
Decalin	BO	Aliphatics
Glycidol	BO	Aliphatics
Mesityl oxide	BO	Aliphatics
Pyrrolidine	BO	Aliphatics
1,4-diethylbenzene	BO	Aromatics
1h-indene	BO	Aromatics
2-butylfuran	BO	Aromatics
Cumene	BO	Aromatics
Isobutylbenzene	BO	Aromatics
P-isopropenylmethylbenzene	BO	Aromatics
Prehnitene	BO	Aromatics
Styrene	BO	Aromatics
Tetralin	BO	Aromatics
2-thujene	BO	Monoterpenes
Beta-angelica lactone	BO	Monoterpenes
Bornane	BO	Monoterpenes
Isobornyl acetate	BO	Monoterpenes
Verbenone	BO	Monoterpenes
Ethyl 2-methylbutyrate	BO	Others
Tolylacetate	BO	Others
Farnesane	BO	Sesquiterpenes
1-hepten-3-one	RH	Aliphatics
2,4-dimethyl-3-pentanone	RH	Aliphatics
2-hydroxyacetamide	RH	Aliphatics
3-methylbutanal	RH	Aliphatics
3-pentanone, 2-methyl -	RH	Aliphatics
Epsilon-caprolactam	RH	Aliphatics
6-phenyldodecane	RH	Aromatics
Musk gx	RH	Aromatics
Beta-myrcene	RH	Monoterpenes
Beta-ocimene	RH	Monoterpenes
Beta-phellandrene	RH	Monoterpenes
Gamma-terpinyl acetate	RH	Monoterpenes
Isoamyl acetate	RH	Others
N,n-diethyl-m-toluamide	RH	Others
Sulfide, allyl methyl	RH	Others
Alpha-eudesmol	RH	Sesquiterpenes
Alpha-selinene	RH	Sesquiterpenes
Beta-selinene	RH	Sesquiterpenes
Cedrene	RH	Sesquiterpenes
Copaene	RH	Sesquiterpenes
Gamma-cadinene	RH	Sesquiterpenes
Longifolene	RH	Sesquiterpenes
Valencene	RH	Sesquiterpenes
4-terpineol	RH	Terpenoids
Isophorone	RH	Terpenoids
Linalyl acetate	RH	Terpenoids
Sulcatone	RH	Terpenoids
2-octanone	UD	Aliphatics
Decan-2-one	UD	Aliphatics
Nonadecane	UD	Aliphatics
Nonan-2-one	UD	Aliphatics

2,6-dimethylnaphthalene	**UD**	**Aromatics**
Diphenylmethane	**UD**	**Aromatics**
Cyclohex-2-enone	**UD**	**Others**
Pentan-1-ol	**UO**	**Aliphatics**
1,4-dimethoxybenzene	**UO**	**Aromatics**
3,3-dimethyl-2-oxobutanoic..	**UO**	**Others**
Alpha-campholenaldehyde	**UO**	**Terpenoids**
1-hexadecene	**WA**	**Aliphatics**
Butan-1-ol	**WA**	**Aliphatics**
Cyclohexene	**WA**	**Aliphatics**
Nonan-1-ol	**WA**	**Aliphatics**
2,5-di-tert-butylphenol	**WA**	**Aromatics**
2-pentylfuran	**WA**	**Aromatics**
2-phenylpropan-2-ol	**WA**	**Aromatics**
Benzothiazole	**WA**	**Aromatics**
Phenol	**WA**	**Aromatics**
Car-3-ene	**WA**	**Monoterpenes**
Lauryl acrylate	**WA**	**Others**
Neophytadiene	**WA**	**Terpenoids**
1-pentadecene	**WY**	**Aliphatics**
2,3-dimethylpentane	**WY**	**Aliphatics**
2,6,8-trimethyldecane	**WY**	**Aliphatics**
Cyclopentanone	**WY**	**Aliphatics**
Pentanal	**WY**	**Aliphatics**
Alpha-phellandrene	**WY**	**Monoterpenes**
Isoterpinolene	**WY**	**Monoterpenes**
Thymol	**WY**	**Monoterpenes**
Butyl acetate	**WY**	**Others**
Butyl propionate	**WY**	**Others**
Monomethyl succinate	**WY**	**Others**
Borneol	**WY**	**Terpenoids**
Carvacrol	**WY**	**Terpenoids**
Fenchone	**WY**	**Terpenoids**

Table S 7: Top-ranking compounds from PLS-DA analyses with explicit directionality. VIP scores (> ranked top 10) identify compounds contributing most strongly to group separation. Group-wise mean abundances are provided to support directional interpretation.

Comparison	Direction	Compound	VIP	Loading	Mean_Group1	Mean_Group2
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	P-xylene (Aromatics)	2.296	-0.148	0.186	0.367
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	Ethylbenzene (Aromatics)	2.268	-0.146	0.108	0.209
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	Isooctane (Aliphatics)	2.206	-0.142	0.035	0.133
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	2,3,4-trimethylpentane (Aliphatics)	2.150	-0.138	0.032	0.087
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	Toluene (Aromatics)	2.138	-0.137	0.239	0.424
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	3-methylhexane (Aliphatics)	2.121	-0.136	0.064	0.136
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	Methylcyclohexane (Aliphatics)	2.065	-0.133	0.065	0.131
WAH/WYH/UDH/RHH v BOH/UOH	> in WAH/WYH/UDH/RHH	2-dodecene (Aliphatics)	2.060	0.132	0.198	0.000
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	1-ethyl-2-methylbenzene (Aromatics)	2.014	-0.129	0.027	0.110
WAH/WYH/UDH/RHH v BOH/UOH	> in BOH/UOH	3-methylheptane (Aliphatics)	2.005	-0.129	0.041	0.074
WAH/WYH/UDH v RHH	> in RHH	Beta-ocimene (Monoterpenes)	2.285	-0.147	0.000	0.072
WAH/WYH/UDH v RHH	> in RHH	Dihydromyrcenol (Terpenoids)	2.281	-0.147	0.000	0.105
WAH/WYH/UDH v RHH	> in RHH	Sabinene (Monoterpenes)	2.273	-0.146	0.005	0.374
WAH/WYH/UDH v RHH	> in RHH	Beta-pinene (Monoterpenes)	2.244	-0.144	0.033	0.287
WAH/WYH/UDH v RHH	> in RHH	Pinocarvone (Terpenoids)	2.232	-0.143	0.000	0.043
WAH/WYH/UDH v RHH	> in RHH	Hydrindane (Aliphatics)	2.216	-0.142	0.000	0.028
WAH/WYH/UDH v RHH	> in RHH	Alpha-pinene (Monoterpenes)	2.196	-0.141	0.104	0.376
WAH/WYH/UDH v RHH	> in RHH	Copaene (Sesquiterpenes)	2.195	-0.141	0.000	0.048
WAH/WYH/UDH v RHH	> in RHH	4-terpineol (Terpenoids)	2.195	-0.141	0.000	0.034
WAH/WYH/UDH v RHH	> in RHH	Beta-phellandrene (Monoterpenes)	2.169	-0.139	0.000	0.045
BOH v UOH/UDH	> in BOH	2-thujene (Monoterpenes)	2.300	-0.148	0.055	0.000
BOH v UOH/UDH	> in BOH	Prehnitene (Aromatics)	2.288	-0.147	0.035	0.000
BOH v UOH/UDH	> in BOH	Farnesane (Sesquiterpenes)	2.284	-0.147	0.035	0.000
BOH v UOH/UDH	> in BOH	P-isopropenylmethylbenzene (Aromatics)	2.274	-0.146	0.020	0.000
BOH v UOH/UDH	> in BOH	Bromodichloromethane (Others (Esters, N…)	2.268	-0.146	0.013	0.000
BOH v UOH/UDH	> in BOH	1,4-diethylbenzene (Aromatics)	2.259	-0.145	0.045	0.000
BOH v UOH/UDH	> in UOH/UDH	Tricyclene (Monoterpenes)	2.251	0.145	0.000	0.076
BOH v UOH/UDH	> in BOH	2,2,4,6,6-pentamethylheptane (Aliphatics)	2.201	-0.142	0.099	0.062
BOH v UOH/UDH	> in BOH	Bornyl acetate (Terpenoids)	2.183	-0.140	0.017	0.000
BOH v UOH/UDH	> in BOH	Propylbenzene (Aromatics)	2.146	-0.138	0.093	0.009