A walk in the park – identifying healthy biomes using scent snapshots

William T. Kay [a,*], Anya Lindström Battle [a], Mahal Humberstone [a], Molly Tucker [a],

Kieran E. Storer [a], Geoffrey Kite [b], Katherine Willis [a]

[a] Department of Biology, University of Oxford, South Parks Road, Oxford OX1 3RB, UK

[b] Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK.

* Corresponding author: william.kay@biology.ox.ac.uk

1   Abstract

Biogenic volatile organic compounds (bVOCs) are carbon-based secondary metabolites emitted by all kingdoms of life, often playing roles in stress protection, growth modulation, and inter-organism communication. In plants, bVOCs such as terpenes (e.g., alpha-pinene, d-limonene) are highly prevalent and have been linked to diverse health benefits including stress reduction, enhanced cognitive function, and neuroprotection. In contrast, urban environments contain anthropogenic VOCs (aVOCs) produced from vehicles and industrial activity. Examples of aVOCs such as benzene, toluene, ethylbenzene and p-xylene (BTEX compounds) are all associated with serious health risks. Using Tenax™ filters and GCMS analysis, this study assessed the presence and relative amounts of bVOCs and aVOCs in the air across six urban greenspaces in Oxford. Sampling protocols were standardised using a self-contained sampling box (VOKSBOX), and data were processed using workflows optimised for large metabolomics-style datasets. Our methods resulted in 175 unique compounds of biological interest being identified over our six sampling sites – a significant increase from those identified in previous open-air studies. Using PCA, PLS-DA and HCA analysis we show that sites significantly and consistently separate by volatile profile, and that the spread of compounds emitted can be used to assess the potential healthiness of varied Oxford greenspaces. Target 12 of the Global Biodiversity Framework calls for expanding high-quality, health-promoting urban greenspaces. By quantifying both beneficial and harmful VOCs, our study proposes a novel, health-oriented metric for site evaluation. These findings support the integration of VOC profiling into urban planning and public health strategies to enhance the well-being of urban populations.

2   Keywords

VOC, aVOC, bVOC, volatiles, health, terpenes, BTEX

3   Introduction

Biogenic volatile organic compounds (bVOCs), often characterised by their specific scent, are a chemically diverse group of carbon-based secondary metabolites produced by all kingdoms of life (Luo et al., 2025, Peñuelas et al., 2014, Amann et al., 2014). Many bVOCs have primary roles within the organisms which produce them, these include stress/disease protection, growth modulation, and cell-cell communication (Peñuelas and Munné-Bosch, 2005, Bartwal et al., 2013, Roberts, 2007). Once their solution equilibrium is exceeded, bVOCs enter the air and can then act as agents of communication at both intra- and inter-species levels (Weisskopf et al., 2021, Bergman et al., 2025). Examples include compounds whose scents act as attractants for mates/pollinators (Kessler et al., 2019, Xu and Turlings, 2018, Xu et al., 2014, Xu et al., 2017a, Boncan et al., 2020), or deterrents of pests/pathogens (Heil and Ton, 2008, Hammerbacher et al., 2019, Brambilla et al., 2022). Previous reviews have estimated that in plants the hemiterpene isoprene (C_5H_8), chiefly produced to combat abiotic stress in plants, constitutes approximately 50 % of bVOC emissions, with simple C1 and C2 compounds and more complex isoprenoids such as monoterpenes, sesquiterpenes and oxygenated terpenes comprising close to the other 50 % (Guenther et al., 2012, Antonelli et al., 2020).

Plant-produced bVOCs such as the terpenes alpha -pinene, beta-pinene, and d-limonene have been the focus of most studies on ambient air to date (Walker et al., 2023, Lahr et al., 2015, Seco et al., 2015, Sanaei et al., 2023, Ma et al., 2025, Guenther et al., 2012, Guenther et al., 1995, Guenther et al., 1993, Wang et al., 2025). This is due to research highlighting the varied temporary and lasting human health benefits imparted by these compounds when inhaled. Short term benefits include physiological and psychological reductions in stress via lowering cortisol, blood pressure and heart rate (Park et al., 2010, Sousa et al., 2018), as well as enhancing cognitive function through increases in working memory and reaction time (Saenen et al., 2023, Zeng et al., 2022, Bratman et al., 2015, Shibata and Suzuki, 2002, Xu et al., 2017b, Farimani et al., 2024). Terpenes have also been shown to act in a neuroprotective manner via promoting cell survival and inhibiting neuronal apoptosis, even reversing hippocampal cell injury in mice (Xu et al., 2022, Xu et al., 2017b). As reviewed elsewhere (Siddiqui et al., 2024, Masyita et al., 2022, Ren et al., 2017), further health benefits include antimicrobial, antioxidant, immunomodulatory, sedative, respiratory, anti-inflammatory, and anticancer effects (Li, 2010, Ben Ammar, 2023, Kim et al., 2020, Del Prado-Audelo et al., 2021, Jo et al., 2021).

Conversely, anthropogenic volatiles (aVOCs) such as from plastics, industrial outputs, or transport related activity can dominate in urban greenspaces (McDonald et al., 2018, Gao et al., 2023). The compounds benzene, toluene, ethylbenzene, and p-xylene (BTEX compounds) are emitted by vehicle exhausts and as such are prevalent urban pollutants (Liu et al., 2025, Gao et al., 2023, Hoque et al., 2008). aVOCs such as these are directly damaging to human health via increasing risks of cancer, respiratory conditions, and cardiovascular disease (Fazakas et al., 2024, He et al., 2024, Saeedi et al.,

2024, Muda et al., 2024). Additionally, the reaction of aVOCs with specific bVOCs leads to the production of ozone and secondary organic aerosols (Ren et al., 2017, Cao et al., 2022, Wang et al., 2025). This effect is especially problematic in sunlit, polluted, urban environments, leading to chronic exposure for urban residents.

The potential impact of volatiles in urban environments means that VOC monitoring is highly relevant to national and international health policy. Target 12 of the Kunming-Montréal Global Biodiversity Framework, adopted by COP 15 in December 2022 and now signed by 189 Parties, commits to: "significantly increase the area and quality, and connectivity of, access to, and benefits from green and blue spaces in urban and densely populated areas… improving human health and well-being and connection to nature…" (Kunming-Montreal, 2022). Yet this target assumes all greenspaces are equal. Specifically relating to air quality and greenspace, questions such as (i) which VOCs contribute to a healthy/unhealthy urban greenspace, and (ii) where healthy urban scentscapes are most likely to be found, need to be answered. Greater knowledge of these aspects will allow for more nuanced improvements to urban greenspace design and subsequent functional nature prescriptions (Nguyen et al., 2023, Bratman et al., 2015, White et al., 2019).

The primary aim of our study was therefore to understand the relative distribution of healthy and unhealthy VOCs in six urban greenspaces in Oxford, UK. Although similar methods have been developed for headspace sampling (Yin et al., 2022, de Carvalho Couto et al., 2024, Reyrolle et al., 2024), none have yet been published for the analysis of ambient air. Using our methods we demonstrate significant differences in the VOC profiles and show that site location and ecology are significant drivers of volatile these differences. Using this information, predictions can then be made as to the characteristics of greenspaces which are likely to have positive impacts on human health.

1   Materials and Methods

1.1   Site locations and sampling strategy

Six urban greenspace sites around Oxford were chosen as sampling sites due to their (i) public accessibility, (ii) different ecologies, and (iii) contrasting distance from the city centre. Five of the chosen sites were outdoor greenspaces which could be compared directly to each other to test the hypotheses that a sites location and ecology are significant drivers of volatile difference. These outdoor greenspaces, as shown in and marked in Figure 1 were Botanic Garden Outdoors, University Parks Dense, University Parks Open, Warneford Meadow, and Wytham Woods. The sixth site chosen was the Rainforest Glasshouse at the Oxford Botanical Gardens. This site was included to test the hypothesis that an indoor, plant rich, and semi enclosed environment may contain a higher number and relative amount of beneficial bVOCs compared to potentially damaging aVOCs. All samples used in this study were collected in December 2024.

To investigate the characteristic bVOC profiles of the selected sites (Figure 1), we employed a

collection protocol adapted from Walker et al (2023) with the addition of housing the pump and

apparatus within a bespoke metal box (VOKSBOX, Figure 2B). Ambient air was captured using a

GilAir® pump connected simultaneously to 4 replicate Markes Tenax® TA tubes using a flow rate of

400ml/min for 2 hours (12 L of air per tube). Samples were collected at a height of 1.5m – based on

the approximate level of the adult human nose while walking - between the hours of 10:00 - 15:00

GMT. Tenax® tubes were capped immediately after collection and stored at - 20 °C for 14 days.

Figure 1: A: Map of the sampling locations used in this study. (Botanic Garden Outdoors; Botanic Garden

Rainforest Glasshouse; University Parks Dense; University Parks Open; Warneford Meadow; Wytham Woods).

Numbers on roads indicate Annual Average Daily Traffic (AADF) for 2023 as representative data for each site

1.2   GC-MS protocol for identification of VOC profiles

For identification of VOC profiles, we employed gas chromatography mass spectrometry (GC-MS) analysis at the Jodrell laboratory, Royal Botanic Gardens, Kew. Samples stored in Tenax® TA tubes were removed from the freezer and stored at room temperature for at least 2 hours before starting the thermal desorption protocol. For this, tubes were desorbed for 10 mins at 230 °C using a TD100-xr automated thermal desorption (TD) unit from Markes. Samples were prefocused on a Tenax® TA packed cold trap at 4 °C for 0.5 mins followed by heating from 4 °C to 300 °C over 5 mins at a purge flow rate of 50ml/min onto a J&W DB-5ms GC column (30 m × 0.25 mm × 0.25 μm) within an Agilent 8890 GC System. Chromatography was achieved by heating the column at an initial temperature of 40 °C at a rate of 3 °C/min to 220 °C at a flow rate of 1 ml/min Helium. Data were collected in electron ionization (EI) mode using an Agilent 5977C GC/ MSD over a mass range of 50–400 amu.

1.3   GC-MS Data Analysis

The full GC-MS data analysis pipeline developed in this work is presented in Figure 2 and includes the following key steps:

Library Development: The GC-MS library used for identification of compounds was a modified version of the NIST20 Wiley Registry 12th edition mass spectral library. Using NIST-Lib2nist and Mass Hunter Library Editor (Agilent), the complete library was reduced by selecting only compounds which exist in the Chemical Entities of Biological Interest (ChEBI) database (Degtyarenko et al., 2007). To assist in identification, only compounds with entries for retention indexes (RI) on semi-standard non-polar columns (to match the J&W DB-5ms column used) were selected for the library. Compounds missing a RI were either searched for and inputted manually using PubChem data or removed if no values were found. To avoid mismatches, isomers with the same RI were combined (e.g. (−)-α-pinene / (+)-α-pinene, became α-pinene). The final library used (W12N20MAIN_chebi_WK_edit.MSP) contained 3,027 unique compounds.

Pre-Analysis: To aid in the correct identification of compounds, a non-isothermal Kovats index (Kovats, 1958) was built using retention times from C8 – C20 alkane standards (Merck, 04070-1ML). This index uses known compounds to produce accurate estimations of where all compounds will elute from the GC column (retention times) thus allowing precise identification. We validated this index using 60 pure terpene standards (Restek, 34142 and 34143). All standard compounds tested were confirmed to appear at the estimated retention times.

Compound Identification: Sample analysis was performed using the Automated Mass Spectral Deconvolution and Identification System (AMDIS) version 2.73. Compound identification relied on

spectral and RI matching, with a minimum match factor threshold of 70 (Net), a RI window of +/- 10,

and a very strong match factor penalty. Deconvolution settings were: component width: 25; adjacent

peak subtraction: two; shape requirements: medium; sensitivity: high; and resolution: medium. The

analysis was restricted to a single identification per compound.

Post Processing: As outlined in Figure 2E compounds deemed GC-MS column bleed were removed

from the analysis. Using the Kovats index, falsely identified compounds in our dataset were removed

if Outdoors the estimated retention time (maximum RI deviation of 10). Best-match compounds for

each retention time and replicate tube were then chosen based on Net match score. Peak areas of

any compounds identified in blank control tubes were removed from the peak areas of all samples

within a GC-MS run. The relative amount (%) was then recalculated as 'edited amount' based on the

Total Ion Count (TIC) of the sample / modified peak areas. Only compounds which were still found to

be present in >= 50% of technical replicate tubes within each experimental location were retained.

Finally, outlier sample tubes - such as those potentially affected by TD cap failures - were identified

and removed using a two-step statistical procedure. First, a Principal Component Analysis (PCA) was

performed on compound abundance data to produce the Euclidean distance of each technical

replicate from the centroid of its corresponding sampling group. Tubes were flagged as outliers if

greater than 1.5 × the median distance (IQR) for that group.

161
162 **Figure 2: B: Sampling box (VOKSBOX) developed for this work on a tripod at 1.5m height. Air from the GilAir®**

163 **sample pump is split 6 ways using brass splitters with silicone tubing. C-E: A flow chart of the methods used in**

164 **this paper for acquisition, identification, and processing of compounds. Red text shows number of compounds**

165 **identified after each processing step.**

1.4   Statistics and data visualisation

167 To discern the volatile compounds found in the ambient air samples from the different study

168 locations, means of abundance data (relative amount) were calculated using 4 technical replicates for

169 all compounds identified per sampling location.

Principal Component Analysis (PCA): PCA analysis using all 175 compounds was carried out for an initial exploratory study into the structure and relationships of the overall volatile profile of the 6 sampling sites (Figure 3A). A Hellinger transformation (square root of the relative amounts) was first applied to the raw compound abundance data to better account for zero inflation and to make the data suitable for Euclidean-based ordination methods (Legendre and Legendre, 2012, 2013, Chang and Lin, 2011). For visualisation, the PCA data of all 175 compounds identified were first centred (subtracting the mean of each compound) and scaled (dividing by the standard deviation) to ensure comparability across compounds. The analysis was then carried out using the FactoMineR (Lê et al., 2008) and factoextra (Kassambara and Mundt, 2017) packages in R. To confirm differences in compound composition between sample groups, a PERMANOVA (Permutational Multivariate Analysis of Variance) was performed using the adonis2 function from the vegan R package (Oksanen, 2015).

Partial Least Squares Discriminant Analysis (PLS-DA): The PLS-DA (Figure 3B, C & D). is a supervised classification (discriminant analysis) and is used for finding components which maximize group separation while explaining variance. This was carried out using pls and mixOmics statistical (Rohart et al., 2017) packages in R. This approach centres and scales the data (as per the PCA) before plotting.

Hierarchical Cluster Analysis (HCA): To explore overall similarities in compound composition between sampling sites, an HCA was carried out using the same Hellinger-transformed compound abundance data (Figure 4). HCA is an unsupervised method that groups samples based on their multivariate dissimilarity. Clustering was performed using Ward's linkage method, which favours the formation of compact, internally coherent clusters, and Euclidean distance as the dissimilarity metric. PLS-DA was also run on the 3 major clades found in the HCA and the top 10 Variable Importance in Projection (VIP) scores for each clade are labelled in Figure 4.

Grouping by Compound Classes: All identified compounds were categorised into 6 major functional classes - aliphatics, aromatics, monoterpenes, sesquiterpenes, terpenoids (oxygenated terpenes) and others (esters, nitrogen- and sulphur-containing compounds). These classes allowed us to assess the differences in volatile composition within the sampling sites. An initial PCA was carried out using Hellinger-transformed data of the 6 compound classes within each sampling site. Further to this, one-way ANOVAs were run on summed abundances within each functional class to find differences between sampling locations. Post-hoc TUKEY tests were used to determine which locations differed significantly (Figure 5).

Dominant and Unique Compounds: Using mean abundance data, the top 5 compounds from each site were calculated (Figure 6A-F).

Analysis of compounds linked to human health: To focus on health-relevant compounds, terpenes and terpenoids only were selected from the full dataset. A hierarchical heatmap (Figure 7) was

generated with colour intensity representing compound abundance scaled within individual
locations.

Analysis of BTEX compounds and terpene/BTEX ratio: The relative amounts of all
terpenes/terpenoids were compared as a ratio to the sum of the anthropogenic BTEX compounds:
Figure 8, benzene, toluene, ethylbenzene and p-xylene (∑terpenes/∑BTEX) (Malik et al., 2025). This
enabled us to make simple comparisons between locations based on searching for areas expected to
be most amenable to human health.

All statistics were run using R version 4.4.1 (2024-06-14 ucrt) within RStudio 2024.09.0+375
"Cranberry Hibiscus" release for windows. All graphs were generated using the ggplot2 (Wickham,
2011) package.

1.5 Measuring sampling site ecology

To assess the ecology of our 6 sampling sites we employed 3 metrics. The first was to use plant
surveys to determine the total number of species found within a 10 metre radius of our sampling
sites (Table S3). The species of all visible plants was recorded as being present regardless of size. For
the OxBG Outdoors site, plant labels were used for species identifications. Plant species Identities
from the Rainforest glasshouse were provided by staff from the Oxford Botanic Garden (University of
Oxford Botanic Garden & Arboretum Plant Records Database, 2024). For other sites, species were
identified using the Seek app and cross-referenced with a physical key (Poland, J., et al., 2020). The
second metric was a measure of canopy cover by trees over the same 10m radius. Google maps was
used to plot the areas and measurements were made using masking in Adobe Photoshop®. Finally,
using estimated Annual Average Daily Traffic (AADF) for 2023 as representative data (GOV.UK, 2023),
the busiest road within 1km of the sampling site was recorded**.**

2 Results

2.1 Distinctive volatile fingerprints of six urban greenspaces

Our analysis pipeline resulted in 175 compounds being identified within our 6 Oxford sampling sites
(Table S1). A PCA which used all identified compounds as predictor variables revealed significant
variation between sites, each showing distinctive site-specific clustering (Figure 3A and PERMANOVA,
$F_{5,16}$ = 35.713, p < 0.001). The first two principal components accounted for 53.4 % of the variance
(PC1: 34.2%, PC2: 19.2%). Further analysis using a PLS-DA showed very similar separation of sampling
sites (Figure 3 B, C, D) with terpenes/terpenoids (generally biogenic, e.g. beta-phellandrene, beta-
ocimene, beta-pinene, and 3-carene), and aromatic compounds (generally anthropogenic, e.g.
benzene, ethylbenzene, p-xylene, and toluene) exerting a dominant influence on both axes. This is
consistent with sampling in both a plant dense greenhouse, and outdoor urban areas.

Figure 3: Compound richness, PCA, PLS-DA loading plot, and HCA of volatile profiles across sampling locations.

A: PCA was performed on full VOC profiles (175 compounds, n = 4 replicates per sampling site). Larger symbol

represents centroid of replicates. Ellipses represent 95% confidence intervals. Distinct clustering indicates

chemically distinct VOC profiles among sampling sites (PERMANOVA, $F_{5,16}$ = 35.713, p = 0.001). B, C & D: PLS-

DA loading plot showing the top 20 compounds contributing to sample separation. Vectors indicate the

magnitude and direction of each compound's influence on PLS1 and PLS2.

2.2 Volatile fingerprints separate into distinctive clades

Using HCA to analyse all volatile compounds (175) found in the 6 sampling sites, major clades appear

(Figure 4). First is the split of University Parks Open (UO) / Botanical Gardens Outdoors (BO) group

and the other sites (Figure 4A). PLS-DA VIP analysis, shown in the lower segment of Figure 4, showed

this separation was due to many vehicle transport emission-related compounds such as 2,3,3-

trimethylpentane, p-xylene, 2-ethylhexan-1-ol, indane, and ethylbenzene. Secondary clustering

appears between Rainforest Glasshouse (RH) and University Parks Dense (UD), Wytham Woods (WY),

and Warneford Meadow (WA) - Figure 4B. This was found to be mostly due to the higher relative

amount of plant related compounds such as beta-ocimene, dihydromyrcenol, sabinene, diacetone

alcohol, beta-pinene, pinocarvone. The third most significant clade (Figure 4C) splits University Parks

Open (UO) and Botanical Gardens Outdoors (BO) which is notably due to the proportion of terpenes

such as p-cymene, 2-thujene, tricyclene. All replicates of sampling locations form tight groupings with

no overlap over other locations with one exception - the presence Wytham Woods (WY) in the

Warneford Meadow (WA) clade. This highlights the close similarity of the two sampling sites.

Figure 4 TOP: HCA conducted using Ward's linkage and Euclidean distances using all technical replicate samples

over 6 urban greenspaces: Botanical Gardens Outdoors (BO), OxBG Rainforest Glasshouse (RH), University

Parks Dense (UD), #University Parks Open (UO), Warneford Meadow (WA), Wytham Woods (WY). BOTTOM:

Top ten compounds responsible for differences between sites based on VIP scores from PLS-DA analysis.

Letters represent the three major clades identified.

2.3   Urban greenspaces show distinctive mix of compound classes

Results from a PCA which used the 6 classes of identified compounds (aliphatics, aromatics,

monoterpenes, sesquiterpenes, others (esters, N, S and halogenated compounds), and terpenoids) as

the predictor for site difference led to significant separation of the six locations sampled (Figure 5).

Using the total number of compounds identified in each compound class (Figure 5), results show that

the number of all six compounds groups differed between sites. The number of identified terpenoids

was found to be highest in the Rainforest Glasshouse, whereas monoterpenes and sesquiterpenes

were found to be significantly higher in both the Rainforest Glasshouse and Botanical Gardens

Outdoors compared to the other 4 sites (one-way ANOVAs followed by Tukey's HSD post-hoc test $p <$

0.05). In terms of relative abundance within the compound classes (Figure 5), results strengthen the

total count data by showing two major differences: Firstly, the Botanic Garden Outdoors contained a

significantly higher proportion of aromatic compounds compared to other sites suggesting a strong influence from anthropogenic sources. Secondly, the Rainforest Glasshouse showed significantly higher proportion of terpenoids whereas the highest relative proportions of monoterpenes, and sesquiterpenes were shared by the Rainforest Glasshouse and Botanical Gardens Outdoors (One-way ANOVAs followed by Tukey's HSD post-hoc test p < 0.05).

Figure 5: Composition of volatile compounds across sampling locations separated by compound class: A: PCA of the six sampling sites based on compound class. PCA was performed on VOC profiles using Hellinger transformed data (n = 4). Ellipses represent 95% confidence intervals. B: Total number (means) of detected volatile compounds per site, categorized by compound class. Results show that the number of all compounds groups differed between sites. C Relative abundance (means) of detected volatile compounds split by classes across sites, expressed as a percentage of the total sample. A and B: Bars represent the mean (n = 4 technical replicate sampling tubes per site). Letters indicate statistically distinct sampling locations within chemical classes based on one-way ANOVA followed by Tukey's HSD post-hoc test (p < 0.05).

2.4   Dominant VOCs identified at sampling sites

Within the top 5 compounds identified (by relative amount) at each site, the Rainforest Glasshouse (Figure 6B) exhibited the greatest predominance of plant-derived terpenes/terpenoids (green), which aligns with its dense and varied vegetation cover. In all other sampling sites, aromatics (orange) and aliphatics (grey) dominated. The aromatic BTEX compounds appear in the top 5 compounds of all outdoor greenspaces except for Warneford Meadow, reinforcing a strong traffic-related influence within outdoor urban sites. Indeed, the Botanical Gardens Outdoors featured all 4 BTEX compounds within the top 5 compounds (Figure 6A) likely due to being the closest of the sample sites to a busy city centre road (Figure 1A). The other outdoor greenspaces - University Parks Open, University Parks Dense, Warneford Meadow, and Wytham Forest (Figure 6C-F) – mostly included a mix of aromatic hydrocarbons (orange) and aliphatics (grey), with a single monoterpene (alpha pinene) appearing at the University Parks Open sampling site.

Figure 6: Dominant volatile compounds across sampling locations. A-F: Bar plots show the top 5 VOCs ranked by relative abundance at each site. Data were collected from n = 4 technical replicate samples per location. Error bars indicate standard error of the mean (SEM). Compound categories are color-coded based on 3 chemical classes: terpenes/terpenoids (green), aliphatics (grey), and aromatics (orange).

2.5   Presence/absence of plant-based terpenes and terpenoids

Terpenes/terpenoids were found at all sampling locations. A total of 34 were identified in this work. Six compounds - alpha-pinene, limonene, 3-carene, camphene, camphor and 1-8 cineole (eucalyptol) - were detected at every site, with a further five – alpha-calacorene, beta-pinene, menthol, p-cymene and tricyclene – detected in five of the six sampling sites. Again, the Rainforest Glasshouse exhibited the largest number of identified terpenes/terpenoid compounds overall (27 compounds), with other sites containing fewer: Botanical Gardens Outdoors (19), University Parks Dense (11), University Parks Open (10), Warneford Meadow (12) and Wytham Forest (12). Seventeen terpene/terpenoid

compounds were found at single sites, with most detected in the Rainforest glasshouse (13) and Botanical Gardens Outdoor site (5). A full breakdown of unique compounds per site is listed in Table S4.

Figure 7: Terpenes/terpenoid compounds identified across sampling locations. Heatmap showing the scaled relative amount (light green > dark green) or absence (grey) of any terpenes/terpenoid VOCs detected across six sampling locations. Scaling was carried out within single locations rather than over the whole dataset. For presence, each compound was found to be present in at least 2 of 4 technical replicate tubes.

2.6   BTEX compounds as predictors of pollution

The ratio of biogenic terpenes/terpenoids to anthropogenic BTEX compounds was significantly higher in the Rainforest Glasshouse compared to the other sites. This is in part due to the increased prevalence of terpenes/terpenoids and in part to the lower proportion of BTEX compounds in this indoor space. The outdoor sites all had a significantly lower ratios with significantly higher amounts of BTEX compounds identified ranging from Warneford (12.6%) to the Botanical Gardens Outdoors (45.4%).

Figure 8: A: Terpene: BTEX ratio of sampling sites (∑terpenes/∑BTEX). Relative amounts of all terpenes/terpenoids as a ratio to the anthropogenic BTEX compound. B: BTEX as a % of all compounds identified. Both graphs display mean values of 4 technical replicates with error bars representing SEM.

2.7   Sampling site ecology

As explained in methods section 1.5, we employed 3 metrics to assess the ecology of our sites. Total plant species was found to be significantly higher within a 10m radius of the sampling site in the OxBG Rainforest Glasshouse (208). The outdoor sites ranged from 43 species in the Botanical Gardens Outdoors site down to just 7 in University Parks Open. Assessments of the localised canopy cover showed that again, OxBG Rainforest Glasshouse was at one end of the spectrum, being completely covered in canopy. Canopy cover was more varied in the outdoor sites ranging from 68% in University Parks Dense to 0% at University Parks Open. Finally, recordings of road traffic within a 1km radius of the sites showed that 4 sites (both Botanical Gardens and University Parks sites) were close to busy roads with more than 12,000 motor vehicles estimated to pass each day. Warneford and Wytham ware more isolated, particularly the latter which only has occasional visitor traffic.

Table 1: A: A summary of the total species counts in our 6 sampling locations. B: Localised Tree cover (% in 20m dia.) measured using Google maps. C: Most busy road within 1km radius of sampling site using AADF data (GOV.UK, 2023).

Location	A: Total plant species in 10m rad.	B: Localised canopy cover (% in 10m rad.)	C: Most busy road in 1km radius (AADF)
Botanical Gardens Outdoors	43	37%	13,871
Rainforest Glasshouse	208	100%	13,871
University Parks Dense	15	68%	12,524
University Parks Open	7	0%	12,524
Warneford Meadow	15	30%	6,731
Wytham Forest	21	51%	100 (est.)

3 Discussion

This study aimed to develop a robust and repeatable protocol for measuring the presence and relative amounts of a wide range of volatiles of biological interest. Our work took place in six urban greenspaces in Oxford - Rainforest Glasshouse at the Oxford Botanic Gardens, Oxford Botanic Garden Outdoors, University Parks Dense, University Parks Open, Warneford Meadow, and Wytham Woods. Using our protocol, this work looked to assess two main hypotheses: (i) that a sites location (and ecology) is a significant driver of volatile difference enabling the separation of greenspaces by chemodiversity, and (ii) that by assessing presence and abundance of both biogenic volatiles (bVOCs) and anthropogenic volatiles (aVOCs), predictions could be made on which greenspaces are more healthy/unhealthy for human recreation.

VOC diversity in ambient air is known to be vast and highly location-dependent, often comprising thousands of compounds. Many air sampling studies have historically employed targeted approaches that significantly limit the number of compounds identified. For instance, previous studies have used between 15 and 66 VOCs across various forest and urban environments (2023, 2025, 2024). In contrast, our study adopted an untargeted metabolomics-style approach commonly used in biological matrices to enhance the depth and resolution of VOC profiling (Fernie et al., 2011, Van den Berg et al., 2006, Rosenthal et al., 2024) to confidently identify 175 unique compounds (Table S1) across these sampling sites. This improved identification capacity is likely attributable to several key enhancements in our analytical pipeline: greater technical replication, the use of a Kováts retention index, a more focused (ChEBI-based) compound library, a slow GC temperature ramp, and AMDIS deconvolution for resolving remaining co-eluting peaks.

Using all 175 identified compounds we can conclude that site location is a significant driver of volatile difference. This mirrors previous work who have used a PCA of volatiles to explain up to 70% of the variance between rural and urban sites in other studies (2023, Pripdeevech et al., 2025, Hewitt et al., 2022). Further PLS-DA and VIP analysis showed that terpenes/terpenoids and aromatic compounds exerted a dominant influence on both axes, consistent with sampling in both a plant dense

378 greenhouse, and outdoor urban areas (Pugliese et al., 2023, Jones et al., 2011, Cao et al., 2022, Wang

379 et al., 2025, Janson, 1993, Noe et al., 2012).

3.1 The role of aVOCs in site separation

To look for more precise trends in our data we employed HCA analysis. This highlighted the

separation of a major clade (Figure 4, Clade A), due to the relative proportion of aVOCs present

between our sampling sites. Specifically, two outdoor sites - Botanical Gardens Outdoors and

University Parks Open – separated from the other 4 sites based on containing significantly higher

relative amounts of the traffic-related BTEX compounds (Figure 8). Botanical Gardens Outdoors (45%

BTEX) and University Parks Open (40% BTEX) are very close to a busy city centre roads (Figure 1)

suggesting that traffic is the main contributor. This conclusion is strengthened by measurements

from the other outdoor sites Wytham Woods and Warneford Meadow. These sites are more isolated

from traffic and had much lower relative amounts of BTEX presence (14% and 13% respectively),

aligning with other published results Hoque et al (2008).

This trend is particularly important given the well-documented negative health effects of BTEX

compounds, even at relatively low concentrations. For example, chronic toluene inhalation is

associated with cognitive deficits in learning, memory and executive function, and can lead to white-

matter leukoencephalopathy (central nervous system damage) in humans and animal models

(Woodward and Braunscheidel, 2023, Thetkathuek et al., 2015). Chronic occupational exposure to

xylene vapor has been shown to impair pulmonary function - causing respiratory irritation, decreased

lung capacity - and also contribute to neurological symptoms including memory and concentration

deficits (Prevention, 2007). Interestingly, despite its proximity to University Parks Open (~80 m),

University Parks Dense showed markedly lower BTEX levels (41% to 28%). Upmanis et al. (2001) also

reported that concentrations of benzene and toluene were up to one-third lower just 40m into urban

parkland within two large Scandinavian cities. A third study also showed decreases in particulate

matter (PM_{10} and $PM_{2.5}$) of up to 50% at a distance of 200 m inside urban parkland in Madrid. One

possible explanation is the presence of dense localised tree cover at University Parks Dense

compared to the Open site (Table 1, 68% compared to 0% within a 10m radius). Previous studies

have shown that tree canopy structure and urban geometry can significantly influence VOC retention

and diffusion. For example, Jeanjean et al (Jeanjean et al., 2016) showed trees had a significant effect

on reducing particulate matter spread at wind speeds over 2m/s – it is likely a similar buffering of

localised volatiles is happening in our tree dense sampling sites.

Another contributing factor may be the presence of exposed soil and leaf litter layer in University

Parks Dense compared to the grassy University Parks Open site. Soil has been shown to increase

degradation of BTEX compounds through microbial metabolism. A 2012 study, for example, showed

that by adding a mixed bacterial consortium to soil, BTEX compounds were more rapidly degraded

(Mukherjee and Bordoloi, 2012). Additionally, degradation improved when supplemented with

phosphorous and nitrogen - the soil at our University Parks Dense site is likely more organically rich

due to the presence of a higher diversity of plants and as such this may be increasing microbial

metabolism.

3.2   The role of bVOCs in site separation

The second major separation of our sites using our HCA analysis (Figure 4, Clade B) was due to

biogenic volatile presence – particularly plant emitted terpenes such as beta-ocimene, sabinene, and

beta-pinene found in the OxBG Rainforest Glasshouse compared with the other 3 outdoor sites. In

total across all sampling sites, we identified 34 unique terpenes/terpenoids. This is significantly

higher than any studies on ambient air so far published (Walker et al., 2023, Sanaei et al., 2023, Mula

et al., 2024). The Rainforest Glasshouse contained the highest number of terpenes (27) with

considerable spread across the outdoor sampling (10 to 19). The specificity and spread of these

compounds may allow us to be more specific about which cited health benefits could be acquired

from which locations. For example, over all sampling sites we found a ubiquitous presence of Alpha-

pinene, Limonene, 3-Carene, Camphene, Camphor and 1-8 cineole - compounds which are

consistently found in air samples elsewhere using similar methods (Gu et al., 2024, Bryant et al.,

2022, Walker et al., 2023, Hellén et al., 2024, Jones et al., 2011, Hakola et al., 2003, Detournay et al.,

2011). These compounds alone are cited to have anti-inflammatory, anxiolytic, antimicrobial,

immune-boosting, antidepressant, anaesthetic, gastroprotective, antioxidant, anti-cancer (*in vitro*),

immune-modulatory, sedative, hypolipidemic, analgesic, decongestant, bronchodilatory, mucolytic,

and cognitive enhancing effects (Lee et al., 2022, Vallianou and Hadzopoulou-Cladaras, 2016, Woo et

al., 2019, Sun, 2007, Salehi et al., 2019, Jo et al., 2021, Oliveira et al., 2024). It could be argued,

therefore, that spending time in any of these sites could impart significant health gains. However, the

relationships between these compounds and their benefits are likely concentration dependent,

therefore further work is needed to quantify these compounds with the aim of making precise

recommendations such as when to visit specific greenspaces and how long to spend in these

locations to maximise benefits.

3.3   The combination of aVOCs and bVOCs

As discussed above, our sampling sites showed mixed profiles of volatiles suggesting they are shaped

both by biogenic and anthropogenic factors. The result that all our Oxford based sampling locations

contain terpenes and terpenoids is suggestive that recreational time in all sites would provide

benefits to human health, however, the presence of aVOCs mean the opposite could be argued.

Using aVOCs or bVOCs to make conclusions about our greenspaces is clearly a useful metric,

however, if we are to establish links to healthy spaces, both must be accounted for. Perhaps a more

useful metric to separate our sites is a ratio of biogenic to anthropogenic compounds. An example of

this can be found in study which assessed air over Covid lockdown in 2020 at an urban site in India

(Malik et al., 2025). This found a sharp increase in the $\sum\alpha+\beta$-pinene/$\sum$BTEX ratios during the strict

lockdown phase and hypothesised that this was due to a reduction in traffic related activity. We

mirrored this study and found that, on a particular day, the Botanical Gardens Outdoors and

University Parks Open had significantly lower terpene/BTEX ratio compared to the other sampling

sites suggesting that these were the least unhealthiest spaces (Figure 8).

Interestingly, aVOCs were found to be significantly higher in indoor sites than outdoor sites in the

vicinity of an optoelectronics industrial park in 2006–2007 (Chang et al., 2019). This result shows that

indoor spaces likely amplify those compounds found in the outdoor air and that simply being inside is

little protection from external sources of pollution. In our work, the indoor OxBG Rainforest

Glasshouse has a much higher ratio between terpenes and BTEX compounds (5.2) when compared to

the other sites (all below 0.5). This glasshouse is plant rich and enclosed and as such likely acts as an

enclosure for those VOCs produced within, suggesting that plant rich greenhouses may be the most

beneficial place to spend time. The result that BTEX compounds represented a smaller amount of the

total volatiles in the glasshouse may partly be due to restricting the entry of pollutant volatiles, but

also due to breakdown by plants and microbes (Mukherjee and Bordoloi, 2012) as well as a relative

dilution due to the high amounts of other biogenic volatiles present.

3.4   Correlating ecology using volatile diversity

The third most significant separation of our sites based on our HCA analysis (Figure 4, Clade C)

showed that the two outdoor sites closest to busy roads (Botanical Gardens Outdoors and University

Parks Open) are further separated by their relative amounts of both biogenic (e.g. 2-thujene, p-

cymene) and anthropogenic (e.g. benzene derivatives) volatiles. Faster terpene decay occurs in urban

greenspaces due to the increased presence of highly reactive pollutants (Luo et al., 2025, Salvador et

al., 2020), so the presence of such a high number of terpenes in the Botanical Gardens Outdoors site

is suggestive of a plant rich environment. One of the hypotheses of this paper was to test if site

ecology has a significant baring on its volatile profile. By carrying out plant surveys across our sites

we can correlate the presence of health related compounds with plant diversity. Table 1 shows a

summary of the total and unique number of species counts in our 6 sampling locations where the

Botanical Gardens Outdoors (43 species) site contains significantly more plant species than the other

outdoor sampling sites (7 - 21 species).

These findings highlight the potential of site-specific volatile profiles - particularly terpenes - as a

basis for mapping biodiversity. Notably, the Rainforest Glasshouse emitted the highest number of

terpenes unique to a single site (13), while the Botanical Gardens Outdoors site was again the most

terpene-rich open-air location, containing five distinct compounds (2-thujene, bornane, prehnitene,

verbenone, and farnesane). Among the remaining sites, only Wytham Woods emitted a uniquely

identified terpene - α-thujone. Future research will seek to trace the biological sources of these

distinctive volatiles, advancing efforts to develop non-invasive, scent-based indicators of ecological

diversity.

4 Author Contributions

William Kay was responsible for methods development, data collection and all data analyses. William

Kay and Anya Lindstrom Battle were both involved in the writing and editing of the manuscript. Kathy

Willis conceived of the initial ideas and methodologies and was also involved in the writing and

editing of the manuscript. Mahal Humberstone, Molly Tucker and Kieran Storer all contributed to in-

field sample collection and the vegetation surveys. Geoff Kite was involved in the collection and

collation of data via GCMS, as well as providing useful guidance. All authors contributed critically to

the drafts and gave final approval for publication.

5 Funding

495 This work was funded by the Leverhulme Centre for Nature Recovery, made possible thanks to the

496 generous support of the Leverhulme Trust - Grant number RC-2021-76.

6 Acknowledgments

498 We would firstly like to thank the University of Oxford Botanic Garden and Wytham Woods for

499 permission to sample on their sites. We would also like to thank Dr Ben Martin who provided helpful

500 advice for sampling and analysis, and Dr Luciana Carvalho who always had time to share thoughts

501 and ideas relating to the project. Finally, we thank Dr Pedro Bota for his generosity of availability and

502 equipment.

7 Data Availability Statement

504 Raw AGILENT files, AMDIS outputs, Vegetation Surveys, and final R scripts used to analyse data are

505 available here: https://unioxfordnexus-

506 my.sharepoint.com/:f:/r/personal/dops0827_ox_ac_uk/Documents/KW_GCMS_Oxford/_Papers/Pap

507 er%201%20-%20Kay%20et%20al.%20A%20Walk%20in%20the%20Park?csf=1&web=1&e=Z9CtQU

8 Conflicts of Interest

509 The authors declare no conflicts of interest.

9   References

Amann, A., de Lacy Costello, B., Miekisch, W., Schubert, J., Buszewski, B., Pleil, J., Ratcliffe, N. & Risby, T. 2014. The human volatilome: volatile organic compounds (VOCs) in exhaled breath, skin emanations, urine, feces and saliva. *Journal of breath research,* 8**,** 034001. https://doi.org/10.1088/1752-7155/8/3/034001.

Antonelli, M., Donelli, D., Barbieri, G., Valussi, M., Maggini, V. & Firenzuoli, F. 2020. Forest volatile organic compounds and their effects on human health: A state-of-the-art review. *International Journal of Environmental Research and Public Health,* 17**,** 6506. https://doi.org/10.3390/ijerph17186506.

Bartwal, A., Mall, R., Lohani, P., Guru, S. & Arora, S. 2013. Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. *Journal of plant growth regulation,* 32**,** 216-232. https://doi.org/10.1007/s00344-012-9272-x.

Ben Ammar, R. 2023. Potential effects of geraniol on cancer and inflammation-related diseases: A review of the recent research findings. *Molecules,* 28**,** 3669. https://doi.org/10.3390/molecules28093669.

Bergman, M. E., Huang, X.-Q., Baudino, S., Caissard, J.-C. & Dudareva, N. 2025. Plant volatile organic compounds: Emission and perception in a changing world. *Current Opinion in Plant Biology,* 85**,** 102706. https://doi.org/10.1016/j.pbi.2025.102706.

Boncan, D. A. T., Tsang, S. S., Li, C., Lee, I. H., Lam, H.-M., Chan, T.-F. & Hui, J. H. 2020. Terpenes and terpenoids in plants: Interactions with environment and insects. *International Journal of Molecular Sciences,* 21**,** 7382. https://doi.org/10.3390/ijms21197382.

Brambilla, A., Sommer, A., Ghirardo, A., Wenig, M., Knappe, C., Weber, B., Amesmaier, M., Lenk, M., Schnitzler, J.-P. & Vlot, A. C. 2022. Immunity-associated volatile emissions of β-ionone and nonanal propagate defence responses in neighbouring barley plants. *Journal of Experimental Botany,* 73**,** 615-630. https://doi.org/10.1093/jxb/erab520.

Bratman, G. N., Daily, G. C., Levy, B. J. & Gross, J. J. 2015. The benefits of nature experience: Improved affect and cognition. *Landscape and urban planning,* 138**,** 41-50. https://doi.org/10.1016/j.landurbplan.2015.02.005.

Bryant, D. J., Nelson, B. S., Swift, S. J., Budisulistiorini, S. H., Drysdale, W. S., Vaughan, A. R., Newland, M. J., Hopkins, J. R., Cash, J. M. & Langford, B. 2022. Biogenic and anthropogenic sources of isoprene and monoterpenes and their secondary organic aerosol in Delhi, India. *Atmospheric Chemistry and Physics Discussions,* 2022**,** 1-35. https://doi.org/10.5194/acp-23-61-2023.

Cao, J., Situ, S., Hao, Y., Xie, S. & Li, L. 2022. Enhanced summertime ozone and SOA from biogenic
volatile organic compound (BVOC) emissions due to vegetation biomass variability during
1981–2018 in China. *Atmos. Chem. Phys.,* 22**,** 2351-2364. https://doi.org/10.5194/acp-22-
2351-2022.

Chang, C.-C. & Lin, C.-J. 2011. LIBSVM: a library for support vector machines. *ACM transactions on*
intelligent systems and technology (TIST), 2**,** 1-27.
https://doi.org/10.1145/1961189.1961199.

Chang, T.-Y., Liu, C.-L., Huang, K.-H. & Kuo, H.-W. 2019. Indoor and Outdoor Exposure to Volatile
Organic Compounds and Health Risk Assessment in Residents Living near an Optoelectronics
Industrial Park. *Atmosphere,* 10**,** 380. https://doi.org/10.3390/atmos10070380.

de Carvalho Couto, C., Chávez, D. W. H., Oliveira, E. M. M., Freitas-Silva, O. & Casal, S. 2024. SPME-
GC-MS untargeted metabolomics approach to identify potential volatile compounds as
markers for fraud detection in roasted and ground coffee. *Food Chemistry,* 446**,** 138862.
https://doi.org/10.1016/j.foodchem.2024.138862.

Degtyarenko, K., De Matos, P., Ennis, M., Hastings, J., Zbinden, M., McNaught, A., Alcántara, R.,
Darsow, M., Guedj, M. & Ashburner, M. 2007. ChEBI: a database and ontology for chemical
entities of biological interest. *Nucleic acids research,* 36**,** D344-D350.
https://doi.org/10.1093/nar/gkm791.

Del Prado-Audelo, M. L., Cortés, H., Caballero-Florán, I. H., González-Torres, M., Escutia-Guadarrama,
L., Bernal-Chávez, S. A., Giraldo-Gomez, D. M., Magaña, J. J. & Leyva-Gómez, G. 2021.
Therapeutic Applications of Terpenes on Inflammatory Diseases. *Front Pharmacol,* 12**,**
704197. https://doi.org/10.3389/fphar.2021.704197.

Detournay, A., Sauvage, S., Locoge, N., Gaudion, V., Leonardis, T., Fronval, I., Kaluzny, P. & Galloo, J.-
C. 2011. Development of a sampling method for the simultaneous monitoring of straight-
chain alkanes, straight-chain saturated carbonyl compounds and monoterpenes in remote
areas. *Journal of Environmental Monitoring,* 13**,** 983-990.
https://doi.org/10.1039/c0em00354a.

Farimani, F. D., Hosseini, M., Amirahmadi, S., Akbarian, M., Shirazinia, M., Barabady, M. & Rajabian,
A. 2024. Cedrol supplementation ameliorates memory deficits by regulating neuro-
inflammation and cholinergic function in lipopolysaccharide–induced cognitive impairment in
rats. *Heliyon,* 10https://doi.org/10.1016/j.heliyon.2024.e30356.

Fazakas, E., Neamtiu, I. A. & Gurzau, E. S. 2024. Health effects of air pollutant mixtures (volatile
organic compounds, particulate matter, sulfur and nitrogen oxides) – a review of the
literature. *Reviews on Environmental Health,* 39**,** 459-478. https://doi.org/10.1515/reveh-
2022-0252.

Fernie, A. R., Aharoni, A., Willmitzer, L., Stitt, M., Tohge, T., Kopka, J., Carroll, A. J., Saito, K., Fraser, P.
D. & DeLuca, V. 2011. Recommendations for reporting metabolite data. *The Plant Cell,* 23**,**
2477-2482. https://doi.org/10.1105/tpc.111.086272.

Gao, X., Wang, Y., Wu, L., Zheng, F., Sun, N., Liu, G., Liu, Y., Meng, P., Sun, L. & Jing, B. 2023. The
Impact of Anthropogenic VOC Emissions on Atmospheric Pollution: A Case Study of a Typical
Industrialized Area in China. *Atmosphere,* 14**,** 1586. https://doi.org/10.3390/atmos14101586.

GOV.UK. 2023. Road traffic statistics - Local authority Oxfordshire. Available:
https://roadtraffic.dft.gov.uk/local-authorities/142.

Gu, S., Luo, W., Charmchi, A., McWhirter, K. J., Rosenstiel, T., Pankow, J. & Faiola, C. L. 2024.
Limonene Enantiomeric Ratios from Anthropogenic and Biogenic Emission Sources. *Environ*
Sci Technol Lett, 11**,** 130-135. https://doi.org/10.1021/acs.estlett.3c00794.

Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau,
M. & McKay, W. 1995. A global model of natural volatile organic compound emissions.
Journal of Geophysical Research: Atmospheres, 100**,** 8873-8892.
https://doi.org/10.1029/94JD02950.

Guenther, A., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T. a., Emmons, L. & Wang, X. 2012.
The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2. 1): an
extended and updated framework for modeling biogenic emissions. *Geoscientific Model*
Development, 5**,** 1471-1492. https://doi.org/10.5194/gmd-5-1471-2012.

Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K. & Fall, R. 1993. Isoprene and
monoterpene emission rate variability: model evaluations and sensitivity analyses. *Journal of*
Geophysical Research: Atmospheres, 98**,** 12609-12617. https://doi.org/10.1029/93JD00527.

Hakola, H., Tarvainen, V., Laurila, T., Hiltunen, V., Hellén, H. & Keronen, P. 2003. Seasonal variation of
VOC concentrations above a boreal coniferous forest. *Atmospheric Environment,* 37**,** 1623-
1634. https://doi.org/10.1016/S1352-2310(03)00014-1.

Hammerbacher, A., Coutinho, T. A. & Gershenzon, J. 2019. Roles of plant volatiles in defence against
microbial pathogens and microbial exploitation of volatiles. *Plant, Cell & Environment,* 42**,**
2827-2843. https://doi.org/10.1111/pce.13602.

He, Y., Qiu, H., Wang, W., Lin, Y. & Ho, K. F. 2024. Exposure to BTEX is associated with cardiovascular
disease, dyslipidemia and leukocytosis in national US population. *Science of The Total*
Environment, 919**,** 170639. https://doi.org/10.1016/j.scitotenv.2024.170639.

Heil, M. & Ton, J. 2008. Long-distance signalling in plant defence. *Trends in Plant Science,* 13**,** 264-
272. https://doi.org/10.1016/j.tplants.2008.03.005.

Hellén, H., Tykkä, T., Schallhart, S., Stratigou, E., Salameh, T. & Iturrate-Garcia, M. 2024.
Measurements of atmospheric C 10–C 15 biogenic volatile organic compounds (BVOCs) with

sorbent tubes. *Atmospheric Measurement Techniques,* 17**,** 315-333.

https://doi.org/10.5194/amt-17-315-2024.

Hewitt, M. J., Belluomo, I., Zuffa, S., Boshier, P. R. & Myridakis, A. 2022. Variation of volatile organic

compound levels within ambient room air and its impact upon the standardisation of breath

sampling. *Scientific Reports,* 12**,** 15887. https://doi.org/10.1038/s41598-022-20365-7.

Hoque, R. R., Khillare, P. S., Agarwal, T., Shridhar, V. & Balachandran, S. 2008. Spatial and temporal

variation of BTEX in the urban atmosphere of Delhi, India. *Science of The Total Environment,*

392**,** 30-40. https://doi.org/10.1016/j.scitotenv.2007.08.036.

Janson, R. W. 1993. Monoterpene emissions from Scots pine and Norwegian spruce. *Journal of*

geophysical research: atmospheres, 98**,** 2839-2850. https://doi.org/10.1029/92JD02394.

Jeanjean, A. P. R., Monks, P. S. & Leigh, R. J. 2016. Modelling the effectiveness of urban trees and

grass on PM2.5 reduction via dispersion and deposition at a city scale. *Atmospheric*

Environment, 147**,** 1-10. https://doi.org/10.1016/j.atmosenv.2016.09.033.

Jo, H., Cha, B., Kim, H., Brito, S., Kwak, B. M., Kim, S. T., Bin, B. H. & Lee, M. G. 2021. Alpha-pinene

enhances the anticancer activity of natural killer cells via ERK/AKT Pathway. *Int J Mol Sci,*

22https://doi.org/10.3390/ijms22020656.

Jones, C. E., Hopkins, J. & Lewis, A. 2011. In situ measurements of isoprene and monoterpenes within

a south-east Asian tropical rainforest. *Atmospheric chemistry and Physics,* 11**,** 6971-6984.

https://doi.org/10.5194/acp-11-6971-2011.

Kassambara, A. & Mundt, F. 2017. Package 'factoextra'. *Extract and visualize the results of*

multivariate data analyses [Online], 76. Available: https://rpkgs.datanovia.com/factoextra/.

Kessler, D., Bing, J., Haverkamp, A. & Baldwin, I. T. 2019. The defensive function of a pollinator-

attracting floral volatile. *Functional Ecology,* 33**,** 1223-1232. https://doi.org/10.1111/1365-

2435.13332.

Kim, T., Song, B., Cho, K. S. & Lee, I. S. 2020. Therapeutic potential of volatile terpenes and terpenoids

from forests for inflammatory diseases. *Int J Mol Sci,*

21https://doi.org/10.3390/ijms21062187.

Kovats, v. E. 1958. Gas-chromatographische charakterisierung organischer verbindungen. Teil 1:

retentionsindices aliphatischer halogenide, alkohole, aldehyde und ketone. *Helvetica Chimica*

Acta, 41**,** 1915-1932. https://doi.org/10.1002/hlca.19580410703.

Kunming-Montreal. 2022. Kunming-Montreal Global Biodiversity Framework: Targets. *Convention on*

Biological Diversity [Online]. Available: https://www.cbd.int/gbf/targets [Accessed 2025].

Lahr, E. C., Schade, G. W., Crossett, C. C. & Watson, M. R. 2015. Photosynthesis and isoprene

emission from trees along an urban–rural gradient in Texas. *Global Change Biology,* 21**,** 4221-

4236. https://doi.org/10.1111/gcb.13010.

Lê, S., Josse, J. & Husson, F. 2008. FactoMineR: an R package for multivariate analysis. *Journal of*
statistical software, 25**,** 1-18. https://doi.org/10.18637/jss.v025.i01.

Lee, S. H., Kim, D. S., Park, S. H. & Park, H. 2022. Phytochemistry and Applications of Cinnamomum
camphora Essential Oils. *Molecules,* 27https://doi.org/10.3390/molecules27092695.

Legendre, P. & Legendre, L. 2012. *Numerical ecology*, Elsevier.

Li, Q. 2010. Effect of forest bathing trips on human immune function. *Environmental health and*
preventive medicine, 15**,** 9-17. https://doi.org/10.1007/s12199-008-0068-3.

Liu, X., Zhang, X., Dufresne, M., Wang, T., Wu, L., Lara, R., Seco, R., Monge, M., Yáñez-Serrano, A. M.,
Gohy, M., Petit, P., Chevalier, A., Vagnot, M. P., Fortier, Y., Baudic, A., Ghersi, V., Gille, G.,
Lanzi, L., Gros, V., Simon, L., Héllen, H., Reimann, S., Le Bras, Z., Müller, M. J., Beddows, D.,
Hou, S., Shi, Z., Harrison, R. M., Bloss, W., Dernie, J., Sauvage, S., Hopke, P. K., Duan, X., An,
T., Lewis, A. C., Hopkins, J. R., Liakakou, E., Mihalopoulos, N., Zhang, X., Alastuey, A., Querol,
X. & Salameh, T. 2025. Measurement report: Exploring the variations in ambient BTEX in
urban Europe and their environmental health implications. *Atmos. Chem. Phys.,* 25**,** 625-638.
https://doi.org/10.5194/acp-25-625-2025.

Luo, R., Lun, X., Gao, R., Wang, L., Yang, Y., Su, X., Habibullah-Al-Mamun, M., Xu, X., Li, H. & Li, J.
2025. A Review of Biogenic Volatile Organic Compounds from Plants: Research Progress and
Future Prospects. *Toxics,* 13**,** 364. https://doi.org/10.3390/toxics13050364.

Ma, J., Wang, S., Chen, G., Zhu, S., Wang, P., Chen, J. & Zhang, H. 2025. Estimating emissions of
biogenic volatile organic compounds from urban green spaces and their contributions to
secondary pollution. *Environmental Science: Atmospheres,* 5**,** 129-141.
https://doi.org/10.1039/d4ea00099d.

Malik, T. G., Gupta, M., Tripathi, N. & Sahu, L. K. 2025. Change in monoterpene concentrations during
winter-to-summer transition period and impact of COVID-19 lockdown at an urban site in
India. *Atmospheric Environment,* 350**,** 121141.
https://doi.org/10.1016/j.atmosenv.2025.121141.

Masyita, A., Mustika Sari, R., Dwi Astuti, A., Yasir, B., Rahma Rumata, N., Emran, T. B., Nainu, F. &
Simal-Gandara, J. 2022. Terpenes and terpenoids as main bioactive compounds of essential
oils, their roles in human health and potential application as natural food preservatives. *Food*
Chem X, 13**,** 100217. https://doi.org/10.1016/j.fochx.2022.100217.

McDonald, B. C., de Gouw, J. A., Gilman, J. B., Jathar, S. H., Akherati, A., Cappa, C. D., Jimenez, J. L.,
Lee-Taylor, J., Hayes, P. L., McKeen, S. A., Cui, Y. Y., Kim, S.-W., Gentner, D. R., Isaacman-
VanWertz, G., Goldstein, A. H., Harley, R. A., Frost, G. J., Roberts, J. M., Ryerson, T. B. &
Trainer, M. 2018. Volatile chemical products emerging as largest petrochemical source of
urban organic emissions. *Science,* 359**,** 760-764. https://doi.org/10.1126/science.aaq0524.

Muda, I., Mohammadi, M. J., Sepahvad, A., Farhadi, A., Obaid, R. F., Taherian, M., Alali, N.,
Chowdhury, S. & Farhadi, M. 2024. Associated health risk assessment due to exposure to
BTEX compounds in fuel station workers. *Reviews on Environmental Health,* 39**,** 435-446.
https://doi.org/10.1515/reveh-2023-0012.

Mukherjee, A. K. & Bordoloi, N. K. 2012. Biodegradation of benzene, toluene, and xylene (BTX) in
liquid culture and in soil by Bacillus subtilis and Pseudomonas aeruginosa strains and a
formulated bacterial consortium. *Environ Sci Pollut Res Int,* 19**,** 3380-8.
https://doi.org/10.1007/s11356-012-0862-8.

Mula, V., Bogdanov, J., Stanoeva, J. P., Zeneli, L. & Zdravkovski, Z. 2024. Monitoring Volatile Organic
Compounds in Air Using Passive Sampling: Regional Cross-Border Study between N.
Macedonia and Kosovo. *Aerosol and Air Quality Research,* 24**,** 230170.
https://doi.org/10.4209/aaqr.230170.

Müller, A., Faubert, P., Hagen, M., Zu Castell, W., Polle, A., Schnitzler, J.-P. & Rosenkranz, M. 2013.
Volatile profiles of fungi–chemotyping of species and ecological functions. *Fungal genetics*
and biology, 54**,** 25-33. https://doi.org/10.1016/j.fgb.2013.02.005.

Nguyen, P.-Y., Astell-Burt, T., Rahimi-Ardabili, H. & Feng, X. 2023. Effect of nature prescriptions on
cardiometabolic and mental health, and physical activity: a systematic review. *The Lancet*
Planetary Health, 7**,** e313-e328. https://doi.org/10.1016/s2542-5196(23)00025-6.

Noe, S. M., Hüve, K., Niinemets, Ü. & Copolovici, L. 2012. Seasonal variation in vertical volatile
compounds air concentrations within a remote hemiboreal mixed forest. *Atmos. Chem.*
Phys., 12**,** 3909-3926. https://doi.org/10.5194/acp-12-3909-2012.

Oksanen, J. 2015. Vegan: community ecology package. *R package version* [Online], 2. Available:
https://vegandevs.github.io/vegan/.

Oliveira, C. D., Cardoso, M. d. G., Batista, L. R., Alves, E., Rosa, M. B. P., Ferreira, V. R. F., de Souza, L.,
Pineda, M., Fernandes, A. I. & Nelson, D. L. 2024. The antibacterial, antioxidant, and
insecticidal activities of essential oils from Thymus vulgaris L., Salvia officinalis L., and
Ocimum basilicum L. *Journal of Food Safety,* 44**,** e13145. https://doi.org/10.1111/jfs.13145.

Park, B. J., Tsunetsugu, Y., Kasetani, T., Kagawa, T. & Miyazaki, Y. 2010. The physiological effects of
Shinrin-yoku (taking in the forest atmosphere or forest bathing): evidence from field
experiments in 24 forests across Japan. *Environmental health and preventive medicine,* 15**,**
18-26. https://doi.org/10.1007/s12199-009-0086-9.

Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., Piechulla, B. & Schnitzler, J. 2014.
Biogenic volatile emissions from the soil. *Plant, cell & environment,* 37**,** 1866-1891.
https://doi.org/10.1111/pce.12340.

Peñuelas, J. & Munné-Bosch, S. 2005. Isoprenoids: an evolutionary pool for photoprotection. *Trends*
in plant science, 10**,** 166-169. https://doi.org/10.1016/j.tplants.2005.02.005.

Prevention, C. f. D. C. a. 2007. Toxicological Profile for Xylene. Available:
https://www.atsdr.cdc.gov/toxprofiles/tp71-c2.pdf.

Pripdeevech, P., Janta, R., Sripahco, T., Meesang, W., Aiyathiti, C., Prabamroong, T., Mahatheeranont,
S., Poshyachinda, S., Pongpiachan, S. & Khruengsai, S. 2025. Seasonal volatile organic
compound dynamics in urban and forest environments in Thailand: Implications for air
quality and secondary pollutants. *Environmental Pollution,* 367**,** 125565.
https://doi.org/10.1016/j.envpol.2024.125565.

Pugliese, G., Ingrisch, J., Meredith, L. K., Pfannerstill, E. Y., Klüpfel, T., Meeran, K., Byron, J., Purser, G.,
Gil-Loaiza, J. & Van Haren, J. 2023. Effects of drought and recovery on soil volatile organic
compound fluxes in an experimental rainforest. *nature communications,* 14**,** 5064.
https://doi.org/10.1038/s41467-023-40661-8.

Ren, Y., Qu, Z., Du, Y., Xu, R., Ma, D., Yang, G., Shi, Y., Fan, X., Tani, A., Guo, P., Ge, Y. & Chang, J.
2017. Air quality and health effects of biogenic volatile organic compounds emissions from
urban green spaces and the mitigation strategies. *Environmental Pollution,* 230**,** 849-861.
https://doi.org/10.1016/j.envpol.2017.06.049.

Reyrolle, M., Desauziers, V., Pigot, T., Gautier, L. & Le Bechec, M. 2024. Comparison of Untargeted
and Markers Analysis of Volatile Organic Compounds with SIFT-MS and SPME-GC-MS to
Assess Tea Traceability. *Foods,* 13**,** 3996. https://doi.org/10.3390/foods13243996.

Roberts, S. C. 2007. Production and engineering of terpenoids in plant cell culture. *Nature chemical*
biology, 3**,** 387-395. https://doi.org/10.1038/nchembio.2007.8.

Rohart, F., Gautier, B., Singh, A. & Lê Cao, K.-A. 2017. mixOmics: An R package for 'omics feature
selection and multiple data integration. *PLoS computational biology,* 13**,** e1005752.
https://doi.org/10.1371/journal.pcbi.1005752.

Rosenthal, K., Lindley, M., Turner, M., Ratcliffe, E. & Hunsicker, E. 2024. Current data processing
methods and reporting standards for untargeted analysis of volatile organic compounds
using direct mass spectrometry: a systematic review. *Metabolomics,* 20**,** 42.
https://doi.org/10.1007/s11306-024-02104-3.

Saeedi, M., Malekmohammadi, B. & Tajalli, S. 2024. Interaction of benzene, toluene, ethylbenzene,
and xylene with human's body: Insights into characteristics, sources and health risks. *Journal*
of Hazardous Materials Advances, 16**,** 100459.
https://doi.org/10.1016/j.hazadv.2024.100459.

Saenen, N. D., Nawrot, T. S., Hautekiet, P., Wang, C., Roels, H. A., Dadvand, P., Plusquin, M. & Bijnens,
E. M. 2023. Residential green space improves cognitive performances in primary

schoolchildren independent of traffic-related air pollution exposure. *Environmental Health,* 22**,** 33. https://doi.org/10.1186/s12940-023-00982-z.

Salehi, B., Upadhyay, S., Erdogan Orhan, I., Kumar Jugran, A., LD Jayaweera, S., A. Dias, D., Sharopov, F., Taheri, Y., Martins, N. & Baghalpour, N. 2019. Therapeutic potential of α-and β-pinene: A miracle gift of nature. *Biomolecules,* 9**,** 738. https://doi.org/10.3390/biom9110738.

Salvador, C. M., Chou, C. C.-K., Ho, T.-T., Tsai, C.-Y., Tsao, T.-M., Tsai, M.-J. & Su, T.-C. 2020. Contribution of Terpenes to Ozone Formation and Secondary Organic Aerosols in a Subtropical Forest Impacted by Urban Pollution. *Atmosphere,* 11**,** 1232. https://doi.org/10.3390/atmos11111232.

Sanaei, A., Herrmann, H., Alshaabi, L., Beck, J., Ferlian, O., Fomba, K. W., Haferkorn, S., van Pinxteren, M., Quaas, J. & Quosh, J. 2023. Changes in biodiversity impact atmospheric chemistry and climate through plant volatiles and particles. *Communications Earth & Environment,* 4**,** 445. https://doi.org/10.1038/s43247-023-01113-9.

Seco, R., Karl, T., Guenther, A., Hosman, K. P., Pallardy, S. G., Gu, L., Geron, C., Harley, P. & Kim, S. 2015. Ecosystem-scale volatile organic compound fluxes during an extreme drought in a broadleaf temperate forest of the Missouri Ozarks (central USA). *Global Change Biology,* 21**,** 3657-3674. https://doi.org/10.1111/gcb.12980.

Shibata, S. & Suzuki, N. 2002. Effects of the foliage plant on task performance and mood. *Journal of environmental psychology,* 22**,** 265-272. http://dx.doi.org/10.1006/jevp.2002.0232.

Siddiqui, T., Khan, M. U., Sharma, V. & Gupta, K. 2024. Terpenoids in essential oils: Chemistry, classification, and potential impact on human health and industry. *Phytomedicine Plus,* 4**,** 100549. https://doi.org/10.1016/j.phyplu.2024.100549.

Sousa, F. S. S., Birmann, P. T., Balaguez, R., Alves, D., Brüning, C. A. & Savegnago, L. 2018. α-(phenylselanyl) acetophenone abolishes acute restraint stress induced-comorbid pain, depression and anxiety-related behaviors in mice. *Neurochemistry international,* 120**,** 112-120. https://doi.org/10.1016/j.neuint.2018.08.006.

Sun, J. 2007. D-Limonene: safety and clinical applications. *Alternative medicine review* [Online], 12. Available: https://www.anaturalhealingcenter.com/documents/Thorne/articles/Limonene12-3.pdf.

Thetkathuek, A., Jaidee, W., Saowakhontha, S. & Ekburanawat, W. 2015. Neuropsychological Symptoms among Workers Exposed to Toluene and Xylene in Two Paint Manufacturing Factories in Eastern Thailand. *Adv Prev Med,* 2015**,** 183728. https://doi.org/10.1155/2015/183728.

Upmanis, H., Eliasson, I. & Andersson-Sköld, Y. 2001. Case Studies of the Spatial Variation of Benzene
and Toluene Concentrations in Parks and Adjacent Built-up Areas. *Water, Air, and Soil*
Pollution, 129**,** 61-81. https://doi.org/10.1023/A:1010357914047.

Vallianou, I. & Hadzopoulou-Cladaras, M. 2016. Camphene, a Plant Derived Monoterpene, Exerts Its
Hypolipidemic Action by Affecting SREBP-1 and MTP Expression. *PloS one,* 11**,** e0147117.
https://doi.org/10.1371/journal.pone.0147117.

Van den Berg, R. A., Hoefsloot, H. C., Westerhuis, J. A., Smilde, A. K. & Van der Werf, M. J. 2006.
Centering, scaling, and transformations: improving the biological information content of
metabolomics data. *BMC genomics,* 7**,** 1-15. https://doi.org/10.1186/1471-2164-7-142.

Walker, H., Jena, A., McEwan, K., Evans, G. & Campbell, S. 2023. Natural volatile organic compounds
(NVOCs) are greater and more diverse in UK forests compared with a public garden. *Forests,*
14**,** 92. https://doi.org/10.3390/f14010092.

Wang, H., Li, Y., Liu, Y., Lu, X., Zhang, Y., Fan, Q., Shen, C., Lai, S., Zhou, Y., Zhang, T. & Yue, D. 2025.
Underappreciated contributions of biogenic volatile organic compounds from urban green
spaces to ozone pollution. *Atmos. Chem. Phys.,* 25**,** 5233-5250. https://doi.org/10.5194/acp-
25-5233-2025.

Weisskopf, L., Schulz, S. & Garbeva, P. 2021. Microbial volatile organic compounds in intra-kingdom
and inter-kingdom interactions. *Nature Reviews Microbiology,* 19**,** 391-404.
https://doi.org/10.1038/s41579-020-00508-1.

White, M. P., Alcock, I., Grellier, J., Wheeler, B. W., Hartig, T., Warber, S. L., Bone, A., Depledge, M. H.
& Fleming, L. E. 2019. Spending at least 120 minutes a week in nature is associated with good
health and wellbeing. *Scientific reports,* 9**,** 7730. https://doi.org/10.1038/s41598-019-44097-
3.

Wickham, H. 2011. ggplot2. *Wiley interdisciplinary reviews: computational statistics,* 3**,** 180-185.

Woo, J., Yang, H., Yoon, M., Gadhe, C. G., Pae, A. N., Cho, S. & Lee, C. J. 2019. 3-Carene, a Phytoncide
from Pine Tree Has a Sleep-enhancing Effect by Targeting the GABA A benzodiazepine
Receptors. *Experimental Neurobiology,* 28**,** 593-601.
https://doi.org/10.5607/en.2019.28.5.593.

Woodward, J. J. & Braunscheidel, K. M. 2023. The Effects of the Inhalant Toluene on Cognitive
Function and Behavioral Flexibility: A Review of Recent Findings. *Addict Neurosci,*
5http://doi.org/10.1016/j.addicn.2022.100059.

Xu, B., Bai, L., Chen, L., Tong, R., Feng, Y. & Shi, J. 2022. Terpenoid natural products exert
neuroprotection via the PI3K/Akt pathway. *Frontiers in Pharmacology,* 13**,** 1036506.
https://doi.org/10.3389/fphar.2022.1036506.

Xu, H., Desurmont, G., Degen, T., Zhou, G., Laplanche, D., Henryk, L. & Turlings, T. C. 2017a.

Combined use of herbivore-induced plant volatiles and sex pheromones for mate location in

braconid parasitoids. *Plant, cell & environment,* 40**,** 330-339.

https://doi.org/10.1111/pce.12818.

Xu, H. & Turlings, T. C. 2018. Plant volatiles as mate-finding cues for insects. *Trends in Plant Science,*

23**,** 100-111. https://doi.org/10.1016/j.tplants.2017.11.004.

Xu, H., Veyrat, N., Degen, T. & Turlings, T. C. 2014. Exceptional use of sex pheromones by parasitoids

of the genus Cotesia: males are strongly attracted to virgin females, but are no longer

attracted to or even repelled by mated females. *Insects,* 5**,** 499-512.

https://doi.org/10.3390/insects5030499.

Xu, P., Wang, K., Lu, C., Dong, L., Gao, L., Yan, M., Aibai, S., Yang, Y. & Liu, X. 2017b. Protective effects

of linalool against amyloid beta-induced cognitive deficits and damages in mice. *Life Sciences,*

174**,** 21-27. https://doi.org/10.1016/j.lfs.2017.02.010.

Yin, X.-L., Fu, W.-J., Chen, Y., Zhou, R.-F., Sun, W., Ding, B., Peng, X.-T. & Gu, H.-W. 2022. GC-MS-

based untargeted metabolomics reveals the key volatile organic compounds for

discriminating grades of Yichang big-leaf green tea. *LWT,* 171**,** 114148.

https://doi.org/10.1016/j.lwt.2022.114148.

Zeng, W., Liu, H., Hou, S., Qiu, X., Chen, X., Liu, M., Wu, D. & Liu, L. 2022. Explore the Benefits of

Natural Air: New Insights from Field and Chamber Tests on Cognitive Performance.

Atmosphere, 13**,** 1006. https://doi.org/10.3390/atmos13071006.

839