Evidence for Genetic Nurture Effects on Substance Use

Authors: Mannan Luo*[1], Victória Trindade Pons[1], Nathan A. Gillespie[2], Hanna M. van Loo[1]

Affiliations:

[1] Department of Psychiatry, University Medical Center Groningen, University of Groningen,

Groningen, the Netherlands

[2] Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University,

Richmond, VA, USA

*Corresponding author

Address:

Hanzeplein 1

9713 GZ Groningen

Netherlands

Corresponding author:

Mannan Luo (m.luo@umcg.nl)

ABSTRACT

Substance use runs in families. Beyond genetic transmission, parental genetics can indirectly influence offspring substance use through the rearing environment, known as "genetic nurture". This study utilized transmitted and non-transmitted polygenic scores to investigate genetic nurture effects on tobacco, alcohol and cannabis use in up to 15,863 adults with at least one genotyped parent from the Lifelines cohort. Genetic nurture significantly influenced smoking quantity (cigarettes per day: β=0.03; pack-years: β=0.02), accounting for 18.8% and 28.6% of the corresponding effects of genetic transmission. However, it had minimal impact on tobacco or cannabis initiation, suggesting a stage-specific pattern. Maternal and paternal genetic nurture contributed equally to offspring smoking quantity, especially for pack-years. Mediation analyses revealed that both maternal and paternal smoking partially explained these effects, with higher mediation proportions observed for maternal smoking quantity. These findings highlight the importance of considering stage-specific and parent-specific effects when investigating genetic nurture in substance use.

INTRODUCTION

Substance use, including tobacco, alcohol and cannabis use, runs in families[1,2]. Parental substance use is one of the most salient familial risks for offspring's substance use behavior and disorders[3-5]. Understanding the mechanisms underlying parent-offspring similarities is important for addressing intergenerational transmission of substance use. These similarities are attributed to both *nature* (the genes passed from parents to offspring) and *nurture* (the family environment created by parents)[6-8]. However, disentangling these influences remains challenging, particularly due to unmeasured genetic confounding in risk factors traditionally considered "environmental"[9]. Many environmental risk factors, including parental substance use, are partially genetic[10], further complicating our understanding of substance use transmission across generations.

Advances in molecular genetics have made it possible to distinguish between transmitted and non-transmitted parental genotypes[11,12]. Parental transmitted alleles influence offspring outcomes both directly through inheritance and indirectly through the environment, while non-transmitted alleles affect offspring only indirectly through the rearing environment – a process known as "genetic nurture" [11,12]. For example, parents with a higher genetic risk for substance use may be more likely to engage in substance use themselves, thereby increasing the likelihood of offspring substance use indirectly through the rearing environment (e.g., modeling behavior, and accessibility of substances)[2,13]. To distinguish genetic transmission from genetic nurture, researchers calculate polygenic scores for transmitted (PGS_T) and non-transmitted alleles (PGS_{NT}), each score representing genetic predisposition for a given trait[11,12]. By examining PGS_T and PGS_{NT}, we can determine how parental genetics shape both inherited risk and the rearing environments that predispose offspring to substance use.

While robust genetic nurture effects have been demonstrated for educational outcomes[14],

research on its role in substance use remains scarce. This study aims to disentangle genetic

transmission and genetic nurture on substance use in adulthood. Specifically, we address three key

questions: i) Do genetic nurture effects contribute to offspring substance use in adulthood? ii) Do

these effects differ between mothers and fathers (i.e., parent-of-origin effects)? iii) To what extent

are these effects mediated by familial environmental factors, such as parental substance use?

First, we investigated whether parental PGS_T and PGS_{NT} (aggregated across both parents)

predict offspring tobacco, alcohol, and cannabis use in adulthood. Previous studies have identified

genetic nurture effects, primarily in adolescence and young adulthood[15] or in high-risk samples[16]

with a narrow focus of substance types (e.g., problematic alcohol use). However, it remains unclear

whether these effects i) persist into adulthood, even after offspring leave the family household[17];

and ii) generalize to the broader population and across different substances. To address these gaps,

we examined genetic nurture on a set of substance use measures in adult offspring from Lifelines,

a general population cohort in the Netherlands. We hypothesized that PGS_{NT} would be associated

with offspring substance use outcomes, providing evidence of genetic nurture, while PGS_T would

explain a larger proportion of the variance.

Additionally, prior research has highlighted that relying solely on complete trio data (both

parents and offspring genotyped) may introduce selection bias and limit generalizability[18]. To

mitigate this, we utilized a validated haplotype-based approach[19] to construct PGS_T and PGS_{NT} for

smoking initiation (SmkInit), cigarettes per day (CigDay), alcoholic drinks per week (DrnkWk),

and lifetime cannabis use (CanU). This method enables the inclusion of both parent-offspring pairs

(one parent and their offspring are genotyped) and complete trios, increasing sample size and

enhancing generalizability.

Second, we examined parent-of-origin effects with separate maternal and paternal PGS_T

and PGS_{NT}. Although both parents contribute to genetic nurture, their effects may differ in

magnitude. Prior research suggests that paternal and maternal genetics similarly influence

offspring educational attainment[11,14,19], but mothers have stronger genetic nurture effects than

fathers on health-related traits[11,20] and attention problems[21]. However, no study has investigated

parent-of-origin effects on substance use. This study directly addresses this gap by partitioning

maternal and paternal genetic nurture effects, providing a more nuanced understanding of the roles

of mothers and fathers in the intergenerational transmission of substance use. Given the absence of

prior research in this area, our analysis was exploratory and did not include a specific hypothesis.

Third, we explored whether and to what extent parental substance use represents a

mediating pathway through which genetic nurture contributes to offspring outcomes. While

identifying genetic nurture reveals that parental genetics influences offspring outcomes through

environmental pathways, it does not specify which environmental factors are involved. Mediation

analysis helps to clarify these pathways by examining how parental genetic liability influences

offspring outcomes through specific environmental factors. Parental substance use is a particularly

promising mediator due to its role in offspring substance use and its partially genetic basis[3,6].

Previous studies have identified parental substance use as a mediator of genetic nurture in

adolescents and young adults[15] or high-risk families[16]. This study extends prior work by: i)

distinguishing maternal and paternal mediating pathways, and ii) explicitly modeling parallel

mediation pathways for both transmitted and non-transmitted genetics. We hypothesized that these

genetic effects would be partially mediated by parental substance use.

RESULTS

Population characteristics

We utilized data from a total of 19,233 genotyped adult offspring with at least one parent genotyped

from the Lifelines cohort, comprising 15,966 parent-offspring pairs and 3,267 trios. Of these, up

to 15,863 participants (mean age=31.66 years; 61.9% female), consisting of 13,411 pairs and 2,452

trios, completed assessments for tobacco and alcohol use at baseline (2006–2013) and cannabis use

during the second wave (2014–2017). Descriptive statistics for these participants are presented in

Table 1. Notably, offspring with both parents genotyped (trios) reported significantly lower

prevalence of smoking initiation, fewer cigarettes per day and pack-years, compared to those with

only one genotyped parent (pairs). Correlations among PGSs and offspring outcomes are provided

in **Supplemental Table S1.**

Genetic nurture effects on substance use

We combined maternal and paternal PGS_T and PGS_{NT} for statistical power (see **Supplemental**

Information). Linear mixed regression models were used to examine overall effects of PGS_T and

PGS_{NT} (**Table 2**). Following Kong et al.[11], we estimated the effect of direct genetic transmission

(β_{DGT}) by subtracting the effect of PGS_{NT} from that of PGS_T (i.e., $\beta_T - \beta_{NT}$). This approach isolates

"true" genetic transmission by removing the influence of genetic nurture, given that PGS_T may

influence offspring directly through genetic inheritance and indirectly via genetic nurture.

As expected, each PGS_T was significantly associated with its corresponding substance use

outcomes, with odds ratios between 1.3 and 1.9 for binary outcomes and standardized coefficients

ranging from 0.09 to 0.19 for continuous outcomes. For genetic transmission effects, β_{DGT} was 0.16

for cigarettes per day, 0.07 for pack-years, and 0.10 for daily alcohol intake. For genetic nurture

effects, parental PGS_{NT_CigDay} showed relatively smaller but significant associations with offspring

cigarettes per day ($\beta_{NT} = 0.03$) and pack-years ($\beta_{NT} = 0.02$) after false discovery rate (FDR)

correction. These genetic nurture effects equaled approximately 18.8% and 28.6% of β_{DGT} for

cigarettes per day and pack-years, respectively. While parental PGS_{NT_DrnkWk} was significantly

associated with offspring daily alcohol intake, this effect did not survive FDR correction. No

associations found between parental PGS_{NT} and smoking initiation or lifetime cannabis use.

Overall, these findings indicate modest genetic nurture effects on smoking quantity. In

contrast, initial tobacco and cannabis use, and daily alcohol intake were predominantly influenced

by genetic transmission, with minimal contributions from genetic nurture.

Parent-of-origin effects

Given the significant genetic nurture effects observed for smoking quantity, we further examined

parent-specific effects on cigarettes per day and pack-years (**Table 3**). Structural equation

modelling (SEM) was used to simultaneously estimate maternal and paternal PGS_T and PGS_{NT}

effects, while accounting for potential genetic assortative mating on tobacco use[22]. Full-

information maximum-likelihood estimation (FIML)[23] was applied to handle missing data,

enabling the inclusion of all 19,233 offspring from the full genotyped family sample.

Both maternal and paternal PGS_{T_CigDay} significantly predicted offspring smoking quantity,

with paternal PGS_{T_CigDay} showing slightly greater effects. Wald tests indicated that the magnitude

of PGS_{T_CigDay} did not differ significantly between mothers and fathers ($\Delta\chi^2 = 3.35$, $p = .07$ for

cigarettes per day, $\Delta\chi^2 = 2.49$, $p = .12$ for pack-years).

For genetic nurture effects, both maternal and paternal PGS_{NT_CigDay} were significantly

associated with offspring pack-years, with similar effect sizes (maternal $\beta = 0.027$, 95% CI [0.001,

0.052], paternal $\beta = 0.034$, 95% CI [0.001, 0.065]). Wald test confirmed no significant difference

between maternal and paternal effects ($\Delta\chi^2 = 0.01$, $p = 0.91$). For cigarettes per day, maternal and

paternal PGS_{NT_CigDay} were also nearly identical in magnitude, but the association was significant

only for mothers (maternal 95% CI [0.008, 0.069], paternal 95% CI [-0.001, 0.071]). No significant

associations were found between maternal and paternal PGSs (**Supplemental Table S2**),

indicating little evidence of genetic assortative mating for smoking quantity. This suggests that

estimates of genetic nurture effects were unlikely to be inflated by genetic similarity between

parents.

Mediation by parental tobacco use

To explore the extent to which parental tobacco use mediated the genetic nurture effects identified

above, we conducted mediation analyses separately for mothers and fathers (**Supplemental**

Information).

169   As shown in **Figure 1(A, B)**, maternal smoking quantity significantly mediated the effects

of both PGS_{T_CigDay} and PGS_{NT_CigDay} on offspring smoking outcomes, with mediated proportions

of 23.2% (cigarettes per day: $\beta_{mediation} = 0.029$, $\beta_{total} = 0.125$) and 17.86% (pack-years: $\beta_{mediation} =$

0.015, $\beta_{total} = 0.084$) for PGS_T, and 74.42% (cigarettes per day: $\beta_{mediation} = 0.032$, $\beta_{total} = 0.043$) and

74.07% (pack-years: $\beta_{mediation} = 0.020$, $\beta_{total} = 0.027$) for PGS_{NT}. In contrast, paternal pack-years

had a lower mediated proportion (**Figure 1C**), mediating 7.48% of PGS_{T_CigDay} ($\beta_{mediation} = 0.008$,

SE = 0.002; $\beta_{total} = 0.107$) and 26.47% of PGS_{NT_CigDay} ($\beta_{mediation} = 0.009$; $\beta_{total} = 0.034$) on offspring

pack-years. These findings suggest that parental smoking, especially maternal smoking, partially

mediates genetic nurture effects on offspring smoking behaviors.

DISSCUSION

This study aimed to disentangle genetic nurture and genetic transmission effects on substance use during adulthood, utilizing transmitted and non-transmitted polygenic scores. We examined these effects on various substance use traits, including across smoking initiation, smoking quantity (cigarettes per day, pack-years), daily alcohol intake, and lifetime cannabis use. Additionally, we investigated parent-of-origin effects and the mediating role of parental substance use to provide a nuanced understanding of intergenerational transmission underlying substance use. We highlight three key findings. First, genetic nurture significantly influenced smoking quantity, but not smoking initiation, daily alcohol intake, or lifetime cannabis use. As expected, genetic transmission played a substantial role across all measured outcomes. Second, parent-of-origin analyses revealed that the magnitudes of genetic nurture effects on smoking quantity were nearly identical for mothers and fathers. Finally, parental smoking quantity partially mediated the effects of both transmitted and non-transmitted genetic risk on offspring smoking outcomes in adulthood, with higher mediation proportions for maternal smoking quantity.

This study extends previous research by exploring genetic nurture on multiple substance use traits during adulthood within a large population-based cohort. While Saunders et al.[15] reported genetic nurture effects on cigarettes per day at age 24 but not at age 29, our results indicate these effects persist into late adulthood, even after offspring have likely left the parental home. This discrepancy likely reflects differences in study design (twin study versus population-based study), sample age ranges (ages 17-29 versus ages 18-67), and the temporary dynamics of genetic nurture. Additionally, our study found no significant genetic nurture on smoking initiation or lifetime cannabis use, suggesting that such effects might be minimal on the initiation of substance use in adults. Several explanations may underlie this finding. First, the dichotomous measure used for

initial use may be underpowered to detect genetic nurture effects, despite the large sample size for

smoking initiation ($N = 15,853$). Continuous and cumulative metrics, such as cigarettes per day

and pack-years, potentially enhance sensitivity to capture the persistence of genetic nurture into

adulthood. Second, genetic nurture on substance use initiation may be more pronounced during

adolescence, a critical period for the onset of such behaviors[24], but these effects likely diminish by

adulthood. This aligns with studies showing genetic and environmental contributions change over

time [25,26] and across different stages of substance use[6,7,27,28], such as initiation, quantity/persistence,

and dependence. For instance, shared environmental factors have consistently been shown to

explain substantial variance in smoking initiation during early adolescence but contribute

minimally to young adulthood[26,29]. Conversely, genetic factors explain an increasing proportion of

variance in substance use as individuals age[25,30]. As genetic nurture partly mirrors these shared

environment effects[31], its effects may be limited for traits with low shared environmental estimates.

We found no evidence for genetic nurture on daily alcohol intake. This could be due to our

short-term measurement of alcohol use (quantity and frequency over the past 30 days) which may

not adequately capture long-term drinking patterns[32]. One study[16] in a high-risk sample identified

genetic nurture for alcohol use disorder, suggesting that genetic nurture may exist for pathological

alcohol use but not for daily alcohol intake. Alcohol consumption, particularly measured in a

general population cohort, may vary depending on social situations or specific life events (e.g.,

being a college student, celebrations or holidays). These contextual factors may contribute to why

genetic nurture plays a less important role in daily alcohol intake.

Together, these findings provide valuable insights into the influence of genetic nurture on

various types and stages of substance use. Genetic nurture appears to have a stronger influence on

sustained or problematic substance use, such as smoking quantity and alcohol dependence

involving regulation and reinforcement over time[33,34]. Conversely, its impact on initial

experimentation, such as tobacco or cannabis initiation, may be minimal in adulthood. These

observed patterns of genetic nurture, particularly their persistence into adulthood, align with

developmental models[35] emphasizing long-term parental influences on substance use behaviors.

Future research should leverage longitudinal data to explore how genetic nurture effects unfold

across the lifespan and across stages of substance use.

Another key contribution of this study is the investigation of parent-of-origin effects,

revealing that maternal and paternal genetic nurture contribute equally to offspring smoking

quantity. Although paternal genetic nurture was not statistically significant for cigarettes per day,

likely due to reduced power from a smaller sample size, its overall effect was comparable to

maternal genetic nurture. Our result aligns with previous findings of similar genetic nurture effects

from mothers and fathers for other complex traits, including educational attainment[11,14] and birth

weight[36]. One plausible explanation is that both parents are equally important in shaping shared

family environments and parenting practices, which in turn influence offspring smoking outcomes.

Consistent with this, previous evidence suggests that substance use intervention efforts may benefit

from considering both maternal and paternal influences[37,38].

Interestingly, despite maternal and paternal genetic nurture effects on pack-years were

nearly equal at the genetic level, our mediation analyses showed maternal smoking quantity

constitutes a more substantial mediating pathway in both relative terms (mediating approximately

74% versus 26% of genetic nurture effects) and absolute magnitude (mediation effect of 0.020

versus 0.009). This suggests that the mechanisms through which genetic nurture acts on offspring

may differ between parents. Our findings highlight the importance of dissecting parent-specific

mediation pathways, as maternal and paternal genetic nurture may impact offspring behaviors

through distinct mechanisms. Mothers may exert genetic nurture effects on offspring smoking

primarily through accumulative smoking exposure (e.g., prenatal and postnatal), aligning with

research linking such exposure to offspring substance use[3,39]. Fathers' influence may operate

through other pathways, such as paternal psychiatric disorders[40] and father-child relationship[41],

beyond direct smoking exposure, which were not included in our mediation model. Future research

should explore alternative mediation pathways to gain a more comprehensive understanding of

intergenerational transmission of substance use. Such findings might inform interventions aimed

at reducing substance use risks through targeted parental support strategies.

This study has several strengths, including a large, population-based cohort and the use of

a unique dataset integrating genotypic data with various substance use measures from mothers,

fathers, and offspring. Unlike much existing research that combines parental PGS, this study

partitioned maternal and paternal genetic effects, allowing for the investigation of parent-of-origin

genetic nurture. Additionally, by differentiating between transmitted and non-transmitted alleles,

we were able to identify genetic nurture effects unconfounded by direct genetic transmission.

However, several limitations remain. First, our sample included only participants of European

descent, potentially limiting generalizability. Second, PGSs explain only a small proportion of

genetic liability to substance use, and the modest genetic nurture effect sizes observed reflect this

limitation. Third, paternal analyses may be underpowered due to limited availability of fathers with

both genotypic and phenotypic data. Fourth, retrospective self-reports on substance use, such as

pack-years, may be subject to recall bias or underreporting[42]. However, measures like cigarettes

per day and pack-years have demonstrated strong reliability and validity in assessing lifetime

smoking exposure[43]. Fifth, we were unable to examine genetic nurture effects across different age

groups due to sample size constraints, despite evidence suggesting that such effects may vary with

age[15,44-46]. Finally, our hypothetical mediation models were theory-driven but not exhaustive.

Comparisons with saturated models (**Supplemental Table S3**), which perfectly fit the data by

estimating all possible paths, showed better statistical fit for the saturated models across all

analyses ($p < .01$). While this suggests potentially unmodeled paths, these comparisons should be

interpreted in light of our large sample size, which increases sensitivity to detect even minor

modelling misspecifications. Notably, the differences in sample-size-insensitive fit indices were

minimal (ΔRMSEA = 0.02, ΔCFI = -0.01 to -0.004), suggesting our theoretically-grounded

mediation models maintain acceptable fit while offering greater parsimony.

CONCLUSIONS

To conclude, we provide evidence for genetic nurture on smoking behaviors in adulthood with a

stage-specific pattern, while direct genetic transmission remains the primary driver of substance

use. We found that maternal and paternal genetic nurture contribute equally to offspring smoking

quantity, particularly pack-years. Moreover, parental substance use mediates genetic nurture

effects, with maternal smoking quantity potentially playing a larger mediating role. Our study

provides a nuanced understanding of genetic nurture effects on substance use, which might operate

in stage-specific and parent-specific ways. These findings, particularly the persistence of genetic

nurture effects into adulthood, highlight the importance of both maternal and paternal influences

in research and interventions aimed at reducing the intergenerational transmission of substance use.

METHODS

Participants

Lifelines is a multi-disciplinary prospective population-based cohort study examining in a unique

three-generation design the health and health-related behaviors of 167,729 persons living in the

North of the Netherlands. It employs a broad range of investigative procedures in assessing the

biomedical, socio-demographic, behavioral, physical and psychological factors which contribute

to the health and disease of the general population, with a special focus on multi-morbidity and

complex genetics[47,48]. Data collection was accomplished at three general assessments, along with

additional assessments. The baseline assessment took place between 2007 and 2013, followed by

a second wave between 2014 and 2017, and a third wave between 2019 and 2023. The design and

sample characteristics of Lifelines have been described in detail elsewhere[47,48].

Measurements

Substance use

Substance use was measured by self-report questionnaires at baseline for adult offspring and their

parents, except for cannabis use which was measured at the second wave. Smoking initiation was

defined as having smoked for one year or longer with the question "Have you ever smoked for a

full year in your lifetime?". Participants who smoked for less than a year were not considered as a

smoker. In participants who smoked for a year or more, smoking quantity was assessed in cigarettes

per day and pack years. Cigarettes per day was defined as the lifetime average number of cigarettes

smoked per day, either as a current or former smoker. Pack-years was calculated by multiplying

the average amount smoked per day (different types of tobacco, including cigarettes, cigarillos,

cigars and pipes) by the number of years the person smoked in their lifetime (1 pack-year equals

20 cigarettes per day for one year). Pack-years is a cumulative indicator of lifetime smoking

behavior (not only singular tobacco use variables like cigarettes per day, but also other tobacco

types).

Alcohol use was assessed with a food frequency questionnaire developed by Wageningen

University[49]. Two questions referred to the frequency and quantity of alcohol consumed in the past

month: "How often did you drink alcoholic drinks in the past month?" (ranging from 'not this

month' to '6-7 days per week'), and "On days that you drank alcohol, how many glasses did you

drink on average?" (from '1' to '12 or more'). These questions were split up for different alcoholic

groups (beer, alcohol-free beer, red wine/rose, white wine, sherry, distilled wine, other alcoholic

beverages). Based on these questions, an average daily alcohol consumption in grams per day was

calculated[50]. This composite index of daily alcohol intake provides a more comprehensive measure

of overall alcohol use than a single measure of frequency (e.g., number of drinking days per month)

or quantity (e.g., glasses per day).

Lifetime cannabis use was defined using two questions: i) "Have you ever used drugs?",

and if yes, ii) "Have you ever used cannabis, such as weed, marijuana, hashish?". The answer

categories were recoded to ever (1) versus never (0) used cannabis.

Genotyping and imputation

A total of 79,988 participants were genotyped across three batches in Lifelines. Quality control

(QC) of markers and samples was performed separately per batch. Detailed pre-imputation QC

criteria is described in **Supplemental Information**.

In brief, markers that were duplicated and monomorphic, markers with a low call rate or

low minor allele frequency, and markers that deviated significantly from Hardy-Weinberg

equilibrium were removed. Post QC data from each array was imputed through the Sanger

Imputation Service with the Haplotype Reference Consortium v1 panel. We selected overlapping

imputed markers with quality scores ≥0.8 across arrays to create a common set of markers for all

genotyped parents and offspring in any arrays. Samples with a low call rate, heterozygosity outliers

or mix-ups on sex and familial relationship were filtered out. Samples were restricted to

participants of European ancestry, determined through principal component analysis with the 1000

Genomes reference, to control for population stratification.

Non-transmitted alleles inference

We applied our newly developed haplotype-based approach to differentiate transmitted and non-

transmitted alleles in genotyped parent-offspring pairs and trios[19]. By including parent-offspring

pairs, rather than restricting the analysis to trios, this approach reduces sample attrition and

improves statistical power. The development and validation of this method have been described in

detail elsewhere[19]. Briefly, we used SHAPEIT5 to estimate haplotypes including pedigree

information[51]. Offspring haplotypes were then compared to parental haplotypes using tiles of 150

adjacent markers on each chromosome. The best match between the parent and offspring tiles,

taking recombination spots into account, was used to determine which parental tiles were

transmitted to the offspring. The remaining non-transmitted alleles were recorded in a separate

dataset, and for parent-offspring pairs, the non-transmitted alleles of the parent who was not

genotyped were set as missing. This method was validated by comparison with standard software

in parent-offspring trios and found a concordance rate for the non-transmitted alleles of 99.8%.

Furthermore, the identification of non-transmitted alleles was confirmed to be unaffected by

missing parental data through simulations of pairs from trios.

Polygenic scores

We calculated PGS_T and PGS_{NT} based on summary statistics from previous genome-wide

association studies (GWAS) for SmkInit[52], CigDay[52], DrnkWk[52], and CanU[53] (**Supplementary**

Table S4). These GWAS were chosen because they were based on the largest sample sizes

currently available for each corresponding phenotype in Lifelines.

To increase the variance explained by each PGS, SNP effects were re-weighted using the

'auto' setting from LDpred2[54], a Bayesian method that adjusts the effect estimates from GWAS

summary statistics by incorporating trait-specific genetic architecture (e.g., SNP-based heritability

and polygenicity measured as the fraction of causal variants) and linkage disequilibrium (LD) data

from UK Biobank reference panel for European ancestry[55]. For each offspring, PGSs were created

based on transmitted and non-transmitted datasets. To estimate overall genetic nurture and genetic

transmission effects, parental PGS_T and PGS_{NT} were defined as the sum of the PGS based on

paternal and maternal transmitted and non-transmitted haplotypes, respectively. The value of the

missing PGS_{NT} in parent-offspring pairs was imputed with the average PGS_{NT} of the observed

parents (**Supplementary Information**). To estimate parent-of-origin effects, we separated four

maternal and paternal PGS_T and PGS_{NT} respectively. To control for population structure and batch

effects across arrays, we standardized PGS residuals within each array after regressing out the first

ten genetic principal components.

Statistical analysis

All analyses were conducted in R[56]. We applied a stepwise approach in which genetic nurture

effects had to be statistically significant to continue to the next analysis.

Genetic nurture and genetic transmission on substance use

First, we used mixed-effects regression models to examine associations between parental

PGS_T and PGS_{NT} with offspring substance use outcomes, including smoking initiation, smoking

quantity (cigarettes per day, pack-years), daily alcohol intake, and lifetime cannabis use.

Continuous outcomes were analyzed using mixed-effects linear regression with the '*lme4*'

package[57], while dichotomous outcomes were analyzed using mixed-effects logistic regression

with the '*GLMMadaptive*' package[58]. Each model was specified as follows: $Y_i = Intercept_Y +$

$\beta PGS_T + \beta PGS_{NT} + sex + age + 1|FamilyID + e_i$. Family ID was included as a random effect

(intercept) to account for the relatedness among siblings, along with sex and age as covariates.

False discovery rate (FDR) corrections[59] (5 tests, $\alpha = 0.05$) were applied to control for multiple

testing across the two mixed-effects logistic regression models (smoking initiation, lifetime

cannabis use) and three mixed-effects linear regression models (cigarettes per day, pack-years, and

daily alcohol intake).

Parent-of-origin effects

If genetic nurture effects remained significant after FDR, we examined parent-of-origin

effects on offspring's substance use outcomes using structural equation modeling (SEM) in

'*Lavaan*' package[60]. SEM enabled us to handle missing data with Full Information Maximum

Likelihood(FIML), which utilizes all available data to estimate parameters and standard errors

without imputing missing values[23]. For each SEM model, $Y_i = Intercept_Y + \beta PGS_{T\_mother} +$

$\beta PGS_{NT\_mother} + \beta PGS_{T\_father} + \beta PGS_{NT\_father} + sex + age + e_i$, all paths and covariances

will be freely estimated. To account for non-normality and non-independence of observation[61], we

generated standard errors and 95% confidence intervals (CIs) using bootstrapping with 1,000

replications. To assess whether the maternal and paternal effects differed significantly, we

compared the equality of standardized regression coefficients using a Wald test[62], via *'lavTestWald'*

function in Lavaan.

Mediation pathways via parental substance use

Finally, we conducted mediation analysis using SEM (*Lavaan* package) to assess the extent

to which parental substance use mediates the associations of PGS_T and PGS_{NT} with offspring

substance use outcomes. FIML was used to handle missing data and reduce the likelihood of biased

parameter estimates. Standardized mediation effects were estimated via bootstrapping with 1,000

replications, with statistical significance determined by 95% bootstrap CIs excluding zero.

Additionally, we calculated the proportion mediated, which represents the percentage of the total

effect on the outcome explained by mediation.

To assess the appropriateness of mediation model, we compared our hypothetical mediation

models with saturated models, which perfectly fit the data by estimating all variances, covariances,

and means of the observed variables (**Supplementary Information**).

426 **DATA AVAILABILITY**

427 Data may be obtained from a third party and are not publicly available. Researchers can apply to

428 use the Lifelines data used in this study. More information about how to request Lifelines data and

429 the conditions of use can be found on their website (https://www.lifelines.nl/researcher/how-to-

430 apply).

ACKNOWLEDGEMENTS

This work was supported by grants from the United States National Institutes of Health, National Institute on Drug Abuse (R01DA052453, R00DA023549). The work of HvL was supported by a VENI grant from the Talent Program of the Netherlands Organization of Scientific Research (NWO-ZonMW 09150161810021). The Lifelines initiative has been made possible by subsidy from the Dutch Ministry of Health, Welfare and Sport, the Dutch Ministry of Economic Affairs, the University Medical Center Groningen (UMCG), Groningen University and the Provinces in the North of the Netherlands (Drenthe, Friesland, Groningen). We acknowledge the services of the Lifelines Cohort Study, the contributing research centers delivering data to Lifelines, and all the study participants. We also thank Prof. Jean-Baptiste Pingault for his valuable input on the statistical analysis.

444 **COMPETING INTERESTS**

445 The authors declare no competing interests.

447 **ETHICS STATEMENT**

448 The Lifelines protocol has been approved by the UMCG Medical ethical committee under number

449 2007/152.

REFERENCES

Merikangas, K. R. *et al.* Familial transmission of substance use disorders. *Archives of general psychiatry* **55**, 973-979 (1998).

Kendler, K. S., Abrahamsson, L., Ohlsson, H., Sundquist, J. & Sundquist, K. Cross-generational transmission of genetic risk for alcohol and drug use disorders: the impact of substance availability on the specificity of genetic risk. *Psychological Medicine* **53**, 5109-5118 (2023). https://doi.org/10.1017/S0033291722002549

McGovern, R. *et al.* The association between maternal and paternal substance use and child substance use, internalizing and externalizing problems: a systematic review and meta-analysis. *Addiction* **118**, 804-818 (2023). https://doi.org/https://doi.org/10.1111/add.16127

Mellentin, A. I. *et al.* The risk of offspring developing substance use disorders when exposed to one versus two parent (s) with alcohol use disorder: A nationwide, register-based cohort study. *Journal of psychiatric research* **80**, 52-58 (2016).

Madras, B. K. *et al.* Associations of Parental Marijuana Use With Offspring Marijuana, Tobacco, and Alcohol Use and Opioid Misuse. *JAMA Network Open* **2**, e1916015-e1916015 (2019). https://doi.org/10.1001/jamanetworkopen.2019.16015

Rhee, S. H. *et al.* Genetic and Environmental Influences on Substance Initiation, Use, and Problem Use in Adolescents. *Archives of General Psychiatry* **60**, 1256-1264 (2003). https://doi.org/10.1001/archpsyc.60.12.1256

Verweij, K. J. *et al.* Genetic and environmental influences on cannabis use initiation and problematic use: a meta‐analysis of twin studies. *Addiction* **105**, 417-430 (2010).

Li, M. D., Cheng, R., Ma, J. Z. & Swan, G. E. A meta-analysis of estimated genetic and environmental effects on smoking behavior in male and female adult twins. *Addiction* **98**, 23-31 (2003). https://doi.org/10.1046/j.1360-0443.2003.00295.x

Kendler, K. & Baker, J. Genetic influences on measures of the environment: a systematic review. *Psychol Med* **37**, 615-626 (2007).

Barr, P. B. *et al.* Clinical, environmental, and genetic risk factors for substance use disorders: characterizing combined effects across multiple cohorts. *Molecular Psychiatry* **27**, 4633-4641 (2022). https://doi.org/10.1038/s41380-022-01801-6

Kong, A. *et al.* The nature of nurture: Effects of parental genotypes. *Science* **359**, 424-428 (2018). https://doi.org/10.1126/science.aan6877

Bates, T. C. *et al.* The Nature of Nurture: Using a Virtual-Parent Design to Test Parenting Effects on Children's Educational Attainment in Genotyped Families. *Twin Res Hum Genet* **21**, 73-83 (2018). https://doi.org/10.1017/thg.2018.11

Broman, C. L. The Availability of Substances in Adolescence: Influences in Emerging Adulthood. *J Child Adolesc Subst Abuse* **25**, 487-495 (2016). https://doi.org/10.1080/1067828x.2015.1103346

Wang, B. *et al.* Robust genetic nurture effects on education: A systematic review and meta-analysis based on 38,654 families across 8 cohorts. *The American Journal of Human Genetics* **108**, 1780-1791 (2021). https://doi.org/https://doi.org/10.1016/j.ajhg.2021.07.010

Saunders, G. R. B., Liu, M., Vrieze, S., McGue, M. & Iacono, W. G. Mechanisms of parent-child transmission of tobacco and alcohol use with polygenic risk scores: Evidence

for a genetic nurture effect. *Dev Psychol* **57**, 796-804 (2021).
https://doi.org/10.1037/dev0001028
16   Thomas, N. S. *et al.* Genetic nurture effects for alcohol use disorder. *Molecular*
psychiatry **28**, 759-766 (2023).
17   Sipilä, P. N., Keski-Rahkonen, A., Lindbohm, J. V., Rose, R. J. & Kaprio, J. Paternal and
Maternal Problem Drinking and Lifetime Problem Drinking of Their Adult Children.
Twin Research and Human Genetics **26**, 152-163 (2023).
https://doi.org/10.1017/thg.2023.12
18   Martin, J. *et al.* Investigating Direct and Indirect Genetic Effects in Attention-
Deficit/Hyperactivity Disorder Using Parent-Offspring Trios. *Biol Psychiatry* **93**, 37-44
(2023). https://doi.org/10.1016/j.biopsych.2022.06.008
19   Trindade Pons, V., Claringbould, A., Kamphuis, P., Oldehinkel, A. J. & van Loo, H. M.
Using parent-offspring pairs and trios to estimate indirect genetic effects in education.
Genetic Epidemiology **n/a** https://doi.org/https://doi.org/10.1002/gepi.22554
20   Tubbs, J. D., Hwang, L.-D., Luong, J., Evans, D. M. & Sham, P. C. Modeling Parent-
Specific Genetic Nurture in Families with Missing Parental Genotypes: Application to
Birthweight and BMI. *Behavior Genetics* **51**, 289-300 (2021).
https://doi.org/10.1007/s10519-020-10040-w
21   Hegemann, L. *et al.* Direct and indirect genetic effects on early neurodevelopmental traits.
Journal of Child Psychology and Psychiatry **n/a**
https://doi.org/https://doi.org/10.1111/jcpp.14122
22   Horwitz, T. B., Balbona, J. V., Paulich, K. N. & Keller, M. C. Evidence of correlations
between human partners based on systematic reviews and meta-analyses of 22 traits and
UK Biobank analysis of 133 traits. *Nature Human Behaviour* **7**, 1568-1583 (2023).
https://doi.org/10.1038/s41562-023-01672-z
23   Lee, T. & Shi, D. A comparison of full information maximum likelihood and multiple
imputation in structural equation modeling with missing data. *Psychological Methods* **26**,
466 (2021).
24   Nuyts, P. A. W., Kuipers, M. A. G., Willemsen, M. C. & Kunst, A. E. Trends in age of
smoking initiation in the Netherlands: a shift towards older ages? *Addiction* **113**, 524-532
(2018). https://doi.org/https://doi.org/10.1111/add.14057
25   Kendler, K. S., Schmitt, E., Aggen, S. H. & Prescott, C. A. Genetic and environmental
influences on alcohol, caffeine, cannabis, and nicotine use from early adolescence to
middle adulthood. *Arch. Gen. Psychiatry* **65**, 674-682 (2008).
https://doi.org/10.1001/archpsyc.65.6.674
26   Koopmans, J. R., Slutske, W. S., Heath, A. C., Neale, M. C. & Boomsma, D. I. The
Genetics of Smoking Initiation and Quantity Smoked in Dutch Adolescent and Young
Adult Twins. *Behavior Genetics* **29**, 383-393 (1999).
https://doi.org/10.1023/A:1021618719735
27   Vink, J. M., Willemsen, G. & Boomsma, D. I. Heritability of smoking initiation and
nicotine dependence. *Behav. Genet.* **35**, 397-406 (2005).
28   Stallings, M. C., Hewitt, J. K., Beresford, T., Heath, A. C. & Eaves, L. J. A twin study of
drinking and smoking onset and latencies from first use to regular use. *Behav Genet* **29**,
409-421 (1999). https://doi.org/10.1023/a:1021622820644
29   Bares, C. B., Kendler, K. S. & Maes, H. H. Developmental Changes in Genetic and
Shared Environmental Contributions to Smoking Initiation and Subsequent Smoking

Quantity in Adolescence and Young Adulthood. *Twin Research and Human Genetics* **18**,
497-506 (2015). https://doi.org/10.1017/thg.2015.48

30 Sullivan, P. F. & Kendler, K. S. The genetic epidemiology of smoking. *Nicotine &*
Tobacco Research **1**, S51-S57 (1999). https://doi.org/10.1080/14622299050011811

31 McAdams, T. A., Cheesman, R. & Ahmadzadeh, Y. I. Annual Research Review: Towards
a deeper understanding of nature and nurture: combining family-based quasi-experimental
methods with genomic data. *J Child Psychol Psychiatry* **64**, 693-707 (2023).
https://doi.org/10.1111/jcpp.13720

32 Agrawal, A. *et al.* Measuring alcohol consumption for genomic meta-analyses of alcohol
intake: opportunities and challenges12345. *The American Journal of Clinical Nutrition*
95, 539-547 (2012). https://doi.org/https://doi.org/10.3945/ajcn.111.015545

33 Neale, M. C., Bustamante, D., Zhou, Y. & Gillespie, N. A. in *Genetics of Substance Use:*
Research and Clinical Aspects (ed Michael M. Vanyukov) 99-117 (Springer
International Publishing, 2022).

34 Caspi, A., Bem, D. J. & Elder Jr., G. H. Continuities and Consequences of Interactional
Styles Across the Life Course. *Journal of Personality* **57**, 375-406 (1989).
https://doi.org/https://doi.org/10.1111/j.1467-6494.1989.tb00487.x

35 Elam, K. K., Lemery-Chalfant, K. & Chassin, L. A gene-environment cascade theoretical
framework of developmental psychopathology. *J Psychopathol Clin Sci* **132**, 287-296
(2023). https://doi.org/10.1037/abn0000546

36 Ghatan, S. *et al.* Genetic Nurture: Estimating the direct genetic effects of pediatric
anthropometric traits. *medRxiv*, 2024.2012.2010.24318796 (2024).
https://doi.org/10.1101/2024.12.10.24318796

37 Stover, C. S., Carlson, M., Patel, S. & Manalich, R. Where's dad? The importance of
integrating fatherhood and parenting programming into substance use treatment for men.
Child abuse review **27**, 280-300 (2018).

38 McGovern, R., Newham, J., Addison, M., Hickman, M. & Kaner, E. The effectiveness of
psychosocial interventions at reducing the frequency of alcohol and drug use in parents:
findings of a Cochrane Review and meta‐analyses. *Addiction* **117**, 2571-2582 (2022).

39 Duko, B. *et al.* Prenatal tobacco exposure and the risk of tobacco smoking and
dependence in offspring: a systematic review and meta-analysis. *Drug and Alcohol*
Dependence **227**, 108993 (2021).

40 Sarala, M. *et al.* Parental smoking and young adult offspring psychosis, depression and
anxiety disorders and substance use disorder. *European journal of public health* **32**, 254-
260 (2022).

41 Yoon, S. *et al.* Vulnerability or resilience to early substance use among adolescents at
risk: The roles of maltreatment and father involvement. *Child Abuse & Neglect* **86**, 206-
216 (2018). https://doi.org/https://doi.org/10.1016/j.chiabu.2018.09.020

42 Bernaards, C. M., Twisk, J. W., Snel, J., Van Mechelen, W. & Kemper, H. C. Is
calculating pack‐years retrospectively a valid method to estimate life‐time tobacco
smoking? A comparison between prospectively calculated pack‐years and
retrospectively calculated pack‐years. *Addiction* **96**, 1653-1661 (2001).

43 Brigham, J. *et al.* Test-retest reliability of web-based retrospective self-report of tobacco
exposure and risk. *Journal of medical Internet research* **11**, e1248 (2009).

44   Demange, P. A. *et al.* Estimating effects of parents' cognitive and non-cognitive skills on
offspring education using polygenic scores. *Nature Communications* **13**, 4801 (2022).
https://doi.org/10.1038/s41467-022-32003-x
45   Armstrong-Carter, E. *et al.* The Earliest Origins of Genetic Nurture: The Prenatal
Environment Mediates the Association Between Maternal Genetics and Child
Development. *Psychol Sci* **31**, 781-791 (2020).
https://doi.org/10.1177/0956797620917209
46   de Zeeuw, E. L. *et al.* Intergenerational Transmission of Education and ADHD: Effects of
Parental Genotypes. *Behavior Genetics* **50**, 221-232 (2020).
https://doi.org/10.1007/s10519-020-09992-w
47   Scholtens, S. *et al.* Cohort Profile: LifeLines, a three-generation cohort study and
biobank. *International journal of epidemiology* **44**, 1172-1180 (2015).
48   Sijtsma, A. *et al.* Cohort Profile Update: Lifelines, a three-generation cohort study and
biobank. *International journal of epidemiology* **51**, e295-e302 (2022).
49   Brouwer-Brolsma, E. M. *et al.* Development and external validation of the 'Flower-FFQ':
a FFQ designed for the Lifelines Cohort Study. *Public Health Nutr* **25**, 225-236 (2022).
https://doi.org/10.1017/s1368980021002111
50   Mangot-Sala, L., Smidt, N. & Liefbroer, A. C. The association between unemployment
trajectories and alcohol consumption patterns. Evidence from a large prospective cohort
in The Netherlands. *Advances in Life Course Research* **50**, 100434 (2021).
https://doi.org/https://doi.org/10.1016/j.alcr.2021.100434
51   Hofmeister, R. J., Ribeiro, D. M., Rubinacci, S. & Delaneau, O. Accurate rare variant
phasing of whole-genome and whole-exome sequencing data in the UK Biobank. *Nature
Genetics* **55**, 1243-1249 (2023). https://doi.org/10.1038/s41588-023-01415-w
52   Saunders, G. R. B. *et al.* Genetic diversity fuels gene discovery for tobacco and alcohol
use. *Nature* **612**, 720-724 (2022). https://doi.org/10.1038/s41586-022-05477-4
53   Pasman, J. A. *et al.* GWAS of lifetime cannabis use reveals new risk loci, genetic overlap
with psychiatric traits, and a causal effect of schizophrenia liability. *Nature Neuroscience*
21, 1161-1170 (2018). https://doi.org/10.1038/s41593-018-0206-1
54   Privé, F., Arbel, J. & Vilhjálmsson, B. J. LDpred2: better, faster, stronger. *Bioinformatics*
36, 5424-5431 (2021). https://doi.org/10.1093/bioinformatics/btaa1029
55   Privé, F., Albiñana, C., Arbel, J., Pasaniuc, B. & Vilhjálmsson, B. J. Inferring disease
architecture and predictive ability with LDpred2-auto. *The American Journal of Human
Genetics* **110**, 2042-2055 (2023).
https://doi.org/https://doi.org/10.1016/j.ajhg.2023.10.010
56   R Core Team. *R: A language and environment for statistical computing.* (R Foundation
for Statistical Computing, Vienna, Austria, 2024).
57   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software* **67**, 1 - 48 (2015).
https://doi.org/10.18637/jss.v067.i01
58   D, R. GLMMadaptive: Generalized Linear Mixed Models using Adaptive Gaussian
Quadrature. (2023).
59   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B
(Methodological)* **57**, 289-300 (1995). https://doi.org/https://doi.org/10.1111/j.2517-
6161.1995.tb02031.x

60   Rosseel, Y. lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical*
Software **48**, 1 - 36 (2012). https://doi.org/10.18637/jss.v048.i02
61   Cheung, G. W. & Lau, R. S. Testing mediation and suppression effects of latent variables:
Bootstrapping with structural equation models. *Organizational Research Methods* **11**,
296-325 (2008). https://doi.org/10.1177/1094428107300343
62   Klopp, E. A tutorial on testing the equality of standardized regression coefficients in
structural equation models using Wald tests with lavaan. *The Quantitative Methods for*
Psychology **16**, 315-333 (2020). https://doi.org/10.20982/tqmp.16.4.p315
642

Table 1. Descriptive characteristics of Lifelines participants, including the subsamples of adult offspring from genotyped family parent-offspring pairs and trios

	All		Pairs		Trios		p*
	N	M±SD or n (%)	*N*	M±SD or n (%)	*N*	M±SD or n (%)	
Age (years) at assessment	15871	31.66 ± 8.61	13417	30.59 ± 8.02	2454	31.86 ± 8.70	
Sex, female	15871	9827 (61.9)	13417	8376 (62.4)	2454	1451 (59.1)	
Tobacco use							
Smoking initiation, yes	15853	6560 (41.4)	13403	5620 (41.9)	2450	940 (38.4)	<.01
Cigarette per day	5972	10.24 ± 6.00	5134	10.34 ± 6.06	838	9.59 ± 5.57	<.01
Pack-years	6276	7.03 ± 6.73	5382	7.19 ± 6.85	894	6.04 ± 5.88	<.001
Alcohol use							
Daily alcohol intake (grams/day)	15863	6.74 ± 8.46	13411	6.73 ± 8.47	2452	6.81 ± 8.41	0.68
Cannabis use							
Lifetime cannabis use, yes	9097	2066 (22.7)	7608	1710 (22.5)	1489	359 (24.1)	0.18

Note: *N*= Sample size, M= Mean, SD = Standard deviation. *Chi-square test for binary outcomes and t-test for continuous outcomes when comparing the pairs versus trios on substance use. Pack-years of smoking were calculated by multiplying the amount smoked per day (of different types of tobacco products, including cigarettes/roll-ups, cigarillos, cigars. grams of pipe tobacco) by the number of years the person has smoked.

Table 2. Regression coefficients of parental transmitted, non-transmitted polygenic scores, and direct genetic effects on offspring substance use in adulthood

	N	β_T (SE)/OR	95% CI	p	p_{FDR}	β_{NT} (SE)/OR	95% CI	p	p_{FDR}	β_T - β_{NT}
		PGS$_T$				**PGS$_{NT}$**				**DGT**
PGS$_{SmkInit}$										
Smoking initiation	15853	**1.853**[a]	1.767, 1.944	<.0001	<.0001	.989 [a]	.950, 1.031	.612	.612	–
PGS$_{CigDay}$										
Cigarettes per day	5972	**.185 (.012)**	.162, .208	<.0001	<.0001	**.033 (.012)**	.011, .057	.004	.020	.16
Pack-years	6276	**.084 (.006)**	.071, .098	<.0001	<.0001	**.017 (.007)**	.004, .031	.014	.035	.07
PGS$_{DrnkWk}$										
Daily alcohol intake	15863	**.120 (.007)**	.106, .134	<.0001	<.0001	.015 (.007)	.001, .031	.034	.057	.10
PGS$_{CanU}$										
Lifetime cannabis use	9097	**1.274**[a]	1.188, 1.365	<.0001	<.0001	1.033[a]	.967, 1.104	.328	.410	–

Note. Mixed-effects regression models include up to 15,863 adult offspring with at least one parent genotyped and available data on substance use outcomes. Age and sex were included as covariates, and sibling effect was controlled for by including family ID as a random effect into models. β_T and β_{NT}: standardized coefficients of the polygenic scores calculated for the transmitted and non-transmitted alleles, respectively, when they are analyzed jointly. DGT= β_T - β_{NT}: estimated effect of genetic transmission of the polygenic score. SE=standard error, OR = odds ratio, 95% CI = 95% confidence interval, p = unadjusted p value, p_{FDR} = p value after FDR correction, SmkInit = smoking initiation, CigDay = cigarettes per day, DrnkWk = drink per week, CanU = lifetime cannabis use.
[a] the estimated odds ratio for smoking initiation and lifetime cannabis use from logistic regression models.

Table 3. Parent-of-origin effects on offspring smoking quantity: comparison of transmitted and non-transmitted polygenic scores for cigarettes per day split by paternal and maternal haplotypes

	Maternal PGS$_{T\_CigDay}$			Maternal PGS$_{NT\_CigDay}$			Paternal PGS$_{T\_CigDay}$			Paternal PGS$_{NT\_CigDay}$		
	β (SE)	95% CI	p	β (SE)	95% CI	p	β (SE)	95% CI	p	β (SE)	95% CI	p
Cigarettes per day	**.124 (.012)**	.100, .145	<.0001	**.039 (.015)**	.010, .069	.008	**.155 (.012)**	.131, .179	<.0001	.036 (.020)	.000, .076	.066
Pack-years	**.082 (.011)**	.062, .104	<.0001	**.027 (.013)**	.001, .052	.043	**.107 (.011)**	.085, .127	<.0001	**.034 (.016)**	.001, .065	.036

Note. Structural equation modeling was used to assess the effects of maternal and paternal transmitted (PGS$_T$) and non-transmitted (PGS$_{NT}$) polygenic scores on offspring smoking outcomes. As the analyses used a full information maximum likelihood (FIML) approach to handle missing data, retaining all available data from the full genotyped family sample (N=19,233), there was no list-wise N for the sample. Standardized coefficients (β), bootstrapped standard errors (SE), bootstrapped 95% confidence intervals (CIs) and p value were reported. FDR correction was not applied to the parent-of-origin analyses given the high correlation between outcome measures (cigarettes per day and pack-years), which would make such correction overly conservative for non-independent tests.

Figure 1. Mediation analysis using structural equation modeling: maternal/paternal smoking quantity as a mediator of the associations between transmitted (PGS_T) and non-transmitted (PGS_{NT}) polygenic scores and offspring smoking outcomes.

Note. In all panels, aT path is PGS_T on maternal/paternal smoking quantity; aNT path is PGS_{NT} on maternal/paternal smoking quantity; b path is maternal/paternal smoking quantity on offspring smoking outcomes; c'T path is the direct effect of PGS_T on offspring smoking outcomes, while accounting for the mediator; and c'NT path is the direct effect of PGS_{NT} on offspring smoking outcomes, while accounting for the mediator. Mediation effects are presented below each mediation model for aT*b and aNT*b. Sex and age were included as covariates in all models. Solid line and bold indicate a significant pathway, and dashed line represent a non-significant ($p > .05$) pathway. **A. Maternal cigarettes per day** (CigDay). Maternal CigDay mediated the influence of both PGS_T ($\beta_{total} = 0.125$, SE = 0.012, 95% CI: 0.102 to 0.148) and PGS_{NT} ($\beta_{total} = 0.043$, SE = 0.016, 95% CI: 0.011 to 0.073) on offspring CigDay. **B. Maternal pack-years**. Maternal pack-years mediated the influence of both PGS_T ($\beta_{total} = 0.084$, SE = 0.011, 95% CI: 0.062 to 0.103) and PGS_{NT} ($\beta_{total} = 0.027$, SE = 0.013, 95% CI: 0.000 to 0.052) on offspring pack-years. **C. Paternal pack-years**. Paternal pack-years mediated the influence of both PGS_T ($\beta_{total} = 0.107$, SE = 0.011, 95% CI: 0.085 to 0.128) and PGS_{NT} ($\beta_{total} = 0.034$, SE = 0.016, 95% CI: 0.003 to 0.065) on offspring pack-years.