Data-driven profiling of accelerometer-determined physical behaviors and health outcomes in adults: A systematic review

Vahid Farrahi[1], PhD., Esmaeil Farhang[1], MSc

[1]Research Group of Data Analytics in Sport Science, Institute for Sports and Sport Science, TU Dortmund University

Corresponding author

Prof. Dr. Vahid Farrahi, PhD

Institute for Sports and Sport Science

Otto-Hahn-Straße 3, 44227 Dortmund, Germany

E-mail: vahid.farrahi@tu-dortmund.de

ABSTRACT

Objective: Data-driven population segmentation analysis are analytical approaches applicable
for profiling physical behaviors from wearable accelerometry data. These methods rely on data
rather than predefined, knowledge-driven hypotheses to first identify multiple subgroups
within a sample population and then evaluate their relationships with health outcomes. This
systematic review aims to describe and synthesize multidimensional physical behavior profiles
derived from accelerometry data across adult populations, as identified using data-driven
population segmentation analysis.

Method: Three electronic databases were searched for relevant articles published up to July
2025. Peer-reviewed journal articles that applied data-driven population segmentation analysis
to accelerometer-monitored physical behaviors in adult participants (>18 years) to create
profiles of physical behaviors, and that examined the associations of the profiles with a health
outcome, were considered. Studies conducted with clinical or specific sample populations were
not considered.

Results: Of the 16,289 publications retrieved, 40 were included. The most commonly
employed technique for physical behavior profiling was the machine learning K-means
clustering algorithm (n=18), followed by latent profile analysis (n=8). A diverse set of
descriptor variables was derived from accelerometer signals and utilized. The review of profiles
of physical behavior revealed several hypothesis-generating, preliminary evidence about how
different components and/or aspects of physical behaviors could cluster together and influence
health outcomes.

Conclusion: Data-driven population segmentation analysis are viable analytical approaches
increasingly employed in accelerometry studies to drive physical behavior profiles. The
application of these techniques to accelerometer-measured physical behaviors has generated
data-driven findings regarding how various physical behavior profiles may differentially relate
to health outcomes.

Keywords: actigraphy, clustering, movement behavior profiles, activity behaviors

INTRODUCTION

Over the past few decades, cohort and epidemiological studies have increasingly relied on wearable accelerometers for measurement of physical activity (PA), sedentary behavior, and sleep, collectively known as physical behaviors (Evenson et al., 2022; Migueles et al., 2022). Modern wearable accelerometers can nowadays capture acceleration signals over multiple days, enabling the quantification of various descriptors that reflect the amount, accumulation patterns, timing, and inter-day and intra-day variations of physical behaviors, among others (Backes et al., 2022; Giurgiu et al., 2025). This enhanced ability to measure physical behaviors in greater detail raises questions about how to best model and interpret the relationships between physical behaviors and health outcomes (Farrahi & Rostami, 2024; Migueles et al., 2022; Pedišić et al., 2017; Rosenberger et al., 2019).

Currently, emerging evidence highlights that physical activities, sedentary behaviors, and sleep do not independently influence health outcomes; rather, these behaviors often interact, cluster together, and exhibit codependent relationships with various health markers and indicators (Chastin, Palarea-Albaladejo, et al., 2015; Migueles et al., 2022; Pedišić, 2014; Pedišić et al., 2017). Further complicating this, accumulating evidence suggests that the timing of PA (H. Feng et al., 2023), the regularity and stability of the rest-activity rhythmicity (Makarem et al., 2024), and patterns of accumulation of sedentary and activity behaviors (Herold et al., 2025) are also important characteristics of one's physical behavior profile that may be linked to health outcomes. Given these complexities, proper modeling of the relationships between accelerometer-measured physical behaviors and health outcomes requires careful consideration of the analytical approaches and their adequacy for exploring such potentially sophisticated relationships (Migueles et al., 2022; Pedišić, 2014; Pedišić et al., 2017).

Historically, the dominant approach for exploring the relationships between measures of

physical behaviors and health outcomes was to employ classical regression approaches

(Pedišić, 2014; Pedišić et al., 2017). Recently, new evidence has highlighted a complex

interplay between different components and dimensions of physical behaviors (Rollo et al.,

2020; Rosenberger et al., 2019), leading to the development of innovative analytical

techniques. Analytical approaches, such as compositional data analysis (Dumuid et al., 2019;

McGregor et al., 2020), isotemporal substitution methods (Mekary et al., 2009; Migueles et al.,

2022), multivariate pattern analysis (Aadland et al., 2019), and functional data analysis

(Goldsmith et al., 2015) are now frequently used for exploring the health impacts of physical

behaviors (Blodgett et al., 2024; Farrahi et al., 2023; Leviäkangas et al., 2024; Niemelä et al.,

2023; Seppänen et al., 2025). These novel techniques are generally known to have improved

capacity for better modeling of the complex relationships between PA, sedentary behaviors,

and sleep with health outcomes over classical statistical approaches (Migueles et al., 2022).

From a methodological perspective, however, such analytical approaches can be considered as

variable-centered methods. A variable-centered method refers to techniques that focus on

exploring the relationships between one or more variables of interest and an outcome variable

(Howard & Hoffman, 2017; Woo et al., 2024), assessing how changes in one or more

independent variables (e.g., physical activity, sedentary behavior, sleep) impact the outcome

variable (e.g., weight).

Another type of analytical approach often used for modeling the health influence of

accelerometer-measured physical behaviors is cluster analysis (Kebede et al., 2024; Migueles

et al., 2022). Cluster analysis divide the study population into distinct smaller groups typically

based on a diverse set of descriptive variables (Ezugwu et al., 2022; Howard & Hoffman, 2017;

Liu et al., 2023; Weller et al., 2020; Woo et al., 2024; Xu & Tian, 2015) and are often referred

to as data-driven population segmentation analysis when applied to individual-level data and

characteristics (i.e., to distinguish it from applications such as image clustering and

segmentation) (Liu et al., 2023; Yan et al., 2018). From a methodological perspective, data-

driven population segmentation analysis are conceptually different from variable-centered

approaches (Woo et al., 2024). Rather than focusing on relationships among variables, these

methods use input descriptors to form multiple subgroups in a data-oriented manner,

maximizing within-group similarity and between-group dissimilarity based on individual

characteristics (Howard & Hoffman, 2017; Liu et al., 2023). Such methods are considered data-

driven approach because the subgroups are formed solely based on input descriptor variables,

with subsequent analysis focusing on understanding the differential associations between these

derived subgroups and health outcomes. In practice, several clustering algorithms are used for

data-driven population segmentation analyses (Ezugwu et al., 2022; Woo et al., 2024; Xu &

Tian, 2015), with k-means clustering, hierarchical clustering, and latent profile analysis being

among the most commonly applied methods in health and medical research (Liu et al., 2023;

Yan et al., 2018).

When applied to characteristics of physical behaviors, the subgroups identified and formed

using data-driven population segmentation analysis can be referred to as physical behavior

profiles (Kebede et al., 2024). These profiles reflect various behavioral patterns within the

sample population. Profiles of individuals grouped by similar physical behaviors can then be

compared with other profiles in terms of how the variables combine to create these profiles and

how these combinations relate differentially to health outcomes. Despite the growing reliance

on data-driven population segmentation analysis to derive multidimensional profiles of

physical behaviors from wearable accelerometry data across various populations, it remains

unclear what types of profiles would emerge in different adult populations and how these might

relate to health outcomes. Additionally, the potential of data-driven population segmentation

analysis as an analytical approach for exploring the health associations in physical behavior

research needs further clarification.

This study sought to systematically review the studies that created and identified profiles of

physical behaviors among adult participants with sample populations that were not restricted

to specific health conditions, using data-driven population segmentation analysis (i.e.,

clustering analysis) and wearable accelerometry data. The objectives of this review were to (1)

identify and describe physical behavior profiles emerged in studies applying data-driven

methods on wearable accelerometer-determined physical behaviors and (2) to examine whether

and how these physical behavior profiles relate differentially to health outcomes.

MATERIALS AND METHODS

This systematic review was registered with the International Prospective Register of Systematic

Reviews (ID: CRD42024612460) and performed in accordance with the Preferred Reporting

Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Page et al., 2021).

Search strategy

Three electronic databases (PubMed, Scopus, and Web of Science) were used to identify

relevant articles. The initial search was conducted in June, 2024 and repeated in July 2025. The

search strings were reported in the Supplementary File. The search results were limited for

human subjects and English-language articles. These were not limited by date and included

papers from all years. Additional articles were identified by searching the references of relevant

articles identified by the search strategy.

Eligibility criteria

Full-text original peer-reviewed journal articles applying data-driven population segmentation

analysis to accelerometer-monitored physical behaviors in adult participants (18 years or older)

to create profiles of physical behaviors and examining the associations of the formed profiles

with a health outcome were considered. We defined "health outcome" as any health-related

problem that could be influenced by components (PA intensities, sedentary behaviors, and

sleep) or any of the dimensions (e.g., timing, temporal distribution, accumulation patterns, or

daily/weekly frequencies, etc.) of physical behaviors. Given that data-driven approaches can

be used for different purposes on the same dataset, studies performed with the same analytical

sample were included.

Studies were excluded if they met at least one of the following criteria: (1) physical behaviors

were not determined using wearable accelerometers; (2) data-driven profiles were created by

combining measures of accelerometer-determined physical behaviors with nonphysical

behavior measures (e.g., diet, background information, etc.); (3) profiles were formed based on

knowledge-driven criteria without employing a data-driven approach (e.g., (Länsitie et al.,

2021)); (4) the associations between identified physical behavior profiles and health outcomes

were not statistically examined; (5) the study was not conducted with adult participants; or (6)

the study was conducted with clinical or other specific populations.

Study selection

Covidence review management software was used for the screening and selection of records

retrieved from the databases. One person (VF) performed the initial search and screened the

titles and abstracts of all articles identified by the search strings to exclude articles that were

not relevant according to the eligibility criteria. Articles identified in the title and abstract were

first assessed for eligibility and then examined in full text independently by the authors (VF

and EF). The reviewers were not blinded to authors or journals. Conflicting decisions were

resolved by discussion between the authors until full consensus was achieved. Reasons for

exclusion of articles after full-article screening were recorded. Included articles were then

reviewed and data from the studies were extracted by one of the authors (VF).

Data extraction

The extracted data items on the characteristics of the included studies were first author, year of

publication, sample population, country, study design (cross-sectional or prospective design),

number of participants included in profile analyses, age of participants, proportion of women,

and the primary health outcomes. For profile analysis, extracted data items included input

descriptor variables used for profile analysis and the method used for driving the profiles. The

identified profiles and the number of participants within each profile were recorded.

Statistically significant associations observed between physical behavior profiles and health

outcomes were also recorded. Data items related to accelerometer data collection methods and

characteristics included the type of device used (uniaxial or triaxial), device placement, wear

time protocol, minimum valid hours and days required for inclusion in analyses, and criteria

for nonwear time.

Assessment of bias

Risk of bias was assessed using a combination of two tools, adapted to accommodate both the

observational nature of the studies and the wearable-accelerometer measurements. Four items

from the Newcastle–Ottawa Quality Assessment Scale (Wells et al., 2000) and eight items

based on recommended reporting practices for accelerometer studies were utilized (Montoye

et al., 2018). These items evaluated the quality of selection strategies, outcome measures, and

reporting methods related to accelerometers. A total quality score was determined, with a

maximum of 11 points for cross-sectional studies and 12 points for prospective cohort studies

(low quality [1–5 points], moderate quality [6–10 points], and high quality [10–11/12 points]).

Synthesis of results

The derived profiles could be largely attributed to the input descriptor variables used for profile

analysis. For better synthesis of the results regarding the health implications of identified

profiles, the publications were categorized into five groups based on the input descriptors and

types of profiles identified in each study. These categories were PA profiles, sedentary behavior

profiles, sleep profiles, rest-activity profiles, and joint profiles. Joint profiles refer to those

profiles that were created by incorporating descriptors from at least two components of physical

behaviors (i.e., any combination of descriptors of PA, sedentary behavior, and sleep).

We presented the overview of profiles in color-coded figures. The input descriptors used for

profile analysis, the derived profiles, statistically significant differences in health outcomes

among profiles, and statistical modeling used for examining the associations were presented in

tables**.** A persistent challenge in wearable accelerometer-based measurement is the

incomparability of study results because of large variability in decisions related to

accelerometer data cleaning and processing (Pedišić & Bauman, 2015). To enable a better

comparison among physical behavior profiles identified, profiles with comparable overall

characteristics that were seen in *two or more distinct* sample populations were identified and

marked.

The common approach to examining health associations with data-driven profiles involves

assessing differences relative to a referent profile. When selecting a reference profile, it is

typically chosen based on its representation of either the unhealthiest or healthiest physical

behavior profile. Comparing the differences in the health outcomes among the derived profiles

with respect to one another could also be performed. When statistically significant associations

or differences were observed between a profile and a health outcome, both the direction of the

association and the referent group were indicated. Differences were denoted using upward

arrows (↑) to indicate higher values in the health outcome relative to the referent group.

Downward arrows (↓) were used to indicate lower values. Only statistically significant

associations were reported. For analyses that were gender-stratified, statistical significance was

only noted if the observed associations were significant for both men and women. If multiple

unadjusted and covariate-adjusted models were built, the significant association from the fully

adjusted models was extracted and reported.

The characteristics and health associations of profiles identified across two or more sample

populations were compared and synthesized narratively. Other key emerging profiles in the

studies, together with the health associations linked to the key characteristics, were also

described and summarized narratively.

RESULTS

A flow diagram of the included publications at each stage is presented in Figure 1. Of the

16,289 records initially retrieved (3,244 from PubMed, 7,390 from Scopus, and 5,655 from

Web of Science), 7,156 duplicates were removed. Following the screening of titles and

abstracts, 60 articles were identified as potentially relevant. After conducting a full-text review

of these 60 publications, 38 were deemed eligible for inclusion. Additionally, two more articles

were identified through hand/citation searches, resulting in a final total of 40 included studies.

210
Figure 1: Flow diagram showing the selection of studies for inclusion in the systematic

review

Study design, participant characteristics, and publication characteristics

The main characteristics of the included studies are summarized in Table 1. Publication dates

for the 40 included articles ranged from 2013 to 2025, with the majority published since 2019

(n = 36). Of these studies, 25 were conducted using a cross-sectional design (Aqeel et al., 2021;

Biswas et al., 2022, 2023; Choi et al., 2025; Farrahi et al., 2021, 2022, 2024; Fukuoka et al.,

2018; Full et al., 2019; Graves et al., 2021; Gupta et al., 2020, 2025; Hibbing et al., 2022; P. H.

Lee et al., 2013; Leskinen et al., 2023; Ma et al., 2023; Marschollek, 2016; Matricciani et al.,

2021; Nawrin et al., 2024; Niemelä et al., 2019; O'Regan et al., 2021; Palmberg et al., 2024;

Paolillo et al., 2023; Smagula et al., 2022; Yamamoto et al., 2023) while 15 studies had a

prospective study design (Albalak et al., 2023; Bai, Ning, et al., 2024; Bai, Shao, et al., 2024;

Bai, Zhou, et al., 2024; Chung et al., 2024; Evenson et al., 2017; Fabrizio et al., 2024; Q. Feng

et al., 2025; Li et al., 2024; Reuter et al., 2020; Shim et al., 2023; von Rosen et al., 2020;

Yamamoto et al., 2025; Yerramalla et al., 2024; Zhang et al., 2024).

Among all included studies, the number of participants in the sample population used for

physical behavior profiling ranged between 124 (Paolillo et al., 2023) to 102,323 (Bai, Zhou,

et al., 2024). Sample populations were drawn from 13 countries. Sixteen studies were

performed with a sample population from the United States, seven from the United Kingdom,

four from Finland, two from Canada, two from Japan, and two from Denmark. Other sample

populations were from Sweden, the Netherlands, Switzerland, Ireland, Australia Korea, and

China. A large number of studies analyzed accelerometer data from the National Health and

Nutrition Examination Survey study (NHANES, n = 10).

Table 1: Main characteristics of 40 studies included in the review, listed by study design, publication year, and primary health outcome.

Study, first author (year)	Sample population / Country	Study design	Age, mean (SD)	Number of participants (female%)	Profiling method	Profile type	Primary health outcomes
(Yerramalla et al., 2024)	Whitehall II study / UK	Prospective	69.4 (5.7)	3,991(26%)	K-means	Joint	All-cause mortality
(Zhang et al., 2024)	NHANES 2011–2014 / USA	Prospective	49.7 (17.4)	3,423 (51%)	K-means	Physical activity / sleep	All-cause mortality
(Bai, Zhou, et al., 2024)	UK biobank / UK	Prospective	56.1 (7.8)	102,323 (56%)	K-means	Physical activity	All-cause mortality, CVD mortality, CVD incidence
(Fabrizio et al., 2024)*	CoLaus-PsyCoLaus / Switzerland	Prospective	61.2 (9.7)	2,465 (55%)	K-means	Physical activity	CVD events
(Chung et al., 2024)	MESA / USA	Prospective	68.2 (9.0)	1,759 (54%)	K-means	Sleep	All-cause mortality
(Shim et al., 2023)	NHANES 2011-2014 / USA	Prospective	Median (IQR): 51 (36–65)	7,297 (53%)	Hierarchical clustering approach using Ward's linkage algorithm	Physical activity	All-cause mortality
(von Rosen et al., 2020)	Sweden Attitude Behaviour and Change / Sweden	Prospective	53 (NR)	851 (56%)	Latent class analysis	Joint	All-cause mortality
(Evenson et al., 2017)	NHANES, 2003-2006 cycles / USA	Prospective	56.5 (NR)	4,510 (54%)	Latent profile analysis	Physical activity / sedentary behavior	All-cause mortality
(Yamamoto et al., 2025)	Toon Health Study / Japan	Prospective	56.2 (11.7)	794 (100%)	Latent profile analysis	Physical activity	Metabolic syndrome onset
(Albalak et al., 2023)	UK biobank / UK	Prospective	61.6 (7.8)	86,657 (58%)	K-means	Physical activity	CVD incidence
(Bai, Ning, et al., 2024)	UK biobank / UK	Prospective	55.9 (7.8)	96,687 (56%)	K-means	Physical activity	17 types of site-specific cancer incidence
(Q. Feng et al., 2025)	UK biobank / UK	Prospective	61.7 (7.9)	89,439 (57%)	K-means	Physical activity	Type 2 diabetes incidence
(Bai, Shao, et al., 2024)	UK biobank / UK	Prospective	56.0 (7.8)	99,532 (57%)	K-means	Physical activity	Type 2 diabetes incidence
(Li et al., 2024)	UK biobank / UK	Prospective	61.2 (7.9)	70,617 (60%)	K-means	Physical activity	Incident hypertension
(Reuter et al., 2020)	OPACH / USA	Prospective	79 (7)	6,204 (100%)	K-means for longitudinal data	Sedentary behavior	Physical functioning

					combined with hierarchical clustering		
(Choi et al., 2025)	KNHANES / Korea	Cross-sectional	Age: ≥65	425 (54%)	Latent profile analysis	Joint	Subjective health, metabolic syndrome, number of chronic diseases
(Gupta et al., 2025)	DPHACTO / Denmark	Cross-sectional	45.1 (9.7)	807 (56%)	Compositional latent class analysis	Joint	Cardiorespiratory fitness, systolic blood pressure, self-rated health, low back pain
(Farrahi et al., 2024)	NHANES, 2003-2006 cycles / USA	Cross-sectional	43.1 (12.5)	1,812 (53%)	Convolutional autoencoders with K-means	Joint	Insulin, triglycerides, homeostasis insulin resistance, plasma glucose, C-reactive protein, total/HDL cholesterol ratio, LDL, WC, BMI
(Leskinen et al., 2023)	Maastricht Study / Netherlands	Cross-sectional	60.2 (8.6)	6,702 (50%)	Group-based multi-trajectory modeling	Physical activity	BMI, waist, HbA1c, fasting glucose, 2-hour glucose, total/HDL cholesterol ratio, triglycerides, systolic BP, diastolic BP
(Ma et al., 2023)	NHANES 2003-2006 cycles / USA	Cross-sectional	Age: ≥20	5,285 (49%)	K-means	Physical activity	BMI and WC
(Biswas et al., 2023)	CHMS, 2007-2017 cycles / Canada	Cross-sectional	38.5 (NR)	8,229 (47%)	Dynamic time warping with hierarchical clustering	Physical activity	WC, systolic BP, diastolic BP, cholesterol
(Yamamoto et al., 2023)	Toon Health Study / Japan	Cross-sectional	57.7 (12.5)	1,850 (66%)	Latent profile analysis	Physical activity	Metabolic syndrome, abdominal obesity, hypertension, hyperglycemia, dyslipidemia
(Biswas et al., 2022)	CHMS, 2007-2017 cycles / Canada	Cross-sectional	42.4 (0.3)	8,909 (47%)	Dynamic time warping with hierarchical clustering	Physical activity	10-year risk for CVD event
(Farrahi et al., 2022)	NFBC1966 / Finland	Cross-sectional	46.6 (0.6)	3,702 (59%)	KmL3D (K-means for joint longitudinal data)	Joint	Fasting serum insulin, 2-hour insulin, fasting glucose, 2-hour glucose, triglycerides, total/HDL cholesterol ratio, LDL/HDL cholesterol ratio, body fat, fat mass, visceral fat area, BMI, WC
(Hibbing et al., 2022)	NHANES 2003-2006 cycles / USA	Cross-sectional	54.3 (17.0)	5,495 (49%)	K-medoid	Sedentary behavior	10-year CVD risk
(Aqeel et al., 2021)	NHANES, 2003-2006 cycles / USA	Cross-sectional	Age range: 20-65	1,999 (48%)	Dynamic time warping and kernel K-means	Physical activity	BMI, WC, hemoglobin A1c, Total cholesterol, triglycerides, fasting plasma glucose, HDL cholesterol, systolic BP, diastolic BP
(Farrahi et al., 2021)	NFBC1966 / Finland	Cross-sectional	46.6 (0.6)	4,439 (57%)	K-means	Sedentary behavior	Fasting serum insulin, 2-hour insulin, fasting glucose, 2-hour glucose, triglycerides, total/HDL cholesterol ratio, LDL/HDL cholesterol ratio, body fat, fat mass, visceral fat area, BMI, WC
(Matricciani et al., 2021)	CheckPoint / Australia	Cross-sectional	44.0 (5.1)	1,137 (87%)	K-means	Sleep	SBP, DPB, BMI, glycoprotein acetyls, apolipoprotein, metabolic syndrome severity score
(Gupta et al., 2020)	DPHACTO / Denmark	Cross-sectional	45.1 (9.7)	807 (56%)	Compositional latent class analysis	Joint	BMI, WC, body fat percentage

(Niemelä et al., 2019)	NFBC1966 / Finland	Cross-sectional	46 (NR)	4582 (58%)	X-means	Physical activity	10-year CVD risk score, HDL cholesterol, total cholesterol, systolic BP, being diabetic
(Marschollek, 2016)	NHANES 2005-2006 cycles / USA	Cross-sectional	NR	7,334 (NR)	X-means	Physical activity	HDL cholesterol, C-reactive protein, BMI
(P. H. Lee et al., 2013)	Hong Kong Jockey Club FAMILY / China	Cross-sectional	43.2 (16.4)	1,714 (54%)	K-means	Physical activity	Having chronic health condition, body fat percentage, BMI
(Fukuoka et al., 2018)	mPED / USA	Cross-sectional	52.4 (11.2)	219 (100%)	K-means	Physical activity	Depression symptoms, BMI, WC, hip circumference, resting systolic BP, resting diastolic BP, cholesterol, triglycerides, LDL, HDL, hemoglobin A1c, fasting plasma glucose
(Nawrin et al., 2024)	NHANES 2011-2012 cycles / USA	Cross-sectional	NR	6,613 (NS)	K-means	Physical activity	Depression
(Smagula et al., 2022)	NHANES 2011-2014 cycles / USA	Cross-sectional	72.9 (7.3)	1,800 (57%)	Finite normal mixture modelling	Rest-activity	Depression symptoms and cognitive performance
(O'Regan et al., 2021)	Move for Life / Ireland	Cross-sectional	Age: ≥45	485 (79%)	Hierarchical clustering	Joint	Having mental illness, obesity, having complex multimorbidity
(Palmberg et al., 2024)	AGNES / Finland	Cross-sectional	75, 80, and 85 years old	441 (60%)	K-means	Joint	Indicators of physical capacity, life-space mobility, depression symptom
(Graves et al., 2021)	MOVEUP and DECOS / USA	Cross-sectional	71.3 (6.7)	181 (79%)	K-means	Rest-activity	Perceived physical fatigability
(Full et al., 2019)	Convenient sample from four US cities / USA	Cross-sectional	55.4 (10.2)	372 (100%)	Latent profile analysis	Joint	Physical functioning, BMI, self-rated health, number of health conditions
(Paolillo et al., 2023)	Community-dwelling older adults / USA	Cross-sectional	72.2 (8.37)	124 (55%)	Latent profile analysis	Joint	Executive functioning, Memory, Vascular burden

*: Profiles were created based on data collected from two follow-ups; the items reported here correspond to the profiles identified during the first follow-up and were prospectively examined in relation to cardiovascular disease events. Abbreviations: NR = Not reported; BP = Blood pressure; BMI = Body mass index; CVD = Cardiovascular diseases; HDL = High-density lipoprotein; LDL = Low-density lipoprotein; WC = Waist circumference; CHMS = Canadian Health Measures Survey; DECOS = Developmental Epidemiologic Cohort Study; DPHACTO = Danish PHysical ACTivity cohort with Objective measurements; KNHANES = Korea National Health and Nutrition Examination Survey; MESA = Multi-Ethnic Study of Atherosclerosis; MOVEUP = Mobility and Vitality Lifestyle Program; mPED = Mobile Phone Based Physical Activity Education; NFBC1966 = Northern Finland Birth Cohort 1966; NHANES = National Health and Nutrition Examination Survey; OPACH = Objective Physical Activity and Cardiovascular Health Study.

Quality score, data collection methods, and accelerometer characteristics

Details of the accelerometer data collection measures are summarized in Supplementary Table S1. Across the studies, 12 required participants to wear the accelerometer during waking hours for 7 days (Aqeel et al., 2021; Biswas et al., 2022, 2023; Choi et al., 2025; Evenson et al., 2017; Farrahi et al., 2024; Fukuoka et al., 2018; Hibbing et al., 2022; Leskinen et al., 2023; Marschollek, 2016; von Rosen et al., 2020; Yamamoto et al., 2023, 2025), while 11 had continuous 7-day measurement including sleep (Albalak et al., 2023; Bai, Ning, et al., 2024; Bai, Shao, et al., 2024; Bai, Zhou, et al., 2024; Q. Feng et al., 2025; Graves et al., 2021; Li et al., 2024; Nawrin et al., 2024; Reuter et al., 2020; Shim et al., 2023; Zhang et al., 2024). The number of valid accelerometer measurement days deemed sufficient across the studies ranged from 1 to 7 consecutive days.

Individual publication quality scores are shown in Supplementary Table S2. Twelve studies with a prospective study design (Albalak et al., 2023; Bai, Ning, et al., 2024; Bai, Shao, et al., 2024; Evenson et al., 2017; Fabrizio et al., 2024; Q. Feng et al., 2025; Li et al., 2024; Reuter et al., 2020; Shim et al., 2023; von Rosen et al., 2020; Yerramalla et al., 2024; Zhang et al., 2024) and 20 studies with a cross-sectional study design were rated as high quality (Aqeel et al., 2021; Biswas et al., 2022, 2023; Choi et al., 2025; Farrahi et al., 2021, 2022, 2024; Fukuoka et al., 2018; Graves et al., 2021; Gupta et al., 2025, 2020; Hibbing et al., 2022; P. H. Lee et al., 2013; Leskinen et al., 2023; Ma et al., 2023; Nawrin et al., 2024; Niemelä et al., 2019; O'Regan et al., 2021; Palmberg et al., 2024; Yamamoto et al., 2023). Three studies with a prospective study design (Bai, Zhou, et al., 2024; Chung et al., 2024; Yamamoto et al., 2025) and five studies with a cross-sectional study design were rated as moderate quality (Full et al., 2019; Marschollek, 2016; Matricciani et al., 2021; Paolillo et al., 2023; Smagula et al., 2022). No study was rated as low quality.

Health outcomes

The health outcomes assessed in the included studies are shown in Table 1. In most prospective studies, the primary health outcome (n = 7) was all-cause mortality (Bai, Zhou, et al., 2024; Chung et al., 2024; Evenson et al., 2017; Shim et al., 2023; von Rosen et al., 2020; Yerramalla et al., 2024; Zhang et al., 2024). Other primary health outcomes in prospective studies were cardiovascular disease (CVD) incidence (Albalak et al., 2023; Bai, Zhou, et al., 2024; Fabrizio et al., 2024), type-2 diabetes incidence (Bai, Shao, et al., 2024; Q. Feng et al., 2025), hypertension incidence (Li et al., 2024), onset of metabolic syndrome (Yamamoto et al., 2025), site-specific cancer incidence (Bai, Ning, et al., 2024), and deterioration in physical functioning (Reuter et al., 2020). Primary health outcomes in most studies with a cross-sectional design (n = 16) were indicators of cardiometabolic health, CVD risk factors, metabolic syndrome, and/or obesity indicators (Aqeel et al., 2021; Biswas et al., 2022, 2023; Choi et al., 2025; Farrahi et al., 2021, 2022, 2024; Fukuoka et al., 2018; Gupta et al., 2020; Hibbing et al., 2022; P. H. Lee et al., 2013; Leskinen et al., 2023; Ma et al., 2023; Marschollek, 2016; Matricciani et al., 2021; Niemelä et al., 2019; Yamamoto et al., 2023). Other primary health outcomes in cross-sectional studies were depression or mental health outcomes (Nawrin et al., 2024; O'Regan et al., 2021; Smagula et al., 2022); indicators of physical function capacity, or frailty (Full et al., 2019; Graves et al., 2021; Palmberg et al., 2024); low back pain (Gupta et al., 2025); or indicators of cognitive health (Paolillo et al., 2023).

Profiling method

Of the 40 included studies (see Table 1), 18 used the K-means clustering algorithm to derive physical behavior profiles (Albalak et al., 2023; Bai, Ning, et al., 2024; Bai, Shao, et al., 2024; Bai, Zhou, et al., 2024; Chung et al., 2024; Fabrizio et al., 2024; Farrahi et al., 2021; Q. Feng et al., 2025; Fukuoka et al., 2018; Graves et al., 2021; P. H. Lee et al., 2013; Li et al., 2024; Ma

et al., 2023; Matricciani et al., 2021; Nawrin et al., 2024; Palmberg et al., 2024; Yerramalla et

al., 2024; Zhang et al., 2024). Seven of these applied different variations and forms of K-means,

including K-means for longitudinal data (Farrahi et al., 2022; Reuter et al., 2020), dynamic

time warping and kernel K-means (Aqeel et al., 2021), convolutional autoencoders with K-

means (Farrahi et al., 2024), X-means (Marschollek, 2016; Niemelä et al., 2019), and K-

medoid (Hibbing et al., 2022). Nine studies used latent profile analysis (Choi et al., 2025;

Evenson et al., 2017; Full et al., 2019; Gupta et al., 2020, 2025; Paolillo et al., 2023; von Rosen

et al., 2020; Yamamoto et al., 2023, 2025), four employed hierarchical clustering (Biswas et

al., 2022, 2023; O'Regan et al., 2021; Shim et al., 2023), one applied finite mixture modelling

(Smagula et al., 2022), and one used group-based multi-trajectory modeling (Leskinen et al.,

2023).

Profile types

Among the 40 included studies, 22 identified profiles of PA, four identified profiles of

sedentary behavior, three identified profiles of sleep, and two identified profiles of rest-activity

rhythmicity. One study reported both PA and sleep profiles (Zhang et al., 2024) and another

reported both PA and sedentary profiles (Evenson et al., 2017); these were counted in both

respective categories. Joint profiling was performed in 11 studies.

Profiles of physical activity

An overview of the profiles identified in studies creating PA profiles is shown in Figure 2. The

number of profiles identified ranged from two to seven. The input descriptor variables, the

derived PA profiles, and their associations with health outcomes in each study are listed in

Table 2.

All but one (Bai, Zhou, et al., 2024) of the studies creating PA profiles (n = 21), specified the

accelerometer-derived descriptors used as input for profile analysis. PA profiling was most

commonly conducted using hourly metrics derived from accelerometer outputs, although the

specific summary metrics varied across studies. Of the 21 studies, 12 utilized average hourly

summary metrics to drive PA profiles (Albalak et al., 2023; Aqeel et al., 2021; Bai, Ning, et al.,

2024; Bai, Shao, et al., 2024; Fabrizio et al., 2024; Q. Feng et al., 2025; P. H. Lee et al., 2013;

Li et al., 2024; Ma et al., 2023; Nawrin et al., 2024; Shim et al., 2023; Zhang et al., 2024).

These 12 studies computed the summary metrics for each hour of the 24-hour day. The

summary metrics included hourly average acceleration (Albalak et al., 2023; Bai, Ning, et al.,

2024; Bai, Shao, et al., 2024; Fabrizio et al., 2024; Q. Feng et al., 2025; Li et al., 2024), hourly

monitor-independent movement summary (MIMS) unit (Shim et al., 2023; Zhang et al., 2024),

hourly activity counts (Aqeel et al., 2021; Nawrin et al., 2024), and hourly moderate-to-

vigorous PA (MVPA) minutes (Ma et al., 2023). One study used hourly activity counts

calculated separately over weekdays and weekend days (P. H. Lee et al., 2013). Four other

studies computed summary metrics that were aggregated over different time intervals than on

an hourly basis (Biswas et al., 2022, 2023; Leskinen et al., 2023; Niemelä et al., 2019).

Key profile characteristics and their relationships with health outcomes

Of the 22 studies identifying PA profiles, 14 identified PA profiles that were characterized by

having the lowest level of PA relative to other PA profiles in the sample study population. These

profiles are depicted in orange in Figure 2. Conversely, of the 22 studies identifying PA profiles,

13 identified PA profiles that were distinguished by having the highest level of PA relative to

other identified profiles in the sample study population. These profiles are shown in black in

Figure 2.

When a low PA profile emerged, it was generally one of the largest profiles in terms of

participant numbers, with few exceptions (Bai, Ning, et al., 2024; Bai, Shao, et al., 2024; Bai,

Zhou, et al., 2024). In contrast to low PA profiles, high PA profiles represented a smaller group

in terms of participant numbers (see Figure 2). High PA profiles were found to have the most

favorable health outcomes, exhibiting reduced risks of all-cause mortality (Bai, Zhou, et al.,

2024; Evenson et al., 2017; Shim et al., 2023), incident CVD and CVD mortality (Bai, Zhou,

et al., 2024), incident type 2 diabetes (Bai, Shao, et al., 2024), and certain types of cancer (Bai,

Ning, et al., 2024), and better markers of cardiometabolic health and obesity (Biswas et al.,

2022, 2023; P. H. Lee et al., 2013; Leskinen et al., 2023; Marschollek, 2016; Niemelä et al.,

2019; Yamamoto et al., 2023).

Across all studies performing PA profiling, 17 identified at least one profile whose PA peaked

at a certain time within a 24-hour cycle. Thirteen of these studies performed with six unique

sample populations identified a profile that could be characterized by having a relatively higher

PA peak at some time during morning. These profiles are highlighted in green in Figure 2.

Profiles of afternoon or evening PA characterized by having a greater proportion of PA in the

afternoon or evening emerged in 10 studies performed with six unique sample populations.

These profiles are highlighted in yellow in Figure 2. Five studies, conducted across four distinct

sample populations, identified profiles characterized by relatively higher physical activity

during the day (Biswas et al., 2022, 2023; Q. Feng et al., 2025; Ma et al., 2023; Zhang et al.,

2024). These profiles are highlighted in gray in Figure 2.

There were substantial variations among the studies, making it difficult to compare the profiles

with respect to the exact timing of peak PA. Those studies identifying morning profiles

observed that PA in morning profiles generally peaked *approximately* at some time in between

5:00 and 10:00 (Albalak et al., 2023; Biswas et al., 2022, 2023; Ma et al., 2023; Zhang et al.,

2024) or 9:00 and 12:00 (Albalak et al., 2023; Aqeel et al., 2021; Bai, Ning, et al., 2024; Bai,

Shao, et al., 2024; Bai, Zhou, et al., 2024; Fukuoka et al., 2018; Leskinen et al., 2023; Li et al.,

2024; Nawrin et al., 2024). The PA peak in afternoon and evening profiles *approximately*

occurred at some time in between 18:00 and 21:00 (Albalak et al., 2023; Q. Feng et al., 2025;

Fukuoka et al., 2018; Ma et al., 2023; Zhang et al., 2024), or showed a steady increase starting

sometime between 12:00 and 16:00 and continuing up to nearly midnight (Biswas et al., 2022,

2023; Nawrin et al., 2024; Niemelä et al., 2019). Studies identifying day profiles generally

reported relatively higher physical activity from morning until the end of day around 17:00–

18:00.

The association of morning, day, afternoon, and evening profiles with health outcomes was

somewhat mixed and different, depending on the health outcomes (see Table 2). One study

reported that the "Early morning" profile, but not the "Late afternoon," had a higher mortality

risk compared with the "Midday" profile (Zhang et al., 2024). Two studies were performed

with the same sample populations and identified the same profile (Albalak et al., 2023; Q. Feng

et al., 2025). One of these reported that both "Early morning peak" and "Late morning peak"

had a lower incidence of coronary artery disease compared with "Average pattern,"

characterized by having average midday PA (Albalak et al., 2023). In the same study (Albalak

et al., 2023), "Late morning peak" was also associated with a lower incidence of stroke and

ischemic stroke compared with the "Average pattern." In the other study (Q. Feng et al., 2025),

"Early morning peak," but not "Late morning," was associated with a lower incidence of type

2 diabetes compared with "Midday average pattern." Another study reported that "Morning

engaged" had higher depression symptoms compared with "Afternoon engaged," while both

"Morning engaged" and "Afternoon engaged" had lower depression symptoms compared with

the "Unengaged" profile (Fukuoka et al., 2018).

Three studies identified a profile in the same sample population whose PA peaked during the

day at around 9:00–10:00 (Bai, Ning, et al., 2024; Bai, Shao, et al., 2024; Bai, Zhou, et al.,

2024). These reported that the single peak profile was associated with lower all-cause mortality,

CVD mortality, stroke, and ischemic heart disease incidence compared with the profile with

the lowest PA (Bai, Zhou, et al., 2024), a lower incidence of type 2 diabetes compared with the

profile with minimal PA (Bai, Shao, et al., 2024), and a lower incidence of breast, kidney, and

myeloma cancer incidence compared with the profile with the lowest PA (Bai, Ning, et al.,

2024).

One study reported that "High daytime activity," but not "Moderate evening activity," was

associated with lower waist circumference and cholesterol levels when compared with the

lowest activity profile (Biswas et al., 2023). In contrast, another study using the same sample

population found no associations for "High daytime activity" relative to the lowest activity

profile, while "Moderate evening activity" was linked to a lower composite 10-year CVD score

(Biswas et al., 2022). In another study, the "Evening active" profile did not have differences in

terms of cardiometabolic health markers or composite 10-year CVD score when compared with

all other profiles, including the profile with the lowest activity level (Niemelä et al., 2019).

However, one study reported that "Evening (MVPA)" had higher BMI and waist circumference

compared with both "Midday (MVPA)" and "Morning (MVPA)" (Ma et al., 2023). Another

study identifying an "Evening dominant" profile observed higher depression symptoms for this

profile when compared with the "Morning dominant" profile (Nawrin et al., 2024).

Study (first author, year)	Derived profiles (name or main characteristics)					
▲,1(Zhang et al., 2024)	Early morning (2,159)	Late afternoon (1,677)	Midday (2,837)			
▲,2(Bai, Zhou, et al., 2024)	Persistently low (6,388)	Moderate and stable (36,678)	Single increase (29,975)	Double increase (17,291)	Vigorous activity patterns (11,991)	
▲(Fabrizio et al., 2024)	10 a.m. – 2 p.m. PA peak (386)	7 a.m. – 12 a.m. PA peak (364)	8 a.m. – 1 p.m. PA peak (670)	No clear PA peak (639)		
▲,1(Shim et al., 2023)	High activity (1,628)	Low activity (2,670)	Mild circadian rhythm disruption (1,256)	Severe circadian rhythm disruption (558)	Very low activity (1,185)	
▲,3,*(Evenson et al., 2017)	Class 6 [Most active] (41)	Class 5 (216)	Class 4 (66)	Class 3 (748)	Class 2 (1,892)	Class 1 [Least active] (1,547)
▪,4(Yamamoto et al., 2025)	Group-A [low PA] (407)	Group-B (81)	Group-C (248)	Group-D [high PA] (58)		
▲,2(Albalak et al., 2023)	Average pattern (27,552)	Early morning peak (19,843)	Late morning peak (22,648)	Evening peak (16,614)		
▲,2(Q. Feng et al., 2025)	Mid-day average pattern (27,681)	Early morning peak (18,614)	Late morning peak (23,299)	Early evening peak (19,845)		
▲,2(Bai, Shao, et al., 2024)	Consistently low (13,764)	Single peak (47,500)	Double peak (18,681)	Intense (19,587)		
▲,2(Li et al., 2024)	PA peak at around 10 a.m. every day (37,216)	Peaks at around 8 a.m. and 6 p.m. daily (20,045)	A relatively uniform distribution of physical intensity throughout the day (12,666)	Even daily distribution of PA intensity (690)		
▲,3(Ma et al., 2023)	Morning [MVPA] (642)	Midday [MVPA] (2,456)	Evening [MVPA] (2,187)			
▲,5(Biswas et al., 2023)	Lowest activity (2,808)	Moderate consistent activity (3,219)	Fluctuating moderate activity (1,194)	High daytime activity (713)	Moderate evening activity (228)	Highest activity 750
4(Yamamoto et al., 2023)	Group-A [low PA] (1,047)	Group-B (261)	Group-C (418)	Group-D [high PA] (124)		
▲,5(Biswas et al., 2022)	Lowest activity (2,808)	Moderate consistent activity (3,219)	Fluctuating moderate activity (1,194)	High daytime activity (713)	Moderate evening activity (225)	Highest activity (750)
▲(Leskinen et al., 2023)	Consistently inactive (1,301)	Consistently low active (2,498)	Active on weekdays (881)	Early birds (138)	Consistently moderately active (449)	Weekend warriors (453) / Consistently highly active (352)
▲,3(Aqeel et al., 2021)	Lowest activity counts from 6:00–23:00 (783)	High activity counts with a decline after 18:00 (247)	Lower activity counts between 6:00–12:00 (336)	Higher activity counts between 8:00–11:00 (633)		
▲(Niemelä et al., 2019)	Inactive (1,881)	Evening active (802)	Moderately active (1,297)	Very active (602)		
▪,3(Marschollek, 2016)	High overall physical with irregular high-intensity activity (92)	Regularly and extensively active (2,230)	Irregularly and less extensively active (2,990)	Inactive (2,010)		
▲(P. H. Lee et al., 2013)	Active (480)	Less active (1,234)				
▲(Fukuoka et al., 2018)	Morning engaged (65)	Afternoon engaged (48)	Unengaged (102)			
▲,1(Nawrin et al., 2024)	Morning dominant (1,964)	All-day dominant (1,792)	All-day or morning dominant (1,855)	Evening dominant (1,002)		
▲,2(Bai, Ning, et al., 2024)	Persistent low (14,037)	Single peak (46,100)	Double peak (17,763)	Vigorous (18,787)		

Comparable profiles:

■ Highest activity profile
■ Lowest activity profile
■ Morning peak
■ Afternoon/evening peak
■ Day peak

Figure 2: Overview of derived profiles in 22 studied identifying physical activity (PA) profiles. The numbers shown in () are the number of participants in the profile. The profiles are specified according to the names given to them in the study or a short outline of the main characteristics. To facilitate comparison across studies, profiles with comparable overall characteristics *seen in at least two or more distinct sample populations* are highlighted using distinct background colors. Note that the color-coded profiles are not necessarily exactly identical or directly comparable, because of the underlying heterogeneities related to profile analysis and/or accelerometer data cleaning/processing. 1, 2, 3, 4, 5: studies marked with the same number were performed with the same sample population. ▲: High quality study. ▪: Moderate quality study. Individual publication quality scores are shown in Supplementary Table S2. *: Five models were created based on different input descriptor variables into the profiling approach; profiles reported here are for the profiles identified on the basis of average activity counts/minute.

Table 2: Input descriptor variables, profiles identified, and the statistically significant associations observed among the profiles identified in 22 studies performing physical activity (PA) profiling.

Study (first author, year)	Input descriptor variables	Derived profiles, name or main characteristics	Number of participants	Statistically significant associations	Statistical approach used to examine associations
(Zhang et al., 2024)	Hourly proportion of total MIMS units accumulated in each hour over a 24-hour period	Early morning	2,159	All-cause mortality risk (vs. Ref.)↑	Cox proportional hazards models
		Late afternoon	1,677	No statistically significant association (vs. Ref.)	
		Midday	2,837	Ref.	
(Bai, Zhou, et al., 2024)	Not reported clearly	Persistently low	6,388	Ref.	Cox proportional hazards models
		Moderate and stable	36,678	All-cause mortality (vs. Ref.)↓, CVD mortality (vs. Ref.)↓, Incident ischemic heart disease (vs. Ref.)↓, Incident stroke (vs. Ref.)↓	
		Single increase	29,975	All-cause mortality (vs. Ref.)↓, CVD mortality (vs. Ref.)↓, Incident ischemic heart disease (vs. Ref.)↓, Incident stroke (vs. Ref.)↓	
		Double increase	17,291	All-cause mortality (vs. Ref.)↓, CVD mortality (vs. Ref.)↓, Incident ischemic heart disease (vs. Ref.)↓, Incident stroke (vs. Ref.)↓	
		Vigorous activity patterns	11,991	All-cause mortality (vs. Ref.)↓, CVD mortality (vs. Ref.)↓, Incident ischemic heart disease (vs. Ref.)↓, Incident stroke (vs. Ref.)↓	
(Fabrizio et al., 2024)	Mean hourly acceleration over a 24-hour period standardized by average 24-hour acceleration across all measurement days	10 a.m. – 2 p.m. PA peak	386	Ref.	Cox proportional hazards models
		7 a.m. – 12 a.m. PA peak	364	CVD event (vs. Ref.)↑	
		8 a.m. – 1 p.m. PA peak	670	No statistically significant difference (vs. Ref.)	
		No clear PA peak	639	CVD event (vs. Ref.)↑	
(Shim et al., 2023)	Mean hourly MIMS units over a 24-hour period across all measurement days, with recursive feature selection to remove irrelevant input features	High activity	1,628	Ref.	Cox proportional hazards models
		Low activity	2,670	No statistically significant difference (vs. Ref.)	
		Mild circadian rhythm disruption	1,256	All-cause mortality risk (vs. Ref.)↑	
		Severe circadian rhythm disruption	558	No statistically significant difference (vs. Ref.)	
		Very low activity	1,185	All-cause mortality risk (vs. Ref.)↑	
[1](Evenson et al., 2017)	Average activity counts/minute on every day of the week	Class 6 [Most active]	41	No death observed	Cox proportional hazards models
		Class 5	216	All-cause mortality (vs. Ref.)↓	
		Class 4	66	No death observed	
		Class 3	748	All-cause mortality (vs. Ref.)↓	
		Class 2	1,892	All-cause mortality (vs. Ref.)↓	
		Class 1 [Least active]	1,547	Ref.	

(Yamamoto et al., 2025)	Number of steps, LPA, MVPA, and the ratio of MVPA to total activity	Group-A [low PA]	407	Ref.	Logistic regression analysis
		Group-B	81	No statistically significant difference (vs. Ref.)	
		Group-C	248	Metabolic syndrome onset (Ref.)↓	
		Group-D [high PA]	58	No statistically significant difference (vs. Ref.)	
(Albalak et al., 2023)	Mean hourly relative acceleration over a 24-hour period across all measurement days	Average pattern	27,552	Ref.	Cox proportional hazards models
		Early morning peak	19,843	Incident coronary artery disease (vs. Ref.)↓	
		Late morning peak	22,648	Incident coronary artery disease (vs. Ref.)↓, Incident stroke (vs. Ref.)↓, Incident ischemic stroke (vs. Ref.)↓	
		Evening peak	16,614	No statistically significant difference (vs. Ref.)	
(Q. Feng et al., 2025)	Mean hourly relative acceleration over a 24-hour period across all measurement days	Mid-day average pattern	27,681	Ref.	Cox proportional hazards models
		Early morning peak	18,614	No statistically significant difference (vs. Ref.)	
		Late morning peak	23,299	Incident type 2 diabetes (vs. Ref.)↓	
		Early evening peak	19,845	No statistically significant difference (vs. Ref.)	
(Bai, Shao, et al., 2024)	Mean hourly acceleration over a 24-hour period across all measurement days	Consistently low	13,764	Ref.	Cox proportional hazards models
		Single peak	47,500	Incident type 2 diabetes (vs. Ref.)↓	
		Double peak	18,681	Incident type 2 diabetes (vs. Ref.)↓	
		Intense	19,587	Incident type 2 diabetes (vs. Ref.)↓	
(Li et al., 2024)	Mean hourly acceleration over a 24-hour period across all measurement days	PA peak at around 10 a.m. every day	37,216	Incident ischemic heart disease (vs. Ref.)↓	Cox proportional hazards models
		Peaks at around 8 a.m. and 6 p.m. daily	20,045	Incident ischemic heart disease (vs. Ref.)↓	
		A relatively uniform distribution of physical intensity throughout the day	12,666	Ref.	
		Even daily distribution of PA intensity	690	No statistically significant difference (vs. Ref.)	
(Ma et al., 2023)	Mean hourly proportion of MVPA over a 24-hour period	Morning [MVPA]	642	Ref.	Linear regression models
		Midday [MVPA]	2,456	BMI (vs. Ref.)↑, WC (vs. Ref.)↑	
		Evening [MVPA]	2,187	BMI (vs. Ref.)↑, WC (vs. Ref.)↑	
(Biswas et al., 2023)	Average counts per minute over 10-minute intervals throughout the 24-hour period	Lowest activity	2,808	Ref.	Ordinary least squares regression models
		Moderate consistent activity	3,219	WC (vs. Ref.)↓	
		Fluctuating moderate activity	1,194	WC (vs. Ref.)↓, Systolic BP↓, Cholesterol (vs. Ref.)↓	
		High daytime activity	713	WC (vs. Ref.)↓, Cholesterol (vs. Ref.)↓	
		Moderate evening activity	225	No differences vs. Ref.	
		Highest activity	750	WC (vs. Ref.)↓, Systolic BP (vs. Ref.)↓, Diastolic BP (vs. Ref.)↓, Cholesterol (vs. Ref.)↓	
(Yamamoto et al., 2023)	Number of steps, LPA, MVPA, and the ratio of MVPA to total activity	Group-A [low PA]	1,047	Ref.	Logistic regression models
		Group-B	261	No statistically significant difference (vs. Ref.)	
		Group-C	418	Metabolic syndrome (vs. Ref.)↓, Abdominal obesity (vs. Ref.)↓	

		Group-D [high PA]	124	Metabolic syndrome (vs. Ref.)↓, Abdominal obesity (vs. Ref.)↓	
(Biswas et al., 2022)	Average counts per minute over 10-minute intervals throughout the 24-hour period.	Lowest activity	2,808	Ref.	Robust Poisson regression models
		Moderate activity	3,219	10-year risk for atherosclerotic CVD event (vs. Ref.)↓	
		Fluctuations of moderate activity	1,194	10-year risk for atherosclerotic CVD event (vs. Ref.)↓	
		High daytime activity	713	No statistically significant difference (vs. Ref.)	
		Moderate evening activity	225	10-year risk for atherosclerotic CVD event (vs. Ref.)↓	
		Highest activity	750	10-year risk for atherosclerotic CVD event (vs. Ref.)↓	
[2](Leskinen et al., 2023)	Hourly stepping minutes of two consecutive hours	Consistently inactive	1,301	BMI (vs. Ref.)↑, Waist (vs. Ref.)↑, 2-hour glucose (vs. Ref.)↑, Total/HDL ratio (vs. Ref.)↑, Triglycerides (vs. Ref.)↑	Analysis of covariance (ANCOVA)
		Consistently low active	2,498	BMI (vs. Ref.)↑, Waist (vs. Ref.)↑, 2-hour glucose (vs. Ref.)↑, Total/HDL ratio (vs. Ref.)↑, Triglycerides (vs. Ref.)↑	
		Active on weekdays	881	BMI (vs. Ref.)↑, Waist (vs. Ref.)↑	
		Early birds	138	BMI (vs. Ref.)↑, Waist (vs. Ref.)↑, Systolic BP (vs. Ref.)↑, Diastolic BP (vs. Ref.)↑	
		Consistently moderately active	449	No statistically significant difference (vs. Ref.)	
		Weekend warriors	453	BMI (vs. Ref.)↑, Waist (vs. Ref.)↑	
		Consistently highly active	352	Ref.	
(Aqeel et al., 2021)	Summation of hourly activity counts and time stamps	Lowest activity counts from 6:00–23:00	783	Ref.	Linear regression models
		High activity counts with a decline after 18:00	247	BMI (vs. Ref.)↓, WC (vs. Ref.)↓	
		Lower activity counts between 6:00–12:00	336	BMI (vs. Ref.)↓, WC (vs. Ref.)↓, Hemoglobin A1c (vs. Ref.)↓	
		Higher activity counts between 8:00–11:00	633	BMI (vs. Ref.)↓, WC (vs. Ref.)↓	
[3](Niemelä et al., 2019)	Average MET (metabolic equivalent) over 10-minute intervals throughout the 24-hour period	Inactive	1,881	10-year CVD risk (vs. Very active)↑, HDL cholesterol (vs. Very active)↑	Analysis of variance (ANOVA)
		Evening active	802	No statistically significant difference (vs. any of the profiles)	
		Moderately active	1,297	HDL cholesterol (vs. Very active)↓	
		Very active	602	10-year CVD risk (vs. Inactive)↓, HDL cholesterol (vs. Inactive)↓, HDL cholesterol (vs. Moderately active)↑	
(Marschollek, 2016)		Cluster 1: High overall physical with regular low-intensity	92	BMI (vs. Cluster 3)↓	Analysis of variance (ANOVA)

	PA intensity, mean duration of PA per 1000 min, and PA regularity	activity and more irregular high-intensity activity			
		Cluster 2: Regularly and extensively active	2,230	C-reactive protein (vs. Cluster 3 and Cluster 4)↓, BMI (vs. Cluster 3 and Cluster 4)↓	
		Cluster 3: Irregularly and less extensively active	2,990	C-reactive protein (vs. Cluster 4)↓, BMI (vs. Cluster 4)↓, C-reactive protein (vs. Cluster 2)↑, BMI (vs. Cluster 1 and Cluster 2)↑	
		Cluster 4: Inactive	2,010	C-reactive protein (vs. Cluster 2 and Cluster 3)↑, BMI (vs. Cluster 1, Cluster 2, and Cluster 3)↑	
(P. H. Lee et al., 2013)	Hourly average counts per minute over a 24-hour period on weekdays and weekends	Active	480	Ref.	Logistic regression models
		Less active	1,234	Having been diagnosed with a chronic condition (vs. Ref.)↑, Asthma (vs. Ref.)↑	
(Fukuoka et al., 2018)	Raw minute-level MET for 7 consecutive days	Morning engaged	65	Depression symptom (vs. Afternoon)↑; Depression symptom (vs. Unengaged)↓	Chi-square or analysis of variance (ANOVA)
		Afternoon engaged	48	Depression symptom (vs. Morning engaged)↓	
		Unengaged	102	Depression symptom (vs. Afternoon and Morning engaged)↑	
(Nawrin et al., 2024)	Hourly activity counts over a 24-hour period	Morning dominant	1,964	Ref.	Logistic regression models
		All-day dominant	1,792	No apparent differences in health outcomes	
		All-day or morning dominant	1,855	No apparent differences in health outcomes	
		Evening dominant	1,002	Depressive symptoms↑	
(Bai, Ning, et al., 2024)	Mean hourly acceleration over a 24-hour period across all measurement days	Persistent low	14,037	Ref.	Cox proportional hazards models
		Single peak	46,100	Incident breast cancer (vs. Ref.)↓, Incident kidney cancer (vs. Ref.)↓, Incident myeloma cancer (vs. Ref.)↓	
		Double peak	17,763	Incident colorectal cancer (vs. Ref.)↓, Incident kidney cancer (vs. Ref.)↓	
		Vigorous	18,787	Incident bladder cancer (vs. Ref.)↓, Incident breast cancer (vs. Ref.)↓, Incident kidney cancer (vs. Ref.)↓, Incident lung cancer (vs. Ref.)↓, Incident myeloma cancer (vs. Ref.)↓, Incident oral cavity and pharynx cancer (vs. Ref.)↓	

Only statistically significant associations observed in the studies are reported. Upward arrow (↑) indicates higher values/incidence in the health outcome relative to the referent group, and downward arrow (↓) lower values/incidence in the health outcome relative to the referent group. [1]: Five models were created based on different input variables into the profiling approach; associations reported here are for the profiles identified on the basis of average activity counts/minute. [2]: The analyses were stratified by type 2 diabetes status; significant associations reported here are those that were seen for no diabetes group [3]: Analysis were stratified by gender; significant differences reported here are those that were seen in both males and females. Abbreviations: Ref. = Referent profile; BMI = Body mass index; BP = Blood pressure; CVD = Cardiovascular diseases; HDL = High-density lipoprotein; LDL = Low-density lipoprotein; LPA= Light-intensity physical activity; MET = Metabolic equivalents; MIMS = Monitor Independent Movement Summary unit; MVPA = Moderate-to-vigorous physical activity; PA = Physical activity; WC = Waist circumference.

Profiles of sedentary behavior

The overview of profiles identified in four studies creating sedentary behavior profiles is shown in Figure 3. The number of profiles that emerged in these studies ranged from three to five. The input descriptor variables used for profile analysis, the derived sedentary profiles, and their associations with health outcomes in each study are listed in Table 3.

In one study (Evenson et al., 2017), the daily percentage of total sedentary time from total accelerometer wear time was used to create and identify sedentary behavior profiles. Two studies created sedentary behavior profiles using several variables describing how sedentary time was accumulated and interrupted (Farrahi et al., 2021; Hibbing et al., 2022), but the descriptor variables used in these studies differed. In the other study, sedentary behavior profiles were created using mean hourly sedentary time computed for up to 14 hours after waking (Reuter et al., 2020). This approach allowed the assessment of diurnal patterns of sedentary time.

Key profile characteristics and their relationships with health outcomes

Of the four studies creating sedentary profiles, two identified sedentary behavior profiles in distinct sample populations based on how sedentary time was accumulated and interrupted (Farrahi et al., 2021; Hibbing et al., 2022). Both studies identified profiles characterized by having more frequent interruptions in sedentary time compared with other emerging profiles. These profiles, labeled as "Interrupted" (Hibbing et al., 2022) and "Breakers" (Farrahi et al., 2021) are highlighted in green in Figure 3 (part Sedentary profiles). In contrast, both studies also identified profiles characterized by prolonged periods of uninterrupted sedentary time, labeled as "Prolonged" (Hibbing et al., 2022) and "Prolonged sitters" (Farrahi et al., 2021). These profiles are highlighted in orange in Figure 3 (part Sedentary profiles). A comparable threshold for defining prolonged sedentary bouts emerged in both studies (i.e., >20 min and

>15–30 min). One of these two studies identified two additional unique profiles: "Shortened

sitters," characterized by sedentary time accumulated in relatively short bouts of 15–30

minutes, and "Couch potatoes," distinguished by the highest overall levels of sedentary time

among all identified profiles (Farrahi et al., 2021).

As shown in Table 3, the "Prolonged" profile was associated with a higher 10-year CVD risk

compared with the "Interrupted" profile (Hibbing et al., 2022). In the other study (Farrahi et

al., 2021), no statistically significant differences in cardiometabolic health markers were

observed between the "Couch potatoes" and "Prolonged sitters." However, both the "Shortened

sitters" and "Breakers" demonstrated better cardiometabolic health compared with the "Couch

potatoes" (Farrahi et al., 2021). One study identified sedentary behavior profiles based on

sedentary time as a proportion of total daily wear time across all days of the week and found

no association between these profiles and mortality risk (Evenson et al., 2017).

Profiles of sleep

The overview of profiles identified in three studies creating sleep profiles is shown in Figure

3. The input descriptor variables used for profile analysis, the derived sleep profiles, and their

associations with health outcomes in each study are listed in Table 3. The number of profiles

that emerged in these three data-driven sleep profiling studies was two (Chung et al., 2024),

three (Zhang et al., 2024) and four (Matricciani et al., 2021). One of these studies utilized

hourly proportions of accumulated sleep minutes within each hour over a 24-hour period as

descriptor variables for profile analysis (Zhang et al., 2024). Although not identical, the other

two studies employed descriptor metrics related to sleep duration, timing, and variability

(Chung et al., 2024; Matricciani et al., 2021).

Key profile characteristics and their relationships with health outcomes

The three studies identifying sleep profiles were conducted with distinct sample populations

(Chung et al., 2024; Matricciani et al., 2021; Zhang et al., 2024). Among these, a profile

characterized by late bedtime was identified in two studies, as illustrated in green in Figure 3

(part Sleep profile). In one of these two studies, the profiles identified were "Irregular sleep",

"Morning lark", and "Night owl" (Zhang et al., 2024). "Irregular sleep" was characterized by

varying times of sleep onset and awakening throughout the day, "Morning lark" by sleeping

early at night and waking early in the morning, and "Night owl" by going to bed late and

waking up late (Zhang et al., 2024). "Irregular sleep" was associated with a higher mortality

risk compared with the "Morning lark." In the other study (Matricciani et al., 2021), the

identified profiles were "Overall good sleepers," "Late to bed," "Long sleepers", and "Short

sleepers." Among these profiles (Matricciani et al., 2021), the "Late to bed" profile was

associated with higher BMI and metabolic syndrome scores compared with the "Overall good

sleepers" profile (see Table 3).

Profiles of rest-activity rhythmicity

The overview of profiles identified in the two studies that created rest-activity rhythmicity

profiles is shown in Figure 3. The input descriptor variables used for profile analysis, the

derived rest-activity rhythmicity profiles, and their associations with health outcomes in each

study are listed in Table 3. Four profiles emerged in both studies that created and identified

profiles of rest-activity rhythmicity (Graves et al., 2021; Smagula et al., 2022). The rest-activity

rhythm descriptors used as input for profile analysis in both studies largely overlapped and

were derived from an extended cosine model (Graves et al., 2021; Smagula et al., 2022).

Key profile characteristics and their relationships with health outcomes

The two studies identified rest-activity profiles in distinct populations (Graves et al., 2021;

Smagula et al., 2022). "Early rising" profile, highlighted in green in Figure 3 (part Rest-activity

rhythmicity), was identified in both studies. In one of the studies (Smagula et al., 2022),

"Shorter active period/less modelable" and "Shorter active period/very weak," characterized

by starting their daily activities later and settling down earlier, with a relatively less rhythmic

24-hour rest-activity pattern, exhibited higher depression symptoms, compared with "Earlier

rising/robust," which was characterized by an estimated activity onset before 7 AM and a more

rhythmic 24-hour rest-activity pattern (Smagula et al., 2022). In the other study (Graves et al.,

2021), "Later rest activity rhythms (RARs)," characterized by having a later activity onset, had

higher perceived fatiguability compared with "Earlier risers," characterized by a relatively

earlier activity onset. The "Less activity/robust" profile, characterized by having relatively a

more rhythmic rest-activity pattern, exhibited higher perceived fatiguability compared with

"Earlier risers" (Graves et al., 2021).

Joint profiles of physical activity, sedentary behavior, and/or sleep

The overview of profiles identified in 11 studies creating joint profiles is shown in Figure 3.

The number of profiles that emerged in these studies ranged from three to five. The input

descriptor variables used for profile analysis, the derived joint profiles, and their associations

with health outcomes in each study are listed in Table 3. Although durations of physical activity

and sedentary behavior were commonly used as descriptor variables, there was overall

heterogeneity in the descriptors used as input for identifying and creating joint profiles (see

Table 3).

Of the 11 studies creating joint profiles, seven incorporated a combination of accelerometer-

derived PA and sedentary behavior descriptors for profile analyses (Choi et al., 2025; Farrahi

et al., 2022; Gupta et al., 2025; Palmberg et al., 2024; Paolillo et al., 2023; von Rosen et al.,

2020; Yerramalla et al., 2024) and three incorporated a combination of accelerometer-derived

PA, sedentary behavior, and sleep descriptors (Full et al., 2019; Gupta et al., 2020; O'Regan et

al., 2021). One study used transformed accelerometer outputs into what was named *activity*

images and used the activity images for profile analysis (Farrahi et al., 2024).

Key profile characteristics and their relationships with health outcomes

Due to the heterogeneity and variety of input descriptor variables in profile analysis, comparing

joint profiles across the included studies in terms of similarities and differences was more

challenging than comparing other types of profiles. Most commonly, profiles distinguished by

low levels of PA combined with high levels of sedentary time emerged among the studies

identifying joint profiles (n = 7) (Choi et al., 2025; Farrahi et al., 2024; Full et al., 2019;

O'Regan et al., 2021; Paolillo et al., 2023; von Rosen et al., 2020; Yerramalla et al., 2024).

These profiles are marked in orange in Figure 3 (part Joint profiles). Compared to profiles with

low levels of PA combined with high levels of sedentary time, relatively more active profiles

having lower levels of sedentary time and higher levels of PA, demonstrated reduced risks of

mortality (von Rosen et al., 2020; Yerramalla et al., 2024), lower amount of chronic illness and

multimorbidity (O'Regan et al., 2021), better cardiometabolic health and obesity indicators

(Farrahi et al., 2024; Full et al., 2019), and better cognitive health (Paolillo et al., 2023) and

physical capacity outcomes (Paolillo et al., 2023).

Profiles most distinguished from other profiles by having high sedentary time combined with

having sufficient PA were seen in two studies conducted with distinct sample populations

(Farrahi et al., 2022; Yerramalla et al., 2024). These profiles are marked in green in Figure 3

(part Joint profiles). As shown in Table 3, in one of these studies, the "Active sitters," most

distinguished by having sufficient MVPA but also high sedentary time, were shown to have a

higher mortality risk compared with the "Active" profile, characterized by having both lower

sedentary time and higher MVPA (Yerramalla et al., 2024). The other study identified an

"Active couch potato" profile, most distinguished by having sufficient MVPA coupled with

high sedentary time. Compared with this profile, "Sedentary light movers" who engaged in

sufficient PA but had relatively less sedentary time and more light-intensity PA were found to

have better cardiometabolic health (Farrahi et al., 2022).

Sedentary profiles (n = 4)

Study (first author, year)	Derived profiles (name or main characteristics)				
▲, 1 (Hibbing et al., 2022)	Interrupted (1,740)	Intermediate (2,453)	Prolonged (1,302)		
▲ (Farrahi et al., 2021)	Couch potatoes (1,222)	Prolonged sitters (1,222)	Shortened sitters (1,529)	Breakers (509)	
▲, 1, * (Evenson et al., 2017)	Class 1 [Most sedentary] (451)	Class 2 (1,286)	Class 3 (1,640)	Class 4 (872)	Class 5 [Least sedentary] (261)
▲ (Reuter et al., 2020)	High-SB-throughout-the-day (2,239)	Moderate-SB-with-lower-morning-SB (1,536)	Moderate-SB-with-higher-morning-SB (1,137)	Low-SB-throughout-the-day (1,292)	

Comparable profiles:
- 🟧 Relatively longer sedentary bouts
- 🟩 Frequent sedentary interruptions

Sleep profiles (n = 3)

Study (first author, year)	Derived profiles (name or main characteristics)			
▲, 3 (Zhang et al., 2024)	Irregular sleep (2,159)	Morning lark (1,677)	Night owl (2,837)	
• (Chung et al., 2024)	Irregular-insufficient (744)	Regular-optimal (1,015)		
• (Matricciani et al., 2021)	Short sleepers (347)	Late to bed (269)	Long sleepers (297)	Overall good sleepers (424)

Comparable profiles:
- 🟩 Late sleep

Rest-activity profiles (n = 2)

Study (first author, year)	Derived profile (name or main characteristics)			
• (Smagula et al., 2022)	Earlier rising/robust (677)	Shorter active period/less modellable (587)	Shorter active period/very weak (177)	Later settling/very weak (359)
▲ (Graves et al., 2021)	Earlier risers (70)	More active/Robust (37)	Later RARs (63)	Less active/Robust (14)

Comparable profiles:
- 🟩 Early rising

Joint profiles (n = 11)

Study (first author, year)	Derived profiles (name or main characteristics)				
▲ (Yerramalla et al., 2024)	Active (726)	Active sitters (890)	Light movers (1,033)	Prolonged sitters (1,040)	Most sedentary (302)
▲ (von Rosen et al., 2020)	Low active (147)	Average active (397)	High active (307)		
▲ (Choi et al., 2025)	High lifestyle physical activity (64)	High MVPA (37)	Low overall physical activity (108)	Average physical activity (219)	
▲, 1 (Gupta et al., 2025)	Chimpanzees (226)	Lions (179)	Koalas (158)	Ants (244)	
▲ (Farrahi et al., 2024)	Lowest activity (351)	Early morning movers (595)	Least sedentary movers (552)	Highest activity (314)	
▲ (Farrahi et al., 2022)	Active couch potatoes (1,173)	Sedentary light movers (1,199)	Sedentary exercisers (694)	Movers (636)	
▲, 1 (Gupta et al., 2020)	Chimpanzees (226)	Lions (179)	Koalas (158)	Ants (244)	
▲ (O'Regan et al., 2021)	Inactive-sedentary (50)	Low activity (295)	Active (111)	Very active (29)	
▲ (Palmberg et al., 2024)	Exercisers (79)	Actives (177)	Inactives (185)		
• (Full et al., 2019)	Moderately active poor sleepers (27)	Highly actives (151)	Inactives (34)	Moderately actives (105)	Actives (55)
• (Paolillo et al., 2023)	Low physical activity (49)	Average physical activity (59)	High intensity physical activity (16)		

Comparable profiles:
- 🟧 High sedentary/Low physical activity
- 🟩 Active and sedentary

Figure 3: Overview of derived profiles in studied identifying sedentary (n = 4), sleep (n = 2), rest-activity (n = 3), and joint (n = 11) profiles. The numbers shown in () are the number of participants in the profile. The profiles are specified according to the names given to them in the study or a short outline of the main characteristics. To facilitate comparison across studies, profiles with comparable overall characteristics *seen in at least two or more distinct sample populations* are highlighted with distinct colors. Note that the color-coded profiles are not necessarily exactly identical, because of the underlying heterogeneities related to profile analysis and/or accelerometer data cleaning/processing. 1: studies marked with the same number were performed with the same sample population. ▲: High quality study. •: Moderate quality study. Individual publication quality scores are shown in Supplementary Table S2. *: Two models were created based on different input variables into the profiling approach; profiles reported here are for the profiles identified on the basis of percentage of sedentary behavior out of total wear time per day. Abbreviations: SB = sedentary behavior; MVPA = Moderate-to-vigorous physical activity.

Table 3: Input descriptor variables, profiles identified, and the statistically significant associations observed among the profiles in studies identifying sedentary behavior, sleep, rest-activity, and joint profiles.

Study (first author, year)	Input descriptor variables	Derived profiles, name or main characteristics	Number of participants	Observed associations	Statistical approach used to examine associations
Sedentary profiles (n = 4)					
(Hibbing et al., 2022)	10th-90th percentiles, interquartiled and interdecile range of sedentary bouts, number of sedentary bouts, % of wear time in sedentary	Interrupted	1,740	Ref.	Logistic regression models
		Intermediate	2,453	10-year CVD risk (vs. Ref.)↑	
		Prolonged	1,302	10-year CVD risk (vs. Ref.)↑	
(Farrahi et al., 2021)	65 variables characterizing how sedentary time was accumulated and interrupted.	Couch potatoes	1,222	Ref.	Linear regression models
		Prolonged sitters	1,222	No statistically significant differences (vs. Ref.)	
		Shortened sitters	1,529	2-hour insulin (vs. Ref.)↓, Fasting serum insulin (vs. Ref.)↓, Triglycerides (vs. Ref.)↓, Total/HDL cholesterol ratio (vs. Ref.)↓, LDL/HDL cholesterol ratio (vs. Ref.)↓, Body fat (vs. Ref.)↓, Fat mass (vs. Ref.)↓, Visceral fat area (vs. Ref.)↓, BMI (vs. Ref.)↓, WC (vs. Ref.)↓	
		Breakers	509	Total/HDL cholesterol ratio (vs. Ref.)↓, LDL/HDL cholesterol ratio (vs. Ref.)↓	
[1](Evenson et al., 2017)	Percentage of sedentary behavior out of total wear time per day on every day of the week	Class 1 - Most sedentary	451	No statistically significant difference (vs. Ref.)	Cox proportional hazards models
		Class 2	1,286	No statistically significant difference (vs. Ref.)	
		Class 3	1,640	No statistically significant difference (vs. Ref.)	
		Class 4	872	No statistically significant difference (vs. Ref.)	
		Class 5 – Least sedentary	261	Ref.	
[2](Reuter et al., 2020)	Mean hourly sedentary time up to 14 hours after waking time	High-SB-throughout-the-day	2,239	Rate of physical functioning decline (vs. Moderate-SB-with-lower-morning-SB)↑	Multivariable linear mixed-effects regression
		Moderate-SB-with-lower-morning-SB	1,536	Rate of physical functioning decline (vs. High-SB-throughout-the-day and Low-SB-throughout-the-day)↓	
		Moderate-SB-with-higher-morning-SB	1,137	No statistically significant association	
		Low-SB-throughout-the-day	1,292	Rate of physical functioning decline (vs. Moderate-SB-with-lower-morning-SB)↑	
Sleep profiles (n = 3)					
(Zhang et al., 2024)	Hourly proportions of accumulated sleep minutes accumulated in each hour over a 24-hour period	Irregular sleep	2,159	All-cause mortality risk (vs. Ref.)↑	Cox proportional hazard models
		Morning lark	1,677	Ref.	
		Night owl	2,837	No statistically significant difference (vs. Ref.)	
(Chung et al., 2024)	Total sleep duration, night-to-night timing irregularity, and night-to-night duration irregularity	Irregular-insufficient	744	Ref.	Cox proportional hazard models
		Regular-optimal	1,015	All-cause mortality (vs. Ref.)↓	

(Matricciani et al., 2021)	Sleep period, sleep timing, day-to-day sleep variability, sleep efficiency	Short sleepers	347	No statistically significant difference (vs. Ref.)	Generalized estimating equation
		Late to bed	269	Metabolic syndrome severity score (vs. Ref.)↑, BMI (vs. Ref.)↑	
		Long sleepers	297	No statistically significant differences (vs. Ref.)	
		Overall good sleepers	424	Ref.	

Rest-activity profiles (n = 2)

(Smagula et al., 2022)	Activity onset time, activity offset time, the pseudo–F statistic, interdaily stability, relative amplitude, and intradaily variability	Earlier rising/robust	677	Ref.	Logistic regression models
		Shorter active period/less modellable	587	Depressed mood (vs. Ref.)↑, Verbal fluency (vs. Ref.)↓, Delayed memory (vs. Ref.)↓	
		Shorter active period/very weak	177	Tired/Little energy (vs. Ref.)↑, Guilt (vs. Ref.)↑, Suicidality (vs. Ref.)↑, Attention and processing speed (vs. Ref.)↓, Verbal fluency (vs. Ref.)↓, Delayed memory (vs. Ref.)↓	
		Later settling/very weak	359	Sleep disturbance (vs. Ref.)↑, Tired/Little energy (vs. Ref.)↑, Suicidality (vs. Ref.)↑, Attention and processing speed (vs. Ref.)↓, Verbal fluency (vs. Ref.)↓, Delayed memory (vs. Ref.)↓	
(Graves et al., 2021)	Width of active period, amplitude, acrophase, beta (steepness), mesor, estimated rise time, and the pseudo-F statistic.	Earlier risers	70	Ref.	Multivariate quantile regression
		More active/Robust	37	No statistically significant difference (vs. Ref.)	
		Later RARs	63	Perceived physical fatigability (vs. Ref.)↑	
		Less active/Robust	14	Perceived physical fatigability (vs. Ref.)↑	

Joint profiles (n = 11)

(Yerramalla et al., 2024)	13 variables, representing overall activity level, total duration and frequency, typical duration, fragmentation, activity intensity distribution and timing of activity behaviors.	Active	726	Ref.	Cox proportional hazard model
		Active sitters	890	All-cause mortality (vs. Ref.)↑	
		Light movers	1,033	All-cause mortality (vs. Ref.)↑	
		Prolonged sitters	1,040	All-cause mortality (vs. Ref.)↑	
		Most sedentary	302	All-cause mortality (vs. Ref.)↑	
(von Rosen et al., 2020)	14 measures representing duration of PA intensities and SB, variation of PA intensities and SB, total time of MVPA and sedentary bouts, number of sedentary and MVPA bouts, total counts, and the relative time spent in one behavior relative to the two remaining behaviors	Low active	147	Ref.	Cox proportional hazard model
		Average active	397	All-cause mortality (vs. Ref.)↓	
		High active	307	All-cause mortality (vs. Ref.)↓	
(Choi et al., 2025)	Minutes per week of sedentary behaviors, light PA, lifestyle PA, moderate PA, MVPA, and vigorous PA	High lifestyle physical activity	61	Subjective health (vs. Ref.)↑	Multinomial logistic regression
		High MVPA	37	Metabolic syndrome (vs. Ref.)↓	
		Low overall physical activity	108	No statistically significant difference (vs. Ref.)	
		Average physical activity	219	Ref.	
(Gupta et al., 2025)	Isometric log ratio of time spent in sedentary, standing, LPA, and MVPA in work and leisure and time in bed	Chimpanzees	226	Ref.	Linear regression and logistic regression models
		Lions	179	Systolic BP (vs. Ref.)↑, low back pain (vs. Ref.)↑	
		Koalas	158	Systolic BP (vs. Ref.)↓, cardiorespiratory fitness (vs. Ref.)↑	
		Ants	244	Self-rated health (vs. Ref.)↓, cardiorespiratory fitness (vs. Ref.)↑	

(Farrahi et al., 2024)	Valid minute-by-minute activity counts over 7 measurement days were shown as *'activity images'*	Lowest activity	351	Ref.	Linear regression models
		Early morning movers	595	Insulin (vs. Ref.)↓, Triglycerides (vs. Ref.)↓, Homeostasis insulin resistance (vs. Ref.)↓, Plasma glucose (vs. Ref.)↓	
		Least sedentary movers	552	WC (vs. Ref.)↓	
		Highest activity	314	Insulin (vs. Ref.)↓, Triglycerides (vs. Ref.)↓, Homeostasis insulin resistance (vs. Ref.)↓, Plasma glucose (vs. Ref.)↓, C-reactive protein (vs. Ref.)↓, Total/HDL cholesterol ratio (vs. Ref.)↓	
(Farrahi et al., 2022)	Time spent in sedentary, LPA, and MVPA on each weekday and weekend day	Active couch potatoes	1,173	Ref.	Linear regression models
		Sedentary light movers	1,199	2-hour insulin (vs. Ref.)↓, Fasting serum insulin (vs. Ref.)↓, Triglycerides (vs. Ref.)↓, Body fat (vs. Ref.)↓, Fat mass (vs. Ref.)↓, Visceral fat area (vs. Ref.)↓, BMI (vs. Ref.)↓, WC (vs. Ref.)↓	
		Sedentary exercisers	694	2-hour insulin (vs. Ref.)↓, Fasting serum insulin (vs. Ref.)↓, Triglycerides (vs. Ref.)↓, Total/HDL cholesterol (vs. Ref.)↓, LDL/HDL cholesterol (vs. Ref.)↓, 2-hour glucose (vs. Ref.)↓, Body fat (vs. Ref.)↓, Fat mass (vs. Ref.)↓, Visceral fat area (vs. Ref.)↓, BMI (vs. Ref.)↓, WC (vs. Ref.)↓	
		Movers	636	2-hour insulin (vs. Ref.)↓, Fasting serum insulin (vs. Ref.)↓, Triglycerides (vs. Ref.)↓, Total/HDL cholesterol (vs. Ref.)↓, LDL/HDL cholesterol (vs. Ref.)↓, Body fat (vs. Ref.)↓, Fat mass (vs. Ref.)↓, Visceral fat area (vs. Ref.)↓, WC (vs. Ref.)↓	
(Gupta et al., 2020)	Isometric log ratio of time spent in sedentary, standing, LPA, and MVPA in work and leisure and time in bed	Chimpanzees	226	Ref.	Linear regression models
		Lions	179	BMI (vs. Ref.)↑, WC (vs. Ref.)↑, Body fat percentage (vs. Ref.)↑	
		Koalas	158	WC (vs. Ref.)↑, Body fat percentage (vs. Ref.)↑	
		Ants	244	No statistically significant difference (vs. Ref.)	
(O'Regan et al., 2021)	Average daily hours in bed, average daily waking sedentary hours, average daily standing hours, average daily LPA hours, average daily MVPA minutes and average daily step count	Inactive-sedentary	50	Having mental illness (vs. Ref.)↑, Being obese (vs. Ref.)↑, Having multi-morbidity (vs. Ref.)↑	Logistic regression models
		Low activity	295	Having mental illness (vs. Ref.)↑, Being obese (vs. Ref.)↑, Having multi-morbidity (vs. Ref.)↑	
		Active	111	Having mental illness (vs. Ref.)↑, Being obese (vs. Ref.)↑, Having multi-morbidity (vs. Ref.)↑	
		Very active	29	Ref.	
[3](Palmberg et al., 2024)	13 features describing duration of movement intensities, gait characteristics, sit-to-stand transition, and activity fragmentation	Exercisers	79	6MWT walking speed (vs. Inactives)↑	General linear models
		Actives	177	10-m walking speed (vs. Inactives)↑, 6MWT walking speed (vs. Inactives)↑	
		Inactives	185	10-m walking speed (vs. Actives)↓, 6MWT walking speed (vs. Exercisers and Actives)↓	
(Full et al., 2019)	Nightly sleep time, nightly sleep efficiency, MVPA, and percent time spent sedentary	Moderately active poor sleepers	27	Physical functioning (vs. Moderately Actives)↓	Analysis of variance (ANOVA)
		Highly actives	151	No statistically significant difference (vs. any of the other profiles)	
		Inactives	34	BMI (vs. Moderately actives)↑	
		Moderately actives	105	BMI (vs. Inactives)↓, Physical functioning (vs. Inactives)↑	
		Actives	55	No statistically significant difference (vs. any of the other profiles)	
(Paolillo et al., 2023)	Average daily step count, average daily sedentary time, and average daily high intensity time	Low physical activity	49	Ref.	Linear regression models
		Average physical activity	59	No statistically significant difference (vs. Ref.)	

	High intensity physical activity	16	Executive functioning (vs. Ref.)↑, vascular burden (vs. Ref.)↓

Only statistically significant associations observed in the studies are reported. Upward arrow (↑) indicates higher values/incidence in the health outcome relative to the referent group, and downward arrow (↓) lower values/incidence in the health outcome relative to the referent group. [1]: Two models were created based on different input variables into the profiling approach; associations reported here are for the profiles identified on the basis of percentage of sedentary behavior out of total wear time per day. [2]: Analysis were stratified by low MVPA and high MVPA groups; the associations reported here are for the low MVPA group. [3]: Analysis were stratified by gender; significant differences reported here are those that were seen in both males and females. Abbreviations: Ref. = Referent profile; 6MWT = 6-minute walking test; BMI = Body mass index; BP = Blood pressure; CVD = Cardiovascular diseases; HDL = High-density lipoprotein; LDL = Low-density lipoprotein; LPA= Light-intensity physical activity; MET = Metabolic equivalents; MVPA = Moderate-to-vigorous physical activity; PA = Physical activity; RARs = Rest-activity rhythm; SB= sedentary behavior. WC = Waist circumference.

DISCUSSION

The present study systematically identified and reviewed 40 studies that utilized data-driven population segmentation analysis to drive multidimensional physical behavior profiles from wearable accelerometry data in a sample of adult participants. Machine learning K-means clustering algorithm and latent profile analysis were the most commonly employed methods for profiling of physical behavior from accelerometer data. The review of derived profiles from accelerometer-monitored physical behaviors revealed several hypothesis-generating, preliminary evidence about adult physical behavior profiles. Multiple studies have identified data-driven profiles distinguished by the timing of physical activity across the 24-hour day. Some profiles suggest that adults can be physically active while still spending substantial time in sedentary behaviors. In addition to total sedentary time, the patterns of accumulation and interruption of sedentary behavior may also be associated with health risks and outcomes.

Data-driven population segmentation analysis are widely used in a number of different research fields (Woo et al., 2024). Compared with variable-centered approaches (Migueles et al., 2022), one of the main advantage of algorithms used for cluster analysis and population segmentations is that they can naturally accommodate a large number of input variables (Biswas et al., 2022; Farrahi et al., 2021; Kebede et al., 2024; Liu et al., 2023; Niemelä et al., 2019; Weller et al., 2020). In the domains related to health and medicine, data-driven population segmentation analysis have been predominantly employed for examining how different risk factors in multidimensional data may cluster together (Hafkamp et al., 9900; Kang et al., 2014; Liu et al., 2023; Woo et al., 2024). Consistent with previous research in other domains (Hafkamp et al., 9900; Kang et al., 2014; Liu et al., 2023; Woo et al., 2024), our study highlights the utility of data-driven profiling of accelerometer-measured physical behaviors for addressing questions about how different components and dimensions of physical behaviors cluster together, and analyzing whether and how the relative similarities and dissimilarities.

In general, studies used a wide variety of accelerometer-derived descriptors to characterize

physical behavior profiles, making direct comparison of profiles difficult. Still, the profiles

identified through data-driven methods generally aligned with the well-established evidence

observed in studies employing variable-centered approaches. For example, profiles

distinguished by relatively a low PA combined with high sedentary time frequently emerged in

studies examining how different components of physical behaviors cluster together (Choi et al.,

2025; Farrahi et al., 2024; Full et al., 2019; O'Regan et al., 2021; Paolillo et al., 2023; von

Rosen et al., 2020; Yerramalla et al., 2024). Consistent with previous findings (Ekelund et al.,

2016; Stamatakis et al., 2019), profiles most distinguished by a combination of low PA and high

sedentary time were consistently found to be linked with increased risks of CVD and all-cause

mortality and adversely to several other risk indicators (Choi et al., 2025; Farrahi et al., 2024;

Full et al., 2019; O'Regan et al., 2021; Paolillo et al., 2023; von Rosen et al., 2020; Yerramalla

et al., 2024).

Joint profiles identified in data-driven studies suggest that a segment of the adult population

may exhibit a behavioral pattern characterized by being physically active while simultaneously

engaging in significant amounts of sedentary behavior. Current robust evidence indicates that

although prolonged sedentary behavior is associated with increased risks of various chronic

conditions and mortality (Biswas et al., 2015), high levels of PA can mitigate some of the health

risks linked to excessive sedentary time (Ekelund et al., 2016). In line with our findings, some

population-based studies have reported a high prevalence of people meeting PA

recommendations while also accumulating high average sedentary time (Collings et al., 2022;

Moreno-Llamas et al., 2022). Although further research is needed to better understand how

socioeconomic factors may be related to this physical behavior profile, *active yet sedentary*

profile appears to be common among adults in high-income and European countries (Collings

et al., 2022; Moreno-Llamas et al., 2022). Currently, it remains unclear whether the level of PA

among these individuals is sufficient to offset the health risks posed by their sedentary behavior.

The findings of this review highlight that *active yet sedentary* profile may be present among

adults, underscoring the need for further research to better understand its prevalence and health

implications in adults.

Several studies identifying PA profiles found a profile in which PA peaked at sometime during

morning, day, afternoon, or evening (Albalak et al., 2023; Aqeel et al., 2021; Bai, Ning, et al.,

2024; Bai, Shao, et al., 2024; Bai, Zhou, et al., 2024; Biswas et al., 2022, 2023; Fabrizio et al.,

2024; Q. Feng et al., 2025; Fukuoka et al., 2018; Leskinen et al., 2023; Li et al., 2024; Ma et

al., 2023; Nawrin et al., 2024; Niemelä et al., 2019; Shim et al., 2023; Zhang et al., 2024). The

health implications of these profiles were mixed in terms of whether PA during a certain time

of the 24-hour period is more beneficial than at other times. Currently, whether PA at one time

of day provides greater health benefits than at another is still not fully understood. Although the

optimal time window for PA may be outcome-specific (H. Feng et al., 2023; Keiser et al., 2024),

findings from data-driven studies overall support the hypothesis that the timing of PA may

influence its health benefits.

Low PA profiles most distinguished by having the lowest levels of PA, were repeatedly

identified across studies performing physical activity profiling. These profiles emerged across

several sample populations from various countries and often represented one of the largest

participant groups among the identified profiles. The health risks associated with insufficient

PA are well-documented and extensively studied previously (I.-M. Lee et al., 2012). Current

estimates based on self-reported PA indicate that nearly one-third of adults worldwide do not

meet the recommended levels of PA (Strain et al., 2024). In line with existing literature (I.-M.

Lee et al., 2012), data-driven PA profiles derived from multidimensional accelerometer data

consistently indicate that low PA is associated with increased health risks.

Two studies created sedentary profiles based on how sedentary time is accumulated and

interrupted, both identifying a profile distinguished by spending sedentary time in relatively

longer bouts lasting 15–30 min (Farrahi et al., 2021; Hibbing et al., 2022). Currently, it is well-

established that excessive sedentary time, particularly when accumulated in uninterrupted

bouts, poses a health risk (Biswas et al., 2015). However, it remains unclear at what point

sedentary behavior begins to negatively impact health outcomes (Chastin, Egerton, et al., 2015).

Sedentary profiles derived using data-driven techniques provide preliminary, hypothesis-

generating evidence suggesting that limiting sedentary time to bouts of approximately 15–30

min may be linked to better cardiometabolic health (Farrahi et al., 2021; Hibbing et al., 2022).

This data-driven hypothesis warrants further research to explore whether similar profiles can

be observed across different populations and how limiting sedentary bouts to specific durations

is linked to other health outcomes.

Only a few studies have created and identified rest-activity rhythmicity or sleep profiles. Due

to the limited number of studies and the heterogeneity observed among them, synthesizing

evidence regarding the profiles derived from these studies was challenging. In general, there

has been a tendency to apply data-driven population segmentation analysis primarily to

accelerometer-determined physical behaviors during waking hours. This may be partly because

earlier wearable accelerometers were traditionally used to monitor physical behaviors only

during waking hours, with relatively recent studies adopting a 24-hour wear protocol (Evenson

et al., 2022; Giurgiu et al., 2025). Although evidence remains inconclusive, early findings from

data-driven profiles of sleep suggest that patterns of bedtime and wake-up time may be linked

to mortality risk and cardiometabolic health (Matricciani et al., 2021; Zhang et al., 2024).

Studies identifying rest-activity profiles also provided some support for this hypothesis

indicating that earlier activity onset, compared with later activity onset, may be favorably linked

with depressive symptoms (Smagula et al., 2022) and perceived fatiguability (Graves et al.,

2021). However, further research is warranted to clarify the potential bidirectional relationships

between rest-activity and sleep profiles and health outcomes.

This review study has several notable strengths. An extensive literature search was conducted

and both prospective cohort and cross-sectional studies were included. While a previously

conducted systematic scoping review has described the application of latent profile analysis to

accelerometry data based on a total of 12 included studies (Brusco et al., 2017), our study is

significantly more inclusive. It is the first to systematically review studies that identify physical

behavior profiles from wearable accelerometry data in adult participants using data-driven

population segmentation analysis. Multiple data-driven hypotheses were identified from the

review of physical behavior profiles and summarized their relationships with various health

outcomes.

This study is not without limitations. In our review, title and abstract screening was conducted

by a single reviewer, which may have increased the risk of missed eligible studies. However,

single-reviewer screening is often used to reduce the substantial resources required for

systematic reviews and has been suggested to be sufficient when performed by a senior expert

(Waffenschmidt et al., 2019). In general, profiles derived from accelerometer-measured

physical behaviors in different sample populations seem to be dependent on the descriptor

variables used as input for profile analysis. The decisions made for cleaning and preparing

accelerometer data, as well as measurement specification and accelerometer wear location,

could potentially have influenced the final derived profiles within the included studies

(Migueles et al., 2017). Heterogeneity in accelerometer measurement and data analysis across

existing studies precluded assessment of the influence of data processing decisions and

measurement specifications on the final derived profiles. Currently, it is still unclear whether

and how physical behavior profiles identified in one sample population could differ from those

identified in another population independent of differences attributed to accelerometer data and

data processing decisions. It is also unclear how differences in summary metrics and variables

derived from accelerometer data, such as MIMS and Euclidean Norm Minus One (ENMO)

(Migueles et al., 2019), could have influenced the final profiles identified. The profiling

algorithms used for profile analysis (e.g., K-means, latent profile analysis, etc.) may influence

the profiles ultimately identified (Brusco et al., 2017). However, it remains unclear to what

extent the reported physical behavior profiles depend on the specific algorithm employed.

Further research is needed to better understand the advantages and limitations of different

algorithms for physical behavior profiling.

CONCLUSIONS

Data-driven population segmentation analysis is well suited for exploring the complex and

potentially unknown interplay among multiple components and aspects of physical behaviors,

particularly for investigating how these behaviors could cluster together and synergistically

influence health outcomes and indicators. The use of data-driven techniques on wearable

accelerometer-measured physical behaviors has generated several hypothesis-generating, data-

driven findings that warrant further investigation. Multiple studies have identified data-driven

profiles that could be distinguished by the temporal distribution of PA at specific times during

a 24-hour period. Joint profiles emerging in data-driven studies indicate that being physically

active while simultaneously spending considerable amounts of time in sedentary behaviors may

represent a profile present among the adult population. Beyond total sedentary time, patterns of

accumulation and interruption of sedentary time may also be linked with health risks and

indicators. Although few studies have identified sleep and rest–activity profiles, data-driven

findings suggest that bedtime and wake-up patterns and the timing of activity onset may be

related to health outcomes.

REFERENCES

Aadland, E., Kvalheim, O. M., Anderssen, S. A., Resaland, G. K., & Andersen, L. B. (2019). Multicollinear physical activity accelerometry data and associations to cardiometabolic health: challenges, pitfalls, and potential solutions. *International Journal of Behavioral Nutrition and Physical Activity*, *16*(1), 74. https://doi.org/10.1186/s12966-019-0836-z

Albalak, G., Stijntjes, M., van Bodegom, D., … Noordam, R. (2023). Setting your clock: associations between timing of objective physical activity and cardiovascular disease risk in the general population. *European Journal of Preventive Cardiology*, *30*(3), 232–240.

Aqeel, M., Guo, J., Lin, L., … Eicher-Miller, H. A. (2021). Temporal physical activity patterns are associated with obesity in U.S. adults. *Preventive Medicine*, *148*, 106538. https://doi.org/https://doi.org/10.1016/j.ypmed.2021.106538

Backes, A., Gupta, T., Schmitz, S., Fagherazzi, G., van Hees, V., & Malisoux, L. (2022). Advanced analytical methods to assess physical activity behavior using accelerometer time series: A scoping review. *Scandinavian Journal of Medicine & Science in Sports*, *32*(1), 18–44. https://doi.org/https://doi.org/10.1111/sms.14085

Bai, P., Ning, X., Gao, R., … Yu, P. (2024). Association between circadian physical activity patterns and cancer incidence through regulation of inflammation: A UK biobank study. *Preventive Medicine*, *179*, 107831. https://doi.org/https://doi.org/10.1016/j.ypmed.2023.107831

Bai, P., Shao, X., Chen, L., … Yu, P. (2024). Association between circadian physical activity trajectories and incident type 2 diabetes in the UK Biobank. *Scientific Reports*, *14*(1), 6459. https://doi.org/10.1038/s41598-024-57082-2

Bai, P., Zhou, S., Shao, X., Lin, Y., Liu, H., & Yu, P. (2024). Ideal 24-h physical activity trajectory reduces all-cause, cause-specific mortality and cardiovascular outcomes through aging deceleration and inflammation regulation: A UK biobank study. *International Journal of Cardiology*, *399*, 131770.

https://doi.org/https://doi.org/10.1016/j.ijcard.2024.131770

Biswas, A., Chen, C., Prince, S. A., Smith, P. M., & Mustard, C. A. (2022). Workers' Activity

Profiles Associated With Predicted 10-Year Cardiovascular Disease Risk. *Journal of the*

American Heart Association, *11*(14), e025148.

Biswas, A., Chen, C., Prince, S. A., Smith, P. M., & Mustard, C. A. (2023). Daily

accelerometer-measured physical activity patterns and associations with cardiometabolic

health among Canadian working adults. *Health Reports*, *34*(3), 15–29.

https://doi.org/10.25318/82-003-x202300300002-eng

Biswas, A., Oh, P. I., Faulkner, G. E., … Alter, D. A. (2015). Sedentary time and its

association with risk for disease incidence, mortality, and hospitalization in adults: a

systematic review and meta-analysis. *Annals of Internal Medicine*, *162*(2), 123–132.

Blodgett, J. M., Ahmadi, M. N., Atkin, A. J., … Hamer, M. (2024). Device-measured physical

activity and cardiometabolic health: the Prospective Physical Activity, Sitting, and Sleep

(ProPASS) consortium. *European Heart Journal*, *45*(6), 458–471.

https://doi.org/10.1093/eurheartj/ehad717

Brusco, M. J., Shireman, E., & Steinley, D. (2017). A comparison of latent class, K-means,

and K-median methods for clustering dichotomous data. *Psychological Methods*, *22*(3),

563–580. https://doi.org/10.1037/met0000095

Chastin, S., Egerton, T., Leask, C., & Stamatakis, E. (2015). Meta-analysis of the relationship

between breaks in sedentary behavior and cardiometabolic health. *Obesity*, *23*(9), 1800–

1810. https://doi.org/10.1002/oby.21180

Chastin, S., Palarea-Albaladejo, J., Dontje, M. L., & Skelton, D. A. (2015). Combined effects

of time spent in physical activity, sedentary behaviors and sleep on obesity and cardio-

metabolic health markers: a novel compositional data analysis approach. *PloS One*,

10(10), e0139984.

Choi, J., Park, J., Choi, J.-Y., Lee, T., Jeon, G., & Cho, S. (2025). The Patterns of Physical

716  Activity Using Latent Profile Analysis Applied to Accelerometry and Associated

717  Sociodemographic and Health Factors Among Older Adults in Korea. *Journal of Aging*

718  *and Physical Activity*, 1–11. https://doi.org/10.1123/japa.2024-0296

719 Chung, J., Goodman, M. O., Huang, T., … Redline, S. (2024). Objectively regular sleep

720  patterns and mortality in a prospective cohort: The Multi-Ethnic Study of

721  Atherosclerosis. *Journal of Sleep Research*, *33*(1), e14048.

722  https://doi.org/https://doi.org/10.1111/jsr.14048

723 Collings, P. J., Backes, A., Aguayo, G. A., … group, on behalf of the O.-L. study. (2022).

724  Device-measured physical activity and sedentary time in a national sample of

725  Luxembourg residents: the ORISCAV-LUX 2 study. *International Journal of Behavioral*

726  *Nutrition and Physical Activity*, *19*(1), 161. https://doi.org/10.1186/s12966-022-01380-3

727 Dumuid, D., Pedišić, Ž., Stanford, T. E., … Olds, T. (2019). The compositional isotemporal

728  substitution model: a method for estimating changes in a health outcome for reallocation

729  of time between sleep, physical activity and sedentary behaviour. *Statistical Methods in*

730  *Medical Research*, *28*(3), 846–857.

731  https://doi.org/https://doi.org/10.1177/0962280217737805

732 Ekelund, U., Steene-Johannessen, J., Brown, W. J., … Lee, I.-M. (2016). Does physical

733  activity attenuate, or even eliminate, the detrimental association of sitting time with

734  mortality? a harmonised meta-analysis of data from more than 1 million men and

735  women. *The Lancet*, *388*(10051), 1302–1310. https://doi.org/10.1016/S0140-

736  6736(16)30370-1

737 Evenson, K. R., Herring, A. H., & Wen, F. (2017). Accelerometry-Assessed Latent Class

738  Patterns of Physical Activity and Sedentary Behavior With Mortality. *American Journal*

739  *of Preventive Medicine*, *52*(2), 135–143. https://doi.org/10.1016/j.amepre.2016.10.033

740 Evenson, K. R., Scherer, E., Peter, K. M., Cuthbertson, C. C., & Eckman, S. (2022). Historical

741  development of accelerometry measures and methods for physical activity and sedentary

behavior research worldwide: A scoping review of observational studies of adults. *PloS One*, *17*(11), e0276890. https://doi.org/10.1371/journal.pone.0276890

Ezugwu, A. E., Ikotun, A. M., Oyelade, O. O., … Akinyelu, A. A. (2022). A comprehensive survey of clustering algorithms: State-of-the-art machine learning applications, taxonomy, challenges, and future research prospects. *Engineering Applications of Artificial Intelligence*, *110*, 104743. https://doi.org/https://doi.org/10.1016/j.engappai.2022.104743

Fabrizio, C., Julien, V., Pedro, M.-V., & Vanessa, K. (2024). Associations Between Physical Activity Patterns and Cardiovascular Events and Risk Factors. *JACC: Advances*, *3*(11), 101324. https://doi.org/10.1016/j.jacadv.2024.101324

Farrahi, V., Collings, P. J., & Oussalah, M. (2024). Deep learning of movement behavior profiles and their association with markers of cardiometabolic health. *BMC Medical Informatics and Decision Making*, *24*(1), 74. https://doi.org/10.1186/s12911-024-02474-7

Farrahi, V., Kangas, M., Kiviniemi, A., Puukka, K., Korpelainen, R., & Jämsä, T. (2021). Accumulation patterns of sedentary time and breaks and their association with cardiometabolic health markers in adults. *Scandinavian Journal of Medicine & Science in Sports*, *31*(7), 1489–1507. https://doi.org/https://doi.org/10.1111/sms.13958

Farrahi, V., & Rostami, M. (2024). Machine learning in physical activity, sedentary, and sleep behavior research. *Journal of Activity, Sedentary and Sleep Behaviors*, *3*(1), 5. https://doi.org/10.1186/s44167-024-00045-9

Farrahi, V., Rostami, M., Dumuid, D., … Oussalah, M. (2022). Joint Profiles of Sedentary Time and Physical Activity in Adults and their Associations with Cardiometabolic Health. *Medicine & Science in Sports & Exercise*, *54*(12), 2118–2128. https://doi.org/10.1249/MSS.0000000000003008.

Farrahi, V., Rostami, M., Nauha, L., … Oussalah, M. (2023). Replacing sedentary time with

physical activity and sleep: Associations with cardiometabolic health markers in adults.

Scandinavian Journal of Medicine & Science in Sports, *33*(6), 907–920.

https://doi.org/https://doi.org/10.1111/sms.14323

Feng, H., Yang, L., Liang, Y. Y., … others. (2023). Associations of timing of physical activity

with all-cause and cause-specific mortality in a prospective cohort study. *Nature*

Communications, *14*(930).

Feng, Q., Albalak, G., van Dijk, K. W., … van Heemst, D. (2025). Accelerometer-measured

chronoactivity and type 2 diabetes risk: A prospective study in UK Biobank participants.

Preventive Medicine, *197*, 108312.

https://doi.org/https://doi.org/10.1016/j.ypmed.2025.108312

Fukuoka, Y., Zhou, M., Vittinghoff, E., Haskell, W., Goldberg, K., & Aswani, A. (2018).

Objectively measured baseline physical activity patterns in women in the mPED trial:

cluster analysis. *JMIR Public Health and Surveillance*, *4*(1), e10.

Full, K. M., Moran, K., Carlson, J., … Kerr, J. (2019). Latent profile analysis of

accelerometer-measured sleep, physical activity, and sedentary time and differences in

health characteristics in adult women. *PLOS ONE*, *14*(6), e0218595.

Giurgiu, M., von Haaren-Mack, B., Fiedler, J., … Bussmann, J. B. J. (2025). The wearable

landscape: Issues pertaining to the validation of the measurement of 24-h physical

activity, sedentary, and sleep behavior assessment. *Journal of Sport and Health Science*,

14, 101006. https://doi.org/https://doi.org/10.1016/j.jshs.2024.101006

Goldsmith, J., Zipunnikov, V., & Schrack, J. (2015). Generalized Multilevel Function-on-

Scalar Regression and Principal Component Analysis. *Biometrics*, *71*(2), 344–353.

https://doi.org/10.1111/biom.12278

Graves, J. L., Qiao, Y. S., Moored, K. D., … Glynn, N. W. (2021). Profiles of Accelerometry-

Derived Physical Activity Are Related to Perceived Physical Fatigability in Older Adults.

Sensors, *21*(5), 1718. https://doi.org/10.3390/s21051718

Gupta, N., Hallman, D. M., Dumuid, D., … Holtermann, A. (2020). Movement behavior

profiles and obesity: a latent profile analysis of 24-h time-use composition among

Danish workers. *International Journal of Obesity*, *44*(2), 409–417.

https://doi.org/10.1038/s41366-019-0419-8

Gupta, N., Hallman, D. M., Korshøj, M., Rasmussen, C. L., & Holtermann, A. (2025). From

Single Movement Behaviors to Complete 24-h Behaviors Profiles and Multiple Health

Outcomes—A Cross-Sectional Study Using Accelerometry. *Scandinavian Journal of*

Medicine & Science in Sports, *35*(5), e70060.

https://doi.org/https://doi.org/10.1111/sms.70060

Hafkamp, F., Dekker, L., Tio, R., … van de Ven, R. (9900). Characterizing High Risk Patients

in Heart Failure: A Latent Class Analysis of Rehospitalization and Mortality. *Journal of*

Cardiovascular Nursing.

Herold, F., Zou, L., Theobald, P., … Gronwald, T. (2025). Beyond FITT: addressing density in

understanding the dose–response relationships of physical activity with health—an

example based on brain health. *European Journal of Applied Physiology*.

https://doi.org/10.1007/s00421-025-05858-3

Hibbing, P. R., Bellettiere, J., & Carlson, J. A. (2022). Sedentary Profiles: A New Perspective

on Accumulation Patterns in Sedentary Behavior. *Medicine and Science in Sports and*

Exercise, *54*(4), 696–706. https://doi.org/10.1249/MSS.0000000000002830

Howard, M. C., & Hoffman, M. E. (2017). Variable-Centered, Person-Centered, and Person-

Specific Approaches: Where Theory Meets the Method. *Organizational Research*

Methods, *21*(4), 846–876. https://doi.org/10.1177/1094428117744021

Kang, J., Ciecierski, C. C., Malin, E. L., … Hitsman, B. (2014). A latent class analysis of

cancer risk behaviors among U.S. college students. *Preventive Medicine*, *64*, 121–125.

https://doi.org/https://doi.org/10.1016/j.ypmed.2014.03.023

Kebede, M., Howard, A. G., Ren, Y., … Evenson, K. R. (2024). A systematic scoping review

of latent class analysis applied to accelerometry-assessed physical activity and sedentary

behavior. *PLOS ONE*, *19*(1), e0283884.

Keiser, T., Katz, S., Robson, S. M., … Patterson, F. (2024). Association between time-of-day

for eating, exercise, and sleep with blood pressure in adults with elevated blood pressure

or hypertension: a systematic review. *Journal of Hypertension*, *42*(6), 95.

Länsitie, M., Niemelä, M., Kangas, M., … Korpelainen, R. (2021). Physical activity profiles

and glucose metabolism — A population-based cross-sectional study in older adults.

TRANSLATIONAL SPORTS MEDICINE, *4*(4), 439–446.

https://doi.org/https://doi.org/10.1002/tsm2.237

Lee, I.-M., Shiroma, E. J., Lobelo, F., Puska, P., Blair, S. N., & Katzmarzyk, P. T. (2012).

Effect of physical inactivity on major non-communicable diseases worldwide: an

analysis of burden of disease and life expectancy. *The Lancet*, *380*(9838), 219–229.

https://doi.org/https://doi.org/10.1016/S0140-6736(12)61031-9

Lee, P. H., Yu, Y.-Y., McDowell, I., Leung, G. M., & Lam, T. H. (2013). A cluster analysis of

patterns of objectively measured physical activity in Hong Kong. *Public Health*

Nutrition, *16*(8), 1436–1444. https://doi.org/10.1017/S1368980012003631

Leskinen, T., Lima Passos, V., Dagnelie, P. C., … Koster, A. (2023). Daily Physical Activity

Patterns and Their Associations with Cardiometabolic Biomarkers: The Maastricht

Study. *Medicine & Science in Sports & Exercise*, *55*(5), 837–846.

Leviäkangas, A., Korpelainen, R., Pinola, P., … Farrahi, V. (2024). Associations of

accelerometer-estimated free-living daily activity impact intensities with 10-year

probability of osteoporotic fractures in adults. *Gait & Posture*, *112*, 22–32.

https://doi.org/https://doi.org/10.1016/j.gaitpost.2024.05.002

Li, X., Zeng, J., Chen, B., … Xu, S. (2024). Effects of the Timing of Intense Physical Activity

on Hypertension Risk in a General Population: A UK-Biobank Study. *Current*

Hypertension Reports, *26*(2), 81–90. https://doi.org/10.1007/s11906-023-01278-w

Liu, P., Wang, Z., Liu, N., & Peres, M. A. (2023). A scoping review of the clinical application

of machine learning in data-driven population segmentation analysis. *Journal of the*

American Medical Informatics Association, *30*(9), 1573–1582.

https://doi.org/10.1093/jamia/ocad111

Ma, T., Bennett, T., Lee, C.-D., & Wicklow, M. (2023). The diurnal pattern of moderate-to-

vigorous physical activity and obesity: a cross-sectional analysis. *Obesity*, *31*(10), 2638–

2647. https://doi.org/https://doi.org/10.1002/oby.23851

Makarem, N., German, C. A., Zhang, Z., … Shechter, A. (2024). Rest-Activity Rhythms Are

Associated With Prevalent Cardiovascular Disease, Hypertension, Obesity, and Central

Adiposity in a Nationally Representative Sample of US Adults. *Journal of the American*

Heart Association, *13*(1), e032073. https://doi.org/10.1161/JAHA.122.032073

Marschollek, M. (2016). Associations between sensor-based physical activity behaviour

features and health-related parameters. *Human Movement Science*, *45*, 1–6.

Matricciani, L., Paquet, C., Fraysse, F., … Olds, T. (2021). Sleep and cardiometabolic risk: a

cluster analysis of actigraphy-derived sleep profiles in adults and children. *Sleep*, *44*(7),

zsab014. https://doi.org/10.1093/sleep/zsab014

McGregor, D. E., Palarea-Albaladejo, J., Dall, P. M., Hron, K., & Chastin, S. F. M. (2020).

Cox regression survival analysis with compositional covariates: Application to modelling

mortality risk from 24-h physical activity patterns. *Statistical Methods in Medical*

Research, *29*(5), 1447–1465. https://doi.org/https://doi.org/10.1177/0962280219864125

Mekary, R. A., Willett, W. C., Hu, F. B., & Ding, E. L. (2009). Isotemporal substitution

paradigm for physical activity epidemiology and weight change. *American Journal of*

Epidemiology, *170*(4), 519–527.

Migueles, J. H., Aadland, E., Andersen, L. B., … Ortega, F. B. (2022). GRANADA consensus

on analytical approaches to assess associations with accelerometer-determined physical

behaviours (physical activity, sedentary behaviour and sleep) in epidemiological studies.

British Journal of Sports Medicine, *56*, 376–384.

https://doi.org/http://dx.doi.org/10.1136/bjsports-2020-103604

Migueles, J. H., Cadenas-Sanchez, C., Ekelund, U., … Ortega, F. B. (2017). Accelerometer

data collection and processing criteria to assess physical activity and other outcomes: a

systematic review and practical considerations. *Sports Medicine*, *47*(9), 1821–1845.

https://doi.org/10.1007/s40279-017-0716-0

Migueles, J. H., Cadenas-Sanchez, C., Rowlands, A. V, … Ortega, F. B. (2019).

Comparability of accelerometer signal aggregation metrics across placements and

dominant wrist cut points for the assessment of physical activity in adults. *Scientific*

Reports, *9*(1), 18235. https://doi.org/10.1038/s41598-019-54267-y

Montoye, A. H. K., Moore, R. W., Bowles, H. R., Korycinski, R., & Pfeiffer, K. A. (2018).

Reporting accelerometer methods in physical activity intervention studies: a systematic

review and recommendations for authors. *Br J Sports Med*, *52*(23), 1507–1516.

Moreno-Llamas, A., García-Mayor, J., & De la Cruz-Sánchez, E. (2022). How Europeans

move: a moderate-to-vigorous physical activity and sitting time paradox in the European

Union. *Public Health*, *203*, 1–8.

https://doi.org/https://doi.org/10.1016/j.puhe.2021.11.016

Nawrin, S. S., Inada, H., Momma, H., & Nagatomi, R. (2024). Twenty-four-hour physical

activity patterns associated with depressive symptoms: a cross-sectional study using big

data-machine learning approach. *BMC Public Health*, *24*(1), 1254.

https://doi.org/10.1186/s12889-024-18759-5

Niemelä, M., Kangas, M., Farrahi, V., … Jämsä, T. (2019). Intensity and temporal patterns of

physical activity and cardiovascular disease risk in midlife. *Preventive Medicine*, *124*,

33–41. https://doi.org/https://doi.org/10.1016/j.ypmed.2019.04.023

Niemelä, M., Kiviniemi, A., Ikäheimo, T. M., … Farrahi, V. (2023). Compositional

association of 24-hour movement behavior with incident major adverse cardiac events

and all-cause mortality. *Scandinavian Journal of Medicine & Science in Sports*, *33*(5),

641–650. https://doi.org/https://doi.org/10.1111/sms.14315

O'Regan, A., Hannigan, A., Glynn, L., … Woods, C. B. (2021). A cluster analysis of device-

measured physical activity behaviours and the association with chronic conditions, multi-

morbidity and healthcare utilisation in adults aged 45 years and older. *Preventive*

Medicine Reports, *24*, 101641.

https://doi.org/https://doi.org/10.1016/j.pmedr.2021.101641

Page, M. J., McKenzie, J. E., Bossuyt, P. M., … Moher, D. (2021). The PRISMA 2020

statement: an updated guideline for reporting systematic reviews. In *BMJ* (Vol. 372, p.

n71). https://doi.org/10.1136/bmj.n71

Palmberg, L., Löppönen, A., Hyvärinen, M., … Karavirta, L. (2024). Physical Behavior

Profiles Among Older Adults and Their Associations With Physical Capacity and Life-

Space Mobility. *Journal of Aging and Physical Activity*, *32*(4), 472–479.

https://doi.org/10.1123/japa.2023-0225

Paolillo, E. W., Lee, S. Y., VandeBunte, A., … Casaletto, K. B. (2023). Data-driven physical

actigraphy patterns relate to cognitive and vascular health in older adults. *Experimental*

Gerontology, *178*, 112231. https://doi.org/https://doi.org/10.1016/j.exger.2023.112231

Pedišić, Ž. (2014). Measurement issues and poor adjustments for physical activity and sleep

undermine sedentary behaviour research—the focus should shift to the balance between

sleep, sedentary behaviour, standing and activity. *Kinesiology*, *46*(1), 135–146.

Pedišić, Ž., & Bauman, A. (2015). Accelerometer-based measures in physical activity

surveillance: current practices and issues. *British Journal of Sports Medicine*, *49*, 219–

223. https://doi.org/10.1136/bjsports-2013-093407

Pedišić, Ž., Dumuid, D., & S Olds, T. (2017). Integrating sleep, sedentary behaviour, and

physical activity research in the emerging field of time-use epidemiology: definitions,

concepts, statistical methods, theoretical framework, and future directions. *Kinesiology*,

49(2), 252–269.

Reuter, C., Bellettiere, J., Liles, S., … Natarajan, L. (2020). Diurnal patterns of sedentary

behavior and changes in physical function over time among older women: a prospective

cohort study. *International Journal of Behavioral Nutrition and Physical Activity*, *17*(1),

88. https://doi.org/10.1186/s12966-020-00992-x

Rollo, S., Antsygina, O., & Tremblay, M. S. (2020). The whole day matters: Understanding

24-hour movement guideline adherence and relationships with health indicators across

the lifespan. *Journal of Sport and Health Science*, *9*(6), 493–510.

https://doi.org/https://doi.org/10.1016/j.jshs.2020.07.004

Rosenberger, M. E., Fulton, J. E., Buman, M. P., … Haskell, W. L. (2019). The 24-hour

activity cycle: a new paradigm for physical activity. *Medicine and Science in Sports and*

Exercise, *51*(3), 454–464. https://doi.org/10.1249/MSS.0000000000001811

Seppänen, M., Lankila, T., Niemelä, M., … Farrahi, V. (2025). Compositional associations of

24-h physical activities, sedentary time and sleep with depressive symptoms in urban and

rural residents: a cross-sectional study. *BMC Medicine*, *23*(219).

https://doi.org/https://doi.org/10.1186/s12916-025-04051-9

Shim, J., Fleisch, E., & Barata, F. (2023). Wearable-based accelerometer activity profile as

digital biomarker of inflammation, biological age, and mortality using hierarchical

clustering analysis in NHANES 2011–2014. *Scientific Reports*, *13*(1), 9326.

https://doi.org/10.1038/s41598-023-36062-y

Smagula, S. F., Zhang, G., Gujral, S., … Krafty, R. T. (2022). Association of 24-Hour Activity

Pattern Phenotypes With Depression Symptoms and Cognitive Performance in Aging.

JAMA Psychiatry, *79*(10), 1023–1031. https://doi.org/10.1001/jamapsychiatry.2022.2573

Stamatakis, E., Gale, J., Bauman, A., Ekelund, U., Hamer, M., & Ding, D. (2019). Sitting

time, physical activity, and risk of mortality in adults. *Journal of the American College of*

Cardiology, *73*(16), 2062–2072. https://doi.org/10.1016/j.jacc.2019.02.031

Strain, T., Flaxman, S., Guthold, R., … Zoma, L. R. (2024). National, regional, and global

trends in insufficient physical activity among adults from 2000 to 2022: a pooled

analysis of 507 population-based surveys with 5.7 million participants. *The Lancet*

Global Health, *12*(8), e1232–e1243. https://doi.org/10.1016/S2214-109X(24)00150-5

von Rosen, P., Dohrn, I.-M., & Hagströmer, M. (2020). Latent profiles analysis of physical

activity and sedentary behaviour with mortality risk: a 15-year follow-up. *Scandinavian*

Journal of Medicine & Science in Sports, *30*(10), 1949–1956.

https://doi.org/https://doi.org/10.1111/sms.13761

Waffenschmidt, S., Knelangen, M., Sieben, W., Bühn, S., & Pieper, D. (2019). Single

screening versus conventional double screening for study selection in systematic

reviews: a methodological systematic review. *BMC Medical Research Methodology*,

19(1), 132. https://doi.org/10.1186/s12874-019-0782-0

Weller, B. E., Bowen, N. K., & Faubert, S. J. (2020). Latent Class Analysis: A Guide to Best

Practice. *Journal of Black Psychology*, *46*(4), 287–311.

https://doi.org/10.1177/0095798420930932

Wells, G. A., Shea, B., O'Connell, D., … others. (2000). *The Newcastle-Ottawa Scale (NOS)*

for assessing the quality of nonrandomised studies in meta-analyses.

Woo, S. E., Hofmans, J., Wille, B., & Tay, L. (2024). Person-Centered Modeling: Techniques

for Studying Associations Between People Rather than Variables. *Annual Review of*

Organizational Psychology and Organizational Behavior, *11*, 453–480.

https://doi.org/https://doi.org/10.1146/annurev-orgpsych-110721-045646

Xu, D., & Tian, Y. (2015). A Comprehensive Survey of Clustering Algorithms. *Annals of Data*

Science, *2*(2), 165–193. https://doi.org/10.1007/s40745-015-0040-1

Yamamoto, N., Maruyama, K., Saito, I., … Osawa, H. (2023). Latent profile analysis

approach to the relationship between daily ambulatory activity patterns and metabolic

syndrome in middle-aged and elderly Japanese individuals: The Toon Health Study.

Environmental Health and Preventive Medicine, *28*, 57.

https://doi.org/10.1265/ehpm.23-00110

Yamamoto, N., Maruyama, K., Saito, I., … Osawa, H. (2025). Patterns of daily ambulatory

activity and the onset of metabolic syndrome in middle-aged and older Japanese women:

the Toon Health Study. *Environmental Health and Preventive Medicine*, *30*, 11.

https://doi.org/10.1265/ehpm.24-00313

Yan, S., Kwan, Y. H., Tan, C. S., Thumboo, J., & Low, L. L. (2018). A systematic review of

the clinical application of data-driven population segmentation analysis. *BMC Medical*

Research Methodology, *18*(1), 121. https://doi.org/10.1186/s12874-018-0584-9

Yerramalla, M. S., Chen, M., Dugravot, A., van Hees, V. T., & Sabia, S. (2024). Association

between profiles of accelerometer-measured daily movement behaviour and mortality

risk: a prospective cohort study of British older adults. *BMJ Open Sport & Exercise*

Medicine, *10*(2). https://doi.org/10.1136/bmjsem-2023-001873

Zhang, Y., Kivimäki, M., Carrillo-Larco, R. M., … Xu, X. (2024). Diurnal patterns of

accelerometer-measured physical activity and sleep and risk of all-cause mortality: a

follow-up of the National Health and Nutrition Examination Surveys (NHANES).

International Journal of Behavioral Nutrition and Physical Activity, *21*(120).

https://doi.org/10.1186/s12966-024-01673-9

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

No datasets were generated or analysed during the current study.

Competing interests

The authors have no competing interests to declare.

Funding

This work was conducted without any financial support from external sources. Authors are

supported by TU Dortmund University and performed this study as part of their work.

Authors' contributions

VF conceived the idea, conducted the literature search, reviewed the titles and abstracts, and

prepared the manuscript draft. EF contributed to assessing the eligibility of articles identified

during the title and abstract screening and revised the manuscript.

Acknowledgments

None.