The genetic architecture of postoperative delirium after major surgery and its

relationship with non-postoperative neurocognitive conditions

Richard A. Armstrong,[1*] Paul Yousefi,[1,2] Ben Gibbison,[3] Golam M Khandaker,[1,2,4,5]

Tom R Gaunt[1,2]

1   MRC Integrative Epidemiology Unit, Bristol Medical School, University of

Bristol, Bristol, UK.

2   NIHR Bristol Biomedical Research Centre, University Hospitals Bristol and

Weston NHS Foundation Trust and University of Bristol, Bristol, UK.

3   Bristol Medical School, University of Bristol, Bristol, UK.

4   Avon and Wiltshire Mental Health Partnership NHS Trust, Bristol, UK.

5   Centre for Academic Mental Health, Population Health Sciences, Bristol

Medical School, University of Bristol, Bristol, UK.

*Correspondence to: Richard A Armstrong

Email: ra17848@bristol.ac.uk

X: @drrichstrong

Short title: Genetic architecture of postoperative delirium

Abstract

Background

Postoperative delirium is the most common postoperative complication in older

individuals. Genome-wide association studies (GWAS) can provide insights into how

genetic factors influence postoperative risk. We examined the genetic architecture of

postoperative delirium after major surgery and its relationship with related cognitive

conditions (delirium of any type and Alzheimer's disease, including the *APOE* ε4

allele).

Methods and Findings

A case-control GWAS was performed in UK Biobank to identify genetic variants

associated with postoperative delirium, adjusted for age, sex, genetic chip and the

first ten principal components. These results were then used in genetic correlation

and polygenic risk score analyses to investigate shared genetic risk between

postoperative delirium and a) delirium of all causes, and b) Alzheimer's disease.

The GWAS (1016 cases, 139,148 controls) identified seven Single Nucleotide

Polymorphisms (SNPs) that mapped to four genes (*APOE, TOMM40, APOC1* and

PVRL2); $p < 5 \times 10^{-8}$. Five SNPs remained significant after excluding pre-existing

dementia, and two after excluding subsequent dementia. The lead SNP was

rs429358, a missense variant of *APOE*. Genetic correlation and polygenic risk score

analyses revealed evidence of shared genetic architecture and risk between

postoperative delirium and Alzheimer's disease (rho 0.68, 95% CI [0.46,0.81];

p<0.001). After adjustment for age and sex, the *APOE* ε4 isoform had a dose-

response effect on risk (odds ratios for one and two copies: 1.75, 95% CI [1.53,2.0],

and 4.19, 95% CI [3.25,5.41], respectively; p <0.001). The main limitations of the

study include the reliance upon clinical coding for outcome definition and limited

statistical power to detect small or modest genetic effects.

Conclusions

We identified genetic variants associated with increased risk of postoperative

delirium. We also found evidence of shared genetic liability with Alzheimer's disease

via *APOE*, complementing recent large-scale studies in all-cause delirium. If

validated the findings have potential clinical applications including preoperative risk

stratification and early identification of pre-clinical Alzheimer's disease risk.

Author Summary

Why was this study done?

• Postoperative delirium is the most common postoperative complication in
older individuals.

• Little is known about the genetic basis of postoperative delirium.

• Knowledge of the genetic risk factors could help treatment or prevention in the
future.

What did the researchers do and find?

• We performed a genome-wide association study of postoperative delirium
including 1016 cases and 139,148 controls.

• We found genetic variants associated with increased postoperative delirium
risk.

• We also found genetic similarity between postoperative delirium, all-cause
delirium, and Alzheimer's disease.

What do these findings mean?

• Postoperative delirium has shared biology with Alzheimer's disease and may
represent unmasking of pre-clinical disease.

• Individuals with genetic risk for Alzheimer's disease have increased risk of
postoperative delirium and vice versa.

• Due to the relatively small number of cases and the fact that we may not have
detected all the true positive diagnoses in the dataset, we were unable to
detect small or modest genetic effects in this study.

Introduction

Postoperative delirium is the most common postoperative complication in older

individuals, with an incidence of up to 65% [1,2]. It has substantial health and

resource implications including increased morbidity, mortality and healthcare costs

[1,3]. Hypotheses of the underlying mechanisms span a range of pathways including

release of excitotoxic neurotransmitters, neuronal injury and inflammation,

physiological stressors, metabolic derangements, electrolyte disorders and genetic

factors [1,2,4–8]. Postoperative delirium has been associated with an increased risk

of cognitive decline and incident dementia [9] but there is conflicting evidence on

how the acute occurrence of postoperative delirium interacts with longer-term

cognitive decline and neurodegeneration [10].

Whilst the aging global population and increasing numbers of older adults

undergoing surgery have made perioperative brain health an increasing focus of

research and clinical initiatives [11], the ability of clinical teams to produce

personalised assessments of clinical risk for individuals remains limited. Improved

detection of underlying biological risk in conjunction with established clinical risk

factors could aid in preoperative risk prediction, facilitate better-informed shared-

decision making and allow targeting of multicomponent interventions to prevent

delirium [11] with potential reductions in morbidity, mortality and resource utilisation.

Population-based genome-wide association studies (GWAS) are a powerful

approach for identifying genetic risk factors associated with complex conditions of

multifactorial aetiology. Their results can aid prediction but also identify underlying

biological mechanisms and potential targets for intervention. They have been

extensively applied to a wide range of clinical phenotypes [12], but rarely to

postoperative delirium specifically. Previous GWAS that have included postoperative

neurocognitive complications have failed to identify any genome-wide significant

associations [13–15] - most likely because they suffered from limited sample sizes

and heterogeneity of outcomes.

This study aimed to evaluate the association between genetic variants genome-wide

and postoperative delirium following major surgery, and to explore whether

postoperative delirium shares genetic risk factors with related conditions, specifically

delirium of all causes and Alzheimer's disease. The limitations of previous studies

are addressed by defining a major surgery cohort in a large scale national biobank

and focusing on a single outcome phenotype.

Methods

Study populations

The data source for the primary analysis was UK Biobank, a long-term prospective

cohort study that provides a large-scale biomedical database and research resource

containing genetic, lifestyle and health information from half a million UK participants

recruited between 2006 and 2010 (http://www.ukbiobank.ac.uk). The UK Biobank

study was approved by the North-West Multi-centre Research Ethics Committee and

all participants provided written informed consent [16]. Additionally, publicly available

GWAS summary statistics from the FinnGen study were used as an external cohort

for genetic correlation analyses. FinnGen is a large-scale genomics initiative that has

analysed over 500,000 Finnish biobank samples and correlated genetic variation

with health data to understand disease mechanisms and predispositions. The project

is a collaboration between research organisations and biobanks within Finland and

international industry partners [17].

Genotyping

UK Biobank genotyping was performed by Affymetrix using two purpose-designed

arrays. Genotype data were quality controlled, phased and ~96 million genotypes

were imputed using the Haplotype Reference Consortium and UK10K haplotype

resources. FinnGen genotyping used a ThermoFisher Axiom custom chip array with

imputation to a reference dataset derived from Finnish whole-genome sequences.

Further information on the methods and QC pipelines are available [16,18].

Inclusion and exclusion criteria

Eligible participants were those undergoing major, inpatient surgery after their date of

enrolment in UK Biobank. Linked hospital inpatient data is available for all UK

Biobank participants with follow-up to 31 May 2022 for Wales, 31 August 2022 for

Scotland and 31 October 2022 for England [19]. Major surgery was defined by Office

of Population Censuses and Surveys Classification of Interventions and Procedures,

Version 4 (OPCS-4) code using the Bupa Schedule of Procedures and Abbott

classification [20,21] and diagnoses of delirium were identified using International

Classification of Diseases, Tenth Revision (ICD-10) code F05. Exclusion criteria

included planned day case surgery and a previous diagnosis of delirium (see S2

Appendix for more detail on phenotype definition and exclusions).

Outcome definition

The primary outcome was a first diagnosis of delirium within 30 days of surgery,

including diagnoses made during readmission. Controls were defined as those

undergoing major surgery with no delirium diagnosis within 30 days. The 30-day

window was used to align with recommendations on the nomenclature of

postoperative delirium [22] and other widely-used sources [23].

Genome-wide Association Study (GWAS)

Case–control GWAS adjusted for age, sex, genetic chip and the first ten principal

components was performed using regenie (v3.2.2) [24]. Participants with missing

data for these variables were excluded. Quality control and ancestry definition were

performed by previously established workflows and methods [25]. Post-hoc analyses

were conducted using Functional Mapping and Annotation of Genome-Wide

Association Studies (FUMA GWAS [26]). Independent significant Single Nucleotide

Polymorphisms (SNPs) were defined as those meeting a genome-wide significance

threshold of $p < 5 \times 10^{-8}$ and independent of each other at $r^2 < 0.6$. Lead SNPs were

identified from independent significant SNPs at $r^2 < 0.1$. Gene-based analysis was

performed using Multi-marker Analysis of GenoMic Annotation (MAGMA [27]).

Sensitivity analyses

The primary GWAS analysis did not include additional clinical covariates as they may

be mediators of genetic risk or colliders given the selection of a surgical patient

cohort. To explore the robustness of the primary analysis the following sensitivity

analyses were performed: a) after excluding participants who had a previous

dementia diagnosis; b) excluding participants who had a previous dementia

diagnosis or were diagnosed with dementia after the index procedure; c) stratified by

cardiothoracic/non-cardiothoracic surgery; d) adjusted for Charlson Comorbidity

Index (0-1 vs ≥2, [28]); and e) conditional on Apolipoprotein E (*APOE*) ε4 haplotype

count (0, 1 or 2) to identify associated variants independent of *APOE*.

Post-GWAS follow-up analyses

Data-driven exploratory follow-up analyses were informed by the results of the

primary GWAS analysis. Two approaches were taken to formally assess shared

genetic architecture between postoperative delirium and a) delirium of any type; b)

Alzheimer's Disease: genetic correlation analysis and polygenic risk scoring.

Genetic correlation analysis

Genetic correlation analyses were performed between the UKBiobank GWAS results

presented here and the external FinnGen study cohort. GWAS summary statistics

(Freeze 9, https://r9.finngen.fi/) were obtained for a) delirium (any cause); and b)

Alzheimer's disease. Local Analysis of [co]Variant Association (LAVA [29]) was used

to analyse bivariate local genetic correlation between these phenotypes and

postoperative delirium (using the summary statistics from the primary GWAS

analysis in UK Biobank). The genome is partitioned into 2,495 blocks for LAVA

analysis so a Bonferroni-adjusted p-value of 0.05/2,495 (2×10^{-5}) was used to define

statistical significance.

Polygenic risk score analysis

Polygenic risk scores (PRS) calculate a weighted sum of risk based on alleles

associated with a disease or outcome phenotype. The Polygenic Score Catalog

(https://www.pgscatalog.org/) was used to identify polygenic risk scores for

Alzheimer's disease. No polygenic risk scores were identified for delirium. The

pgsc_calc workflow [30] was used to download, variant match and calculate

individual polygenic scores for the primary UK Biobank cohort. Individual scores

were then centred and scaled before being summed and averaged. Odds ratios for

the association between increasing quintile of polygenic risk score for Alzheimer's

disease and postoperative delirium were estimated by logistic regression with

adjustment for age and sex using the *glm* function in R (version 4.4.0). Sensitivity

analyses were performed with additional adjustment for Charlson Comorbidity Score

(0-1 vs ≥2, [28]). A Bonferroni correction for the number of comparisons against the

reference quintile was applied, resulting in a significance threshold of p < 0.0125

(0.05/4).

Apolipoprotein E (APOE) genotype analysis

It has been recommended that the *APOE* region of chromosome 19 should be

excluded from Alzheimer's disease polygenic risk scores, with the *APOE* ε4 allele

included as an additional covariate in modelling, due to its outsize effect relative to

other regions across the genome [31]. The subset of polygenic scores identified

above which did not include the *APOE* region (defined as a 10 kilobase window

either side of region from the start position of *TOMM40*, chr19:45,394,477, to the

stop position of *APOC1*, chr19:45,422,606, build 37/hg19 [31]) were therefore

considered in an additional analysis, adjusted for age, sex and *APOE* ε4 allele count

(0, 1, 2).

This study is reported as per the Strengthening the Reporting of Observational

Studies in Epidemiology (STROBE) guideline (S1 Checklist).

Results

Participant characteristics

A total of 1016 cases of postoperative delirium and 139,148 controls were identified

(Figure 1). Individuals who developed postoperative delirium tended to be older,

male and had higher prevalence of comorbidities (Table 1). Compared to controls, a

higher proportion of patients in the postoperative delirium group underwent

emergency (532 (52%) vs 29,530 patients (21%)) and cardiothoracic surgery (172

(17%) vs 5041 patients (3.6%)).

Table 1 Participant characteristics for postoperative delirium cases and controls.

	Cases n = 1016	Controls n = 139,148	p-value
Age (yr), mean (SD)	72 (6)	64 (8)	<0.001
Sex, number (%)			<0.001
Female	384 (38)	77,244 (56)	
Male	632 (62)	61,904 (44)	
Ethnicity, number (%)			0.2
Asian or Asian British	9 (0.9)	2301 (1.7)	
Black or Black British	10 (1.0)	1644 (1.2)	
Chinese	0 (0)	239 (0.2)	
Mixed	5 (0.5)	675 (0.5)	
Other ethnic group	3 (0.3)	984 (0.7)	
White	978 (97)	132,676 (96)	
Missing	11 (1.1)	629 (0.5)	
Charlson Comorbidity Index, median (IQR)	1 (0–3)	0 (0–1)	<0.001
Missing	25 (2.5)	15,817 (11)	
Admission method, number (%)			<0.001
Elective	434 (43)	106,725 (77)	
Emergency	532 (52)	29,530 (21)	
Maternity	0 (0)	113 (<0.1)	

Not known	0 (0)	29 (<0.1)
Transfer	50 (4.9)	2751 (2.0)
Surgical specialty, number (%)		<0.001
Cardiothoracic surgery	172 (17)	5041 (3.6)
General surgery	196 (19)	26,436 (19)
Neurosurgery	55 (5.4)	5192 (3.7)
Other	186 (18)	36,532 (26)
Trauma and orthopaedics	338 (33)	53,722 (39)
Urology	37 (3.6)	10,628 (7.6)
Vascular	32 (3.1)	1597 (1.1)

Numbers are number (proportion), mean (standard deviation, SD) or median

(interquartile range, IQR). Continuous variables compared using Wilcoxon rank sum

test; categorical variables using Pearon's Chi-squared test or Fisher's exact test for

cell counts of 5 or fewer. Additional details can be found in Table A in S3 Appendix.

Genome-wide significant associations for postoperative delirium

Seven independent, statistically significant SNPs were identified within a single

genetic risk region on chromosome 19 (Figure 2, Table 2). The Q-Q plot showed no

evidence of systematic inflation (genomic inflation factor (λ) 1.01, Figure 3). The lead

SNP was rs429358, a missense variant in the *APOE* gene that, together with rs7412,

defines the $\varepsilon2/\varepsilon3/\varepsilon4$ isoforms linked to Alzheimer's disease risk. A total of 21 genes

were mapped to this region, including four (*APOE*, *APOC1, TOMM40* and *PVRL2*)

that were significantly associated in MAGMA gene-level analysis (Figure 4; Table B,

S3 Appendix). The Q-Q plot showed no evidence of systematic inflation (genomic

inflation factor (λ) 1.0, Figure 5).

Table 2 Independent significant single-nucleotide polymorphisms passing genome-wide significance threshold in case-control GWAS of postoperative delirium.

rsID	Chr:position	Nearest gene	Function	Effect allele	Odds ratio (95% CI)	p
rs429358	19:45411941	*APOE*	exonic	T	0.54 (0.49,0.59)	1.41×10^{-29}
rs157592	19:45424514	*APOC1*	intergenic	A	0.58 (0.53,0.64)	1.77×10^{-23}
rs157582	19:45396219	*TOMM40*	intronic	C	0.65 (0.59,0.71)	3.42×10^{-17}
rs11556505	19:45396144	*TOMM40*	exonic	C	0.62 (0.56,0.69)	3.33×10^{-16}
rs10119	19:45406673	*TOMM40*	UTR3	G	0.7 (0.64,0.77)	3.34×10^{-13}
rs75627662	19:45413576	*APOE*	downstream	C	0.72 (0.65,0.79)	5.66×10^{-10}
rs12691088	19:45418486	*APOC1*	Intronic	G	0.45 (0.35,0.58)	1.65×10^{-8}

rsID: Reference SNP cluster ID; UTR3: 3' untranslated region.

GWAS results after excluding existing or incident dementia

After excluding individuals with a pre-existing diagnosis of dementia (56 cases (5.7%) and 375 controls (0.3%)), 5 of the 6 independent significant SNPs remained significant. However, rs75627662 (a downstream variant of *APOE*) no longer reached genome-wide significance. Three genes, *APOE*, *APOC1* and *TOMM40*, remained significant on MAGMA testing (Figures A and B, S3 Appendix). When patients who went on to be diagnosed with dementia during UK Biobank follow-up were also excluded (141 cases, 13.9%, and 2358 controls, 1.7%), two SNPs remained genome-wide significant, rs429358 and rs157592. The MAGMA results were unchanged (Figures C and D, S3 Appendix).

GWAS results in cardiothoracic and non-cardiothoracic surgery groups

The results for the non-cardiothoracic surgery subgroup (798 cases, 126,973

controls) were similar to those in the overall cohort with five independent significant

SNPs identified in the same risk locus on chromosome 19 and three of the four

genes remained significant on MAGMA testing (*APOE*, *APOC1* and *TOMM40*). The

lead SNP was the same as in the primary GWAS and the other variants either

overlapped with the primary analysis or were in strong linkage disequilibrium (r^2 >

0.97) with them (Table C in S3 Appendix). The cardiothoracic analysis was likely

underpowered and returned no genome-wide significant results at $p < 5 \times 10^{-8}$.

Multiple SNPs reached a lower indicative threshold of significance including the lead

SNP from the primary analysis, rs429358 (p = 3.8 x 10^{-7}; Table D in S3 Appendix).

GWAS results adjusted for Charlson Comorbidity Index

A total of 495 (48.7%) cases and 26,659 (19.2%) controls had a Charlson

Comorbidity Index (CCI) ≥2 (χ^2 = 562.42, p < 0.001). GWAS results after adjustment

for CCI were consistent with the primary analysis, returning the same seven

independent significant SNPs on chromosome 19 (Table E in S3 Appendix).

GWAS results conditional on Apolipoprotein E (APOE) ε4 allele count

APOE ε4 allele counts differed between cases and controls (0, 1 and 2 copies in 566

(57.4%), 348 (35.3%) and 71 (7.2%) cases vs 95,489 (71.6%), 34,754 (26.1%) and

3,082 (2.3%) controls; χ^2 = 159.37, p < 0.001). After adjustment for *APOE* ε4 allele

count, all the independent significant SNPs from the primary GWAS analysis lost

statistical significance (Figure E in S3 Appendix).

Genetic correlation with Alzheimer's disease

Significant local genetic correlation was identified on chromosome 19 between

postoperative delirium and both delirium of all causes and Alzheimer's disease

(Table 3). The correlated region (chr19:45040933-45893307) encompassed the

independent significant SNPs from the primary GWAS.

Table 3 Results of Local Analysis of [co]Variant Association (LAVA) testing between

postoperative delirium GWAS results and FinnGen GWAS summary statistics.

FinnGen phenotype	FinnGen sample size (cases / controls)	Rho (95% CI)	R^2 (95% CI)	p
Delirium (all)	3,039 / 356,660	0.63 (0.46,0.81)	0.40 (0.22,0.66)	3.7×10^{-10}
Alzheimer's disease	13,393 / 363,884	0.68 (0.57,0.80)	0.47 (0.32,0.65)	5.2×10^{-21}

Rho represents the local genetic correlation coefficient. R^2 represents the proportion

of variance in postoperative delirium explained by the comparator phenotype.

Association of genetic predisposition for Alzheimer's disease with postoperative

delirium

Prior to averaging, forty-two PRS for Alzheimer's disease were identified and passed

variant matching (Table F in S3 Appendix). Increased polygenic risk for Alzheimer's

disease was associated with increased risk of postoperative delirium in the UK

Biobank cohort after adjusting for age and sex (adjusted OR PRS quintile 4: 1.49

(95% CI [1.2,1.85]), PRS quintile 5: 2.29 (95% CI [1.87,2.8]); p <0.001; Table 4).

These results were largely unchanged in a sensitivity analysis adjusting for Charlson

Comorbidity Index (Table G in S3 Appendix).

Table 4 Odds ratio for postoperative delirium by Alzheimer's disease polygenic risk

score (PRS) quintile. Results are shown for all 42 polygenic risk scores, adjusted for

age and sex, and the subset which did not include the *APOE* region (15 *APOE*-

independent scores), adjusted for age, sex and *APOE* ε4 genotype.

Term	All scores (n=42)			*APOE*-independent scores (n=15)		
	Unadjusted odds ratio (95% CI)	Adjusted odds ratio (95% CI)	P value	Unadjusted odds ratio (95% CI)	Adjusted odds ratio (95% CI)	P value
PRS quintile 1	1.00 (reference)	1.00 (reference)	-	1.00 (reference)	1.00 (reference)	-
PRS quintile 2	1.06 (0.85,1.34)	1.07 (0.85,1.35)	0.54	1.23 (1,1.52)	1.24 (1,1.53)	0.052
PRS quintile 3	1.25 (1,1.56)	1.25 (1,1.57)	0.047	1.32 (1.07,1.62)	1.32 (1.07,1.62)	0.01
PRS quintile 4	1.46 (1.18,1.81)	1.49 (1.2,1.85)	<0.001	1.24 (1,1.53)	1.24 (1,1.53)	0.049
PRS quintile 5	2.18 (1.79,2.66)	2.29 (1.87,2.8)	<0.001	1.46 (1.19,1.79)	1.50 (1.22,1.84)	<0.001
Age		1.22 (1.2,1.23)	<0.001		1.22 (1.20,1.23)	<0.001
Sex - male		1.82 (1.6,2.07)	<0.001		1.82 (1.60,2.07)	<0.001
APOE ε4 x1		-	-		1.75 (1.53,2.00)	<0.001
APOE ε4 x2		-	-		4.19 (3.25,5.41)	<0.001

Effect of APOE genotype on the association of Alzheimer's disease risk with

postoperative delirium

Fifteen of the polygenic risk scores identified for Alzheimer's disease contained no

variants in the *APOE* region (*APOE*-independent scores, Table F in S3 Appendix).

The effect size of increasing PRS quintile was reduced compared to *APOE*-inclusive

scores and the presence of one or more *APOE* ε4 isoforms was associated with

increased risk of postoperative delirium (Table 4). Similar results were seen with

adjustment for Charlson Comorbidity Index (Table G in S3 Appendix).

Discussion

In this study, using a large national biobank with linked healthcare data, we identified

a genetic risk locus for postoperative delirium on chromosome 19 which included

seven independent significant SNPs and four mapped genes meeting genome-wide

significance. Follow-up analyses in an independent cohort confirmed the presence of

shared genetic risk between postoperative delirium and Alzheimer's disease.

Previous GWAS which have included postoperative delirium as an outcome have

failed to find any genome-wide significant results, but were likely underpowered [13–

15]. Our study, to the best of our knowledge the largest GWAS of postoperative

delirium to date, is substantially better powered than previous analyses and

implicates *APOE* – an established genetic determinant of Alzheimer's disease – in

postoperative delirium risk. The lead SNP identified (rs429358) remained significant

when patients with a pre-existing or subsequent diagnosis of dementia were

excluded, as did the significance of the *APOE* gene on MAGMA testing. We have

identified additional genes associated with postoperative delirium (*APOC1*, *TOMM40*

and *PVRL2*), which were previously reported to be associated with a range of

neurocognitive phenotypes including dementia, cognitive function and decline, brain

biomarkers and cerebral imaging metrics (Table B and Figure F, S3 Appendix).

However, a sensitivity GWAS conditional on *APOE* ε4 allele count resulted in the

loss of significance of all SNPs in the *APOE* region suggesting *APOE* is the sole

contributing factor at this genetic risk locus. We did not replicate the results of an

earlier candidate gene study that identified an association between the cholinergic

genes *CHRM2* and *CHRM4* and postoperative delirium [14] (all three identified

variants showed null effects, Table H in S3 Appendix). However, we used a genome-

wide approach which has methodological advantages including increased genetic

coverage, reduced bias and lower false positive rates and therefore represents a

more robust analysis [32].

Post-GWAS follow-up analyses showed that genetic predisposition for Alzheimer's

disease is associated with risk of perioperative delirium, including a dose-response

effect of *APOE* ε4 allele count. Previous studies investigating the association

between *APOE* ε4 and postoperative delirium have shown mixed results. A recent

study of 19,331 patients (2086 of whom had postoperative delirium) found *APOE* ε4

carriers to be at increased risk of neurocognitive disorders compared with non-

carriers [33]. However, the neurocognitive traits in this study were defined using

Phecodes – which include a wider range of medical diagnoses than the more

specific ones used in our study [34] – and the authors did not report the timepoint at

which outcomes were assessed. Several other studies have found no association

[35–37] whilst one small study (29 cases of postoperative delirium) did report

increased risk with the ε4 allele [38].

Taken together, these data suggest shared genetic architecture between

postoperative delirium, delirium in general, and Alzheimer's disease via *APOE*. This

is supported by a large-scale genetic meta-analysis of all-cause delirium which

identified *APOE* as a strong delirium risk factor and showed that incorporating ε4

status into prediction models improved incident delirium prediction [39]. The study

identified a further four risk loci for delirium which we did not replicate in our

postoperative delirium cohort, likely due to differences in statistical power and

analytical approach. We did, however, find that *APOE*-independent PRS for

Alzheimer's disease were associated with postoperative delirium, mirroring the

smaller contribution made by other genomic regions. A role for APOE-related

pathways in postoperative neurocognitive outcomes has biological plausibility

through effects on neuroinflammation and blood-brain barrier integrity [40]. In the

perioperative setting, *APOE* ε4 carriers have shown greater postoperative decreases

in functional connectivity in key Alzheimer's disease risk regions [41]. Further

validation of these findings may lead to potential clinical applications including

preoperative risk stratification based on associated variants or existing polygenic risk

scores (e.g. for Alzheimer's disease) and the identification of potential therapeutic

targets for the reduction of postoperative delirium risk. Alternatively, the acute

occurrence of postoperative delirium might represent an unmasking of underlying

neurocognitive vulnerability or an undiagnosed chronic condition, e.g. pre-clinical

Alzheimer's disease. If genetic liability for Alzheimer's disease extends to

postoperative delirium, patients found to be at increased risk of delirium may benefit

from additional monitoring long-term.

The strengths of this study include the use of a large prospective cohort with linked

healthcare data that is subject to rigorous quality control and validation procedures.

We defined our exposure using published criteria and employed robust statistical

methods to deal with the large case-control imbalance. We were also able to extend

our findings through a range of post-GWAS analyses using an independent

validation cohort. The study does have some limitations. The diagnosis of delirium

relied upon ICD-10 clinical coding and was not prospectively screened for. Validation

studies have shown that ICD-10 coding in administrative databases has low

sensitivity but excellent specificity for postoperative delirium compared to prospective

clinical screening [42,43]. As such there will be some misclassification bias,

particularly true cases which are included in the control group. This will likely have

attenuated the observed genetic effects and reduced our statistical power to detect

small or modest effects. Whilst is also possible that some cases of delirium preceded

surgery, the risk of this was minimised as far as possible. It is also possible that

some cases in the 30-day follow-up period were related to other factors, for example

infection or hospitalisation, but these occurred in the context of recovery from major

surgery so were included in line with recommendations on the nomenclature of

postoperative cognitive changes. Whilst UK Biobank represents a large cohort of

patients there is a recognised "healthy volunteer" selection bias [44] and so the

results may not be entirely generalisable to the wider UK population. Any secondary

data analysis relying on routinely-collected administrative data is limited by the fields

available, with some potentially relevant factors such as the American Society of

Anesthesiologists (ASA) physical status grade, surgical urgency and type of

anaesthesia not recorded. Finally, whilst rates of some comorbidities and surgical

characteristics differed between cases and controls, these variables may lie on the

causal pathway or act as selection-related colliders in a major surgical cohort. We

therefore did not adjust for them in the primary GWAS but examined their influence

through a range of sensitivity analyses which support the robustness of our findings.

In conclusion, we have used a large national biobank with linked healthcare data to

identify seven SNPs and four genes associated with increased risk of postoperative

delirium. Data from an independent cohort suggests there is shared genetic

architecture and risk between postoperative delirium, delirium more broadly, and

Alzheimer's disease. If validated, these findings have potential clinical benefit

through risk stratification based on genotype or polygenic risk score data and early

identification of future Alzheimer's disease risk in the perioperative setting.

Authors' contributions

RAA: study design, data analysis, preparing first and subsequent drafts of the paper;

PY: study design, drafting paper; BG: study design, drafting paper; GK: study design,

drafting paper; TG: study design, drafting paper.

Acknowledgements

This research has been conducted using the UK Biobank Resource under Application Number 128619. This includes linked data provided by patients and collected by the NHS as part of their care and support and data assets made available by National Safe Haven as part of the Data and Connectivity National Core Study, led by Health Data Research UK in partnership with the Office for National Statistics and funded by UK Research and Innovation (grant ref MC_PC_20058).

Quality Control filtering of the UK Biobank data used the process described by R.Mitchell, G.Hemani, T.Dudding, L.Corbin, S.Harrison, L.Paternoster[25].

This work was carried out using the computational and data storage facilities of the Advanced Computing Research Centre, University of Bristol (http://www.bristol.ac.uk/acrc/).

Competing interests

I have read the journal's policy and the authors of this manuscript have the following competing interests: TRG receives funding from GlaxoSmithKline, Biogen and Roche for unrelated research. The other authors have declared that no competing interests exist.

Financial disclosure

This study was funded by a Wellcome Trust GW4-CAT PhD Programme for Health Professionals PhD Fellowship awarded to RAA [316275/Z/24/Z]. RAA, PY, GMK and TRG are supported by the Medical Research Council Integrative Epidemiology Unit at the University of Bristol (RA, TG: MC_UU_00032/3; PY: MC_UU_00032/4; GMK: MC_UU_0032/6). GMK acknowledges additional funding from the Wellcome Trust (grant numbers: 201486/Z/16/Z and 201486/B/16/Z), the Medical Research Council (grant numbers: MR/W014416/1; MR/S037675/1; MR/Z50354X/1; and MR/Z503745/1. PY, TRG, and GMK are also supported by the UK National Institute for Health and Care Research (NIHR) Bristol Biomedical Research Centre (grant number: NIHR 203315). The views expressed are those of the authors and not necessarily those of the UK NIHR or the Department of Health and Social Care. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Evered L, Atkins K, Silbert B, Scott DA. Acute peri-operative neurocognitive disorders: a narrative review. Anaesthesia. 2022;77: 34–42. doi:10.1111/anae.15613

2. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. The Lancet. 2014;383: 911–922. doi:10.1016/S0140-6736(13)60688-1

3. Gou RY, Hshieh TT, Marcantonio ER, Cooper Z, Jones RN, Travison TG, et al. One-Year Medicare Costs Associated With Delirium in Older Patients Undergoing Major Elective Surgery. JAMA Surg. 2021;156: 462–470. doi:10.1001/jamasurg.2020.7260

4. MacLullich AMJ, Ferguson KJ, Miller T, de Rooij SEJA, Cunningham C. Unravelling the pathophysiology of delirium: A focus on the role of aberrant stress responses. J Psychosom Res. 2008;65: 229–238. doi:10.1016/j.jpsychores.2008.05.019

5. Cerejeira J, Firmino H, Vaz-Serra A, Mukaetova-Ladinska EB. The neuroinflammatory hypothesis of delirium. Acta Neuropathol (Berl). 2010;119: 737–754. doi:10.1007/s00401-010-0674-1

6. Rump K, Adamzik M. Epigenetic Mechanisms of Postoperative Cognitive Impairment Induced by Anesthesia and Neuroinflammation. Cells. 2022;11: 2954. doi:10.3390/cells11192954

7. Swarbrick CJ, Partridge JSL. Evidence-based strategies to reduce the incidence of postoperative delirium: a narrative review. Anaesthesia. 2022;77: 92–101. doi:10.1111/anae.15607

8. Schenning KJ, Deiner SG. Postoperative Delirium in the Geriatric Patient. Anesthesiol Clin. 2015;33: 505–516. doi:10.1016/j.anclin.2015.05.007

9. Nadelson MR, Sanders RD, Avidan MS. Perioperative cognitive trajectory in adults. Br J Anaesth. 2014;112: 440–451. doi:10.1093/bja/aet420

10. Goldberg TE, Chen C, Wang Y, Jung E, Swanson A, Ing C, et al. Association of Delirium With Long-term Cognitive Decline: A Meta-analysis. JAMA Neurol. 2020;77: 1373–1381. doi:10.1001/jamaneurol.2020.2273

11. Peden CJ, Miller TR, Deiner SG, Eckenhoff RG, Fleisher LA, Fleisher L, et al. Improving perioperative brain health: an expert consensus review of key actions for the perioperative care team. Br J Anaesth. 2021;126: 423–432. doi:10.1016/j.bja.2020.10.037

12. Abdellaoui A, Yengo L, Verweij KJH, Visscher PM. 15 years of GWAS discovery: Realizing the promise. Am J Hum Genet. 2023;110: 179–194. doi:10.1016/j.ajhg.2022.12.011

13. Rickenbacher M, Reinbold CS, Herms S, Hoffmann P, Cichon S, Wueest AS, et al. Genome-wide Association Study of Postoperative Cognitive Dysfunction in Older Surgical Patients. J Neurosurg Anesthesiol. 2022;34: 248. doi:10.1097/ANA.0000000000000749

14. Heinrich M, Sieg M, Kruppa J, Nürnberg P, Schreier PH, Heilmann-Heimbach S, et al. Association between genetic variants of the cholinergic system and postoperative delirium and cognitive dysfunction in elderly patients. BMC Med Genomics. 2021;14: 248. doi:10.1186/s12920-021-01071-1

15. Westphal S, Stoppe C, Gruenewald M, Bein B, Renner J, Cremer J, et al. Genome-wide association study of myocardial infarction, atrial fibrillation, acute stroke, acute kidney injury and delirium after cardiac surgery - a sub-analysis of the RIPHeart-Study. BMC Cardiovasc Disord. 2019;19: 26. doi:10.1186/s12872-019-1002-x

16. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank resource with deep phenotyping and genomic data. Nature. 2018;562: 203–209. doi:10.1038/s41586-018-0579-z

17. Kurki MI, Karjalainen J, Palta P, Sipilä TP, Kristiansson K, Donner KM, et al. FinnGen provides genetic insights from a well-phenotyped isolated population. Nature. 2023;613: 508–518. doi:10.1038/s41586-022-05473-8

18. FinnGen. Genetic data. FinnGen. [cited 1 September 2025]. Available from: https://www.finngen.fi/en/genetic_data

19. UK Biobank. Data providers and dates of data availability. UK Biobank. [cited 1 September 2025]. Available from: https://biobank.ndph.ox.ac.uk/showcase/exinfo.cgi?src=Data_providers_and_dates

20. Abbott TEF, Fowler AJ, Dobbs TD, Harrison EM, Gillies MA, Pearse RM. Frequency of surgical treatment and related hospital procedures in the UK: a national ecological study using hospital episode statistics. BJA Br J Anaesth. 2017;119: 249–257. doi:10.1093/bja/aex137

21. Bupa. Schedule of procedures. Bupa. [cited 1 September 2025]. Available from: https://codes.bupa.co.uk/procedures

22. Evered L, Silbert B, Knopman DS, Scott DA, DeKosky ST, Rasmussen LS, et al. Recommendations for the nomenclature of cognitive change associated with anaesthesia and surgery—2018. Br J Anaesth. 2018;121: 1005–1012. doi:10.1016/j.bja.2017.11.087

23. American College of Surgeons National Surgical Quality Improvement Programme. About the ACS risk calculator. American College of Surgeons. [cited 7 Jan 2026]. Available from: https://riskcalculator.facs.org/RiskCalculator/about.html

24. Mbatchou J, Barnard L, Backman J, Marcketta A, Kosmicki JA, Ziyatdinov A, et al. Computationally efficient whole-genome regression for quantitative and

binary traits. Nat Genet. 2021;53: 1097–1103. doi:10.1038/s41588-021-00870-7

25. Mitchell R, Hemani G, Dudding T, Corbin L, Harrison S, Paternoster L. UK Biobank genetic data: MRC-IEU quality control, version 2. In: data.bris [Internet]. 22 Jan 2019 [cited 1 September 2025]. doi:10.5523/bris.1ovaau5sxunp2cv8rcy88688v

26. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. Nat Commun. 2017;8: 1826. doi:10.1038/s41467-017-01261-5

27. Leeuw CA de, Mooij JM, Heskes T, Posthuma D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. PLOS Comput Biol. 2015;11: e1004219. doi:10.1371/journal.pcbi.1004219

28. Gleason LJ, Schmitt EM, Kosar CM, Tabloski P, Saczynski JS, Robinson T, et al. Effect of Delirium and Other Major Complications on Outcomes After Elective Surgery in Older Adults. JAMA Surg. 2015;150: 1134–1140. doi:10.1001/jamasurg.2015.2606

29. Werme J, van der Sluis S, Posthuma D, de Leeuw CA. An integrated framework for local genetic correlation analysis. Nat Genet. 2022;54: 274–282. doi:10.1038/s41588-022-01017-y

30. Lambert SA, Wingfield B, Gibson JT, Gil L, Ramachandran S, Yvon F, et al. Enhancing the Polygenic Score Catalog with tools for score calculation and ancestry normalization. Nat Genet. 2024; 1–6. doi:10.1038/s41588-024-01937-x

31. Ware EB, Faul JD, Mitchell CM, Bakulski KM. Considering the APOE locus in Alzheimer's disease polygenic scores in the Health and Retirement Study: a longitudinal panel study. BMC Med Genomics. 2020;13: 164. doi:10.1186/s12920-020-00815-9

32. Menon DK, Rosand J. Finding a Place for Candidate Gene Studies in a Genome-Wide Association Study World. JAMA Netw Open. 2021;4: e2118594. doi:10.1001/jamanetworkopen.2021.18594

33. Thedim M, Hu J, Maher M, Wiener-Kronish J, Saxena R, Vacas S. Perioperative polygenic and APOE-based genetic risk assessment for neurocognitive disorders: a biobank study. Br J Anaesth. 2025;0. doi:10.1016/j.bja.2025.05.014

34. Denny JC, Bastarache L, Ritchie MD, Carroll RJ, Zink R, Mosley JD, et al. Systematic comparison of phenome-wide association study of electronic medical record data and genome-wide association study data. Nat Biotechnol. 2013;31: 1102–1111. doi:10.1038/nbt.2749

35. Abelha FJ, Fernandes V, Botelho M, Santos P, Santos A, Machado JC, et al. Apolipoprotein E e4 allele does not increase the risk of early postoperative

delirium after major surgery. J Anesth. 2012;26: 412–421. doi:10.1007/s00540-012-1326-5

36. Vasunilashorn S, Ngo L, Kosar CM, Fong TG, Jones RN, Inouye SK, et al. Does Apolipoprotein E Genotype Increase Risk of Postoperative Delirium? Am J Geriatr Psychiatry Off J Am Assoc Geriatr Psychiatry. 2015;23: 1029. doi:10.1016/j.jagp.2014.12.192

37. Tagarakis GI, Tsolaki-Tagaraki F, Tsolaki M, Diegeler A, Tsilimingas NB, Papassotiropoulos A. The Role of Apolipoprotein E in Cognitive Decline and Delirium after Bypass Heart Operations. Am J Alzheimers Dis Dementias®. 2007;22: 223–228. doi:10.1177/1533317507299415

38. Leung JM, Sands LP, Wang Y, Poon A, Kwok P, Kane JP, et al. Apolipoprotein E e4 Allele Increases the Risk of Early Postoperative Delirium in Older Patients Undergoing Noncardiac Surgery. Anesthesiology. 2007;107: 406–411. doi:10.1097/01.anes.0000278905.07899.df

39. Raptis V, Bhak Y, Cannings TI, MacLullich AMJ, Tenesa A. Dissecting the genetic and proteomic risk factors for delirium. Nat Aging. 2025; 1–17. doi:10.1038/s43587-025-01018-6

40. Yang LG, March ZM, Stephenson RA, Narayan PS. Apolipoprotein E in lipid metabolism and neurodegenerative disease. Trends Endocrinol Metab. 2023;34: 430–445. doi:10.1016/j.tem.2023.05.002

41. Browndyke JN, Wright MC, Yang R, Syed A, Park J, Hall A, et al. Perioperative neurocognitive and functional neuroimaging trajectories in older *APOE4* carriers compared with non-carriers: secondary analysis of a prospective cohort study. Br J Anaesth. 2021;127: 917–928. doi:10.1016/j.bja.2021.08.012

42. Katznelson R, Djaiani G, Tait G, Wasowicz M, Sutherland AM, Styra R, et al. Hospital administrative database underestimates delirium rate after cardiac surgery. Can J Anesth Can Anesth. 2010;57: 898–902. doi:10.1007/s12630-010-9355-8

43. Kim DH, Lee J, Kim CA, Huybrechts KF, Bateman BT, Patorno E, et al. Evaluation of algorithms to identify delirium in administrative claims and drug utilization database. Pharmacoepidemiol Drug Saf. 2017;26: 945–953. doi:10.1002/pds.4226

44. Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, et al. Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. Am J Epidemiol. 2017;186: 1026–1034. doi:10.1093/aje/kwx246

Figure legends

Figure 1 Study flow diagram.

Figure 2 Manhattan plot of case-control GWAS of postoperative delirium results (adjusted for age, sex, chip and first ten principal components). The red line represents a genome-wide significant p value of 5 x 10^{-8}.

Figure 3 Q-Q plot of case-control GWAS of postoperative delirium results (primary analysis). Genomic inflation factor (λ) 1.01.

Figure 4 Manhattan plot for MAGMA gene testing from case-control GWAS of postoperative delirium (adjusted for age, sex, chip and first ten principal components). The red line represents a genome-wide significant p value of 2.5 x 10^{-6}.

Figure 5 Q-Q plot for MAGMA gene testing from case-control GWAS of postoperative delirium results (primary analysis). Genomic inflation factor (λ) 1.0.

Supporting Information files and legends

S1 Checklist. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist

S2 Appendix. Supplementary methods

S3 Appendix. Tables and figures

Table A Additional participant characteristics for case-control GWAS of postoperative delirium. Comorbidities assessed at time of surgery. Operative categories are listed as Abbott/Bupa classification.

Table B Mapped genes from primary case-control GWAS of postoperative delirium. minGwasP represents the minimum p-value associated with the gene; IndSigSNPs lists the independent significant SNPs tagged to that gene.

Table C Independent significant SNPs in non-cardiothoracic surgery case-control GWAS of postoperative delirium (sensitivity analysis). rsID: Reference SNP cluster ID.

Table D Top 10 SNPs from case-control GWAS of postoperative delirium in cardiothoracic cohort (sensitivity analysis). rsID: Reference SNP cluster ID.

Table E Independent significant single-nucleotide polymorphisms passing genome-wide significance in case-control GWAS of postoperative delirium after adjustment for age, sex, chip, first 10 PCs and Charlson Comorbidity Index (sensitivity analysis). rsID: Reference SNP cluster ID.

Table F Polygenic risk scores from PGS Catalog passing matching and included in polygenic risk score analyses. *APOE-independent scores.

Table G Odds ratio for postoperative delirium by Alzheimer's disease polygenic risk score (PRS) quintile. Results are shown for all 42 polygenic risk scores, adjusted for age and sex, and the subset which did not include the APOE region (15 APOE-independent scores), adjusted for age, sex, APOE ε4 genotype and Charlson Comorbidity Index (0-1 vs >=2).

Table H Results from primary analysis (case-control GWAS of postoperative delirium) for variants identified in candidate gene study by Heinrich et al. rsID: Reference SNP cluster ID.

Figure A Manhattan plot for case-control GWAS of postoperative delirium after previous dementia diagnoses excluded. The red line represents a genome-wide significant p value of 5×10^{-8}.

Figure B Manhattan plot for MAGMA testing for case-control GWAS of postoperative delirium after previous dementia diagnoses excluded.

Figure C Manhattan plot for case-control GWAS of postoperative delirium after previous or subsequent dementia diagnoses excluded. The red line represents a genome-wide significant p value of 5×10^{-8}.

Figure D Manhattan plot for MAGMA testing for case-control GWAS of postoperative delirium after previous or subsequent dementia diagnoses excluded.

Figure E Manhattan plot for case-control GWAS of postoperative delirium conditional on APOE *ε4* allele count (adjusted for age, sex, chip, first 10 PCs).

Figure F GWAS catalog reported genes for case-control GWAS of postoperative

delirium (primary analysis).