1
2
3
4
5 # A meta-analysis suggests that TMS targeting the hippocampal network
6 # selectively improves episodic memory
7
8

9 **Authors:**
10 Elena Badillo Goicoechea[1†], Phillip F. Agres[2†], Johanna M.H. Rau[2], Arantzazu San Agustín[2], Joel L. Voss[2*]
11
12 **Affiliations:**
13 [1]Department of Public Health Sciences, The University of Chicago; Chicago, IL
14 [2]Department of Neurology, The University of Chicago; Chicago, IL
15 *Corresponding author. Email: joelvoss@uchicago.edu
16 [†]These authors contributed equally to this work.

17
18
19
20 **Impact statement:**
21 Meta-analysis indicates that network-targeted noninvasive brain stimulation consistently enhances memory
22 function supported by the hippocampal network, thus providing robust evidence that specific memory abilities
23 rely on specific modifiable brain networks.
24
25

Abstract:
Episodic memory is critically dependent on the hippocampal network and is frequently impaired in many clinical
disorders. Recent findings highlight Hippocampal Indirectly Targeted Stimulation (HITS) as a promising,
noninvasive transcranial magnetic stimulation (TMS) intervention to enhance episodic memory performance.
Here, we report the first comprehensive meta-analysis of HITS effects on episodic memory, encompassing
both healthy individuals and clinical populations. HITS robustly improved episodic memory, with effects
selective for episodic memory versus other non-memory cognitive domains. Efficacy was significantly greater
when memory performance was assessed using memory tasks sensitive to recollection, which is strongly
linked to hippocampal network function, compared to recognition or other types of episodic memory tasks.
Efficacy was also significantly greater when HITS was delivered before versus after the period of memory
encoding. No serious adverse events were reported. These findings establish HITS as a safe and effective
approach for episodic memory enhancement, with potential for clinical translation in memory disorders.
Selectivity of effects for episodic memory generally and for recollection-format tests in particular indicates
cognitive and mechanistic specificity, supporting the potential for targeted and selective neuromodulation of
hippocampal networks and their associated functions.

INTRODUCTION

Memory for episodes of experience (episodic memory) depends critically on the hippocampus and interactions of hippocampus with a set of brain regions that comprise a hippocampal-centered network [1-8]. Episodic memory impairments due to a variety of neurological and psychiatric conditions have been associated with disrupted functional connectivity of hippocampal-centered networks, as measured via methods such as functional MRI [3, 4, 6, 9-14]. Thus, to the extent that memory impairments result from disrupted network function, modulation of hippocampal network functional connectivity is a potential therapeutic strategy for episodic memory rescue.

Transcranial magnetic stimulation (TMS) induces electrical fields in the human brain with sufficient intensity to trigger focal action potential firing in neocortex [15]. Repetitive TMS (rTMS) protocols influence neuroplasticity of distributed networks of the stimulated location, which can influence functional connectivity [16-21]. This motivates the hypothesis that rTMS of hippocampal network locations could affect network functional connectivity and thereby influence episodic memory. Indeed, Wang et al. [8] applied multi-day rTMS to a parietal-cortex location identified via its high fMRI connectivity with the hippocampus. They reported that stimulation increased fMRI connectivity of the indirectly targeted portion of hippocampus with its network and improved episodic memory performance. We refer to this stimulation method as "Hippocampal Indirectly Targeted Stimulation" (HITS), as the overall goal is to use noninvasive rTMS to indirectly affect hippocampal interactions with its network. It is possible that HITS could be used to enhance memory function, both in healthy individuals and those with clinical memory impairments resulting from abnormal hippocampal network connectivity [22-24]. However, the variety of neuromodulation protocols used by various research groups and variability in reported effectiveness of HITS [25-28] motivate systematic and quantitative evaluation of this potential.

Here, we report results from a systematic review and meta-analysis of studies that investigated the effects of HITS on episodic memory. We reviewed studies in which rTMS was applied to locations of the hippocampal network (**Figure 1**) and that assessed effects of stimulation on objective tests of episodic memory, conducted in healthy young and older adults and in clinical populations with memory impairments. Our main analysis goal was to determine whether HITS affects episodic memory reliably across studies, despite variation in study designs. We also tested whether the effects of HITS were greater on episodic memory than on other cognitive functions measured in the same studies, and whether study design factors modulated the effectiveness of HITS. We focused on hypothesis-driven design factors, including task format, targeting method, and study population, as well as eight factors identified *post hoc* as varying across studies (Table S1).

RESULTS

Sample characteristics

Our search identified 47 studies meeting inclusion criteria, 38 of which provided sufficient data to support meta-analysis (Table 1). These 38 studies included N = 1,009 subjects and reported 253 statistical comparisons (i.e., "effects") of the influence of HITS on performance of episodic-memory (140 effects) and non-memory tasks (113 effects), which we transformed into normalized effects (Hedges' *g*). The mean sample size was n = 23.2 for episodic-memory effects (range = 4 – 68) and n = 23.3 for non-memory effects (range = 8 – 58) (Figure S1). By definition, stimulation was applied to locations of the hippocampal network that were predominantly in left lateral parietal cortex (84% of studies) or in precuneus or other parieto-occipital neocortex locations generally considered as part of the hippocampal network (all remaining studies) (Figure 1).

HITS improved episodic memory performance

A meta-regression model with a random effect by study was used to evaluate the overall effect of HITS on episodic memory performance for the 140 effects derived from episodic-memory tasks. The overall effect was positive and highly significant, indicating that HITS improved episodic memory (Hedges' *g* = 0.44; 95% CI [0.34, 0.54]; $p < 0.001$; Figure 2A). A similar result was obtained when outliers were removed in a sensitivity analysis (Hedges' *g* = 0.38; 95% CI [0.29, 0.46]; $p < 0.001$). Both the main analysis and sensitivity analysis showing lower bounds well above conventional thresholds for small effects ($g \geq 0.2$) indicates robust evidence(*29*) for meaningful improvement in episodic memory performance due to HITS.

Effects of HITS were greater on episodic-memory than non-memory tasks

Meta-regression was used to separately evaluate the overall effect of HITS on performance for 113 effects that concerned tasks assessing cognitive functions other than episodic memory. These included tasks designed to assess attention, working memory, executive functions, and language. The overall effect was almost zero and not significant, indicating that HITS did not reliably affect performance in these tasks (Hedges' *g* = 0.04, 95% CI [-0.01, 0.09], $p = 0.12$; Figure 2B). A similar result was obtained when outliers were removed in a sensitivity analysis (g = 0.01, 95% CI [-0.05 0.06]; $p = 0.79$).

To more rigorously assess whether effects of HITS for tasks that measured episodic memory were significantly greater than for non-memory tasks, we ran a model pooling all 253 memory and non-memory effects and assessing overall moderation by task type. Effects for non-memory tasks were significantly less than for episodic-memory tasks (Hedges' *g* modification = -0.39; 95% CI [-0.48, -0.29]; $p < 0.001$). A similar result was obtained when outliers were removed in a sensitivity analysis (Hedges' *g* modification = -0.34; 95% CI [-0.43 - 0.25]; $p < 0.001$). Thus, the beneficial effects of HITS were significantly greater for tasks measuring episodic memory than other cognitive domains in both the main and sensitivity analyses.

Study factors modulated HITS effects on episodic memory

We categorized episodic-memory effects according to study design factors, based on *a priori* hypotheses as well as *post hoc* considerations of variability in study designs (Table S1). We evaluated which of these factors had the potential for independent modulation of HITS effects on episodic memory via cross-validated lasso regression selection with permutation. Factors that were retained in greater than 50% of the lasso iterations were included in the meta-regression. This led to the exclusion of the TMS Protocol factor, likely because there was strong redundancy of this factor with the Intensity and Sessions factors (Figure S2). As expected, the 10 retained factors were well represented among analyzed effects (Figure 3). In the sensitivity analysis with outlier removal, TMS Protocol was also discarded, as well as the Task Delay factor. Three of the retained factors significantly modulated the effects of HITS on episodic memory performance (Figure 3A), as described next.

Consistent with our *a priori* hypothesis, effects for tasks that measured episodic memory using recollection test formats were significantly greater than for those that used recognition formats (Hedges' *g* modification = 0.22

greater for recollection versus recognition; 95% CI [0.06, 0.37]; p = 0.006; Figure 3A). In contrast, effects for tests that used other episodic memory (non-recollection and non-recognition) formats were not significantly different from recognition (Hedges' g modification = 0.07 less for other-format versus recognition; 95% CI [-0.33, 0.20]; p = 0.62; Figure 3A). The same pattern held in the sensitivity analysis with outliers removed (Hedges' g modification = 0.19 greater for recollection than recognition; 95% CI [0.54, 0.33]; p = 0.006; Hedges' g = 0.02 less for other-format than recognition; 95% CI [-.25, 0.21]; p = 0.84). Thus, among tasks assessing episodic memory, the effects of HITS were significantly greater for recollection.

Regarding the Timing factor, HITS applied before the period of memory encoding (from days to seconds before) was associated with significantly greater effects on episodic memory than HITS applied immediately after the period of memory encoding (Hedges' g modification = 0.69 greater for pre-encoding versus post-encoding; 95% CI [0.34, 1.05]; p < 0.001; Figure 3A). The same pattern held in a sensitivity analysis with outliers removed (Hedge's g modification = 0.61 greater for pre-encoding versus post-encoding; 95% CI [0.33, 0.89]; p < 0.001). The average effect for studies in which HITS was applied post-encoding was nearly zero (Figure 3A), with these being the majority (52.4%) of all observed negative effects (i.e., effects reflecting impairment by HITS). In contrast, effects when HITS was applied pre-encoding were almost all positive (91.5% of pre-encoding effects were positive). Notably, studies in the post-encoding category all followed the same general experiment design, whereby a brief encoding period was immediately followed by a single session of stimulation which was then immediately followed by the test. In contrast, pre-encoding studies included a variety of designs, including those in which brief stimulation trains were given immediately before individual memoranda as well as those in which many days or weeks of stimulation were given for days to weeks before memory encoding (Table S1). Thus, among tasks assessing episodic memory, the effects of HITS were significantly greater when it was applied sometime before the period of memory encoding.

Regarding the Targeting factor, effects differed for studies that used individualized (i.e., subject-specific) measures of functional or structural MRI connectivity to determine the stimulation location for HITS compared to studies that used a similar target for all subjects based on an atlas or *a priori* hypothesis (Hedges' g modification = -0.32; 95% CI [-0.58, -0.07]; p = 0.013; Figure 3A). However, this effect was not significant in a sensitivity analysis with outliers removed (Hedges' g modification = -0.19; 95% CI [-0.38, 0.01]; p = 0.059). Thus, evidence for modification of HITS effects by targeting method was weak, not surviving sensitivity analysis.

None of the other study factors significantly modulated the effects of HITS in either the main or sensitivity analyses (Figure 3B).

Of the 140 effects concerning episodic memory, 17.9% involved both the two significant positive moderating study factors identified in the main and sensitivity analyses (recollection test format and application of HITS before encoding). These effects thus can be considered as reflecting optimized experiment designs. To estimate the expected effects for future studies incorporating both of these factors, we used our meta-regression model to project the expected effect sizes for studies incorporating these factors while accounting for variance associated with other factors, held constant. Their mean effect estimate was g = 0.66 (95% CI [0.38, 0.94]), in contrast to that of all other effects incorporating non-optimized factors of all other effects, g = 0.37 (95% CI [0.07, 0.68]) (Figure 4).

Similar outcomes in studies that were excluded from meta-analysis
Nine studies identified via systematic review were not included in the meta-analysis for reasons summarized in Supplementary Results. Excluded studies reported outcomes from 209 subjects and incorporated study design factors representative of those for included studies (Supplementary Results). Seven of the excluded studies reported results from statistical tests to assess the effects of HITS on episodic memory performance. Of those,

187   five reported statistically significant improvement due to HITS in at least one test of episodic memory and two
188   reported numeric improvements due to HITS that were not statistically significant (Supplementary Results).
189   Thus, excluded studies mostly reported episodic memory improvements due to HITS, with variability in
190   outcomes that is consistent with the variability seen in studies that were included in the meta-analysis. It is
191   therefore unlikely that meta-analysis results would vary significantly if it were possible to include these studies.
192
No serious adverse events were reported
None of the reviewed studies reported any unexpected or serious adverse events. The total sample reviewed
included N=1,238 subjects, including healthy younger and older adults, individuals with mild to moderate
Alzheimer's dementia, and individuals with psychiatric symptoms. This suggests that the incidence of serious
adverse events due to HITS is low, which is consistent with the general evidence for very low incidence of
serious adverse events due to rTMS (*30*).
199
200

DISCUSSION

Enhancement of episodic memory in humans is an important scientific objective with clear implications for the treatment of clinical memory disorders and for enhancing function in healthy individuals, including older adults experiencing normative memory decline. However, interventions for memory enhancement have been elusive. Our findings indicate HITS is a promising method to achieve memory enhancement via noninvasive stimulation. There were similarly positive effects among participant populations, from young healthy adults to those with memory disorders, and for a wide range of assessment delays, from seconds to weeks after subjects received HITS. In contrast, there was essentially no effect on non-memory functions, suggesting that memory enhancement did not come at the cost of any off-target cognitive detriment. There were no reported unexpected or serious adverse events. Findings suggest that HITS is safe and effective for targeted enhancement of episodic memory.

Many have speculated that brain stimulation targeting a given network can have specific effects on the cognitive abilities hypothesized to depend on that network (*17, 25, 26, 31, 32*). Although aspects of this hypothesis for HITS have been supported in isolated experiments (*8, 33, 34*), the current findings are notable in providing meta-analytic support. The hippocampus and its network are strongly implicated in episodic memory, and particularly in tests that that measure recollection compared to recognition (*35-37*). Indeed, we found that effects of HITS were specific to episodic-memory versus other cognitive tests, and were greater for recollection than recognition or other-format memory tests. Although previous individual experiments have shown that HITS affects activity of the hippocampal network in relation to recollection improvements more so than recognition (*23, 34, 38*), the current findings provide strong evidence that across many different experiment parameters, stimulation targeting the hippocampal network specifically affects the memory function thought to depend most heavily on this network.

Specificity of HITS effects on recollection is mechanistically informative. For instance, other interventions such as transcranial electrical stimulation yield nonspecific effects across a variety of cognitive domains (*39*). It is possible that such nonspecific effects could arise via modulation of general non-targeted factors, such as arousal via effects of stimulation on either peripheral or central nervous systems (*40, 41*). Such nonspecific influences could be problematic, as general and/or off-target effects could be ineffective or detrimental when applied to disorders affecting specific brain circuits. Along with neuroimaging evidence for selectivity (*8, 23, 33, 34, 42, 43*), the current findings suggest that HITS is suitable for targeted interventions of hippocampal network contributions to recollection. This could be useful in the treatment of memory disorders, which tend to disproportionately impact recollection relative to recognition (*36*). However, an important caveat is that the tests used as outcomes in the studies we analyzed were primarily designed to measure different cognitive abilities in isolation in the context of neuropsychological diagnosis. Our findings thus do not preclude off-target effects in specialized tasks that deliberately assess the interactions of brain networks. That is, enhancement of episodic memory could affect tests of other cognitive domains when those tests are at least partially sensitive to, or compete with, episodic memory (*44*).

It is notable that HITS had little to no effect in experiments that delivered a single session of HITS immediately after an encoding session and before the corresponding memory test (Figure 3A, HITS Timing factor of Post-Encoding). In contrast, effects were robustly positive when HITS was applied at some point pre-encoding, across a variety of potential stimulation-encoding delay intervals. This suggests that HITS improves memory formation rather than retention of recently learned information or memory retrieval. This is consistent with findings that hippocampus makes distinct contributions to learning versus retrieval (*45*). Regarding HITS as an intervention, it is notable that effects were similar across a variety of delays between when HITS was applied and when memory encoding occurred (Figure 3B, Task Delay factor). Unlike potential interventions that require delivery of stimulation concurrent with cognitive performance (e.g., transcranial electrical stimulation) or with highly specific information about the timing or successfulness of specific memory events (e.g., closed-loop

deep-brain stimulation), this finding suggests that HITS does not require specific knowledge of when memory demands occur following intervention. This is of practical utility, as it is difficult to know *a priori* when memory encoding will be required during ongoing activities of daily living.

It is important to avoid over-interpretation of null findings concerning moderating effects of some study factors (Figure 3B), as many were correlated (Figure S2) and were selected in individual experiments based on *a priori* hypotheses. For instance, based on evidence that stimulation for more consecutive sessions of HITS leads to greater or more persistent effects (*46*), studies that included more HITS sessions also tended to measure outcomes after longer delays (correlation of Sessions and Task Delay factors; Figure S2). Thus, it is possible that studies with more stimulation sessions would have yielded greater effects had they used shorter testing delays, as in the studies with fewer stimulation sessions. Likewise, the Intensity factor was complicated in that studies using lower intensity were more likely to involve theta-burst stimulation than studies using higher intensity, which typically used high-frequency rTMS. These studies were based on hypotheses that hippocampal networks should respond to theta-burst stimulation more than high-frequency stimulation (*25, 43*). But, theta-burst stimulation typically used lower intensity and also fewer pulses of stimulation than high-frequency rTMS. Thus, no differences in outcome based on the Intensity factor thus could support the hypothesis that theta-burst stimulation is more effective than rTMS, as similar effect sizes resulted with fewer and less-intense stimulation pulses. Interpretation of null effect modulation by factors that are highly correlated with other factors is thus ambiguous. Although it was useful to identify those factors that robustly modulated effects of HITS via meta-analysis (Figure 4), future experimental studies will be needed to systematically assess whether effects of HITS vary by study design (Figure 3B), based on specific manipulation of individual factors while holding other factors constant.

A limitation of our approach is that we focused on studies applying HITS to lateral and medial parieto-occipital neocortex targets of the hippocampal network. Other prominent locations of this network include medial frontal and lateral temporal neocortex (*47*), although no studies targeting these locations were identified by our review, potentially because conventional TMS of these locations is not well tolerated. Notably, the hippocampal network includes portions of lateral cerebellar cortex (*48*), and one study identified by our review found episodic memory improvement following theta-burst TMS of this location (*49*). Theta-burst stimulation of lateral cerebellum has also been shown to affect hippocampal network fMRI connectivity (*50*). However, this study was excluded from meta-analysis to avoid heterogeneity in stimulation locations. Future work could systematically evaluate alternative stimulation targets within the hippocampal network, including those that may require stimulation methods other than TMS.

The current findings suggest that HITS has robust effects on episodic memory and that studies including optimized parameters yield especially robust effects (Figure 4). However, further research is needed to test clinical efficacy for specific conditions and to determine mechanisms of action. Although our review included studies of patients with episodic memory impairments due to mild to moderate dementia and psychiatric symptoms, the studies did not assess whether gains by HITS could be maintained for months to years. Further, it is not clear whether HITS will be useful for memory impairments in other disorders, which may involve distinct mechanisms. Implementations for specific disorders will likely require better understanding of how HITS affects neural function, such as could be obtained by directly recording hippocampal activity while HITS is applied (*43, 51*), in order to tailor HITS to target disorder-specific neural impairments. The current findings of HITS efficacy, specificity, and safety encourage such future explorations.

MATERIALS AND METHODS

Eligibility criteria

We included reports published between September 2014 (date of the seminal HITS publication (*8*)) and April 2025 that: (1) used rTMS to stimulate hippocampal network locations and (2) included at least one objectively scored test of episodic memory as an outcome. Studies were considered as stimulating hippocampal network locations if they delivered rTMS to: (1) an individualized/subject-specific location defined by fMRI connectivity analysis (e.g., a neocortical location identified by resting-state fMRI connectivity to hippocampus in an individual), (2) a location based on a group metric of hippocampal fMRI connectivity (e.g., a neocortical location identified in a group-based atlas of hippocampal resting-state fMRI connectivity), and (3) a similar location based on a hypothesis (e.g., a location that is in a group atlas of hippocampal connectivity but that was selected to test a hypothesis about its function). We limited spatial variability among studies by including only those that stimulated parietal and superior occipital neocortex. We excluded studies using other potential hippocampal network locations due to their rarity, including one study with cerebellar cortex stimulation (*49*). Studies were considered as including an objective test of episodic memory if at least one of the reported outcomes used a memory test that allowed unambiguous computation of the effect of stimulation on performance accuracy (i.e., experimenter-controlled stimulus set presented to subjects for encoding, for which memory was subsequently tested after some retention interval).

Search strategy

Two publicly available online scientific databases were searched: PubMed (pubmed.ncbi.nlm.nih.gov) and Scopus (scopus.com). These databases were most recently searched on April 26, 2025. The primary search was conducted with PubMed, using the PubMed Advanced Search Builder. Two search terms of interest were entered: "memory" and "transcranial magnetic stimulation". These terms were selected to be as broad as possible while aiming to exclude studies utilizing stimulation techniques (invasive and non-invasive) other than TMS. These terms were added to the Query Box using 'AND' Boolean operators; all possible fields (i.e., title, abstract) were searched. PubMed automatically included a search for similar terms for each of the terms of interest entered (i.e., 'memory' and 'memories'), and included the Medical Subject Headings (MeSH) terms associated with them. This search returned 104 results. A secondary search was conducted using Scopus after completing the PubMed search, using the same primary search terms using 'AND' Boolean operators and constraining fields to Article title, Abstract, and Keywords and limiting results to peer-reviewed empirical reports by limiting the "Document Type" option to "Article". This search returned 160 results. We also examined the references of publications identified as meeting the eligibility criteria for suggestions about other relevant publications. Publications were selected from the search results when two reviewers (PA and JLV) agreed that they met the two eligibility criteria. Additionally, only studies in human participants were selected.

Outcomes extraction

Following review of the full texts, we extracted data from statistical tests aiming to quantify the effects of rTMS on episodic memory and non-memory outcomes. Extracted data included sample sizes, means, and standard deviations of the outcome variables for experimental and control measurements, as well as results from reported statistical tests. These data were used to calculate effect sizes, using the test statistics or group means and variances following established methods (*52*). Cohen's d was used to measure the effect size of pairwise comparisons. Partial eta squared was used to measure the effect size of comparisons using analysis of variance or other general linear models. We did not include effects from tests that provided "composite" measures spanning multiple cognitive domains (e.g., Montreal Cognitive Assessment, Mini-Mental State Examination, etc.) as these could not be readily classified as reflecting episodic-memory versus various non-memory functions. In tasks with multiple subscales (e.g., Wechsler Memory Scale, NIH Toolbox for Cognition, etc.), individual subscale scores were used and categorized based on the task format (e.g., recall versus recognition) and/or by the intended cognitive domain (e.g., episodic memory versus sustained attention). Effect sizes were computed such that improvement in performance (i.e., higher accuracy, faster response time in

343   speeded non-memory tasks, or fewer errors) were positive and impairments (i.e., lower accuracy, slower
344   responses, or more errors) were negative. Prior to input for meta-regression, all estimates of effect size were
345   converted to Hedges' *g* to minimize potential impact of inflated effect sizes associated with small sample sizes
346   (*52*). Within and between estimation uncertainty around each Hedge's *g* were approximated using its
347   corresponding study effective sample size.
348
Meta-analysis of effect size
A mixed effects univariate meta-regression model estimated via restricted maximum likelihood was used to
estimate the overall effect of rTMS on outcomes, pooling all effects reported in the final sample of studies
collected at the outcome level and weighting them in terms of their within and between estimation uncertainty.
We included a study random effect in the model to account both for inter and intra study variability in our final
estimate, as a single study often reported various effect sizes for different outcomes. Heterogeneity in
individual effect sizes was examined using a generalized Q-test, to assess whether variability in observed
effects exceeds what would be expected from random sampling error alone (*53, 54*).
357
Meta-analysis of effect modifiers
Factors that varied among studies were identified based on *a priori* hypotheses and *post hoc* consideration of
study design details that could influence outcomes (Table S1, Figure 3). We explored the association of effect
size with these suspected modifiers by estimating a multivariate meta-regression model including all study
factors. As with the main effects model, we pooled all effects reported in the final sample of studies collected at
the outcome level and weighted them in terms of their within and between study estimation uncertainty. Since
some study characteristics were *a priori* expected to be highly correlated with one another (e.g., theta-burst
stimulation is typically performed at lower intensity than high-frequency rTMS), we performed a previous-
variable-selection step before assessing effect modification. Indeed, some factors were highly correlated, as
expected (Figure S2). We therefore used a bootstrapped (1000 iterations) lasso model with a cross-validated
penalization term to identify study characteristics that were jointly consistently redundant in terms of statistically
explaining HITS effects (i.e., discarded by lasso with a probability larger than 50% across all iterations), to be
then excluded from the effect modification analysis. Study factors retained after the lasso variable selection
were then simultaneously included in the multivariate mixed effects meta-regression, where their association
with stimulation effects was estimated. Heterogeneity in individual effect sizes after accounting for effect
modifiers was examined using a generalized Q-test, assessing whether significant variability in observed
effects remained, even after accounting for the variation explained by included study characteristics (Q(df =
136) = 307.07, p < 0.001) . Analysis for this step was conducted using the *glmnet* package in R. All meta-
analytic estimates were computed using the *metafor* package in R (*54*).
377
Outlier removal
To guard against the impact that unusually strong reported effects could have had in our findings, we carried
out all meta-analyses, including the lasso variable selection step, after removing outliers identified using the
studentized deleted residuals (*55*). We identified and excluded seven effect outliers in the analysis of episodic
memory outcomes, leaving 133 of 140 effects for that analysis, and seven effect outliers in the analysis of non-
memory outcomes, leaving 106 of 113 effects for that analysis. Categorical variables were included in final
models if at least one level had a coefficient with magnitude ≥ |0.05| or their aggregate coefficients had
magnitude ≥ |0.01|. Variables with all coefficients < |0.01| were not included. We report findings with and
without removal of outliers, which had little impact.
387
Reporting bias assessment
Publication bias was assessed with a funnel plot of individual effect sizes against their standard errors (Figure
S3 and S4). Special attention was paid to the asymmetry of the plot and the absence of effects in the lower-left
corner of the funnel, which indicates the lack of published small-sample studies reporting negative effect sizes,

further confirmed by an Egger's regression test (p <0.001), which is commonly carried out to detect evidence

of publication bias (*56*). We tested for heterogeneity (I^2) to measure the inconsistency of effect sizes across

studies within each cognitive domain. The I^2 statistic represents the percentage of variation in effect sizes

beyond a random sampling error (*57*). Although the interpretation of heterogeneity can depend on multiple

factors, an I^2 of 25 to 75% is typically considered moderate, whereas $I^2 > 75\%$ is considered a substantial

heterogeneity (*57*). Further, we tested for the presence of outliers, and removed them in sensitivity analyses to

assess the robustness of our main findings to outliers. There was no conclusive evidence of publication bias,

with I^2 of 0.53 for episodic memory tests and of 0.0 for non-memory tests (Figure S3 and S4) and with

replication of major findings in the sensitivity analyses.

References

1. F. P. Battaglia, K. Benchenane, A. Sirota, C. M. A. Pennartz, S. I. Wiener, The hippocampus: hub of brain network communication for memory. *Trends in Cognitive Sciences*, S1364661311000891 (2011).
2. R. M. Braga, R. L. Buckner, Parallel Interdigitated Distributed Networks within the Individual Estimated by Intrinsic Functional Connectivity. *Neuron* **95**, 457-471 e455 (2017).
3. B. C. Dickerson, H. Eichenbaum, The episodic memory system: neurocircuitry and disorders. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **35**, 86-104 (2010).
4. M. A. Ferguson *et al.*, A human memory circuit derived from brain lesions causing amnesia. *Nat Commun* **10**, 3497 (2019).
5. D. R. King, M. de Chastelaine, R. L. Elward, T. H. Wang, M. D. Rugg, Recollection-related increases in functional connectivity predict individual differences in memory accuracy. *J Neurosci* **35**, 1763-1772 (2015).
6. M. Ritchey, L. A. Libby, C. Ranganath, Cortico-hippocampal systems involved in memory and cognition: the PMAT framework. *Prog Brain Res* **219**, 45-64 (2015).
7. W. B. Scoville, B. Milner, Loss of recent memory after bilateral hippocampal lesions. *J Neurol Neurosurg Psychiatry* **20**, 11-21 (1957).
8. J. X. Wang *et al.*, Targeted enhancement of cortical-hippocampal brain networks and associative memory. *Science* **345**, 1054-1057 (2014).
9. R. Jalilianhasanpour, E. Beheshtian, G. Sherbaf, S. Sahraian, H. I. Sair, Functional Connectivity in Neurodegenerative Disorders: Alzheimer's Disease and Frontotemporal Dementia. *Top Magn Reson Imaging* **28**, 317-324 (2019).
10. V. Tripathi *et al.*, Default Mode Network Functional Connectivity As a Transdiagnostic Biomarker of Cognitive Function. *Biol Psychiatry Cogn Neurosci Neuroimaging* **10**, 359-368 (2025).
11. M. D. Greicius, G. Srivastava, A. L. Reiss, V. Menon, Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc Natl Acad Sci U S A* **101**, 4637-4642 (2004).
12. A. G. Garrity *et al.*, Aberrant "default mode" functional connectivity in schizophrenia. *Am J Psychiatry* **164**, 450-457 (2007).
13. V. Ives-Deliperi, J. T. Butler, Mechanisms of cognitive impairment in temporal lobe epilepsy: A systematic review of resting-state functional connectivity studies. *Epilepsy Behav* **115**, 107686 (2021).
14. S. A. Joshi, E. R. Duval, B. Kubat, I. Liberzon, A review of hippocampal activation in post-traumatic stress disorder. *Psychophysiology* **57**, e13357 (2020).
15. M. C. Romero, M. Davare, M. Armendariz, P. Janssen, Neural effects of transcranial magnetic stimulation at the single-cell level. *Nat Commun* **10**, 2642 (2019).
16. A. Valero-Cabre, B. R. Payne, J. Rushmore, S. G. Lomber, A. Pascual-Leone, Impact of repetitive transcranial magnetic stimulation of the parietal cortex on metabolic brain activity: a 14C-2DG tracing study in the cat. *Exp Brain Res* **163**, 1-12 (2005).
17. M. D. Fox, M. A. Halko, M. C. Eldaief, A. Pascual-Leone, Measuring and manipulating brain connectivity with resting state functional connectivity magnetic resonance imaging (fcMRI) and transcranial magnetic stimulation (TMS). *Neuroimage* **62**, 2232-2243 (2012).
18. A. C. Chen *et al.*, Causal interactions between fronto-parietal central executive and default-mode networks in humans. *Proc Natl Acad Sci U S A* **110**, 19944-19949 (2013).
19. M. C. Eldaief, M. A. Halko, R. L. Buckner, A. Pascual-Leone, Transcranial magnetic stimulation modulates the brain's intrinsic activity in a frequency-dependent manner. *Proc Natl Acad Sci U S A* **108**, 21229-21234 (2011).
20. C. Gratton, T. G. Lee, E. M. Nomura, M. D'Esposito, The effect of theta-burst TMS on cognitive control networks measured with resting state fMRI. *Front Syst Neurosci* **7**, 124 (2013).
21. C. Liston *et al.*, Default mode network mechanisms of transcranial magnetic stimulation in depression. *Biol Psychiatry* **76**, 517-526 (2014).
22. Y. H. Jung *et al.*, Effectiveness of Personalized Hippocampal Network-Targeted Stimulation in Alzheimer Disease: A Randomized Clinical Trial. *JAMA Netw Open* **7**, e249220 (2024).
23. A. S. Nilakantan *et al.*, Network-targeted stimulation engages neurobehavioral hallmarks of age-related memory decline. *Neurology* **92**, e2349-e2354 (2019).

24. Y. Tang *et al.*, Visuospatial Learning Selectively Enhanced by Personalized Transcranial Magnetic
Stimulation over Parieto-Hippocampal Network among Patients at Clinical High-Risk for Psychosis.
Schizophr Bull **49**, 923-932 (2023).
25. M. Hebscher, J. L. Voss, Testing network properties of episodic memory using non-invasive brain
stimulation. *Curr Opin Behav Sci* **32**, 35-42 (2020).
26. R. F. H. Cash *et al.*, Personalized brain stimulation of memory networks. *Brain Stimulation* **15**, 1300-
1304 (2022).
27. M. V. Freedberg, J. A. Reeves, C. M. Fioriti, J. Murillo, E. M. Wassermann, Reproducing the effect of
hippocampal network-targeted transcranial magnetic stimulation on episodic memory. *Behavioural
Brain Research* **419**, 113707 (2022).
28. M. S. Hermiller, E. Karp, A. S. Nilakantan, J. L. Voss, Episodic memory improvements due to
noninvasive stimulation targeting the cortical-hippocampal network: A replication and extension
experiment. *Brain Behav* **9**, e01393 (2019).
29. G. M. Sullivan, R. Feinn, Using Effect Size-or Why the P Value Is Not Enough. *J Grad Med Educ* **4**,
279-282 (2012).
30. A. J. Lerner, E. M. Wassermann, D. I. Tamir, Seizures from transcranial magnetic stimulation 2012-
2016: Results of a survey of active laboratories and clinics. *Clin Neurophysiol* **130**, 1409-1416 (2019).
31. A. Pascual-Leone, V. Walsh, J. Rothwell, Transcranial magnetic stimulation in cognitive neuroscience--
virtual lesion, chronometry, and functional connectivity. *Curr Opin Neurobiol* **10**, 232-237 (2000).
32. R. Polanía, M. A. Nitsche, C. C. Ruff, Studying and modifying brain function with non-invasive brain
stimulation. *Nature Neuroscience* **21**, 174-187 (2018).
33. K. N. Warren, M. S. Hermiller, A. S. Nilakantan, J. L. Voss, Stimulating the hippocampal posterior-
medial network enhances task-dependent connectivity and memory. *Elife* **8**, (2019).
34. S. Kim *et al.*, Selective and coherent activity increases due to stimulation indicate functional distinctions
between episodic memory networks. *Sci Adv* **4**, eaar2768 (2018).
35. H. Eichenbaum, A. P. Yonelinas, C. Ranganath, The medial temporal lobe and recognition memory.
Annu Rev Neurosci **30**, 123-152 (2007).
36. A. P. Yonelinas, The nature of recollection and familiarity: A review of 30 years of research. *J Mem Lang*
46, 441-517 (2002).
37. A. J. Barnett *et al.*, Intrinsic connectivity reveals functionally distinct cortico-hippocampal networks in
the human brain. *PLoS Biol* **19**, e3001275 (2021).
38. A. S. Nilakantan, D. J. Bridge, E. P. Gagnon, S. A. VanHaerents, J. L. Voss, Stimulation of the Posterior
Cortical-Hippocampal Network Enhances Precision of Memory Recollection. *Current biology: CB* **27**,
465-470 (2017).
39. S. Grover, R. Fayzullina, B. M. Bullard, V. Levina, R. M. G. Reinhart, A meta-analysis suggests that
tACS improves cognition in healthy, aging, and psychiatric populations. *Sci Transl Med* **15**, eabo2044
(2023).
40. L. van Boekholdt, S. Kerstens, A. Khatoun, B. Asamoah, M. Mc Laughlin, tDCS peripheral nerve
stimulation: a neglected mode of action? *Mol Psychiatry* **26**, 456-461 (2021).
41. A. Majdi, B. Asamoah, M. Mc Laughlin, Reinterpreting published tDCS results in terms of a cranial and
cervical nerve co-stimulation mechanism. *Front Hum Neurosci* **17**, 1101490 (2023).
42. M. Freedberg *et al.*, Persistent Enhancement of Hippocampal Network Connectivity by Parietal rTMS Is
Reproducible. *eNeuro* **6**, (2019).
43. M. S. Hermiller, Y. F. Chen, T. B. Parrish, J. L. Voss, Evidence for Immediate Enhancement of
Hippocampal Memory Encoding by Network-Targeted Theta-Burst Stimulation during Concurrent fMRI.
J Neurosci **40**, 7155-7168 (2020).
44. M. V. Freedberg *et al.*, A direct test of competitive versus cooperative episodic-procedural network
dynamics in human memory. *Cereb Cortex* **32**, 4715-4732 (2022).
45. T. Gedankien *et al.*, Cholinergic blockade reveals role for human hippocampal theta in encoding but not
retrieval. *bioRxiv*, (2025).
46. M. Freedberg *et al.*, Optimizing Hippocampal-Cortical Network Modulation via Repetitive Transcranial
Magnetic Stimulation: A Dose-Finding Study Using the Continual Reassessment Method.
Neuromodulation **23**, 366-372 (2020).
47. I. Kahn, J. R. Andrews-Hanna, J. L. Vincent, A. Z. Snyder, R. L. Buckner, Distinct cortical anatomy
linked to subregions of the medial temporal lobe revealed by intrinsic functional connectivity. *J
Neurophysiol* **100**, 129-139 (2008).

48. R. L. Buckner, F. M. Krienen, A. Castellanos, J. C. Diaz, B. T. Yeo, The organization of the human cerebellum estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 2322-2345 (2011).

49. S. Dave, S. VanHaerents, J. L. Voss, Cerebellar Theta and Beta Noninvasive Stimulation Rhythms Differentially Influence Episodic Memory versus Semantic Prediction. *J Neurosci* **40**, 7300-7310 (2020).

50. M. A. Halko, F. Farzan, M. C. Eldaief, J. D. Schmahmann, A. Pascual-Leone, Intermittent theta-burst stimulation of the lateral cerebellum increases functional connectivity of the default network. *J Neurosci* **34**, 12049-12056 (2014).

51. Z. Li *et al.*, Multimodal Evidence for Hippocampal Engagement and Modulation by Functional Connectivity-Guided Parietal TMS. *bioRxiv*, (2025).

52. H. M. I. Turner, Calculating and synthesizing effect sizes. *Contemporary Issues in Communication Science and Disorders* **33**, 42-55 (2006).

53. H. C. van Houwelingen, L. R. Arends, T. Stijnen, Advanced methods in meta-analysis: multivariate approach and meta-regression. *Statistics in Medicine* **21**, 589-624 (2002).

54. W. Viechtbauer, Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software* **36**, 1 - 48 (2010).

55. W. Viechtbauer, M. W. Cheung, Outlier and influence diagnostics for meta-analysis. *Res Synth Methods* **1**, 112-125 (2010).

56. M. Egger, G. Davey Smith, M. Schneider, C. Minder, Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**, 629-634 (1997).

57. J. P. Higgins, S. G. Thompson, Quantifying heterogeneity in a meta-analysis. *Stat Med* **21**, 1539-1558 (2002).

58. G. B. Saturnino *et al.*, in *Brain and Human Body Modeling: Computational Human Modeling at EMBC 2018,* S. Makarov, M. Horner, G. Noetscher, Eds. (Springer, Copyright 2019, The Author(s). Cham (CH), 2019), pp. 3-25.

Author contributions:

Conceptualization: EBG, PFA, JLV

Methodology: EBG

Investigation: PFA, JMHR, ASA, JLV

Writing – original draft: EBG, PFA, JLV

Writing – review & editing: EBG, PFA, JMHR, ASA, JLV

Competing interests: Authors declare that they have no competing interests.

Data availability: Upon acceptance, the primary data will be deposited in a publicly available database (Dryad, datadryad.org) for use by other researchers without any restrictions on access or use. The data made available will include a table of all standardized effect sizes organized by study and by factor, such that reported analyses could be replicated.

Figure 1. Targets and methods used in HITS experiments. (**A**) Hippocampal indirect targets and neocortical stimulation locations are shown overlaid on a template brain with the resting-state fMRI connectivity network of the hippocampal indirect targets displayed in red. Stimulation locations are colorized separately for studies that used individualized targeting versus those that used group/atlas-based targets. For studies using individualized targeting, the average (centroid) location of the targets for all subjects in the study is shown. The hippocampal indirect targets are shown for those studies using individualized targeting. Note that not all stimulation locations fall within the highlighted red hippocampal network, as that specific network was defined based only on those studies with hippocampal indirect targets reported, and is shown for illustrative purposes to highlight proximity of all stimulation targets to a typical hippocampal network. (**B**) Representative electrical field (*58*) induced by TMS of one of the left-parietal stimulation sites at a typical intensity (estimate of 100% MT). The field is thresholded at 66% of its maximum intensity, as stimulation of lower intensity should have negligible effects on neuronal activity (all analyzed studies used stimulation intensity of 70% or above). (**C**) Coronal slices for the indicated positions show the hippocampal indirect targets and stimulation locations in greater detail. K-means clustering indicated that the majority of stimulation locations (84%) comprised a cluster within left parietal cortex, as indicated.

571 **Figure 2. HITS selectively improved episodic memory overall.** (**A**) Forest plot of all effects of HITS on
572 episodic-memory task outcomes, ordered by size. (**B**) Forest plot of all effects of HITS on non-memory task
573 outcomes. The pooled effect is shown for both plots. Red circles indicate outliers that were excluded in
574 sensitivity analyses. Circle size indicates the study's weighted contribution to the meta-regression model.

Figure 3. Study factors that modulated effects of HITS on episodic memory. (**A**) Factors that significantly modulated HITS effects on episodic memory. The pie chart indicates the percentage of effects for each category of each factor. Individual effects are shown in the box plots, grouped by factor levels. (**B**) Factors that did not significantly modulate HITS effects on episodic memory, plotted in the same format. *Indicates an effect modification that was significant in the main analysis but not in the sensitivity analysis.

Figure 4. Greater effects of HITS on memory in studies having optimized designs. Box plots of expected effect sizes projected from the meta-regression model for studies in which HITS was applied before encoding and measured using recollection format tests, versus all other study factors. Circles are individual projected effects, with circle size indicating standard error of the prediction, as indicated.

Study #	First Author	Year	Sample Size	DOI
Included in meta-analyses				
1	Wang	2014	16	10.1126/science.1252900
2	Yazar	2014	69	10.1371/journal.pone.0110414
3	Wang	2015	*	10.1002/hipo.22416
4	Bonnì	2015	30	10.1016/j.bbr.2014.12.032
5	Nilakantan	2017	16	10.1016/j.cub.2016.12.042
6	Yazar	2017	23	10.1016/j.brs.2017.02.011
7	Kim	2018	32	10.1126/sciadv.aar2768
8	Tambini	2018	22	10.1162/jocn_a_01300
9	Bonnici	2018	22	10.1523/JNEUROSCI.1239-18.2018
10	Koch	2018	14	10.1016/j.neuroimage.2017.12.048
11	Wynn	2018	19	10.1101/lm.048033.118
12	Ye	2018	18	10.1523/JNEUROSCI.0660-18.2018
13	Hermiller	2019	24	10.1002/hipo.23054
14	Hermiller	2019	14	10.1002/brb3.1393
15	Nilakantan	2019	15	10.1212/WNL.0000000000007502
16	Hermiller	2020	16	10.1523/JNEUROSCI.0486-20.2020
17	Chen	2020	13	10.18632/aging.202313
18	Gao	2021	32	10.1016/j.brainres.2021.147510
19	Hebscher	2021	20	10.1016/j.cub.2021.01.027
20	Freedberg	2021	23	10.1016/j.neuroimage.2021.118199
21	Velioğlu	2021	15	10.1016/j.nlm.2021.107410
22	Jia	2021	69	10.3389/fnagi.2021.693611
23	Freedberg	2022	29	10.1016/j.bbr.2021.113707
24	Dave	2022	16	10.1016/j.crneur.2022.100030
25	Hua	2022	39	10.3389/fnhum.2022.973298
26	Hermiller	2022	30	10.1016/j.neurobiolaging.2021.09.018
27	Chen	2022	8	10.1073/pnas.2113778119
28	Wei	2022	29	10.1016/j.psychres.2022.114721
29	Tang	2023	89	10.1093/schbul/sbad015
30	Chen	2023	18	10.1111/cns.14177
31	You	2023	34	10.2147/CIA.S416992
32	Jin	2024	11	10.1016/j.bspc.2023.105725
33	Cheng	2025	15	10.1162/jocn_a_02273
34	Jung	2024	30	10.1001/jamanetworkopen.2024.9220
35	Lv	2024	60	10.1016/j.bbr.2024.115117
36	Webler	2024	24	10.1016/j.bpsgos.2024.100309
37	Velioğlu	2024	12	10.29399/npa.28420
38	Zheng	2024	43	10.1093/cercor/bhae460
Excluded from meta-analyses				
39	Zhao	2016	30	10.18632/oncotarget.13060
40	Warren	2019	*	10.7554/eLife.49458
41	Hendrikse	2020	39	10.1016/j.cortex.2020.08.028
42	Wang	2020	8	10.3389/fnhum.2020.541791
43	Cash	2022	*	10.1016/j.brs.2022.09.004
44	Yang	2022	22	10.3233/JAD-215390
45	Chen	2022	12	10.3233/JAD-210661
46	Li	2023	28	10.3390/brainsci13030419
47	Li	2024	38	10.2147/NDT.S468219

592 **Table 1. Studies identified by systematic review.** Study information is listed separately for studies
593 that were included versus excluded from meta-analyses. Sample sizes listed are for the number of
594 subjects contributing to statistical effects that were analyzed or described. *Studies that re-analyzed
595 data from another study and therefore did not contribute independent data are indicated.

Supplementary Materials

A meta-analysis suggests that TMS targeting the hippocampal network selectively improves episodic memory

Elena Badillo Goicoechea, Phillip F. Agres, Johanna M.H. Rau, Arantzazu San Agustín, Joel L. Voss

Contents:

Figure S1. Histogram of sample sizes per analyzed effect, plotted separately for episodic-memory and non-memory effects.

Figure S2. Relationships among factors in episodic memory studies.

Figure S3. Funnel plot of episodic memory HITS effects. Outliers are indicated in red. The I^2 statistic is indicated.

Figure S4. Funnel plot of non-memory HITS effects. Outliers are indicated in red. The I^2 statistic is indicated.

#	Factor	Levels	Description	Rationale
1	Task	Recognition	Episodic memory tests using recognition formats; i.e., memoranda are repeated from study to test and subjects make old/new discrimination judgments	Recollection is thought by many to be more heavily dependent on hippocampus and its network than familiarity, which differentially contribute to tasks using recollection and recognition formats. While Recollection tasks could have been sensitive to many types of memory processing, including familiarity, at least one aspect of performance depended on recollection. Performance of tasks with other formats could be supported by an unknown mix of recollection, familiarity, or other processes.
		Recollection	Episodic memory tests using recollection formats; i.e., free recall, cued recall, or self-reported recollection judgments for tasks using paired associates, item-context associates, or multi-part complex stimuli as memoranda	
		Other	Episodic memory tests using other formats; i.e., temporal order judgments, recall of details from videos, scored visual reproduction, and tests that assess mnemonic pattern separation	
2	Targeting	Individual	The location where TMS was applied was determined by subject-specific analysis of functional or structural MRI connectivity of hippocampus; i.e., a location of parietal/occipital neocortex was targeted based on a maximal connectivity value with hippocampus defined in each subject.	Individualized targeting has been used in many studies based on the assumption that network structure varies across people and that indirect influences of stimulation on the hippocampus are transsynaptic and predicted by connectivity. Group targeting assumes that there is sufficient homogeneity across subjects to reliably affect networks, or was based on hypotheses about specific parietal areas that happen to be among hippocampal network areas typically targeted using individualized targeting approaches.
		Group/atlas	TMS was applied to approximately the same target in each subject, based on functional or structural MRI connectivity in a group/atlas or based on a specific hypothesis about the function of a specific location.	
3	HITS Timing	Pre-encoding	rTMS was delivered sometime before the period of memory encoding, ranging from application during the several seconds before individual memoranda were studied, in the minutes to hours before an encoding block was administered, to for many sessions over several days or weeks before the encoding phase of the memory task was completed.	This is a post hoc factor. Most studies used the "before" design whereas some used the "after" design. Hippocampal involvement in memory could differ for encoding versus retrieval.
		Post-encoding	All of these experiments applied a single session of stimulation after a block of encoding and before the corresponding memory test was given minutes to tens of minutes later.	
4	Population	Younger adult	Young adults were participants (<45 years old).	Populations with age-related memory decline (older adult) or with clinical memory impairments (MCI/AD and potentially Other) could present more opportunity for stimulation to improve memory. Alternatively, network disruptions frequently observed in memory disorders could reduce the ability for TMS to indirectly influence hippocampus due to reduced possibility of transsynaptic influence. Thus, the a priori hypotheses are that effects will differ among populations.
		Older adult	Older adults were participants (>65 years old).	
		MCI / AD	Participants included individuals with diagnoses of Mild Cognitive Impairment (MCI) or mild to moderate Alzheimer's dementia (AD) following clinical standards.	
		Other	Other groups included individuals with first-episode psychosis, individuals at high risk of schizophrenia, and individuals with symptoms of anxiety disorders of post-traumatic stress disorder.	
5	Target	Left parietal	Lateral parietal cortex (LPC) was stimulated in left hemisphere in most studies, at locations defined via either individualized or group targeting methods, as shown in Figure 1.	This is a post hoc factor. The majority of studies targeted areas of lateral, inferior, posterior parietal cortex near the angular gyrus, following the general method of Wang et al. 2014.
		Other	Other stimulation locations included	Others targeted precuneus or other

			precuneus or other parietal-occipital areas defined based on individual or group targeting methods, as shown in Figure 1. These were combined because separating them would have led to very few effects in each separate level.	parietal or anterior occipital areas based on connectivity with hippocampus. All of these areas are considered within the hippocampal network and so no strong a priori hypotheses regarding the effects of HITS were warranted.
6	Task delay	Short	The effects of stimulation were measured on the same day on which stimulation was delivered. Some studies applied short stimulation trains in the seconds before memoranda were presented, others delivered a several-minute stimulation train and then tested effects within one hour, and others used multi-day or multi-week stimulation and tested the effects on the same day following the last stimulation session.	This is a post hoc factor. Generally, the effects of stimulation are thought to decay, albeit at unknown rates. However, studies tended to measure outcomes at test delays aligned with the duration of stimulation (i.e., shorter test delays for protocols involving less stimulation and longer test delays for protocols involving more stimulation over more days; see Figure S2). Thus, as test delay was aligned to measure outcomes appropriately for hypothetical protocol-specific hypothesized decay rates, there was no strong a priori hypothesis to suggest that test delays would be non-equivalent.
		Medium	The effects of stimulation were measured on the following day after stimulation delivery. These studies mostly delivered several consecutive daily sessions of stimulation and measured the effects ~24 hours after the final stimulation session.	
		Long	The effects of stimulation were measured more than one day after the final stimulation session in multi-day or multi-week stimulation protocols, up to ~4 weeks after the end of stimulation.	
7	Sessions	1	Protocols involved only one session of stimulation before outcomes were assessed. This encompasses studies that used short trains of stimulation (seconds) immediately before memoranda were presented, as well as several-minute trains of stimulation before memory was assessed.	This is a post hoc factor. Generally, more sessions of stimulation are thought to produce greater and/or longer-lasting effects. However, as session counts and task delays were correlated (see "Task delay" factor), there were no strong a priori hypotheses about differences among session counts.
		2 – 4	Protocols involved two to four consecutive sessions of stimulation spread out across multiple days.	
		5+	Protocols involved five or more sessions of stimulation spread out across multiple days and/or weeks	
8	Intensity	100+% MT	Stimulation was applied at an intensity of greater than or equal to the resting motor threshold (MT), typically no higher than 120% of the MT.	This is a post hoc factor. Generally, higher intensities would be thought to have greater effects. However, this factor overlapped considerably with TMS protocol (factor 11, which was not retained in the lasso regressions). That is, studies using repetitive TMS trains at high frequencies tended to use intensities at or above motor threshold, whereas studies using patterned theta-burst stimulation tended to use intensities lower than 100% of motor threshold (Figure S2). Thus, there were no strong a priori hypotheses concerning this factor specifically.
		<100% MT	Stimulation was applied at an intensity of less than 100% of the resting MT, typically at 90%, 80%, or 70% of the MT.	
9	Control	Baseline	The comparison for the effects of stimulation was the baseline, pre-stimulation performance value.	This was a post hoc factor. Different control conditions are used to achieve different scientific objectives. As all control conditions would be expected to yield little if any effect on memory performance, any significant variation in effects of HITS on memory by control condition would raise doubts about its robustness to small differences in experiment design
		Other location	The comparison for the effects of stimulation of the hippocampal network was stimulation of some other part of the brain not within the hippocampal network using the same stimulation parameters. Typically, primary or secondary motor or somatosensory areas were used as	

			control locations.	choices.
		Low intensity	The comparison for the effects of stimulation was stimulation of the same location using the same overall stimulation parameters but at a very low intensity, typically <30% of resting motor threshold.	
		Other	The comparison for the effects of stimulation was stimulation of the same location using a different rhythmic pattern that was hypothesized to have less of an effect on memory.	
10	Design	Post - Post	The effects of stimulation on performance were statistically compared for after receiving HITS versus after receiving the control condition (e.g., t test of the performance scores after HITS versus after control).	This was a post hoc factor. Different statistical designs are used to achieve different scientific objectives and to accommodate different practical constrains on experiment design. As the choice of statistical design would be expected to yield little if any effect on memory performance, any significant variation in effects of HITS on memory by analytic design would raise doubts about its robustness to small differences in experiment analysis choices.
		Pre - Post	The statistical comparison of effects considered performance after versus before receiving HITS (e.g., t test of performance scores after HITS versus baseline before HITS).	
		Delta	The statistical comparison of effects on performance was done by computing the pre- versus post-HITS change (delta) in scores and comparing to the pre- versus post-Control delta (e.g., t test of percentage change baseline to post-HITS versus percentage change baseline to post-Control).	
		Interaction	The statistical test of effects on performance treated time (pre- versus post-) and stimulation type (HITS versus control) as factors, as in a RM-ANOVA.	
		Other	More than two stimulation conditions, including the control condition, were compared as factors, as in a RM-ANOVA	
11	TMS protocol	rTMS	High-frequency (10, 15, or 20 Hz) rTMS, applied in either short trains of several seconds, or as a longer, approximately 20-min train per session.	Some evidence suggests that theta burst stimulation has greater effects on hippocampal network activity and episodic memory performance. However, this factor was strongly correlated with Intensity and Sessions factors (Figure S2). NOTE: this factor was not retained in the lasso regressions as it was strongly correlated with other factors, thus warranting no strong a priori hypotheses of its independent contributions.
		TBS	Theta-burst stimulation, including short trains of several seconds of theta burst, or as continuous theta burst (40 s) or intermittent theta burst (5 minutes) sessions.	

Table S1. Description of study factors.

Supplementary Results

Summary of findings from studies that were not included in the meta-analysis

As indicated in the main text, nine studies were excluded from meta-analysis (Table 1). Five of these exclusions were because statistical information provided in the publications were not sufficient for calculation of standardized effect sizes, and authors could not be reached to provide additional information (studies 39, 42, 44, 45, 47 from Table 1). One exclusion (study 40) was made because the sample of subjects analyzed overlapped 94% with subjects that contributed data to a previous publication that was included, and so exclusion was necessary to avoid redundant statistical results in the meta-analysis. Two studies were excluded because they did not report the effects of HITS on performance of an episodic memory test, but instead quantified the correlation between fMRI connectivity and the effects of HITS on episodic memory performance (studies 43 and 46). One study was excluded (study 41) because re-analysis of the same data by the same research team (study 43) reported a different conclusion than the original report, suggesting that stimulation may not have been delivered to intended hippocampal network targets in the original report, and thereby calling into question findings from the original report and making it unclear how to categorize the Targeting or Target factors for meta-analysis.

Of the nine excluded studies, seven reported the effects of HITS on episodic memory performance (studies 39, 40, 41, 42, 44, 45, 47). Five of these seven reported that HITS significantly improved performance in at least one of the episodic memory tasks that were administered (studies 39, 40, 42, 44, 45). One of those studies (study 44) also reported numeric but non-significant decrease in memory performance in a subgroup analysis of subjects with mild Alzheimer's dementia. Two of the seven studies (41 and 47) reported small numeric improvements in memory following HITS that were not statistically significant.

Regarding the study factors that modulated the effects of HITS in the main study (Figure 3A), the seven studies all all included at least one test using a recollection format (Task factor: Recollection) and all administered HITS before encoding (Timing factor: pre-encoding). Two used individualized targeting (studies 40 and 41), with the others using group/atlas-based targeting. Other factors that were not significantly related to memory outcomes following HITS in the main analysis (Figure 3B) varied among these seven excluded studies.

The other two excluded studies (43 and 46) investigated correlations between resting-state fMRI connectivity and the effects of HITS on episodic memory, without directly assessing the impact of HITS on episodic memory. One reported that fMRI connectivity of the hippocampus to the stimulated location predicted episodic memory improvement due to HITS (study 43). The other study reported that HITS changed the correlation between fMRI connectivity and memory performance scores (study 46).

Overall, reported results in the excluded studies were generally consistent with findings from the meta-analysis of included studies. That is, of the seven excluded studies that reported effects of HITS on episodic memory, the majority (5/7) reported statistically significant episodic memory enhancement by HITS, but with variability in effect sizes among studies and a minority (2/7) reporting small but statistically nonsignificant changes in memory due to HITS. Thus, if effects from these studies could have been included in the meta-analysis, it is unlikely that conclusions would have been substantially affected.