1   **Deep learning-based precision phenotyping of spine curvature identifies novel genetic risk**
2   **loci for scoliosis in the UK Biobank**
3

4   Michael Zeosk[1*], Eucharist Kun[1*], Siddharth Reddy[1], Devansh Pandey[1], Liaoyi Xu[1], Joyce Y.
5   Wang[1], Chenfei Li[1], Ryan S. Gray[2,3], Carol A. Wise[4,5], Nao Otomo[4,5] & Vagheesh M.
6   Narasimhan[1,6+]
7
[1] Department of Integrative Biology, The University of Texas at Austin
[2] Department of Pediatrics, Dell Children's Medical Center of Central Texas
[3] Department of Nutritional Sciences, The University of Texas at Austin
[4] Department of Orthopedic Surgery, University of Texas Southwestern Medical Center
[5] Center for Musculoskeletal Research, Texas Scottish Rite Hospital for Children
[6] Department of Statistics and Data Science, The University of Texas at Austin
[*] Contributed equally to this work
[+] To whom correspondence should be addressed

16
17   **Abstract**
18

Scoliosis is the most common developmental spinal deformity, but its genetic underpinnings
remain only partially understood. To enhance the identification of scoliosis-related loci, we
utilized whole body dual energy X-ray absorptiometry (DXA) scans from 57,887 individuals in
the UK Biobank (UKB), and quantified spine curvature by applying deep learning models to
segment then landmark vertebrae to measure the cumulative horizontal displacement of the spine
from a central axis. On a subset of 120 individuals, our automated image-derived curvature
measurements showed a correlation 0.92 with clinical Cobb angle assessments, supporting their
validity as a proxy for scoliosis severity. To connect spinal curvature with its genetic basis we
conducted a genome-wide association study (GWAS). Our quantitative imaging phenotype
allowed us to identify 2 novel loci associated with scoliosis in a European population not seen in
previous GWAS. These loci are in the gene *SEM1/SHFM1* as well as on a lncRNA on chr 3 that
is downstream of *EDEM1* and upstream of *GRM7*. Genetic correlation analysis revealed
significant overlap between our image-based GWAS and ICD-10 based GWAS in both the UKB
and Biobank of Japan. We also showed that our quantitative GWAS had more statistical power
to identify new loci than a case-control dataset with an order of magnitude larger sample size.
Increased spine curvature was also associated with increased leg length discrepancy, reduced
muscle strength and decreased bone density, and increased incidence of knee but not hip
osteoarthritis. Our results illustrate the potential of using quantitative imaging phenotypes to
uncover genetic associations that are challenging to capture with medical records alone and
identify new loci for functional follow-up.

39
40   **Introduction**

Scoliosis is a complex musculoskeletal disorder characterized by an abnormal lateral curvature of the spine[1,2]. This disorder manifests from a variety of causes with a small proportion caused by rare Mendelian mutations or congenital vertebral anomalies that present from birth. However, the most common forms of scoliosis are idiopathic forms, which can manifest in early childhood, adolescence, or adulthood[3,4]. Untreated scoliosis can lead to a range of complications, including chronic pain and increased risk of fractures, significantly impacting the quality of life for affected individuals. Given that scoliosis is often identified only after progression to noticeable spinal deformity, understanding the genetic basis of scoliosis is critical for advancing treatment strategies and enabling earlier identification of the condition. Despite the multifactorial nature of scoliosis, encompassing both genetic and environmental contributions[5], its underlying genetic factors remain poorly understood. Identifying these also has increases the potential for the identification of therapeutic targets to prevent progression and improve patient outcomes.

Genome-wide association studies (GWAS) have provided valuable insights into the genetic architecture of scoliosis. Notably, several GWAS conducted in Japanese populations have identified susceptibility loci associated with adolescent idiopathic scoliosis (AIS) using binary health record data from the Biobank of Japan[6,7]. A recent meta-analysis of these studies has expanded the total number of reported loci to 21[8]. Another meta-analysis of AIS GWAS across multiple populations replicated many of these reported loci as well as discovered a locus that had sexually dimorphic effects for idiopathic scoliosis[9,10]. These genetic discoveries come from a mix of biobank and disease-specific cohorts, however, no GWAS has successfully identified genome wide significant loci from purely cohort studies such as the UK Biobank (UKB) for scoliosis, as the total number of cases of individuals coded as having scoliosis is only 4,851, or less than 1%.

A primary limitation of unselected cohort studies is the use of ICD-10 records to determine disease status. In the clinic, the severity of scoliosis is often classified using the Cobb angle, made by drawing lines along the upper and lower endplates of the most tilted vertebrae in a spinal curve and measuring the angle where these lines intersect[1]. However, this nuance is often lost in the medical record and simply classified as either having scoliosis or not. In addition, these binary case-control assessments derived from health records typically only include individuals that have extreme severe disease, thereby greatly limiting the number of cases in datasets like population biobanks. Furthermore, coding individuals as cases in the medical record could be dependent on clinician experience, the severity they consider constitutes a diagnosis, or the patient's access to care or insurance provided[11]. Recent advancements in deep learning and computer vision provide an opportunity to overcome these limitations by directly assessing disease phenotypes from radiographic images, thus generating quantitative image-derived phenotypes (IDPs). In musculoskeletal diseases, imaging has been the primary method of diagnosis and tracking of disease progression in the clinic for decades[1]. Therefore, deep learning has the potential to directly capture nuanced disease severity in an accurate and replicable way

that can improve gene discovery for such disorders. In previous work, we have shown that
utilizing deep learning models for phenotyping of radiographic images increases the power of
gene discovery of knee osteoarthritis[11], and others have also applied such approaches to
understand the genetic basis of hip joint space[12]. However, in scoliosis, though methods have
been designed to automatically measure spinal curvature[13], these have not been extended to study
the genetic drivers of the disease. In this study, we utilize deep learning and novel image analysis
methods to produce quantitative image-derived phenotypes associated with spine curvature - a
natural continuous measure of scoliosis severity. By training an image segmentation model to
isolate the spine from existing dual energy X-ray absorptiometry (DXA) images and further
developing a method to quantify the cumulative lateral curvature of the spine on ~60,000
individuals, we provide a quantitative measure of the disease over and beyond the ICD-10
scoliosis diagnosis health record data (M41)[11]. While ascertainment of patients with scoliosis is
typically carried out in adolescents, like those utilized in the previous GWAS studies, this
analysis examines spine curvature using DXA scans of adult individuals encompassing the
different periods in which scoliosis can manifest.

Results
Dataset and quality control of DXA imaging and genetic data

To study the genetic basis of spine curvature, we jointly analyzed paired DXA imaging and
imputed genome sequence data of 69,981 individuals in the UKB[14]. We first restricted the
dataset to individuals of white British ancestry. Next, as the bulk imaging data from the UKB
consisted of DXA images that reflect scans of different body parts, we used a deep-learning
approach to subset the imaging dataset to only full body transparent DXA scans[15,16]. We then
removed individuals that had outlier image resolutions or poor quality DXA scans, and padded
images to a standard size for processing. For the genetic data, we applied standard variant and
sample quality control (QC) and analyzed 7.2 million common biallelic SNPs with minor allele
frequency greater than 0.1%. Post quality control, we were left with combined imaging and
genetic data for a total of 57,887 individuals aged between 46 and 81 with a mean age of 64. The
ratio of males to females was 50.2% to 49.8% in this analysis, consistent with the overall
distribution in the UKB.

A deep learning model for image segmentation and quantification of spine curvature

To perform automated measurement of spinal curvature we used a training dataset of 63 images
(50 training, 13 validation, **Supplementary Table 1**) (**Fig. 1a**). On each of these images, we
manually outlined the spine from the top vertebrae to the bottom. We then trained a deep-
learning model based on the U-Net architecture[17,18] with a 34-layer ResNet encoder[19] to perform
semantic segmentation of the spine at pixel-level resolution (**Fig. 1b**). We then applied this
model to all the 57,887 individual full body transparent images and obtained spine segmentation

masks for each DXA image. To measure spine curvature, we obtained landmarks at 20 different
positions along the spine mask reflecting the pixel centroids at each segment of the spine (**Fig.**
1c). Then taking a line between the top landmark and the bottom landmark of the mask, we
rotated the spine image such that the line was in a vertical orientation (**Fig. 1d**). This then
allowed us to calculate the overall cumulative spine curvature simply by examining the
horizontal displacement between subsequent landmarks. Our spine curvature phenotype was
therefore quantified as the sum of the absolute differences between x-coordinates of adjacent
centroids along the spine, providing a continuous measure of lateral deviation (**Fig. 1e**). We
normalized these measurements across images by converting pixel lengths to mm, which we
accomplished by multiplying the pixel lengths by the pixel to mm scaling reported for each
image provided by the scanner.

Figure 1. Deep learning segmentation and quantification of spinal curvature. **a**. Original DXA
scan converted from DICOM format to high resolution JPEG image for downstream analysis. **b**.
Deep-learning-based segmentation of spine from original image using a U-net architecture[17,18]
with a 34-layer ResNet encoder[19]. **c**. Automatic identification of key landmarks throughout the
spine through the calculation of evenly spaced centroids. **d**. Rotation of spine to a vertical
orientation **e**. Measurement of overall spine curvature as the sum of absolute horizontal
displacement between landmarks (marked in yellow).

Evaluation of the spine curvature measurements

We evaluated the performance of the segmentation model in several ways. First, the set accuracy
- the pixel correspondence between the labeled masks and annotation of the trained model on
validation data - was 0.99. Second, the correlation between images taken of the same person
across two imaging visits was 0.68, despite changes in image resolution, scanner, technician, and
back posture of each subject (**Fig. 2a**). We do not expect to see 100% concordance across these
replicates as differences in spinal curvature can arise due to actual differences in posture or due
to scoliosis. Third, we examined the correlation between Cobb angle as assessed manually by
certificated orthopedic surgeons and our spinal curvature measurement in 124 individuals, which
exhibited a significant Pearson correlation of 0.92. Additionally, from these 124 individuals, we
examined differences in spinal curvature based on scoliosis diagnosis from Cobb angle
measurement. Individuals with a Cobb angle exceeding 10 degrees, which is the typical cutoff

for scoliosis diagnosis, had a statistically significant higher spinal curvature than individuals with
a Cobb angle of less than 10 degrees (t-test, $P < 3.4 \times 10^{-8}$). We also confirmed this by
examining the relationship between the spinal curvature phenotype and scoliosis status as
captured in the ICD-10 code M41 case–control data[14] (**Fig. 2b**). As expected, the spinal
curvature phenotype was significantly higher in cases compared to controls (t-test,
$P < 2.2 \times 10^{-16}$). Finally, we examined the relationships between the spine curvature phenotype
and age—which is also known to be highly associated with spinal compression and curvature.
Again, as expected, we found that the average spinal curvature phenotype increased significantly
with age (linear regression, ß = 0.019, $P < 0.0001$) (**Fig. 2c**). Taken together, these suggest that
our spine curvature phenotype was replicable, was highly correlated with Cobb angle – the
clinical measure of scoliosis – and captured a certain proportion of actual inherent deviation in
the vertebral column and not just deviation due to posture at the time the image was taken.

Figure 2. Evaluation of cumulative spine curvature across visits, ICD-10 scoliosis diagnosis
status, and age. **a** Correlation of spine curvature between two visits for individuals who were
imaged multiple times, (R=0.68, $P <0.0001$). **b** Comparison of cumulative spine curvature
between individuals with and without a recorded ICD-10 M41 Scoliosis diagnosis. **c**. Analysis of
the relationship between average cumulative spine curvature and age, with circle size correlating
to the number of individuals in that age group, (beta = 0.019, $P < 0.0001$).

Genetic associations of image derived spine curvature

Having obtained normalized spine curvature measurements on each individual, we linked this
phenotype to its genetic basis. We carried out a common variant GWAS of our spine curvature
phenotype using PLINK[20] on 57,887 white British individuals adjusting for age, sex, and the first
20 genetic principal components. Using the summary statistics of this GWAS, the estimated
heritability for spine curvature using Linkage Disequilibrium Score Regression (LDSC)[21] was
0.071 ± 0.01. In total, our GWAS identified 3 genome-wide significant loci associated with
scoliosis at a threshold of $P = 5 \times 10^{-8}$ while a GWAS performed using the UKB ICD-10
scoliosis diagnosis data on the non-imaged individuals found no genome-wide significant loci
(**Fig. 3a-3b**). One of the SNPs is in a region upstream of the gene *PAX1* that has previously been
implicated in other scoliosis GWAS and functionally validated in Zebrafish models[8,9]. Deletion

of this locus has also been shown to result in tail abnormalities in female mice[22]. However, we
identified two novel loci, never previously reported by other studies of scoliosis or related traits
in the GWAS catalog. The first is near the gene *SEM1/SHFM1*. Deletions in this region have
been shown to be associated with the rare limb developmental disorder, split hand/split foot
malformation type 1[23]. The second locus is on chr 3 and is in a lncRNA that is downstream of
EDEM1 and upstream of *GRM7*.

Figure 3. GWAS and phenotype associations **a**. Manhattan plot for GWAS performed using
ICD-10 scoliosis code alone. **b**. Manhattan plot for GWAS performed using deep learning
derived spinal curvature measurement. **c.** Regressions of bone and muscle phenotypes with
spinal curvature. Linear and logistic regressions are shown separately. Asterisks denote
significant phenotypes that have FDR-adjusted p-values < 0.05.
198
199   **Deep learning-based phenotyping improves power for discovering scoliosis associated loci**
200
Next, we assessed the correspondence between our GWAS to other GWASs that had been
performed on traditionally measured scoliosis phenotypes obtained from the electronic
healthcare record. To do so, we conducted genetic correlation analyses with two case-control
GWAS datasets. First, we utilized the UKB ICD-10 scoliosis diagnosis GWAS that we had
previously carried out on the non-imaged individuals. The genetic correlation between this
binary GWAS and our image based GWAS was 0.46 ($P = 0.006$). While not close to 1 this is
within expectation for genetic correlation between a case-control GWAS and a quantitative
GWAS on the same trait. In contrast, the genetic correlation between arm length, a skeletal trait
that is not associated with the spine, and the UKB ICD-10 scoliosis diagnosis was not significant
($P = 0.11$). This serves as a negative control and suggests that our imaging-based phenotyping
was identifying genetic loci that were highly correlated to the medical record-based diagnosis for
scoliosis. Second, we examined genetic correlation with a case-control GWAS for scoliosis
carried out in a Japanese population[8] using POPCORN[24] to adjust for differences in LD Structure
between Europeans and Japanese. This demonstrated a slightly higher genetic correlation of 0.56
($P = 1.18 \times 10^{-6}$). These findings suggest that our continuous, image-derived phenotypes
effectively capture scoliosis-related genetic signals identified using diagnostic records while
offering improved statistical power.
218
219   **Associations of scoliosis with bone and muscle density and musculoskeletal disease**
220
We then examined the association between spine curvature and bone and muscle strength. After
controlling for age and sex, we found that in line with epidemiological data, hand grip strength (a
proxy for muscular strength) and spine bone density were both negatively associated with spine
curvature (multivariate regression covariate p-value for grip strength: $P = 1.4 \times 10^{-8}$, p-value for
bone mineral density: $P < 2.2 \times 10^{-16}$) (**Fig. 3c**). We next examined the association between leg
length discrepancy and scoliosis. An actual difference in bone length between the legs is thought
to contribute to scoliosis, as the spine bends to accommodate the discrepancy. To examine this
hypothesis, we utilized deep learning approaches to measure the difference in the length of the
left and right leg. Though the difference between the length of the legs is small, only a few
millimeters, we were able to replicably measure this difference in the same individual across two
imaging visits taken more than two years apart. We found that spine curvature was positively
associated with the absolute value of leg length discrepancy after controlling for age, sex and

height (multivariate regression covariate p-value for leg length discrepancy: $P = 2.58 \times 10^{-4}$). We
also found that individuals with increased spine curvature also self-reported back pain at higher
rates, and had higher incidence of knee osteoarthritis, but not hip osteoarthritis in the ICD-10
code (multivariate regression covariate p-value for back pain: $P = 4.57 \times 10^{-2}$, knee OA: $P =$
4.84×10^{-2}, hip OA: $P = 0.82$).

Discussion

Our study demonstrates the potential of using deep learning-derived phenotypes, such as spine
curvature from DXA images, to improve genetic analyses of scoliosis. By moving beyond
traditional case–control phenotypes sourced from EHR records, we provide a quantitative
measure of spinal curvature that captures a continuum of variation, uncovering subtle phenotypic
differences associated with scoliosis risk. This quantitative approach circumvents challenges of
diagnostic inconsistency and variability in clinical definitions, offering a more standardized and
scalable method for phenotyping in biobank-scale settings[11].

One major limitation of our study is our inability to differentiate adult scoliosis from adolescent
idiopathic scoliosis. While adult scoliosis can occur as a continuation or exacerbation of
adolescent scoliosis, other factors such as spine degeneration can also play a role in adult
scoliosis. For reference, though cobb angle and spinal curvature measured in our cohort
displayed a Pearson correlation of 0.92, the heritability of cobb angle measured in 4465
adolescent patients was 0.21 ± 0.10, while the heritability of our spinal curvature measurement in
57,887 adults was 0.07 ± 0.008[25]. To understand if the genetic signals we obtained were likely to
be involved in adolescent scoliosis, we computed genetic correlation with the Japanese AIS
cohort[8] and observed a correlation of $R_g = 0.56$. This is in line with a previously reported genetic
correlation of 0.60 between cobb angle and adolescent idiopathic scoliosis[25]. Additionally, as
adolescent idiopathic scoliosis is more common in the thoracic region and adult scoliosis more
commonly affects the lumbar region, we wanted to see if we could separate the genetic signals
by dividing our spinal curvature assessment into two parts: the top half reflecting the first 10
points and focused more on thoracic vertebrae in our spine segmentation, and the bottom half
focused on lumbar vertebrae. We then recomputed our GWAS on these two halves of the spine.
We found that the genetic correlation between the adolescent cohort in the Japanese biobank and
the bottom half curvature was $R_g = 0.66 \pm 0.15$, while the top half was 0.48 ± 0.10. Thus, we
were unable to detect significant differences using this approach. However, significantly non-
zero genetic correlation suggests that our approach is capturing a significant proportion of the
genetic basis of adolescent scoliosis as well. Similar to findings in other musculoskeletal
conditions like knee osteoarthritis[11], our results suggest that quantitative phenotyping offers
increased statistical power for genetic studies compared to case–control approaches.

Despite its advantages such as improved consistency and scalability, our approach has

limitations. The DXA-based quantitative phenotype remains a derived measure, and its precision relies on accurate segmentation and alignment. As the derived measure is not a cobb angle commonly used to analyze scoliosis severity, singular results can be difficult to interpret out of context. Similarly, while our model reduces bias inherent to health record data, any bias in the underlying imaging data or the segmentation model may propagate through the analysis pipeline. Additionally, as our GWAS focused primarily on individuals of European ancestry, the transferability of our findings to other populations requires further validation. Future efforts should prioritize extending this methodology to more diverse cohorts and exploring its applicability to other imaging modalities, such as MRI or CT scans.

We observed several associations between spine curvature and phenotypes such as leg length discrepancy and bone mineral density. While these associations reflect plausible biologically meaningful correlations such as leg length discrepancy driving the difference in spine curvature, we note that these analyses do not represent causal inference, and additional work is required to examine these effects.

Our study contributes to a growing body of evidence supporting the use of image-derived phenotypes for large-scale genetic research. By providing a proof-of-concept for the utility of quantitative phenotyping in scoliosis, our results underscore the broader potential for applying similar methodologies to other conditions where imaging plays a key diagnostic role. For example, directly measuring spinal curvature could serve as a model for rethinking disease phenotyping in biobanks and accelerating discoveries in complex traits and diseases.

Methods
UKB dataset and participants

All analyses were conducted using data from the UK Biobank (UKB), a prospective, population-based cohort that recruited 500,000 individuals aged 40–80 in the UK between 2006 and 2010[14]. The dataset offers extensive health records and whole-genome sequencing, allowing us to analyze 12.1 million common biallelic SNPs with a minor allele frequency greater than 0.1%. We focused on 402,000 participants who self-identified as white British and had available genetic data and continued consent as of February 22, 2022. Access to the dataset was granted under application number 65439.

DXA imaging data

The UKB provides dual-energy X-ray absorptiometry (DXA) images collected using a Lunar iDXA instrument (GE Healthcare) in DICOM format as part of a bulk data field ID. For this study, we utilized full-body transparent DXA scans, with the original bulk download yielding 75,268 DICOM images from 70,405 unique participants. Images were assessed for poor quality

or incorrect labeling through a previously developed deep learning pipeline created to classify
types of DXA images and subsequently removed from analysis[15,16]. Each scan was associated
with metadata detailing pixel and millimeter-based dimensions. All participants were instructed
to lie flat on the scanner during imaging, resulting in non-weight-bearing images.
The DXA scans exhibited two distinct categories:
1. **Black background images**: 660–816 × 272 pixels
2. **White background images**: 930–950 × 300–382 pixels
To standardize image dimensions and control for scaling effects during deep learning, we applied
padding to bring the images to consistent sizes. Black background images were padded to 816 ×
288 pixels, while white background images were resized to 960 × 384 pixels. This process
involved converting each DICOM image into a NumPy array and appending rows or columns of
pixels as needed using NumPy[26], SciPy[27], and sci-kit image[28]. This resulted in 56,779 images of
816×288 and 18,489 images of 960×384. These images were then converted to high-resolution
JPEGs using the pydicom library[29] for downstream analyses.
327
328 **Image segmentation**
329
330 We manually annotated 63 DXA scans for spine segmentation using the LabelMe software[30],
331 splitting the sample evenly between participants with and without a prior scoliosis diagnosis
332 (ICD-10 code M41). Relevant demographic characteristics of the annotated dataset are
333 summarized in **Table 1**. An example of the result of our manual annotation is shown in
334 **Supplementary Fig. 1**.

335
336 **Supplementary Fig. 1**. Manual labeling of the spine in two individuals shown in red.

Group	Characteristic	Mean (s.d.)

Model Training Data	Participants, total	63
	Age	58 (7.24)
Male	Total (%)	32 (50.8)
	Age	60 (7.17)
Female	Total (%)	31 (49.2)
	Age	57 (7.04)
Scoliosis	Total (%)	38 (60.3)
	Male, total (%)	19 (50)
	Female, total (%)	19 (50)
	Age	58 (7.06)
Non-Scoliosis	Total (%)	25 (39.7)
	Male, total (%)	13 (52)
	Female, total (%)	12 (48)
	Age	58 (7.64)

Supplementary Table 1. Relevant characteristics and distribution of training data.

For segmentation we implemented a U-net architecture[17,18] with a 34-layer ResNet[19] encoder to
perform semantic segmentation of the spine. A separate model was trained for each image size
category. Both models were trained for 12 epochs with data augmentation such as rotation and
zoom applied to the training data to improve generalization. To improve the robustness of the
segmentation models, padded images from the other size category were subject to further image
resizing during training to increase annotated sample size. Upon visual analysis of early results,
we identified 224 model-predicted images for refinement and retrained the model to enhance
precision. Based on human annotation, the per pixel accuracy of the segmentation was 99%. The
final models were deployed on the 75,268 full-body transparent DXA images. Resulting
segmentation mask files (**Supplementary Fig. 2**) were exported as JPEG images using the
pydicom library[29].

Supplementary Fig 2. Example of the pixel mask output of our segmentation model obtained
for further image processing and measurement of spine curvature.

Spine rotation and curvature measurement

To quantify spinal curvature, we developed a custom pipeline in Python, adapted from the
PlantCV package[31], with additional modules from NumPy and OpenCV[32]. Each segmented spine
mask was processed independently and converted to a binary matrix, where pixels representing
the spine were set to 1, and background pixels were set to 0. Mask contours were analyzed, and a
bounding box was created around the spine, whose length represented the height of the spine.
This box was then divided into 20 equal-height segments to ensure consistent spatial sampling
through the spine and across images. Finally, within each segment, pixels were analyzed row by
row to calculate the centroid, representing the geometric center of the segment's pixels.
To assess spinal alignment, we compared the x-coordinates of the topmost and bottommost
centroids. If they did not align vertically, the angle to align the points was computed and we
performed a rotation using bicubic interpolation in OpenCV. The image was iteratively rotated
until the centroids aligned along a vertical axis to then calculate horizontal displacement. Once
the spine was aligned and centroid positions were stored, the curvature was quantified as the
cumulative horizontal displacement between consecutive centroids along the spine. The sum of
the absolute differences in x-coordinates between adjacent centroids provided a metric for
curvature, reflecting total deviation from a straight axis (**Supplementary Fig. 3**).

372
373 **Supplementary Fig. 3.** Cumulative horizontal displacement measurement
374 To address variability in image resolution, we scaled each distance in pixels using the
375 corresponding pixel to millimeter measurements from the metadata. These conversion ratios
376 were validated against previous studies using an alternate measure of computing these by
377 regressing the overall height in pixels[16].
378
379 **Imaging data processing and quality control**
380
To examine issues associated with our segmentation and image processing pipeline, we
examined key metrics, such as top and bottom centroid positions before and after alignment,
cumulative difference squared to emphasize quick changes in curvature, and pixel value counts
to identify segmentation issues were exported for quality control and downstream analyses.
Outliers identified based on unusual curvature values >2 standard deviations from the mean were
removed. After this quality control, the final dataset consisted of 74,667 images from 69,981
individuals, ready for subsequent analyses.
To ensure the data conformed to a normal distribution, the RankNorm function in R was applied.
This approach uses rank-based inverse normal transformation to standardize the data, enhancing
its suitability for downstream genetic correlation analyses while minimizing the influence of
outliers. This normalization is particularly useful when dealing with datasets where a few strong
cases may dominate and cause other curvature data to be obscured by noise, ensuring that
meaningful patterns are not lost.
394
395 **Correlation between Cobb angle measurement and image derived spinal curvature**
396
For a set of 124 images, with clinician guidance, we measured the Cobb angle, which is the angle
formed by lines drawn along the top and bottom vertebrae of the curve and typically used for
scoliosis diagnosis. We used Pearson correlation to examine the association between these
physician derived Cobb angle measurements and our computationally derived spinal curvature

measurements. Additionally, as scoliosis patients are only diagnosed at Cobb angle
measurements greater than 10 degrees, we split our cohort of 124 individuals into scoliosis and
non-scoliosis groups and compared the mean spinal curvature between the two groups through a
t-test.
405
406 **Genetic quality control**
407
For our genome-wide association analyses, we focused on participants of Caucasian ancestry,
self-reported as white British (FID 21000) and confirmed via genetic principal component
analysis (PCA) from the UKB. We excluded individuals whose self-reported sex (FID 31) did
not match their genetic sex (FID 22001), those with evidence of sex chromosome aneuploidy
(FID 22019), individuals with heterozygosity or genotype missingness outliers (based on UKB's
QC procedures for DNA sample processing and genotyping, FID 22027), and those with more
than nine third-degree relatives or any of unknown kinship (FID 22021). This filtering left us
with a sample of 402,000 individuals. From this subset, we further restricted our analysis to
participants who had undergone imaging (FID 20158) and had complete DXA measurements
(FID 12254), leaving 57,887 participants for analysis. We downloaded imputed genetic data for
487,253 individuals from the UKB[14] (covering chromosomes 1 through 22, FID 22828) and
filtered this dataset according to our QC criteria using PLINK v2[20]. We excluded any duplicate
SNPs (--rm-dup "exclude-all"), retained only biallelic SNPs (--snps-only "just-acgt"), limited the
dataset to those with no more than two alleles (--max-alleles 2), a minor allele frequency greater
than 1% (--maf 0.01), individual missingness rates below 2.5% (FID 22005), and a per-SNP
missingness threshold of 5% (--maxMissingPerSnp 0.05). Following these steps, 7,204,508
SNPs remained in the imputed dataset.
425
426 **Genome-wide association studies (GWAS)**
427
We conducted GWAS using PLINK v2[20] with the following filtering thresholds: minor allele
frequency greater than 0.01, SNP missingness below 5%, and individual missingness below
2.5%. We included the first 20 genetic principal components provided by UKB (FID 22009), sex
(FID 31), and age (FID 21022) as covariates. After applying both genetic and imaging QC filters,
57,887 participants remained for the final GWAS analysis, with 7,204,508 SNPs included in
each association study. SNPs identified in the GWAS were further clumped using the PLINK --
clump function with a significance threshold of 5.0×10^{-8}, a secondary significance threshold
of 1.0×10^{-4} for SNPs in linkage disequilibrium (LD), an r^2 threshold of 0.1, a 1 Mb window,
and a 20kb border around each gene. SNPs were assigned to genes using --clump-verbose and --
clump-range with the glist-hg19 reference. We annotated protein-coding genes within 250 kb of
each clumped region and on the GRCH37 genome browser.
439

Validation of Phenotype Consistency through Genetic Correlation Analysis

To assess the genetic similarity of scoliosis phenotypes across two populations and validate that the identified phenotype aligns with a scoliosis outcome, we carried out genetic correlation analysis. For samples of European ancestry from the UK Biobank we used Linkage Disequilibrium Score Regression[21]. For examining genetic correlation across ancestries and adjusting for LD differences, we used the Popcorn software package[24]. We examined genetic correlation between our image derived phenotypes and a GWAS from the UKB with case-control phenotypes from scoliosis diagnosis health records, and also a GWAS from participants in a Japanese population (BBJ) with case-control derived phenotypes. This analysis yielded estimates of heritability (h^2) and genetic correlation (R_g) between the two scoliosis phenotypes, along with associated standard errors, Z-scores, and p-values.

References

1. Janicki, J. A. & Alman, B. Scoliosis: Review of diagnosis and treatment. *Paediatr Child Health* **12**, 771 (2007).
2. Aebi, M. The adult scoliosis. *European Spine Journal* **14**, 925–948 (2005).
3. Janicki, J. A. & Alman, B. Scoliosis: Review of diagnosis and treatment. *Paediatr Child Health* **12**, 771 (2007).
4. Giampietro, P. F. Genetic Aspects of Congenital and Idiopathic Scoliosis. *Scientifica (Cairo)* **2012**, 152365 (2012).
5. Konieczny, M. R., Senyurt, H. & Krauspe, R. Epidemiology of adolescent idiopathic scoliosis. *J Child Orthop* **7**, 3 (2012).
6. Takahashi, Y. *et al.* A genome-wide association study identifies common variants near LBX1 associated with adolescent idiopathic scoliosis. *Nature Genetics 2011 43:12* **43**, 1237–1240 (2011).
7. Kou, I. *et al.* Genetic variants in GPR126 are associated with adolescent idiopathic scoliosis. *Nature Genetics 2013 45:6* **45**, 676–679 (2013).
8. Kou, I. *et al.* Genome-wide association study identifies 14 previously unreported susceptibility loci for adolescent idiopathic scoliosis in Japanese. *Nature Communications 2019 10:1* **10**, 1–9 (2019).
9. Sharma, S. *et al.* A PAX1 enhancer locus is associated with susceptibility to idiopathic scoliosis in females. *Nat Commun* **6**, 6452 (2015).
10. Khanshour, A. M. *et al.* Genome-wide meta-analysis and replication studies in multiple ethnicities identify novel adolescent idiopathic scoliosis susceptibility loci. *Hum Mol Genet* **27**, 3986–3998 (2018).
11. Flynn, B. I. *et al.* Deep learning based phenotyping of medical images improves power for gene discovery of complex disease. *npj Digital Medicine 2023 6:1* **6**, 1–12 (2023).

479   12.   Faber, B. G. *et al.* The identification of distinct protective and susceptibility mechanisms
480   for hip osteoarthritis: findings from a genome-wide association study meta-analysis of
481   minimum joint space width and Mendelian randomisation cluster analyses. *EBioMedicine*
482   **95**, 104759 (2023).
483   13.   Jamaludin, A. *et al.* Automated measurement of size of spinal curve in population-based
484   cohorts: Validation of a method based on total body dual energy X-ray absorptiometry
485   scans. *Bone* **172**, 116775 (2023).
486   14.   Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.
487   *Nature* **562**, 203–209 (2018).
488   15.   Xu, L. *et al.* The genetic architecture of and evolutionary constraints on the human pelvic
489   form. *Science (1979)* **388**, (2025).
490   16.   Kun, E. *et al.* The genetic architecture and evolution of the human skeletal form. *Science*
491   *(1979)* **381**, (2023).
492   17.   Weng, W. & Zhu, X. U-Net: Convolutional Networks for Biomedical Image
493   Segmentation. *IEEE Access* **9**, 16591–16603 (2015).
494   18.   Ronneberger, O., Fischer, P. & Brox, T. 2015-U-Net. *ArXiv* 1–8 (2015).
495   19.   He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image Recognition.
496   *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern*
497   *Recognition* 770–778 (2015) doi:10.48550/arxiv.1512.03385.
498   20.   Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-
499   Based Linkage Analyses. *Am J Hum Genet* **81**, 559 (2007).
500   21.   Bulik-Sullivan, B. *et al.* LD Score regression distinguishes confounding from polygenicity
501   in genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).
502   22.   Ushiki, A. *et al.* Deletion of Pax1 scoliosis-associated regulatory elements leads to a
503   female-biased tail abnormality. *Cell Rep* **43**, (2024).
504   23.   Crackower, M. A. *et al.* Characterization of the split hand/split foot malformation locus
505   SHFM1 at 7q21.3-q22.1 and analysis of a candidate gene for its expression during limb
506   development. *Hum Mol Genet* **5**, 571–579 (1996).
507   24.   Brown, B. C., Ye, C. J., Price, A. L. & Zaitlen, N. Transethnic Genetic-Correlation
508   Estimates from Summary Statistics. *Am J Hum Genet* **99**, 76 (2016).
509   25.   Otomo, N. *et al.* Polygenic Risk Score of Adolescent Idiopathic Scoliosis for Potential
510   Clinical Use. *Journal of Bone and Mineral Research* **36**, 1481–1491 (2021).
511   26.   Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020).
512   27.   Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python.
513   *Nat Methods* **17**, 261–272 (2020).
514   28.   Van Der Walt, S. *et al.* Scikit-image: Image processing in python. *PeerJ* e453 (2014)
515   doi:10.7717/PEERJ.453/FIG-5.
516   29.   Mason, D. *et al.* pydicom/pydicom: pydicom 2.3.0. (2022).
517   30.   Wada, K. Labelme: Image Polygonal Annotation with Python.

31. Gehan, M. A. *et al.* PlantCV v2: Image analysis software for high-throughput plant phenotyping. *PeerJ* **5**, e4088 (2017).

32. Bradski, G. The OpenCV Library. Preprint at (2000).

Data availability

All data used for this study were obtained from the UK Biobank under application number 65439. GWAS summary statistics are currently being uploaded to the GWAS catalog and are available at this Box link: https://utexas.box.com/s/5iqlxvhul0ri9torr89lspb66thmpcdl

Code availability

Deep-learning and image processing tools can be found at https://github.com/reb12345/Scoliosis.

Acknowledgements

V.M.N. was supported on a grant from the Allen Discovery Center program, a Paul G. Allen Frontiers Group advised program of the Paul G. Allen Family Foundation.

Author information

Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA
Michael Zeosk, Eucharist Kun, Siddharth Reddy, Liaoyi Xu, Devansh Pandey, Joyce Wang, Chenfei Li, Vagheesh M. Narasimhan

Department of Nutritional Sciences, The University of Texas at Austin, Austin, TX, USA,
Department of Pediatrics, Dell Medical School, Austin, TX, USA
Ryan Grey

Center for Musculoskeletal Research, Texas Scottish Rite Hospital for Children, Department of Orthopedic Surgery, The University of Texas Southwestern Medical Center, Ryan Grey

Department of Statistics and Data Science, The University of Texas at Austin, Austin, TX, USA
Vagheesh M. Narasimhan

Contributions

M. Z. wrote the paper with input from all co-authors. E. K., S. R., L. X., D. P., J. W., C. L. and N. O. performed the analysis. R.G., C.W. and V. M. N. supervised the analysis.

558
559   **Corresponding authors**
560
561   Correspondence to Vagheesh M. Narasimhan (vagheesh@utexas.edu)

562
563   **Ethics declarations**
564
565   The authors declare no competing interests