Title

Clinical Trajectory Patterns in IgA Nephropathy Identified by Deep Learning and Their Association with Renal Outcomes

Author

Ryunosuke Noda[1], Daisuke Ichikawa[1], Sayuri Shirai[2], Yugo Shibagaki[1], Takashi Yokoo[3], Yusuke Suzuki[4], and the J-IGACS Working Group

[1] Division of Nephrology and Hypertension, Department of Internal Medicine, St. Marianna University School of Medicine, Kanagawa, Japan

[2] Division of Nephrology and Hypertension, Department of Internal Medicine, St. Marianna University School of Medicine, Yokohama City Seibu Hospital, Kanagawa, Japan

[3] Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan

[4] Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan

Corresponding Author

Daisuke Ichikawa

The address of all authors: 2-16-1 Sugao, Miyamae-Ku, Kawasaki, Kanagawa 216-8511, Japan

Telephone number: +81-44-977-8111

Email address: ichikawa6008@gmail.com

Word Count:

Abstract: 277 words

Main Text: 3,436 words (Introduction, Methods, Results, and Discussion)

Number of Figures and Tables:

Figures: 4,

Tables: 4,

Supplemental Material: Yes

Key Points

- Deep learning on clinical trajectories revealed the heterogeneity of IgA nephropathy, identifying three distinct patient subgroups.

- These subgroups, reflecting a spectrum of progression patterns and treatment responses, had distinct long-term renal outcomes.

- This approach may provide a dynamic framework to understand clinical course, moving beyond static, single-point risk assessment.

Abstract

Background The heterogeneous clinical course of IgA nephropathy complicates risk stratification based on static, single-time-point markers. We aimed to classify these clinical trajectories using unsupervised deep learning and to validate the prognostic value of the resulting subgroups for long-term renal outcomes.

Methods We analyzed data from 873 patients with biopsy-proven IgA nephropathy from the Japan IgA Nephropathy Prospective Cohort Study (J-IGACS). A Long Short-Term Memory autoencoder was used to learn low-dimensional representations from the trajectories of hematuria, proteinuria, and estimated glomerular filtration rate (eGFR) over the first 12 months following renal biopsy. Patients were then classified using k-means clustering on these representations. The primary outcome was a 30% decline in eGFR from baseline.

Results Patients were classified into three distinct clusters based on their clinical trajectories. Cluster 1 (n=284) showed rapid improvement in both hematuria and proteinuria with a stable eGFR, and was associated with a favorable renal prognosis. Cluster 2 (n=215) was characterized by persistent severe hematuria with only a modest reduction in proteinuria, and a mild decline in eGFR. Cluster 3 (n=374), despite having mild hematuria, had the lowest baseline eGFR, which declined further during the first year and showed incomplete resolution of proteinuria. Both Cluster 2 and Cluster 3 had less favorable renal outcomes than Cluster 1. After adjusting for baseline clinicopathological factors including the Oxford classification score, membership in Cluster 2 or 3 remained an independent predictor of the primary outcome (adjusted hazard ratio, 2.12; 95% confidence interval, 1.35-3.34).

Conclusions Unsupervised deep learning successfully classified patients with IgA nephropathy into distinct clinical trajectory patterns. These data-driven patterns were independently associated with long-term renal outcomes, providing a new framework for understanding the disease's heterogeneity.

Introduction

IgA nephropathy is the most common primary glomerulonephritis worldwide and has a highly heterogeneous clinical course, ranging from mild cases with spontaneous remission to severe cases progressing to end-stage kidney disease.[1,2] This clinical diversity presents a major obstacle to accurate risk assessment and personalized medicine.[3] However, the patterns of these heterogeneous clinical courses are not well-characterized or classified. Therefore, there is a pressing need to objectively identify distinct clinical course patterns to improve risk stratification.

Historically, predicting renal outcomes in IgA nephropathy has relied on baseline clinical and pathological markers, such as the degree of proteinuria, estimated glomerular filtration rate (eGFR), and histological findings from renal biopsy.[4–6] However, these static, single-time-point assessments fail to capture dynamic prognostic information, such as treatment response or changes in disease activity over time. Although clinical practice is often guided by these dynamic changes, standardized approaches for classifying patients based on their evolving clinical patterns remain unestablished. To overcome this limitation, analyzing clinical marker trajectories after diagnosis has emerged as a promising approach.[7,8]

Previous studies using statistical models for trajectory analysis have revealed associations between proteinuria or hematuria trajectories and renal outcomes.[7,8] However, these approaches often focus on a single clinical marker or analyze multiple markers individually, limiting their ability to capture the interactive dynamics among markers. Furthermore, the application of dynamic deep learning in this area has been limited. For instance, a supervised model was developed to predict renal outcomes but was not optimized for identifying objective patient subgroups.[9]

Based on these limitations, we hypothesized that distinct patient clusters could be identified from longitudinal clinical data using an unsupervised deep learning approach. To test this

hypothesis, we employed a Long Short-Term Memory (LSTM) autoencoder to learn integrated trajectories of hematuria, proteinuria, and eGFR, followed by k-means clustering to identify patient subgroups. This study had two objectives: first, to identify and characterize distinct clinical trajectory clusters in a real-world setting to better understand disease heterogeneity, and second, to validate the association of these clusters with long-term renal outcomes and assess their incremental prognostic value over conventional baseline predictors.

Methods

Study Population

This study used data from the Japan IgA Nephropathy Prospective Cohort Study (J-IGACS), a multicenter prospective observational study conducted at 33 university hospitals and 11 regional core hospitals in Japan.[10,11] J-IGACS is a nationwide, electronic health record-based registry of patients with biopsy-proven primary IgA nephropathy. Patients were enrolled from April 1, 2005, to August 31, 2015, and were followed up through May 31, 2021. Investigators uploaded datasets to the J-IGACS website every six months.

We included patients from the J-IGACS database who met the following criteria: baseline eGFR ≥ 30 mL/min/1.73 m²; at least 12 months of follow-up; and no more than one missing value for urinary red blood cell (RBC) count, proteinuria, or eGFR at the 0, 6, and 12-month time points used for clustering. As this was an exploratory analysis using data from the J-IGACS registry, no prior sample size calculation was performed. The study was approved by the institutional review board of St. Marianna University School of Medicine (Approval No. 6589) and the ethics committees of each participating institution. The study was conducted in accordance with the Declaration of Helsinki and adhered to the Strengthening the Reporting of Observational Studies in Epidemiology guidelines.[12] All patients or their legal guardians provided written informed consent before enrollment.

Variables and Outcomes

Baseline data collected at the time of renal biopsy included demographics (age, sex), clinical parameters (Body Mass Index [BMI], mean arterial pressure [MAP], eGFR, proteinuria, urinary RBC), histopathological findings (Oxford classification score), and initial treatment information (immunosuppressive therapy, tonsillectomy, Renin-Angiotensin System (RAS) inhibitor). MAP was calculated as follows: MAP (mmHg) = 1/3 × pulse pressure + diastolic blood pressure. For patients aged ≥ 20 years, eGFR was calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation modified with a Japanese coefficient.[13] For patients < 20 years old at biopsy, eGFR was calculated using the Uemura formula[14]; subsequent measurements after a patient reached 20 years of age used the modified CKD-EPI equation for Japanese adults. Urinary RBC counts were measured using either an automated urinalysis system or manual microscopic examination of the sediment, according to the standard protocol of the institution. Proteinuria was recorded from 24-hour urine collections. The Oxford classification score[15,16] included mesangial hypercellularity (M: M0/M1), endocapillary hypercellularity (E: E0/E1), segmental glomerulosclerosis (S: S0/S1), tubular atrophy/interstitial fibrosis (T: T0/T1/T2), and cellular/fibrocellular crescents (C: C0/C1/C2), analyzed by five pathologists blinded to clinical data, with scores determined by a consensus of at least three pathologists. Initial treatment was defined as therapy initiated within one year of biopsy. Immunosuppressive therapy included oral or intravenous corticosteroids. RAS inhibitor use was defined as the initiation of an Angiotensin-Converting Enzyme inhibitor or Angiotensin II Receptor Blocker. Longitudinal data for urinary RBC, proteinuria, and eGFR at baseline, 6 months, and 12 months were used for clustering.

The primary outcome was a 30% decline in eGFR from baseline. This outcome is a widely accepted surrogate endpoint for CKD progression that provides statistical efficiency.[17–19]

Secondary outcomes for sensitivity analyses included a 40% decline in eGFR, a 50% decline in eGFR, and kidney failure (eGFR <15 mL/min/1.73 m² or initiation of kidney replacement therapy). Patients were followed from the time of renal biopsy until the loss to follow-up, or the end of the study period on May 31, 2021. Time to event was calculated in months.

Data Preprocessing and Missing Data Handling

For model input, urinary RBC counts were converted to a five-level ordinal variable based on the thresholds commonly used in Japanese clinical practice to assess hematuria severity: 1: <5/HPF (normal/remission), 2: 5-10/HPF (mild), 3: 11-20/HPF (moderate), 4: 21-50/HPF (significant), 5: >50/HPF (severe). Missing values in the time-series data for clustering (urinary RBC, proteinuria, eGFR) were imputed using last observation carried forward or next observation carried backward if no prior value existed. Continuous variables (proteinuria, eGFR) were standardized within each fold of cross-validation using the mean and standard deviation of the training data to prevent information leakage from the test data. Missing baseline data were imputed using multivariate imputation by chained equations.

Trajectory Pattern Identification using Deep Learning

To comprehensively capture the temporal changes of multiple clinical markers following diagnosis and initial therapeutic interventions, an LSTM autoencoder model was constructed using data from the first 12 months following renal biopsy. This deep learning architecture is particularly well-suited for capturing complex patterns in longitudinal data and has been applied for trajectory-based subgrouping in other kidney diseases, such as acute kidney injury and nephrotic syndrome.[20,21] The model's encoder compresses the one-year trajectories of the three biomarkers into a low-dimensional latent representation, and the decoder subsequently reconstructs the original trajectories from this representation. The ordinal urinary RBC variable

was transformed into a low-dimensional vector via an Embedding layer, concatenated with standardized proteinuria and eGFR, and fed into a single-layer LSTM encoder. A detailed description of the model architecture, hyperparameter optimization process, and specific implementation is provided in the **Supplemental Methods**. Finally, the optimized model was retrained on the entire dataset to extract a latent representation for each patient's trajectory.

k-means clustering [22] was applied to the extracted latent representations to group patients into clusters. The optimal number of clusters (K) was determined by evaluating the silhouette score and Calinski-Harabasz index for K ranging from 2 to 6. The identified clusters were visualized in a two-dimensional space using the t-distributed Stochastic Neighbor Embedding (t-SNE) method.[23]

Statistical Analysis

Continuous variables are presented as mean ± standard deviation (SD) or median (interquartile range [IQR]), and categorical variables as numbers (percentages). The trajectories for each cluster were visualized by plotting the bootstrapped mean (1000 resamples) and 95% confidence interval (CI) at each time point.

The association between the clusters and the primary outcome was visualized using Kaplan-Meier curves and compared using the log-rank test. To assess the independent prognostic value of the clusters, we constructed nested Cox proportional hazards models. Model 1 was a univariate model with cluster classification as the sole explanatory variable. Model 2 added key baseline clinical factors (age [per 10 years], MAP [per 10 mmHg]) as covariates. Model 3 added the Oxford classification scores to Model 2. Baseline urinary RBC, proteinuria, and eGFR values were excluded from the Cox models to avoid multicollinearity, as their trajectory information was incorporated into the cluster classification. The proportional hazards assumption was confirmed for all models using Schoenfeld residual tests.

To evaluate the incremental prognostic value of trajectory-based clusters over static predictors, we compared two nested multivariate Cox models: a "base model" including only static baseline predictors (age, MAP, eGFR, proteinuria, urinary RBC, and the Oxford classification score), and a "base + cluster model" that additionally included the cluster classification variable. Urinary RBC was included as dummy variables in both models. Model discrimination was assessed using the C-statistic (concordance index), reclassification using the category-free Net Reclassification Improvement (NRI), and discrimination improvement using the Integrated Discrimination Improvement (IDI). Differences in these metrics and their 95% CIs were calculated using bootstrapping (1000 resamples).

Sensitivity Analysis

We conducted several sensitivity analyses to confirm the robustness of our findings. First, to assess the impact of initial treatments, Model 4 was constructed by adding immunosuppressive therapy, tonsillectomy, and RAS inhibitor to the covariates in Model 3. Second, we repeated the survival analyses for more severe renal outcomes, specifically a 40% or 50% decline in eGFR and progression to kidney failure. Third, we substituted the k-means algorithm with a Gaussian mixture model to test the stability of the clustering results. Finally, a complete case analysis, which excluded all patients with any missing baseline values, was performed to assess the impact of our data imputation methods on the Cox model results. The analyses were performed using Python (version 3.11.13), and a two-sided p-value <0.05 was considered statistically significant. Key libraries included TensorFlow, scikit-learn, Optuna, Pandas, NumPy, Lifelines, and Matplotlib. The source code is available on GitHub (URL: https://github.com/Ryunosuke1219/iga-nephropathy-lstm-autoencoder/tree/main).

Results

Baseline Characteristics

Of the 1,130 patients in the J-IGACS cohort, 873 met the inclusion criteria and were included in the final analysis (**Figure 1**). Exclusions were due to a baseline eGFR < 30 mL/min/1.73 m² or receipt of dialysis (n=54), a follow-up duration of < 12 months (n=92), or having two or more missing values for the clustering variables (n=111).

The baseline characteristics of the study population are presented in **Table 1**. The mean age was 39.0 ± 15.7 years, 50.9% of patients were female, the mean eGFR was 77.0 ± 25.9 mL/min/1.73 m², and the median proteinuria was 0.58 g/day (IQR, 0.29-1.13). Regarding the Oxford classification, 26.1% were M1, 33.6% were E1, 72.6% were S1, 18.7% were T1/T2, and 36.7% were C1/C2. For initial treatment, 65.3% received immunosuppressive therapy, 44.8% underwent tonsillectomy, and 58.3% were treated with RAS inhibitor. During a median follow-up of 78 months (IQR, 42-96), 152 patients (17.4%) experienced the primary outcome: a 30% decline in eGFR from baseline.

Identification of Patient Clusters Based on Clinical Trajectories

Clustering was performed on latent representations extracted by the LSTM autoencoder from the first year of longitudinal data post-biopsy. Based on the silhouette score and Calinski-Harabasz index, K=3 was determined as the optimal number of clusters (**Supplemental Figure S1**). A two-dimensional t-SNE visualization showed that the three clusters were generally well-separated (**Figure 2**). All 873 patients were classified into three clusters: Cluster 1 (n=284), Cluster 2 (n=215), and Cluster 3 (n=374).

Clinical Features of the Identified Clusters

The three clusters displayed distinct clinical trajectories over the first year following renal biopsy, which were associated with baseline clinicopathological features and initial treatment

choices (**Figure 3**, **Supplemental Table S1**).

Cluster 1 was characterized by marked clinical improvement. The proportion with severe hematuria (Grade 5: > 50/HPF) decreased from 46.8% (n=133) at baseline to 0.4% (n=1) at 12 months. Similarly, mean proteinuria fell from 1.15 (95% CI, 0.88-1.37) to 0.23 g/day (95% CI, 0.20-0.25). Their eGFR, the highest at baseline, remained stable, changing negligibly from 89.5 (95% CI, 87.1-91.7) to 90.6 mL/min/1.73 m² (95% CI, 88.4-93.0).

Cluster 2 was characterized by persistent severe hematuria. This hematuria (Grade 5: > 50/HPF) persisted in a substantial portion of patients (26.5%, n=57) at 12 months. In contrast, proteinuria showed a modest reduction, with the mean decreasing from 0.90 (95% CI, 0.73-1.04) to 0.46 g/day (95% CI, 0.39-0.52). Their eGFR declined mildly from a mean of 76.6 (95% CI, 74.0-79.2) to 74.6 mL/min/1.73 m² (95% CI, 71.9-77.2).

Cluster 3 had the mildest baseline hematuria but showed the modest improvement in proteinuria, which only fell from a baseline of 1.05 (95% CI, 0.94-1.14) to 0.53 g/day (95% CI, 0.43-0.62). Their eGFR, which was already the lowest among the clusters at baseline, also declined during the first year from a mean of 67.7 (95% CI, 66.2-69.3) to 64.6 mL/min/1.73 m² (95% CI, 63.1-66.2).

These three trajectory patterns corresponded to differences in baseline characteristics (**Table 2**). Patients in Cluster 1 were younger (mean age 31.8 years), had the highest eGFR (mean 89.4 mL/min/1.73 m²), and the lowest MAP (mean 86.0 mmHg). Pathologically, this cluster had the highest proportion of crescents (C1/C2: 45.5%) and received the highest proportion of immunosuppressive therapy (81.3%) and tonsillectomy (54.6%). Patients in Cluster 2 had intermediate features regarding age (mean 40.4 years), eGFR (mean 76.6 mL/min/1.73 m²), and MAP (mean 90.4 mmHg). The proportion of immunosuppressive therapy (47.4%) and tonsillectomy (39.1%) were the lowest in this group. Patients in Cluster 3 were older (mean 43.6 years) and had the lowest eGFR at baseline (mean 67.8 mL/min/1.73 m²). Their MAP

(mean 93.0 mmHg) and proteinuria (median 0.67 g/day) were the highest. Pathologically, this cluster had the highest proportion of tubular atrophy/interstitial fibrosis (T1/T2: 24.6%) and the highest proportion of RAS inhibitor (67.6%).

Association Between Clusters and Long-Term Renal Prognosis

Kaplan-Meier analysis demonstrated that event-free survival for the primary outcome differed significantly among the three clusters (log-rank p < 0.001) (**Figure 4**). Patients in Cluster 1 had a significantly more favorable renal prognosis than those in Clusters 2 and 3.

The results of the Cox proportional hazards models are presented in **Table 3**. In the unadjusted analysis (Model 1), membership in Cluster 2 or 3 was associated with a significantly higher risk for the primary outcome compared with Cluster 1 (Hazard Ratio [HR], 2.83; 95% CI, 1.83-4.37; p < 0.001). This association remained significant in Model 2, adjusted for age and MAP (adjusted HR, 2.16; 95% CI, 1.38-3.40; p < 0.001), and in Model 3, further adjusted for the Oxford classification score (adjusted HR, 2.12; 95% CI, 1.35-3.34; p = 0.001).

To evaluate the incremental prognostic value of the trajectory-based classification over established static predictors, we compared nested Cox models (**Table 4**). Adding the cluster classification to the base model significantly improved model discrimination. The C-statistic increased from 0.745 (95% CI, 0.707-0.765) for the base model to 0.757 (95% CI, 0.722-0.776) for the base + cluster model (Δ, 0.013; 95% CI, 0.003-0.022). A significant improvement was also observed for the IDI (absolute value, 0.142; 95% CI, 0.028-0.307). However, the category-free NRI was not statistically significant (0.002; 95% CI, -0.026 to 0.262).

Sensitivity Analysis

The independent prognostic value of the clusters was maintained after adjusting for initial treatments (immunosuppressive therapy, tonsillectomy, and RAS inhibitors) (Model 4; adjusted

HR, 1.90; 95% CI, 1.19-3.02) (**Supplemental Table S2**). Furthermore, significant prognostic separation among clusters was observed for more severe renal outcomes—40% eGFR decline, 50% eGFR decline, and kidney failure (log-rank test p < 0.05 for all) (**Supplemental Figures S2-S4**). Using a Gaussian mixture model instead of k-means also yielded prognostically significant clusters (p < 0.001) (**Supplemental Figure S5**). Finally, a complete case analysis yielded consistent results in the Cox model, confirming the independent prognostic ability of the clusters (**Supplemental Table S3**).

Discussion

In this large, real-world cohort study, we applied an unsupervised deep learning approach to one-year trajectories of hematuria, proteinuria, and eGFR in patients with IgA nephropathy. This method identified three distinct patient clusters that were independently associated with long-term renal outcomes. These findings, which reveal subgroups with different clinical dynamics and risks, challenge a uniform management approach and underscore the need for more personalized therapeutic strategies.

The identified clusters likely represent distinct clinicopathological phenotypes within IgA nephropathy. Cluster 1, for example, appears to be a subgroup with highly active, treatment-responsive inflammation. This cluster was characterized by severe hematuria and a high prevalence of crescents, coupled with the highest rates of immunosuppressive therapy and tonsillectomy. This combination may have led to the improvement in hematuria and proteinuria and, consequently, a favorable long-term renal outcome. Prior studies have shown that crescents were a marker of severe, active glomerular inflammation and are associated with a high risk of progression.[24,25] Our findings suggest that this high-risk state is not fixed; instead, the prognosis can be modified by effective therapeutic intervention. This highlights the importance of the interplay between the disease's biology and treatment responsiveness, which cannot be captured

by static predictors alone.

In contrast, Cluster 3 likely represents a patient group in whom established, irreversible chronic changes are the primary driver of progression. This cluster included the oldest age, the lowest baseline eGFR, and the highest proportion of T-lesions, a robust predictor of progression.[16,26] The poor renal outcome in this cluster, despite having the highest rate of RAS inhibitor use, suggests a treatment-resistant pathophysiology for which current standard-of-care is likely insufficient. This aligns with the concept of renal fibrosis as the final common pathway of CKD progression.[27] Therefore, patients with this trajectory may be ideal candidates for novel therapeutic agents, including anti-fibrotic therapies.

Cluster 2, characterized by persistent hematuria, is also a clinically compelling subgroup. While the prognostic value of hematuria has been debated, recent studies suggest that persistent or time-averaged microscopic hematuria is an independent risk factor for progression.[8,28–30] The trajectory of Cluster 2 may therefore indicate a potential blind spot in current guidelines, which primarily focus on proteinuria as the main therapeutic target.[4] Patients in this cluster whose proteinuria might fall below therapeutic thresholds could be at risk of being undertreated, representing a potential unmet medical need.

Our approach builds upon previous work. While prior clustering studies in IgA nephropathy have provided valuable insights, they have relied on static, cross-sectional data.[31,32] Previous trajectory analyses in IgA nephropathy, however, have often focused on single markers or used statistical models that assume predefined trajectory shapes, which may not fully capture clinical complexity.[7,8] Our study addresses these limitations by employing an LSTM autoencoder, a deep learning architecture specifically designed to capture complex patterns from longitudinal data without such rigid assumptions.[33,34] While standard LSTM networks have recently shown efficacy in predicting outcomes in IgA nephropathy[9], the use of an LSTM autoencoder for data-driven subgrouping represents a novel application in this field. To the best of our knowledge,

this is the first study to leverage this approach to classify the diverse clinical trajectories of the disease, offering a new framework for patient stratification. This dynamic classification could move beyond static risk factors to identify clinically distinct phenotypes, such as treatment-resistant progressors or those with persistent inflammation, thereby enabling a more personalized approach to patient monitoring and therapeutic strategies.

This study has several limitations. First, as an observational study, it is susceptible to confounding by indication. The identified clusters could reflect a combination of disease biology and real-world treatment choices, not pure natural history. Accordingly, the main utility of our approach may be less in improving risk prediction and more in providing a framework to understand the diversity of clinical course. Second, this study was limited to a Japanese cohort where both patient backgrounds and clinical management may differ from other regions. Therefore, while our sensitivity analysis confirmed that the identified patterns are methodologically robust to the choice of clustering algorithm, the generalizability of these specific clinical trajectories requires validation in diverse, international cohorts. Third, our analysis was limited to three clinical markers and did not include the trajectories of other potentially important variables, such as blood pressure or specific medication dosages. These factors, combined with the inherent black-box nature of deep learning models and the limited temporal resolution of the data, imply that the precise factors driving the clustering remain to be fully elucidated. Future work should focus on validating these patterns in diverse, international cohorts and refining the model with additional clinical variables.

In conclusion, this study demonstrated that an unsupervised deep learning approach can identify clinically meaningful patient clusters in IgA nephropathy based on their longitudinal clinical data. This data-driven classification represents a critical step toward a more dynamic and personalized risk assessment. Ultimately, this approach could enable clinicians to better tailor monitoring and therapeutic strategies to a patient's evolving clinical course, thereby advancing

the goal of precision medicine in nephrology.

Disclosures

The authors declare no competing interests to disclose.

Funding

This study was partly supported by a Grant-in-Aid for Progressive Renal Diseases Research, Research on Rare and Intractable Disease, from the Ministry of Health, Laboure and Welfare of Japan. This research was supported by the Japan Agency for Medical Research and Development under grant JP19ek0109261.

Acknowledgments

We thank the clinicians in the J-IGACS Working Group for their kind cooperation.

The J-IGACS Working Group: Yusuke Suzuki: Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan. Takashi Yokoo: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Ryosuke Aoki: Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan. Daisuke Ichikawa: Division of Nephrology and Hypertension, Department of Internal Medicine, St. Marianna University School of Medicine, Kanagawa, Japan. Takafumi Ito: Department of Internal Medicine, Nephrology, Teikyo University School of Medicine, Teikyo University Chiba Medical Center, Chiba, Japan. Hiroyuki Ueda: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Maki Urushihara: Department of Pediatrics, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima, Japan. Ritsuko Katafuchi: Kidney Unit, National Hospital Organization, Fukuoka-Higashi Medical Center, Fukuoka, Japan, and

Division of Nephrology, Medical Corporation Houshikai, Kano Hospital, Fukuoka, Japan. Tetsuya Kawamura: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Masao Kikuchi: Division of Cardiovascular Medicine and Nephrology, Department of Internal Medicine, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan. Masao Kihara: Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan. Kentaro Koike: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Ryoko Sakaguchi: Department of Pathology, The Jikei University School of Medicine, Tokyo, Japan. Takaya Sasaki: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Satoru Sanada: Department of Nephrology, Japan Community Healthcare Organization Sendai Hospital, Sendai, Japan. Takanori Shibata: Division of Nephrology, Department of Medicine, Showa Medical University School of Medicine, Tokyo, Japan. Yuko Shima: Department of Pediatrics, Wakayama Medical University, Wakayama City, Wakayama, Japan. Akihiro Shimizu: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Akira Shimizu: Department of Analytic Human Pathology, Nippon Medical School, Tokyo, Japan. Kensuke Joh: Department of Pathology, The Jikei University School of Medicine, Tokyo, Japan. Sayuri Shirai: Division of Nephrology and Hypertension, Department of Internal Medicine, St. Marianna University School of Medicine, Yokohama City Seibu Hospital, Yokohama, Japan. Hitoshi Suzuki: Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan. Kazuo Takahashi: Department of Biomedical Molecular Sciences, School of Medicine, Fujita Health University, Nagoya, Aichi, Japan. Nobuo Tsuboi: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Yasuhiko Tomino: Asian Pacific Renal Research Promotion Office, Medical Corporation

SHOWAKAI, Shinjuku-ku, Tokyo, Japan. Shinya Nakatani: Department of Metabolism, Endocrinology and Molecular Medicine, Osaka Metropolitan University Graduate School of Medicine, Osaka, Japan. Koichi Nakanishi: Department of Child Health and Welfare (Pediatrics), Graduate School of Medicine, University of the Ryukyus, Ginowan, Okinawa, Japan. Masako Nishikawa: Center for Research Promotion, The Jikei University School of Medicine, Tokyo, Japan. Tomoya Nishino: Department of Nephrology, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan.

Yoshihito Nihei: Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan. Akinori Hashiguchi: Department of Pathology, Keio University School of Medicine, Tokyo, Japan. Hiroshi Hataya: Department of Nephrology and Rheumatology, Tokyo Metropolitan Children's Medical Center, Fuchu, Tokyo, Japan. Keita Hirano: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Yusuke Fukao: Department of Nephrology, Juntendo University Faculty of Medicine, Tokyo, Japan. Akihiro Fukuda: Department of Endocrinology, Metabolism, Rheumatology and Nephrology, Faculty of Medicine, Oita University, Oita, Japan. Shouichi Fujimoto: Department of Medical Environment Innovation, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan. Shiko Honma: Department of Pathology, The Jikei University School of Medicine, Tokyo, Japan. Keiichi Matsuzaki: Department of Public Health, Kitasato University School of Medicine, Kanagawa, Japan. Kenichiro Miura: Department of Pediatric Nephrology, Tokyo Women's Medical University, Tokyo, Japan. Yoichi Miyazaki: Division of Nephrology and Hypertension, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan. Kumiko Muta: Advanced Medical Education Center, Nagasaki University School of Medicine, Nagasaki, Japan. Takahito Moriyama: Department of Nephrology, Tokyo Medical University, Tokyo, Japan. Takashi Yasuda: Naruse Kidney Clinic, Tokyo, Japan. Yoshinari Yasuda: Department of Advanced Science in Renal-

Cardio Medicine/Nephrology, Gifu University Graduate School of Medicine, Gifu, Japan.

Shinya Yokote: Department of Nephrology, Kawaguchi Municipal Medical Center, Saitama, Japan.

Author Contributions

Conceptualization: Ryunosuke Noda, Daisuke Ichikawa.

Data curation: Ryunosuke Noda, Daisuke Ichikawa, Sayuri Shirai.

Formal analysis: Ryunosuke Noda.

Funding acquisition: Yusuke Suzuki.

Investigation: Ryunosuke Noda.

Methodology: Ryunosuke Noda.

Supervision: Daisuke Ichikawa, Sayuri Shirai, Yugo Shibagaki.

Writing – original draft: Ryunosuke Noda, Daisuke Ichikawa, Sayuri Shirai.

Writing – review & editing: Ryunosuke Noda, Daisuke Ichikawa, Sayuri Shirai, Yugo Shibagaki, Takashi Yokoo, Yusuke Suzuki.

Data Sharing Statement

The data underlying this article will be shared upon reasonable request to the corresponding author. The source code used for the analysis is publicly available on GitHub at https://github.com/Ryunosuke1219/iga-nephropathy-lstm-autoencoder/tree/main.

References

1. Stamellou E, Seikrit C, Tang SCW, et al. IgA nephropathy. *Nat Rev Dis Primer*. 2023;9(1):67. doi:10.1038/s41572-023-00476-9

2. Julian BA, Waldo FB, Rifai A, Mestecky J. IgA nephropathy, the most common glomerulonephritis worldwide. A neglected disease in the United States? *Am J Med*.

1988;84(1):129-132. doi:10.1016/0002-9343(88)90019-8

3. El Karoui K, Fervenza FC, De Vriese AS. Treatment of IgA Nephropathy: A Rapidly Evolving Field. *J Am Soc Nephrol*. 2024;35(1):103-116. doi:10.1681/ASN.0000000000000242

4. Rovin BH, Adler SG, Barratt J, et al. KDIGO 2021 Clinical Practice Guideline for the Management of Glomerular Diseases. *Kidney Int*. 2021;100(4, Supplement):S1-S276. doi:10.1016/j.kint.2021.05.021

5. Barbour SJ, Coppo R, Zhang H, et al. Evaluating a New International Risk-Prediction Tool in IgA Nephropathy. *JAMA Intern Med*. 2019;179(7):942-952. doi:10.1001/jamainternmed.2019.0600

6. Selvaskandan H, Shi S, Twaij S, Cheung CK, Barratt J. Monitoring Immune Responses in IgA Nephropathy: Biomarkers to Guide Management. *Front Immunol*. 2020;11. doi:10.3389/fimmu.2020.572754

7. Glenn DA, Carver AW, Helmuth ME, et al. Proteinuria Trajectory and Disease Progression in Children and Adults with IgA Nephropathy/Vasculitis. *Clin J Am Soc Nephrol*. Published online March 2025:10.2215/CJN.0000000707. doi:10.2215/CJN.0000000707

8. Yano Y, Nagasu H, Kanegae H, et al. Kidney outcomes associated with haematuria and proteinuria trajectories among patients with IgA nephropathy in real-world clinical practice: The Japan Chronic Kidney Disease Database. *Nephrol Carlton Vic*. 2024;29(2):65-75. doi:10.1111/nep.14250

9. Chen T, Chen T, Xu W, et al. Development and External Validation of a Multidimensional Deep Learning Model to Dynamically Predict Kidney Outcomes in IgA Nephropathy. *Clin J Am Soc Nephrol*. 2024;19(7):898. doi:10.2215/CJN.0000000000000471

10. Kawamura T, Hirano K, Koike K, et al. Associations of corticosteroid therapy and tonsillectomy with kidney survival in a multicenter prospective study for IgA nephropathy. *Sci Rep*. 2023;13(1):18455. doi:10.1038/s41598-023-45514-4

11. Koike K, Kawamura T, Hirano K, et al. Clinicopathological prognostic stratification for proteinuria and kidney survival in IgA nephropathy: a Japanese prospective cohort study. *Clin Kidney J*. 2024;17(1):sfad294. doi:10.1093/ckj/sfad294

12. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of

Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies. *Ann Intern Med*. 2007;147(8):573-577. doi:10.7326/0003-4819-147-8-200710160-00010

13. Matsuo S, Imai E, Horio M, et al. Revised Equations for Estimated GFR From Serum Creatinine in Japan. *Am J Kidney Dis*. 2009;53(6):982-992. doi:10.1053/j.ajkd.2008.12.034

14. Uemura O, Nagai T, Ishikura K, et al. Creatinine-based equation to estimate the glomerular filtration rate in Japanese children and adolescents with chronic kidney disease. *Clin Exp Nephrol*. 2014;18(4):626-633. doi:10.1007/s10157-013-0856-y

15. A Working Group of the International IgA Nephropathy Network and the Renal Pathology Society, Cattran DC, Coppo R, et al. The Oxford classification of IgA nephropathy: rationale, clinicopathological correlations, and classification. *Kidney Int*. 2009;76(5):534-545. doi:10.1038/ki.2009.243

16. Coppo R, Troyanov S, Bellur S, et al. Validation of the Oxford classification of IgA nephropathy in cohorts with different presentations and treatments. *Kidney Int*. 2014;86(4):828-836. doi:10.1038/ki.2014.63

17. Levey AS, Inker LA, Matsushita K, et al. GFR Decline as an End Point for Clinical Trials in CKD: A Scientific Workshop Sponsored by the National Kidney Foundation and the US Food and Drug Administration. *Am J Kidney Dis*. 2014;64(6):821-835. doi:10.1053/j.ajkd.2014.07.030

18. Levey AS, Gansevoort RT, Coresh J, et al. Change in Albuminuria and GFR as End Points for Clinical Trials in Early Stages of CKD: A Scientific Workshop Sponsored by the National Kidney Foundation in Collaboration With the US Food and Drug Administration and European Medicines Agency. *Am J Kidney Dis*. 2020;75(1):84-104. doi:10.1053/j.ajkd.2019.06.009

19. Coresh J, Turin TC, Matsushita K, et al. Decline in Estimated Glomerular Filtration Rate and Subsequent Risk of End-Stage Renal Disease and Mortality. *JAMA*. 2014;311(24):2518. doi:10.1001/jama.2014.6634

20. Tan Y, Huang J, Zhuang J, et al. Identifying acute kidney injury subphenotypes using an outcome-driven deep-learning approach. *J Biomed Inform*. 2023;143:104393. doi:10.1016/j.jbi.2023.104393

21. Kimura T, Yamamoto R, Yoshino M, et al. Deep learning analysis of clinical course of

primary nephrotic syndrome: Japan Nephrotic Syndrome Cohort Study (JNSCS). *Clin Exp Nephrol*. 2022;26(12):1170-1179. doi:10.1007/s10157-022-02256-3

22. MacQueen J. Some methods for classification and analysis of multivariate observations. In: *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Statistics*. Vol 5.1. University of California Press; 1967:281-298. Accessed July 7, 2025. http://projecteuclid.org/euclid.bsmsp/1200512992

23. Maaten L van der, Hinton G. Visualizing Data using t-SNE. *J Mach Learn Res*. 2008;9(86):2579-2605.

24. Trimarchi H, Barratt J, Cattran DC, et al. Oxford Classification of IgA nephropathy 2016: an update from the IgA Nephropathy Classification Working Group. *Kidney Int*. 2017;91(5):1014-1021. doi:10.1016/j.kint.2017.02.003

25. Du Y, Chen S, Wang F, et al. The Significance of Crescents on the Clinical Features and Outcomes of Primary Immunoglobin A Nephropathy. *Front Med*. 2022;9. doi:10.3389/fmed.2022.864667

26. Lv J, Shi S, Xu D, et al. Evaluation of the Oxford Classification of IgA nephropathy: a systematic review and meta-analysis. *Am J Kidney Dis*. 2013;62(5):891-899. doi:10.1053/j.ajkd.2013.04.021

27. Duffield JS. Cellular and molecular mechanisms in kidney fibrosis. *J Clin Invest*. 2014;124(6):2299-2306. doi:10.1172/JCI72267

28. Sevillano AM, Gutiérrez E, Yuste C, et al. Remission of Hematuria Improves Renal Survival in IgA Nephropathy. *J Am Soc Nephrol*. 2017;28(10):3089-3099. doi:10.1681/ASN.2017010108

29. Yu GZ, Guo L, Dong JF, et al. Persistent Hematuria and Kidney Disease Progression in IgA Nephropathy: A Cohort Study. *Am J Kidney Dis*. 2020;76(1):90-99. doi:10.1053/j.ajkd.2019.11.008

30. He P, Wang H, Huang C, He L. Hematuria was a high risk for renal progression and ESRD in immunoglobulin a nephropathy: a systematic review and meta-analysis. *Ren Fail*. 2021;43(1):488-499. doi:10.1080/0886022X.2021.1879852

31. Wang Y, Wen Q, Lian X, et al. Machine learning-based unsupervised phenotypic clustering analysis of patients with IgA nephropathy: Distinct therapeutic responses of different groups. *Chin Med J (Engl)*. Published online February 2025.

doi:10.1097/CM9.0000000000003422

32. Tan J, Yang R, Xiao L, et al. Risk Stratification in Immunoglobulin A Nephropathy Using Network Biomarkers: Development and Validation Study. *J Med Internet Res*. 2025;27(1):e65563. doi:10.2196/65563

33. Yin C, Liu R, Zhang D, Zhang P. Identifying Sepsis Subphenotypes via Time-Aware Multi-Modal Auto-Encoder. In: *Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*. KDD '20. Association for Computing Machinery; 2020:862-872. doi:10.1145/3394486.3403129

34. Liu P, Sun X, Han Y, He Z, Zhang W, Wu C. Arrhythmia classification of LSTM autoencoder based on time series anomaly detection. *Biomed Signal Process Control*. 2022;71:103228. doi:10.1016/j.bspc.2021.103228

Figure Legends

Figure 1. Flowchart of patient selection.

Abbreviations: eGFR, estimated Glomerular Filtration Rate; J-IGACS, Japan IgA Nephropathy Prospective Cohort Study; RBC, red blood cells.

Figure 2. Visualization of patient clusters using t-SNE.

The t-distributed Stochastic Neighbor Embedding (t-SNE) plot visualizes the low-dimensional latent representations learned by the Long Short-Term Memory (LSTM) autoencoder from the first-year clinical trajectories. Each point represents a single patient, colored according to their assigned cluster: Cluster 1 (blue), Cluster 2 (orange), and Cluster 3 (green).

Figure 3. Clinical marker trajectories over the first 12 months for each cluster.

(a) Hematuria trajectories are shown as stacked area plots, representing the proportion of patients within each of the five hematuria grades over 12 months for each cluster. The grades are defined by urinary red blood cells per high-power field (HPF): Grade 1 (<5/HPF), Grade 2 (5-10/HPF), Grade 3 (11-20/HPF), Grade 4 (21-50/HPF), and Grade 5 (>50/HPF). (b) Proteinuria and estimated Glomerular Filtration Rate (eGFR) trajectories are shown for each cluster. Solid lines represent the bootstrapped mean values, and the surrounding shaded areas indicate the 95% confidence intervals.

Figure 4. Kaplan-Meier curves for the primary outcome by cluster.

Kaplan-Meier curves illustrating the event-free survival proportion for the primary outcome, defined as a $\geq 30\%$ decline in estimated Glomerular Filtration Rate (eGFR) from baseline. The p-value was calculated using the log-rank test and showed a significant difference among the clusters ($p < 0.001$).

Table 1. Baseline characteristics of the study population

Characteristic	Overall (n=873)	Missing, n (%)
Age (years)	39.0 ± 15.7	0 (0.0)
Female	444 (50.9%)	0 (0.0)
Body mass index (kg/m²)	22.2 ± 3.8	22 (2.5)
Mean arterial pressure (mmHg)	90.0 ± 13.5	51 (5.8)
eGFR (ml/min/1.73 m²)	77.0 ± 25.9	2 (0.2)
Proteinuria (g/day)	0.58 [0.29-1.13]	2 (0.2)
Urinary RBC (cells/HPF), n (%)		10 (1.1)
Grade 1: <5	128 (14.7%)	
Grade 2: 5-10	120 (13.7%)	
Grade 3: 11-20	165 (18.9%)	
Grade 4: 21-50	197 (22.6%)	
Grade 5: >50	253 (29.0%)	
M, n (%)		37 (4.2)
M1	228 (26.1%)	
E, n (%)		37 (4.2)
E1	293 (33.6%)	
S, n (%)		37 (4.2)
S1	634 (72.6%)	
T, n (%)		37 (4.2)
T1	144 (16.5%)	
T2	19 (2.2%)	
C, n (%)		37 (4.2)
C1	313 (35.9%)	
C2	7 (0.8%)	
Immunosuppressive therapy	570 (65.3%)	0 (0.0)
Tonsillectomy	391 (44.8%)	0 (0.0)
RAS inhibitor	509 (58.3%)	0 (0.0)

Data are presented as mean ± standard deviation (SD), median [interquartile range (IQR)], or n (%). Abbreviations: BMI, Body Mass Index; eGFR, estimated Glomerular Filtration Rate; MAP, Mean Arterial Pressure; RAS, Renin-Angiotensin System; HPF, high-power field. The Oxford classification score consisted of mesangial hypercellularity (M0, mesangial score ≤ 0.5; M1, mesangial score > 0.5), endocapillary hypercellularity (E0, absent; E1,present), segmental glomerulosclerosis (S0, absent; S1, present), tubular atrophy/interstitial fibrosis (T0, ≤ 25%; T1, 26–50%; T2, > 50%), and crescent (C1, cellular/fibrocellular crescents present in at least one glomerulus; C2, cellular/fibrocellular crescents present in > 25% glomerulus).

Table 2. Comparison of baseline characteristics among the identified clusters

Characteristic	Cluster 1 (n=284)	Cluster 2 (n=215)	Cluster 3 (n=374)
Age (years)	31.8 ± 14.6	40.4 ± 16.0	43.6 ± 14.4
Female, n (%)	161 (56.7)	104 (48.4)	179 (47.9)
BMI (kg/m²)	21.3 ± 4.0	22.5 ± 3.5	22.8 ± 3.6
MAP (mmHg)	86.0 ± 12.4	90.4 ± 14.2	93.0 ± 13.2
eGFR (mL/min/1.73 m²)	89.4 ± 26.5	76.6 ± 26.0	67.8 ± 21.1
Proteinuria (g/day)	0.54 [0.28-0.99]	0.52 [0.26-1.03]	0.67 [0.33-1.30]
Urinary RBC (cells/HPF), n (%)			
Grade 1: <5	2 (0.7)	3 (1.4)	123 (32.9)
Grade 2: 5-10	7 (2.5)	9 (4.2)	104 (27.8)
Grade 3: 11-20	46 (16.2)	26 (12.1)	93 (24.9)
Grade 4: 21-50	95 (33.5)	60 (27.9)	42 (11.2)
Grade 5: >50	133 (46.8)	113 (52.6)	7 (1.9)
Oxford Classification, n (%)			
M1	67 (23.6)	61 (28.4)	100 (26.7)
E1	107 (37.7)	68 (31.6)	118 (31.6)
S1	205 (72.2)	149 (69.3)	280 (74.9)
T1/T2	31 (10.9)	40 (18.6)	92 (24.6)
C1/C2	129 (45.5)	71 (33.0)	120 (32.1)
Initial Treatment, n (%)			
Immunosuppressive therapy	231 (81.3)	102 (47.4)	237 (63.4)
Tonsillectomy	155 (54.6)	84 (39.1)	152 (40.6)
RAS inhibitor	118 (41.5)	138 (64.2)	253 (67.6)

Data are presented as mean ± SD, median [IQR], or n (%). Abbreviations: BMI, Body Mass Index; eGFR, estimated Glomerular Filtration Rate; MAP, Mean Arterial Pressure; RAS, Renin-Angiotensin System; RBC, Red Blood Cell; HPF, high-power field.

Table 3. Cox proportional hazards models for the primary outcome (30% decline in eGFR)

	Model 1			Model 2			Model 3		
	HR	95% CI	p-value	HR	95% CI	p-value	HR	95% CI	p-value
Cluster (2,3 vs 1)	2.83	1.83-4.37	<0.001	2.16	1.38-3.40	<0.001	2.12	1.35-3.34	0.001
Age (per 10 years)				1.25	1.11-1.40	<0.001	1.23	1.09-1.39	<0.001
MAP (per 10 mmHg)				1.09	0.96-1.24	0.167	1.05	0.92-1.19	0.467
M1							1.48	1.03-2.11	0.033
E1							1.02	0.70-1.50	0.903
S1							1.1	0.71-1.70	0.666
T1/2							1.65	1.15-2.36	0.007
C1/2							0.95	0.64-1.39	0.78

Model 1: Univariable model with cluster classification.

Model 2: Adjusted for variables in Model 1 plus age and MAP.

Model 3: Adjusted for variables in Model 2 plus Oxford classification scores.

Abbreviations: eGFR, estimated Glomerular Filtration Rate; HR, Hazard Ratio; CI, Confidence Interval; MAP, Mean Arterial Pressure.

Table 4. Improvement in model performance for predicting 30% eGFR decline after adding cluster classification

	C statistic (95%CI)	Change in C statistic (95%CI)	Category-free NRI (95%CI)	Absolute IDI (95%CI)	Relative IDI (95%CI)
Base model	0.745 (0.707-0.765)				
Base + Cluster model	0.757 (0.722-0.776)	0.013 (0.003-0.022)	0.002 (-0.026-0.262)	0.142 (0.028-0.307)	0.151 (0.028-0.347)

The Base model included static baseline factors (age, MAP, eGFR, proteinuria, urinary RBC, and the Oxford classification score). The Base +Cluster model added the cluster classification (Clusters 2 and 3 vs Cluster 1) to the base model. Abbreviations: CI, Confidence Interval; NRI, Net Reclassification Improvement; IDI, Integrated Discrimination Improvement.

Figure 1

Figure 2

Figure 3

Figure 4