1   **Comprehensive Thyroid Assessment in a Prospective Japanese Cohort:**

2   **Epidemiology, Determinants, and Ultrasonographic Reference Ranges**

Satoko Shimazu-Kuwahara[1], Ichiro Yamauchi[2*], Sachiko Kawashima[1], Makiko Tatsumi[1],

Takuro Hakata[2], Yoriko Sakane[2,3], Masahiro Yakami[1], Kosuke Inoue[4], Daisuke Yabe[2],

Mayumi Inoue[1]

*Corresponding author

[1]Preemptive Medicine and Lifestyle Disease Research Center, Kyoto University Hospital

[2]Department of Diabetes, Endocrinology and Nutrition, Kyoto University Graduate School of

Medicine

[3]Sugawa Clinic

[4]Health Promotion and Human Behavior, School of Public Health, Kyoto University

Graduate School of Medicine

Address correspondence to:

Ichiro Yamauchi

Department of Diabetes, Endocrinology and Nutrition

Kyoto University Graduate School of Medicine

54 Kawaharacho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

Phone: +81-75-751-3560

E-mail: ichiroy@kuhp.kyoto-u.ac.jp

Funding

This study was supported by Medical Corporation Midtown Clinic and Japan Foundation for

Applied Enzymology.

Disclosure summary

SSK, SK, MT, MY, and MI are affiliated with the Preemptive Medicine and Lifestyle Disease

Research Center, Kyoto University Hospital, which is funded by Medical Corporation

Midtown Clinic. IY and TH serve as advisors to the Center. SSK, SK, MT, MY, and MI received

research funding from Medical Corporation Midtown Clinic.

ABSTRACT

Context: A comprehensive evaluation of thyroid disease and health through multimodal assessment is warranted.

Objective: To clarify the epidemiology and clinical significance of abnormal findings on thyroid examinations in the general population.

Design: Prospective cohort study conducted between April 1, 2016, and December 31, 2021.

Setting: Japanese adults undergoing health checkups at the Preemptive Medicine and Lifestyle Disease Research Center, Kyoto University Hospital.

Main Outcome Measures: All subjects underwent thyroid function tests, ultrasonography, and ^{18}F-fluorodeoxyglucose-positron emission tomography (FDG-PET); anti-thyroperoxidase antibody (TPOAb) titers were measured in a subset of subjects.

Results: In the original cohort of 4,407 subjects (2,643 males and 1,764 females), the prevalence of thyroid dysfunction, increased blood flow on ultrasonography, diffuse thyroid FDG uptake, and thyroid nodules was 5.81%, 2.45%, 3.43%, and 39.71%, respectively; all were more frequent in females. Among 2,420 subjects with TPOAb measurements, TPOAb positivity was 7.19% and was significantly associated with thyroid dysfunction only at titers ≥ 128 IU/mL. Multivariate analyses identified age, sex, and thyroid volume as major determinants of thyroid function. Using data from 1,840 subjects without any thyroid abnormalities, we established sex-specific reference ranges for thyroid dimensions and found their correlations with age and body size.

Conclusions: This cohort provides epidemiological and physiological insights into thyroid health by integrating findings from thyroid function tests, ultrasonography, FDG-PET, and TPOAb measurements in the general population. Furthermore, the present study highlights associations across abnormal findings, relationships within thyroid physiology, and clinical relevance of high-titer TPOAb.

Introduction

Thyroid diseases are common in the general population. Thyroid dysfunction is clinically important because, if left untreated, it can lead to multiple comorbidities. In the United Kingdom, the incidence of hypothyroidism is 0.41% per year in females and 0.06% per year in males, whereas that of hyperthyroidism is 0.08% per year in females and negligible in males (1). In the United States, the prevalence of overt and subclinical hypothyroidism is 0.3% and 4.3%, respectively, whereas that of overt and subclinical hyperthyroidism is 0.5% and 0.7% (2). When thyroid dysfunction is detected, its cause should be investigated using additional tests. Hashimoto's disease, the most frequent cause of hypothyroidism, is diagnosed by detecting thyroid autoantibodies, such as anti-thyroperoxidase (TPOAb) and anti-thyroglobulin antibodies (TgAb). In the United States, TPOAb and TgAb positivity has been reported in 13.0% and 11.5% of adults, respectively (2).

Thyroid ultrasonography is a key imaging modality for the diagnosis of thyroid diseases. Diffuse hypervascularity on color Doppler, known as thyroid inferno, is a hallmark of Graves' disease, although increased blood flow can also be observed in some cases of Hashimoto's thyroiditis (3). In addition, ultrasonography is essential for the detailed evaluation of thyroid nodules. The prevalence of thyroid nodules is high and increases with age, reaching nearly 50% when assessed using ultrasonography (4). ^{18}F-fluorodeoxyglucose (FDG) positron emission tomography (PET) is widely used in oncological evaluations, and the thyroid gland occasionally exhibits FDG uptake. Diffuse thyroid FDG uptake suggests a high likelihood of Hashimoto's disease (5,6). Although the prevalence of diffuse thyroid FDG uptake has been reported (5,7), its clinical significance remains unclear.

Although thyroid diseases can be evaluated using multiple modalities, data on the relationships among these examinations are limited, particularly regarding ultrasonography and FDG-PET. The Fukushima Health Management Survey reported estimated thyroid

volume by ultrasonography from infancy through adolescence (8), whereas fewer data are

available for adults (9,10). Moreover, in adults, the high prevalence of Hashimoto's

thyroiditis and thyroid nodules complicates the identification of truly disease-free reference

populations. Therefore, a cohort restricted to individuals with healthy thyroid glands is

warranted to investigate thyroid physiology.

In 2016, we launched a prospective cohort study of Japanese adults in a health

checkup setting. This cohort featured comprehensive phenotyping. For thyroid assessment,

all subjects underwent thyroid function tests, thyroid ultrasonography, and FDG-PET;

measurement of TPOAb was incorporated midway through the study. We aimed to clarify the

epidemiology and clinical significance of abnormal findings detected in these thyroid

examinations among the general population.

Methods

Subjects

A flow diagram of the enrollment process is shown in Figure 1. From April 1, 2016,

to December 31, 2021, we enrolled 4,651 adults who underwent their initial health checkup at

the Preemptive Medicine and Lifestyle Disease Research Center, Kyoto University Hospital.

We excluded 5 individuals without blood tests and 111 who did not provide informed

consent. In addition, 128 individuals were excluded because of current or past thyroid

diseases, including Graves' disease (n = 37), levothyroxine replacement for hypothyroidism

(n = 37), thyroid cancer (n = 36), surgery for thyroid nodules (n = 13), and a history of

subacute thyroiditis (n = 5). A total of 4,407 subjects met the inclusion criteria for this study,

and we defined this group as the Original cohort (Figure 1). As TPOAb measurements began

partway through the study, we additionally analyzed the TPOAb cohort consisting of 2,420

subjects with available TPOAb results (Figure 1). A major strength of this study is the strict

definition of subjects without thyroid abnormalities based on multimodal thyroid assessment.

Accordingly, we established the Thyroid-healthy cohort by excluding subjects with abnormal

TSH levels, TPOAb positivity, increased blood flow, diffuse FDG uptake, or thyroid nodules

(Figure 1). The study was approved by the Ethics Committee of Kyoto University Hospital

(approval numbers R0619 and R3604), and written informed consent was obtained from all

subjects.

Evaluation

Serum levels of TSH, free T4 (fT4), and free T3 (fT3) were measured using

electrochemiluminescence immunoassays (Elecsys; Roche Diagnostics, Mannheim,

Germany). Reference ranges were as follows: TSH, 0.500–5.000 µIU/mL; fT4, 0.880–1.620

ng/dL; and fT3, 2.33–4.00 pg/mL. TPOAb measurements were initiated on January 1, 2018,

and were performed using an Elecsys assay (reference: < 16 IU/L). Thyroid ultrasonography

was performed using an Aplio 500 system (Canon Medical System Corporation, Otawara,

Japan) equipped with a linear transducer (PLT-1005BT; 6-cm footprint). We measured the

isthmus thickness and, for each lobe, the transverse, anteroposterior, and longitudinal

diameters. Lobe volume was calculated as the transverse diameter × anteroposterior diameter

× longitudinal diameter × π/6, and total thyroid volume was obtained by summing volumes of

both lobes. Blood flow was evaluated using color Doppler ultrasonography. The entire gland

was systematically surveyed for nodules and cysts. Ultrasound cine loops were reviewed by a

second sonographer for quality assurance. For FDG-PET, we focused on diffuse thyroid

uptake and did not analyze focal uptake confined to nodules. FDG-PET images were

independently reviewed by board-certified radiologists.

Statistical analyses

All statistical analyses were performed using JMP Pro version 18.0 (SAS Institute

Inc., NC, USA). Unless otherwise specified, values are presented as the mean ± SD. As TSH

values followed a log-normal distribution, statistical analyses were performed using log-

transformed values. Continuous variables were compared using Student's *t*-test, and

categorical variables using Fisher's exact test. For trend analyses, we assigned the median

value of each ordered category and modeled it as a continuous variable in regression

analyses. Multivariable linear regression was used to assess independent associations.

Pearson's correlation coefficients were calculated to evaluate correlations between variables.

Significant differences were considered as $p < 0.05$.

Results

Prevalence of abnormal findings on thyroid examinations

The characteristics of the Original cohort are summarized in Table 1. The mean age

was 52.9 ± 11.3 years; 2,643 (59.97%) were males and 1,764 (40.03%) were females. The

serum levels of TSH, fT4, and fT3 were 1.952 ± 2.245 μIU/mL, 1.288 ± 0.200 ng/dL, and

3.08 ± 0.58 pg/mL, respectively. Significant sex-related differences were observed (Table 1).

Males were slightly older and had larger anthropometric measurements than females. TSH

levels were slightly lower in males than in females (1.859 vs. 2.091 μIU/mL, $p = 0.0001$),

whereas fT4 and fT3 levels were higher in males (fT4, 1.313 vs. 1.250 ng/dL, $p < 0.0001$;

fT3, 3.19 vs. 2.93 pg/mL, $p < 0.0001$).

Thyroid dysfunction was identified in 256 subjects (5.81%) (Figure 2A, Table 1).

The most common pattern was subclinical hypothyroidism (n = 118; 2.68%), followed by

subclinical thyrotoxicosis (n = 104; 2.36%), overt thyrotoxicosis (n = 27; 0.61%), and overt

hypothyroidism (n = 7; 0.16%) (Figure 2B, Table 1). The characteristics of each group,

categorized by thyroid dysfunction pattern, are shown in Table 2. Overt thyrotoxicosis was

more common in younger subjects, whereas subclinical hypothyroidism predominated among

older subjects and was more frequent in females.

Increased blood flow on thyroid ultrasonography was detected in 108 subjects

(2.45%) and was significantly more common in females (4.14%) than in males (1.32%)

(Figure 2C, Table 1). Diffuse thyroid FDG uptake was observed in 151 subjects (3.43%) and

was likewise more common in females (6.35%) than in males (1.48%) (Figure 2D, Table 1).

Both increased blood flow and diffuse FDG uptake were often present in the subjects with

overt thyroid dysfunction or subclinical hypothyroidism, especially those with overt

hypothyroidism (Table 2). Thyroid nodules were found in 1,750 subjects (39.71%) (Figure

2E, Table 1). Based on the maximum diameter, 667 subjects (15.14%) had nodules < 5.0 mm,

705 (16.00%) had nodules 5.0–9.9 mm, 315 (7.15%) had nodules 10.0–19.9 mm, and 63

(1.43%) had nodules ≥ 20.0 mm (Figure 2F, Table 1). Females had a higher prevalence of

nodules ≥ 5.0 mm than males (33.79% vs. 18.43%) (Table 1). The prevalence of nodules

increased with age, whereas the prevalence of thyroid dysfunction, increased blood flow, and

diffuse FDG uptake did not show significant age-related variations (Supplemental Table 1).

Association of clinical characteristics and abnormal findings with thyroid function

We subsequently assessed the relationship of clinical characteristics and abnormal

findings with thyroid function. Although fT4 levels showed a significant correlation with log-

transformed TSH levels ($r = -0.3829$, $p < 0.0001$) (Figure 3A), we excluded subjects whose

thyroid function values were outside the main distribution for subsequent analyses (Figure

3B): those with TSH < 0.1 μIU/mL (n = 18), TSH > 10 μIU/mL (n = 13), and one subject

with syndrome of inappropriate secretion of TSH (SITSH, n = 1). In this dataset restricted to

subnormal thyroid function, older age was associated with higher TSH levels in both sexes

(Figure 3C, Supplemental Table 2). Sex differences were observed in fT4 and fT3 trends: in

males, both hormones decreased with age, whereas in females, only a slight age-related

decline in fT3 was observed (Figure 3D and 3E, Supplemental Table 2). Height showed only

a minor association with thyroid function, whereas body weight correlated positively with

fT3 levels (Supplemental Figure 1, Supplemental Table 3 and 4). Body mass index (BMI)

analysis showed that lean subjects had lower fT3 levels and obese subjects had higher fT3

levels in both sexes; no BMI-related trends were observed for TSH or fT4 (Figure 3F–3H,

Supplemental Table 5).

We then examined the relationship between abnormal findings and thyroid function.

Subjects with increased blood flow had higher TSH and lower fT4 levels, with no changes in

fT3 levels (Figure 4A–4C, Supplemental Table 6). Diffuse FDG uptake was also associated

with higher TSH and lower fT4 levels (Figure 4D–4F, Supplemental Table 6). Regarding

thyroid nodules, only minor size-related trends were observed; nodules $\geq$ 20 mm were

associated with lower TSH levels in females (Figure 4G–4I, Supplemental Table 7).

Moreover, the estimated thyroid volume was analyzed using ultrasonography. Due to

technical limitations, mainly the difficulty in measuring the longitudinal diameter, data were

unavailable for 183 subjects, leaving 4,192 evaluable cases. Thyroid volume was negatively

correlated with TSH and positively correlated with both fT4 and fT3 levels (Figure 4J–4L,

Supplemental Table 8).

Association between TPOAb titers and thyroid function

We then analyzed the TPOAb cohort (n = 2,420), whose characteristics were

similar to those of the Original cohort (Supplemental Table 9). TPOAb positivity was found

in 174 subjects (7.19%), with a higher prevalence in females (96 of 938 subjects, 10.23%)

than in males (78 of 1,482 subjects, 5.26%; $p < 0.0001$) (Figure 5A). Subjects with TPOAb

positivity more frequently exhibited abnormal findings on thyroid examinations than those

without TPOAb; thyroid dysfunction (13.22% vs. 4.94%, $p < 0.0001$), increased blood flow

(28.16% vs. 1.11%, $p < 0.0001$), and diffuse FDG uptake (28.74% vs. 1.25%, $p < 0.0001$)

(Figure 5B–5D, Supplemental Table 10). The prevalence of thyroid nodules did not differ

between TPOAb-positive and TPOAb-negative subjects (40.80% vs. 39.45%, $p = 0.7478$)

(Figure 5E, Supplemental Table 10).

We subsequently examined whether the TPOAb titer influenced these associations.

Thyroid dysfunction, particularly hypothyroidism, was more frequent among subjects with

TPOAb titers $\geq$ 128 IU/mL (Table 3). The prevalence of increased blood flow and diffuse

FDG uptake also increased in a titer-dependent manner (Table 3). To further clarify the

relationship between TPOAb positivity and thyroid function, we excluded subjects whose

thyroid function values were outside the main distribution (Figure 5F), as in Figure 3B: TSH

< 0.1 μIU/mL (n = 11), TSH > 10 μIU/mL (n = 10), or SITSH (n = 1). In this dataset,

TPOAb-positive subjects had higher TSH levels than TPOAb-negative subjects, whereas fT4

and fT3 levels remained unchanged (Figure 5G–5I, Supplemental Table 11). Analysis by titer

showed that TSH elevation was observed only in subjects with TPOAb ≥ 128 IU/mL, with no

corresponding changes in fT4 or fT3 levels (Supplemental Figure 2, Supplemental Table 11).

From these analyses, we identified age, sex, BMI, increased blood flow, diffuse FDG

uptake, thyroid volume, and TPOAb titers ≥ 128 IU/mL as potential determinants of thyroid

function. To assess their relative contributions, we conducted multivariable analyses within

the TPOAb cohort after excluding subjects lacking thyroid volume data (n = 130), those with

nodules ≥ 20 mm, and those with outlying thyroid function values (TSH < 0.1 μIU/mL, n = 6;

TSH > 10 μIU/mL, n = 10; SITSH, n = 1). Among the 2,239 subjects analyzed, Model 1,

which included only demographic and anthropometric variables, showed an age-related

increase in TSH and decreases in fT4 and fT3 levels. Males had lower TSH and higher fT4

and fT3 levels than females, and BMI showed a positive correlation exclusively with fT3

levels (Table 4). Model 2, which incorporated all identified factors, reproduced the age-

related changes, but the association between TSH levels and sex was no longer significant.

BMI was weakly associated with TSH and fT3 levels in both sexes. Among the factors

obtained from thyroid examinations, thyroid volume was negatively associated with TSH and

positively associated with fT4 and fT3 levels; increased blood flow and TPOAb ≥ 128 IU/mL

were linked to higher TSH levels; and increased blood flow and diffuse FDG uptake were

associated with lower fT4 levels. In summary, age and sex remain the dominant determinants

of thyroid function, while ultrasonographic findings have a significant influence.

Reference ranges and factors associated with thyroid size

We further investigated factors influencing thyroid size using the Thyroid-healthy

cohort (n = 1,840) without any thyroid abnormalities. The characteristics of this cohort are

presented in Table 5. Subjects with nodules < 10 mm were included, given their high

prevalence (33.37%). All thyroid size parameters, including the isthmus thickness, the

transverse, anteroposterior, and longitudinal diameters of each lobe, were larger in males than

in females (Table 5). Several sex-independent correlations were observed: the longitudinal

diameter was negatively correlated with age (Figure 6A–6D, Supplemental Table 12) and

positively correlated with height (Figure 6E–6H, Supplemental Table 13). Body weight and

BMI were positively correlated with the isthmus thickness and the transverse and

anteroposterior diameters (Figure 6I–6L, Supplemental Figure 3, Supplemental Table 14 and

15). Consequently, thyroid volume was positively correlated with height, body weight, and

BMI (Supplemental Figure 4, Supplemental Table 12–15).

As thyroid nodules measuring 5.0–9.9 mm could potentially affect lobe size in both

sexes (Supplemental Table 16), we performed an additional analysis by excluding 296

subjects with nodules in this size range. Multiple regression analysis confirmed that sex

remained a significant determinant of thyroid size, independent of height and body weight

(Supplemental Table 17). The longitudinal diameter was significantly correlated with age and

height. The transverse and anteroposterior diameters were positively correlated with body

weight. Thyroid volume was strongly correlated with body weight, but its association with

height was no longer significant.

Finally, we calculated reference ranges for thyroid size using data from 1,544

Thyroid-healthy cohort subjects without nodules $\geq$ 5.0 mm (Table 6). Consistent with the

findings in Table 5, the left lobe was smaller than the right lobe in both sexes. For the right

lobe, the transverse diameter measured 16.6 $\pm$ 2.8 mm in males and 14.1 $\pm$ 2.4 mm in

females; for the left lobe, it was 15.9 $\pm$ 2.8 mm and 14.3 $\pm$ 2.2 mm, respectively (p < 0.0001

for males, p = 0.1904 for females). The anteroposterior diameter of the right lobe was 14.7 $\pm$

2.5 mm in males and 12.1 $\pm$ 2.1 mm in females, while the left lobe measured 12.9 $\pm$ 2.6 mm

and 10.4 ± 2.4 mm, respectively (p < 0.0001 for both sexes). The longitudinal diameter of the

right lobe was 50.2 ± 5.3 mm in males and 48.5 ± 5.6 mm in females; for the left lobe, 47.9 ±

5.7 mm and 45.7 ± 6.3 mm, respectively (p < 0.0001 for both sexes). Thus, the left lobe was

consistently smaller than the right lobe in both sexes, particularly in the anteroposterior and

longitudinal diameters.

Discussion

In this prospective cohort study of the Japanese general population, we

comprehensively analyzed data on thyroid examinations, including thyroid function tests,

ultrasonography, FDG-PET, and TPOAb measurements. In addition to documenting the

prevalence of thyroid dysfunction, we present epidemiological findings. As expected, females

were more susceptible to thyroid dysfunction and other abnormalities than males. We also

identified factors associated with thyroid function, demonstrating that age, sex, and thyroid

volume were dominant determinants. By rigorously excluding subjects with thyroid

abnormalities, we defined the Thyroid-healthy cohort and established reference ranges for

thyroid size. Furthermore, we identified sex differences and demonstrated that thyroid

dimensions correlated with age and body size.

Given that racial differences may affect thyroid function and disease prevalence, our

East Asian cohort provides valuable evidence. A meta-analysis of European epidemiological

data reported a prevalence of 4.59% for subclinical hypothyroidism, 0.62% for overt

hypothyroidism, 0.50% for subclinical hyperthyroidism, and 0.35% for overt

hyperthyroidism (11). In the United States, the National Health and Nutrition Examination

Survey (NHANES; 2007–2012) found prevalence of 4.3%, 0.33%, 3.20%, and 0.2%,

respectively (12). In our cohort, the corresponding values were 2.68%, 0.16%, 2.36% and

0.61%. Another Japanese cohort study of healthy individuals reported an overall prevalence

of hypothyroidism of 1.39% (13). These comparisons suggest that hypothyroidism is less

common and thyrotoxicosis is more common in Japan than in Europe or the United States.

Japan is considered an iodine-sufficient to iodine-excess country (14), and excessive iodine

intake is generally associated with an increased risk of developing hypothyroidism (15). In

this context, the observed differences are unlikely to be explained by iodine status and may

instead reflect other factors, such as racial or genetic differences.

298   Our cohort also provides epidemiological insights into FDG-PET, ultrasonography,

and TPOAb titers. Increased blood flow on ultrasonography was observed in 2.45% of the

subjects, with a marked sex difference (1.32% in males vs. 4.14% in females). Schulz *et al.*

reported that increased blood flow can occur in hypothyroidism and is strongly associated with

elevated serum TSH levels (16). To date, its significance has not been evaluated in population-

based cohorts without known thyroid diseases. While ultrasonography is subject to inter-

operator variability, the relevance of our findings is supported by their association with thyroid

dysfunction, which was more common in subjects with increased blood flow. Furthermore,

increased blood flow was associated with TPOAb positivity, which is a well-established risk

factor for hypothyroidism. Taken together, in settings where thyroid ultrasonography is readily

available, the detection of increased blood flow may facilitate early suspicion of Hashimoto's

disease and hypothyroidism.

310   While most previous studies have focused on focal thyroid FDG uptake, we

specifically examined diffuse FDG uptake to clarify its significance in Hashimoto's disease

and thyroid function. Previous studies have reported diffuse FDG uptake in 3.3% of 1102 cases

(5) and 1.8% of 2594 cases (7). In our study, diffuse FDG uptake showed patterns similar to

those of increased blood flow, with an overall prevalence of 3.43% (1.48% in males and 6.35%

in females), which we report here for the first time. Diffuse FDG uptake was more common in

subjects with thyroid dysfunction, particularly overt hypothyroidism, consistent with earlier

reports identifying diffuse FDG uptake as a surrogate marker for Hashimoto's disease (5,6).

Moreover, diffuse FDG uptake has been reported as a predictive factor for immune-related

adverse events (irAEs) involving the thyroid gland in patients treated with immune checkpoint

inhibitors (17,18). Given that female patients are more susceptible to thyroid irAEs (17,19),

the high prevalence of diffuse FDG uptake in females may contribute to this increased risk.

A meta-analysis reported a prevalence of thyroid nodules of 24.83% detected on

thyroid ultrasonography (20). In our cohort, the prevalence of nodules ≥ 5.0 mm was similar

(24.58%). Data on size-specific prevalence are limited; for example, in the Epidemiological

Study of Health Effects in Fukushima Emergency Workers, nodules > 5.0 mm were observed

in 15.5% of participants (21). The difference from our cohort is likely attributable to sex

composition, as the Fukushima study population was predominantly male. In our cohort, the

prevalence of nodules ≥ 5.0 mm was lower in males (18.43%) than in females (33.79%).

Consistent with previous studies (22-24), we observed a higher prevalence in females and an

age-related increase in the frequency of nodules. Moreover, our size-stratified analyses

revealed that relatively large nodules were associated with lower TSH levels in females.

TPOAb was detected in 7.19% of the subjects in our cohort, with a higher prevalence

in females (10.23%) than in males (5.26%). In NHANES, the prevalence was 11.1% (7.2% in

males and 14.8% in females) (12), and in a Japanese health checkup cohort, it was 11.6% (7.2%

in males and 15.0% in females) (25). Thus, TPOAb positivity was less frequent in our study,

which may reflect differences in assay methodology: NHANES used a different immunoassay

and reference range, whereas the Japanese health checkup cohort used a radioimmunoassay kit.

Supporting this interpretation, a Chinese study that used the same manufacturer's assay as ours,

with a cut-off of < 34 IU/mL, reported a prevalence of 7.98% (6.12% in males and 9.90% in

females) (26), closely matching our results.

Importantly, we identified the clinical relevance of high TPOAb titers. As shown in

Table 3, the prevalence of increased blood flow and diffuse FDG uptake elevated in a titer-

dependent manner, whereas the prevalence of thyroid dysfunction increased only in subjects

with TPOAb titers ≥ 128 IU/mL. Similarly, a previous cohort study found no significant

increase in overt hypothyroidism among subjects with TPOAb titers of 35–105 IU/mL, but a

marked increase among those with TPOAb titers of 106–350 IU/mL (27). These findings

suggest that low-titer TPOAb positivity reflects a relatively mild autoimmune response that

may be self-limiting and insufficient to cause thyroid dysfunction. However, given our cross-

sectional design, we cannot exclude the possibility that low-titer positivity may predict future

thyroid dysfunction. Nonetheless, low-titer TPOAb positivity is unlikely to confer a short-term

risk of hypothyroidism.

352   We further investigated the factors associated with thyroid function, incorporating

physiological perspectives. TSH levels were higher in females, whereas fT4 and fT3 levels

were higher in males. Although this tendency has been reported in other cohorts (2,28,29),

detailed statistical analyses are lacking. To address the potential bias from the higher prevalence

of thyroid abnormalities in females, we confirmed sex-related differences in thyroid function

using multivariate regression analyses. Age was another major determinant. TSH levels

increased with age in both sexes, whereas fT4 and fT3 levels declined with age in males, with

minimal associations observed in females. Similar patterns have been reported, with estrogen

proposed as a contributing factor (30).

361   The association between low body weight and low T3 levels is well recognized in

non-thyroidal illness, whereas the relationship between obesity and thyroid function remains

controversial (31). In our cohort, BMI was positively correlated with fT3 levels in both sexes

and TSH levels in females; both associations remained significant in multivariate analyses.

As fT3 elevation was not accompanied by changes in fT4 levels, the involvement of

iodothyronine deiodinases is likely. Type 2 iodothyronine deiodinase, which exerts a greater

influence on circulating fT3 levels than type 1 (32), may mediate this effect, whereas type 3

iodothyronine deiodinase reduces fT3 levels. The clinical importance of type 2 iodothyronine

deiodinase is supported by evidence that its pathogenic variant, Thr92Ala, is associated with

reduced fT3 levels (33). The observed TSH elevation in females is also noteworthy; one

possible mechanism is leptin-mediated stimulation of thyrotropin-releasing hormone

secretion, given that females have higher circulating leptin levels (34,35) and obesity

increases leptin production (36-38).

Multivariate analyses confirmed that these factors independently determined thyroid

function. Abnormal findings on thyroid examinations were also associated with thyroid

function, even within the physiological range. However, diffuse FDG uptake and TPOAb

titers $\geq$ 128 IU/mL showed weaker associations than increased blood flow and thyroid

volume: diffuse FDG uptake was associated only with fT4, TPOAb titers $\geq$ 128 IU/mL only

with TSH, whereas increased blood flow and thyroid volume were associated with both TSH

and fT4 levels. This discrepancy may reflect that ultrasonographic changes indicate structural

alterations in the thyroid gland, whereas immune activity detected by FDG-PET or TPOAb

positivity alone may be insufficient to cause thyroid dysfunction.

Another strength of the present study is the rigorous assessment of thyroid

abnormalities, which enabled us to define the Thyroid-healthy cohort with no abnormal

findings on any thyroid examination. Using this cohort, we performed a detailed analysis of

thyroid size. First, males in our cohort had larger thyroid glands, independent of age, height,

and body weight. Sex differences in thyroid size have been consistently reported (39-41), and

our findings reaffirm this observation. Second, we observed a negative correlation between

age and the longitudinal diameter. Although age-related changes in thyroid size remain

controversial (40-42), this association may explain why we did not detect significant age

effects on other parameters (the isthmus thickness, the transverse diameter, and the

anteroposterior diameter) and why previous studies focusing on thyroid volume may have

failed to detect size changes (43), as thyroid volume is less sensitive to alterations in a single

dimension. Third, we identified a positive correlation between height and the longitudinal

diameter, whereas body weight was positively correlated with the isthmus thickness and the

transverse and anteroposterior diameters. This discrepancy is intriguing and raises the

possibility that the thyroid gland may grow predominantly in the longitudinal dimension in

proportion to height.

Finally, we established reference ranges for individual thyroid dimensions using

ultrasonography. While previous reports have provided regional reference values for thyroid

volume (43), to our knowledge, none have reported ranges for each parameter, namely, the

isthmus thickness, the transverse diameter, the anteroposterior diameter, and the longitudinal

diameter. We determined the reference values using strict criteria, excluding all subjects with

any abnormal findings on thyroid examinations or with nodules ≥ 5.0 mm, thereby ensuring a

robust reference set. We hope that these sex-specific reference values will serve as a useful tool

for evaluating thyroid enlargement in clinical practice.

This study has several limitations. As Japan is an iodine-sufficient country owing to

seaweed consumption (44), our results may differ from those in iodine-deficient regions. As

our cohort was composed almost entirely of Japanese individuals, racial differences should be

considered. Despite the peer review of ultrasonographic parameters, inter-operator variability

cannot be completely excluded. Diffuse FDG uptake was assessed qualitatively rather than

quantitatively, precluding the establishment of cutoff values. TgAb was not measured, and

inclusion of this parameter might have yielded additional insights.

In summary, this prospective cohort study of Japanese adults reported the prevalence

of thyroid dysfunction and abnormal findings on thyroid examinations and integrated

epidemiological data from ultrasonography, FDG-PET, and TPOAb measurements. We found

that TPOAb levels become clinically significant at titers ≥ 128 IU/mL and that age, sex, and

thyroid volume are dominant determinants of thyroid function. The Thyroid-healthy cohort,

defined through our stringent screening, enabled us to establish sex-specific reference ranges

for thyroid dimensions and confirm associations between thyroid size, age, and body size.

These findings provide comprehensive epidemiological and physiological evidence in

thyroid health and research.

Acknowledgments

We are grateful to the staff of the Preemptive Medicine and Lifestyle Disease Research Center for their assistance in collecting the data analyzed in this study. We also thank the ultrasound technologists for their expertise in performing thyroid ultrasonography and for their careful review of the images.

References

1. Vanderpump MP, Tunbridge WM, French JM, Appleton D, Bates D, Clark F, Grimley Evans J, Hasan DM, Rodgers H, Tunbridge F, Young ET. The incidence of thyroid disorders in the community: a twenty-year follow-up of the Whickham Survey. *Clin Endocrinol (Oxf)*. 1995;43(1):55-68.

2. Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer CA, Braverman LE. Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab*. 2002;87(2):489-499.

3. Ceylan I, Yener S, Bayraktar F, Secil M. Roles of ultrasound and power Doppler ultrasound for diagnosis of Hashimoto thyroiditis in anti-thyroid marker-positive euthyroid subjects. *Quant Imaging Med Surg*. 2014;4(4):232-238.

4. Mazzaferri EL. Management of a solitary thyroid nodule. *N Engl J Med*. 1993;328(8):553-559.

5. Yasuda S, Shohtsu A, Ide M, Takagi S, Takahashi W, Suzuki Y, Horiuchi M. Chronic thyroiditis: diffuse uptake of FDG at PET. *Radiology*. 1998;207(3):775-778.

6. Karantanis D, Bogsrud TV, Wiseman GA, Mullan BP, Subramaniam RM, Nathan MA, Peller PJ, Bahn RS, Lowe VJ. Clinical significance of diffusely increased 18F-FDG uptake in the thyroid gland. *J Nucl Med*. 2007;48(6):896-901.

7.   Chen W, Parsons M, Torigian DA, Zhuang H, Alavi A. Evaluation of thyroid FDG

uptake incidentally identified on FDG-PET/CT imaging. *Nucl Med Commun*.

2009;30(3):240-244.

8.   Suzuki S, Midorikawa S, Fukushima T, Shimura H, Ohira T, Ohtsuru A, Abe M,

Shibata Y, Yamashita S, Suzuki S. Systematic determination of thyroid volume by

ultrasound examination from infancy to adolescence in Japan: the Fukushima Health

Management Survey. *Endocr J*. 2015;62(3):261-268.

9.   Hegedüs L, Perrild H, Poulsen LR, Andersen JR, Holm B, Schnohr P, Jensen G,

Hansen JM. The determination of thyroid volume by ultrasound and its relationship to

body weight, age, and sex in normal subjects. *J Clin Endocrinol Metab*.

1983;56(2):260-263.

10.   Gómez JM, Maravall FJ, Gómez N, Gumà A, Soler J. Determinants of thyroid volume

as measured by ultrasonography in healthy adults randomly selected. *Clin Endocrinol*

(Oxf). 2000;53(5):629-634.

11.   Garmendia Madariaga A, Santos Palacios S, Guillén-Grima F, Galofré JC. The

incidence and prevalence of thyroid dysfunction in Europe: a meta-analysis. *J Clin*

Endocrinol Metab. 2014;99(3):923-931.

12.   Zhang X, Wang Y, Wang H, Zhang X. Trends in Prevalence of Thyroid Dysfunction

and its Associations With Mortality Among US Participants, 1988-2012. *J Clin*

Endocrinol Metab. 2024;109(2):e657-e666.

13.   Konno N, Yuri K, Taguchi H, Miura K, Taguchi S, Hagiwara K, Murakami S.

Screening for thyroid diseases in an iodine sufficient area with sensitive thyrotrophin

assays, and serum thyroid autoantibody and urinary iodide determinations. *Clin*

Endocrinol (Oxf). 1993;38(3):273-281.

14.   Katagiri R, Asakura K, Sasaki S, Hirota N, Notsu A, Miura A, Todoriki H, Fukui M,

Date C. Estimation of habitual iodine intake in Japanese adults using 16 d diet records

over four seasons with a newly developed food composition database for iodine. *Br J Nutr*. 2015;114(4):624-634.

15. Teng W, Shan Z, Teng X, Guan H, Li Y, Teng D, Jin Y, Yu X, Fan C, Chong W, Yang F, Dai H, Yu Y, Li J, Chen Y, Zhao D, Shi X, Hu F, Mao J, Gu X, Yang R, Tong Y, Wang W, Gao T, Li C. Effect of iodine intake on thyroid diseases in China. *N Engl J Med*. 2006;354(26):2783-2793.

16. Schulz SL, Seeberger U, Hengstmann JH. Color Doppler sonography in hypothyroidism. *Eur J Ultrasound*. 2003;16(3):183-189.

17. Yamauchi I, Yasoda A, Matsumoto S, Sakamori Y, Kim YH, Nomura M, Otsuka A, Yamasaki T, Saito R, Kitamura M, Kitawaki T, Hishizawa M, Kawaguchi-Sakita N, Fujii T, Taura D, Sone M, Inagaki N. Incidence, features, and prognosis of immune-related adverse events involving the thyroid gland induced by nivolumab. *PLoS One*. 2019;14(5):e0216954.

18. Yamauchi I, Yabe D. Best practices in the management of thyroid dysfunction induced by immune checkpoint inhibitors. *Eur Thyroid J*. 2025;14(1):e240328.

19. Muir CA, Tsang VHM, Menzies AM, Clifton-Bligh RJ. Immune Related Adverse Events of the Thyroid - A Narrative Review. *Front Endocrinol (Lausanne)*. 2022;13:886930.

20. Mu C, Ming X, Tian Y, Liu Y, Yao M, Ni Y, Liu Y, Li Z. Mapping global epidemiology of thyroid nodules among general population: A systematic review and meta-analysis. *Front Oncol*. 2022;12:1029926.

21. Imaizumi M, Sobue T, Taniguchi N, Miyagawa M, Momose T, Yoshinaga S, Yamamoto S, Ohishi W, Kitamura H, Zha L, Okubo T. Protocol and Progress of a Thyroid Study in the Epidemiological Study of Health Effects in Fukushima Emergency Workers. *Disaster Med Public Health Prep*. 2025;19:e108.

22. Brander A, Viikinkoski P, Nickels J, Kivisaari L. Thyroid gland: US screening in a

random adult population. *Radiology*. 1991;181(3):683-687.

23. Kwong N, Medici M, Angell TE, Liu X, Marqusee E, Cibas ES, Krane JF, Barletta

JA, Kim MI, Larsen PR, Alexander EK. The Influence of Patient Age on Thyroid

Nodule Formation, Multinodularity, and Thyroid Cancer Risk. *J Clin Endocrinol*

Metab. 2015;100(12):4434-4440.

24. Li Y, Jin C, Li J, Tong M, Wang M, Huang J, Ning Y, Ren G. Prevalence of Thyroid

Nodules in China: A Health Examination Cohort-Based Study. *Front Endocrinol*

(Lausanne). 2021;12:676144.

25. Kasagi K, Takahashi N, Inoue G, Honda T, Kawachi Y, Izumi Y. Thyroid function in

Japanese adults as assessed by a general health checkup system in relation with

thyroid-related antibodies and other clinical parameters. *Thyroid*. 2009;19(9):937-944.

26. Zhang M, Ni W, Zhang L, Fan K, Sun Y, Liu C, Xu S. Age-specific association

between thyroid autoimmunity and hypothyroidism in Chinese adults aged over 65

years: a cross-sectional study. *Front Endocrinol (Lausanne)*. 2023;14:1216308.

27. Walsh JP, Bremner AP, Feddema P, Leedman PJ, Brown SJ, O'Leary P. Thyrotropin

and thyroid antibodies as predictors of hypothyroidism: a 13-year, longitudinal study

of a community-based cohort using current immunoassay techniques. *J Clin*

Endocrinol Metab. 2010;95(3):1095-1104.

28. Yamada S, Horiguchi K, Akuzawa M, Sakamaki K, Yamada E, Ozawa A, Kobayashi

I, Shimomura Y, Okamoto Y, Andou T, Andou Y, Yamada M. The Impact of Age- and

Sex-Specific Reference Ranges for Serum Thyrotropin and Free Thyroxine on the

Diagnosis of Subclinical Thyroid Dysfunction: A Multicenter Study from Japan.

Thyroid. 2023;33(4):428-439.

29. Meng Z, Liu M, Zhang Q, Liu L, Song K, Tan J, Jia Q, Zhang G, Wang R, He Y, Ren

X, Zhu M, He Q, Wang S, Li X, Hu T, Liu N, Upadhyaya A, Zhou P, Zhang J. Gender

and Age Impacts on the Association Between Thyroid Function and Metabolic

Syndrome in Chinese. *Medicine (Baltimore)*. 2015;94(50):e2193.

30. Völzke H, Alte D, Kohlmann T, Lüdemann J, Nauck M, John U, Meng W. Reference

intervals of serum thyroid function tests in a previously iodine-deficient area. *Thyroid*.

2005;15(3):279-285.

31. Fontenelle LC, Feitosa MM, Severo JS, Freitas TE, Morais JB, Torres-Leal FL,

Henriques GS, do Nascimento Marreiro D. Thyroid Function in Human Obesity:

Underlying Mechanisms. *Horm Metab Res*. 2016;48(12):787-794.

32. Yamauchi I, Sakane Y, Yamashita T, Hakata T, Sugawa T, Fujita H, Okamoto K, Taura

D, Hirota K, Ueda Y, Fujii T, Yasoda A, Inagaki N. Thyroid hormone economy in

mice overexpressing iodothyronine deiodinases. *FASEB J*. 2022;36(2):e22141.

33. Castagna MG, Dentice M, Cantara S, Ambrosio R, Maino F, Porcelli T, Marzocchi C,

Garbi C, Pacini F, Salvatore D. DIO2 Thr92Ala Reduces Deiodinase-2 Activity and

Serum-T3 Levels in Thyroid-Deficient Patients. *J Clin Endocrinol Metab*.

2017;102(5):1623-1630.

34. Rosenbaum M, Nicolson M, Hirsch J, Heymsfield SB, Gallagher D, Chu F, Leibel

RL. Effects of gender, body composition, and menopause on plasma concentrations of

leptin. *J Clin Endocrinol Metab*. 1996;81(9):3424-3427.

35. Ostlund RE, Jr., Yang JW, Klein S, Gingerich R. Relation between plasma leptin

concentration and body fat, gender, diet, age, and metabolic covariates. *J Clin

Endocrinol Metab*. 1996;81(11):3909-3913.

36. Légrádi G, Emerson CH, Ahima RS, Flier JS, Lechan RM. Leptin prevents fasting-

induced suppression of prothyrotropin-releasing hormone messenger ribonucleic acid

in neurons of the hypothalamic paraventricular nucleus. *Endocrinology*.

1997;138(6):2569-2576.

37. Nillni EA, Vaslet C, Harris M, Hollenberg A, Bjørbak C, Flier JS. Leptin regulates

prothyrotropin-releasing hormone biosynthesis. Evidence for direct and indirect

pathways. *J Biol Chem*. 2000;275(46):36124-36133.

38. Harris M, Aschkenasi C, Elias CF, Chandrankunnel A, Nillni EA, Bjøorbaek C,

Elmquist JK, Flier JS, Hollenberg AN. Transcriptional regulation of the thyrotropin-

releasing hormone gene by leptin and melanocortin signaling. *J Clin Invest*.

2001;107(1):111-120.

39. Berghout A, Wiersinga WM, Smits NJ, Touber JL. Determinants of thyroid volume as

measured by ultrasonography in healthy adults in a non-iodine deficient area. *Clin*

Endocrinol (Oxf). 1987;26(3):273-280.

40. Barrère X, Valeix P, Preziosi P, Bensimon M, Pelletier B, Galan P, Hercberg S.

Determinants of thyroid volume in healthy French adults participating in the

SU.VI.MAX cohort. *Clin Endocrinol (Oxf)*. 2000;52(3):273-278.

41. Gómez JM, Maravall FJ, Gómez N, Gumà A, Casamitjana R, Soler J. Pituitary-

thyroid axis, thyroid volume and leptin in healthy adults. *Horm Metab Res*.

2002;34(2):67-71.

42. Guo W, Tan L, Dong S, Jin Y, Zhu M, Wei H, Chen Y, Fan L, Du C, Zhang W. New

Reference Values for Thyroid Volume and a Comprehensive Assessment for

Influencing Factors in Chinese Adults with Iodine Sufficiency. *Eur Thyroid J*.

2021;10(6):447-454.

43. Srivastava S, Dighe M, Möller K, Chammas MC, Dong Y, Cui XC, Dietrich CF.

Ultrasound measurements and normal findings in the thyroid gland. *Med Ultrason*.

2024; doi: 10.11152/mu-4451.

44. Fuse Y, Ito Y, Shishiba Y, Irie M. Current Iodine Status in Japan: A Cross-sectional

Nationwide Survey of Schoolchildren, 2014-2019. *J Clin Endocrinol Metab*.

2022;107(5):e2065-e2079.

Figure Legends

Figure 1. Flowchart illustrating the establishment of cohorts in this prospective study. n, number of subjects; TPOAb, anti-thyroperoxidase antibody; SITSH, syndrome of inappropriate secretion of TSH; FDG: ^{18}F-fluorodeoxyglucose.

Figure 2. Prevalence of abnormal findings on thyroid examinations in the Original cohort. The actual number of subjects is shown in panels A, C, D, and E. The percentage of subjects with abnormal findings is shown in panels B and F. Thyroid nodules were classified by maximum diameter.

Figure 3. Associations between characteristics and thyroid function in the Original cohort. (A) Scatter plot of serum TSH and free T4 (fT4) levels. (B) Illustration of the exclusion criteria for the analyses presented in Figures 3 and 4, overlaid on a scatter plot. (C– E) Trends in thyroid function according to age (see Supplemental Table 2 for the actual values). (F–H) Trends in thyroid function according to body mass index (BMI) (see Supplemental Table 5 for actual values). Gray violin plots show the distribution of the data, and black circles with connecting lines represent mean ± SD. Log-transformed TSH values were used for statistical analyses because of their log-normal distribution. Trend analyses were conducted by assigning the median value of each category and modeling it as a continuous variable in regression analyses.

Figure 4. Associations between thyroid examination results and thyroid function in the Original cohort. (A–C) Differences in thyroid function according to the presence of increased blood flow on thyroid ultrasonography (see Supplemental Table 6 for actual values). (D–F) Differences in thyroid function according to the presence of diffuse FDG uptake (see Supplemental Table 6 for actual values). (G–I) Trends in thyroid function according to thyroid nodule size, classified by maximum diameter (see Supplemental Table 7 for the actual values). (J–L) Trends in thyroid function according to thyroid volume (see Supplemental Table 8 for actual values). Data from 183 subjects were excluded because of

technical issues in volume measurement using ultrasonography. Gray violin plots show the

distribution of the data, and black circles with connecting lines represent mean ± SD.

Student's *t* test was used for panels A–F. Trend analyses were performed for panels G–L.

Figure 5. Associations between TPOAb titers and thyroid examination results in the

TPOAb cohort. (A) Number of TPOAb-positive subjects. (B–E) Prevalence of abnormal

findings in TPOAb-positive subjects; any thyroid dysfunction (B), increased blood flow (C),

diffuse FDG uptake (D), and any thyroid nodules (E). (F) Exclusion criteria for analyses in

panels G–I overlaid on the scatter plot. (G–I) Differences in thyroid function according to

TPOAb positivity (see Supplemental Table 11 for the actual values). Gray violin plots show

the distribution of the data, and black circles with connecting lines represent mean ± SD.

Student's *t* test was used.

Figure 6. Associations between characteristics and thyroid size in the Thyroid-healthy

cohort. (A–D) Trends in thyroid size according to age (see Supplemental Table 12 for the

actual values). (E–H) Trends in thyroid size according to height (see Supplemental Table 13

for the actual values). (I–L) Trends in thyroid size according to body weight (see

Supplemental Table 14 for actual values). Two male subjects weighing < 50.0 kg were

excluded because of the small sample size. Gray violin plots show the distribution of the data,

and black circles with connecting lines represent mean ± SD.

Table 1. Characteristics and prevalence of abnormal findings in the Original cohort

	All (n = 4407)	Male (n = 2643)	Female (n = 1764)	p
Age (year)	52.9 (11.3)	53.2 (11.0)	52.3 (11.9)	0.0114
Height (cm)	166.2 (8.7)	171.5 (6.1)	158.3 (5.5)	< 0.0001
Body weight (kg)	66.0 (14.1)	73.5 (11.5)	54.7 (9.3)	< 0.0001
BMI (kg/m^2)	23.7 (3.8)	25.0 (3.5)	21.8 (3.6)	< 0.0001
Thyroid function				
TSH (μIU/mL)	1.952 (2.245)	1.859 (1.972)	2.091 (2.596)	0.0001
fT4 (ng/dL)	1.288 (0.200)	1.313 (0.180)	1.250 (0.221)	< 0.0001
fT3 (pg/mL)	3.08 (0.58)	3.19 (0.39)	2.93 (0.76)	< 0.0001
Thyroid dysfunction				
No n (%)	4151 (94.19%)	2505 (94.78%)	1646 (93.31%)	0.0485
Yes n (%)	256 (5.81%)	138 (5.22%)	118 (6.69%)	
Overt thyrotoxicosis	27 (0.61%)	15 (0.57%)	12 (0.68%)	
Subclinical thyrotoxicosis	104 (2.36%)	69 (2.61%)	35 (1.98%)	
Subclinical hypothyroidism	118 (2.68%)	52 (1.97%)	66 (3.74%)	
Overt hypothyroidism	7 (0.16%)	2 (0.08%)	5 (0.28%)	
Increased blood flow				
No n (%)	4299 (97.55%)	2608 (98.68%)	1691 (95.86%)	< 0.0001
Yes n (%)	108 (2.45%)	35 (1.32%)	73 (4.14%)	
Diffuse FDG uptake				
No n (%)	4256 (96.57%)	2604 (98.52%)	1652 (93.65%)	< 0.0001
Yes n (%)	151 (3.43%)	39 (1.48%)	112 (6.35%)	
Thyroid nodule				
No n (%)	2657 (60.29%)	1763 (66.71%)	894 (50.68%)	< 0.0001
Yes n (%)	1750 (39.71%)	880 (33.29%)	870 (49.32%)	
< 5.0 mm	667 (15.14%)	393 (14.87%)	274 (15.53%)	
5.0–9.9 mm	705 (16.00%)	329 (12.45%)	376 (21.32%)	
10.0–19.9 mm	315 (7.15%)	133 (5.03%)	182 (10.32%)	
≥ 20.0 mm	63 (1.43%)	25 (0.95%)	38 (2.15%)	

BMI, body mass index; fT4, free T4; fT3, free T3; FDG, [18]F-fluorodeoxyglucose. Continuous variables are expressed as mean (SD). Comparisons between males and females were performed using the Student's t-test or Fisher's exact test, as appropriate. TSH values were log-transformed for statistical analyses due to their log-normal distribution.

Table 2. Characteristics and prevalence of abnormal findings in the Original cohort, classified by thyroid dysfunction

	Euthyroid (n = 4151)	Overt thyrotoxicosis (n = 27)	Subclinical thyrotoxicosis (n = 104)	Subclinical hypothyroidism (n = 118)	Overt hypothyroidism (n = 7)
Thyroid function					
TSH (µIU/mL)	1.805 (0.932)	0.185 (0.195)****	0.382 (0.116)****	7.041 (4.487)****	33.436 (31.148)****
fT4 (ng/dL)	1.285 (0.166)	2.295 (0.970)****	1.333 (0.137)**	1.161 (0.142)****	0.677 (0.194)****
fT3 (pg/mL)	3.07 (0.39)	6.06 (4.79)****	3.18 (0.48)**	2.89 (0.33)****	2.68 (0.47)**
Age	52.8 (11.3)	47.0 (8.2)**	51.7 (10.3)	58.1 (13.3)****	60.0 (11.5)
Sex					
Male n (%)	2505 (60.35%)	15 (55.56%)	69 (66.35%)	52 (44.07%)***	2 (28.57%)
Female n (%)	1646 (39.65%)	12 (44.44%)	35 (33.65%)	66 (55.93%)	5 (71.43%)
Increased blood flow					
No n (%)	4071 (98.07%)	19 (70.37%)****	101 (97.12%)	106 (89.83%)****	2 (28.57%)****
Yes n (%)	80 (1.93%)	8 (29.63%)	3 (2.88%)	12 (10.17%)	5 (71.43%)
Diffuse FDG uptake					
No n (%)	4026 (96.99%)	23 (85.19%)**	100 (96.15%)	106 (89.83%)***	1 (14.29%)****
Yes n (%)	125 (3.01%)	4 (14.81%)	4 (3.85%)	12 (10.17%)	6 (85.71%)
Thyroid nodule					
No n (%)	2527 (60.88%)	18 (66.67%)	45 (43.27%)***	64 (54.24%)	3 (42.86%)
Yes n (%)	1624 (39.12%)	9 (33.33%)	59 (56.73%)	54 (45.76%)	4 (57.14%)

Continuous values are expressed as mean (SD). Statistical analyses were performed for each group in comparison with the euthyroid group. Thyroid function and age were analyzed using the Student's *t*-test. TSH values were log-transformed for statistical analyses due to their log-normal distribution. Sex, FDG uptake, increased blood flow, and thyroid nodules were analyzed using Fisher's exact test. **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

Table 3. Characteristics and prevalence of abnormal findings in the TPOAb cohort, classified by TPOAb titers

	< 16.0 IU/mL (n = 2246)	16.0–31.9 IU/mL (n = 63)	32.0–127.9 IU/mL (n = 41)	≥ 128.0 IU/mL (n = 70)
Thyroid dysfunction				
No	2135 (95.06%)	59 (93.65%)	39 (95.12%)	53 (75.71%)****
Yes	111 (4.94%)	4 (6.35%)	2 (4.88%)	17 (24.29%)
Overt thyrotoxicosis	13 (0.58%)	0 (0.00%)	1 (2.44%)	3 (4.29%)
Subclinical thyrotoxicosis	46 (2.05%)	1 (1.59%)	1 (2.44%)	1 (1.43%)
Subclinical hypothyroidism	51 (2.27%)	3 (4.76%)	0 (0.00%)	10 (14.29%)
Overt hypothyroidism	1 (0.04%)	0 (0.00%)	0 (0.00%)	3 (4.29%)
Age	52.2 (11.2)	50.5 (12.2)	51.3 (10.9)	54.4 (10.7)
Sex				
Male n (%)	1404 (62.51%)	32 (50.79%)*	17 (41.46%)**	29 (41.43%)***
Female n (%)	842 (37.49%)	31 (49.21%)	24 (58.54%)	41 (58.57%)
Increased blood flow				
No n (%)	2221 (98.89%)	57 (90.48%)***	33 (80.49%)****	35 (50.00%)****
Yes n (%)	25 (1.11%)	6 (9.52%)	8 (19.51%)	35 (50.00%)
Diffuse FDG uptake				
No n (%)	2218 (98.75%)	57 (90.48%)***	32 (78.05%)****	35 (50.00%)****
Yes n (%)	28 (1.25%)	6 (9.52%)	9 (21.95%)	35 (50.00%)
Thyroid nodule				
No n (%)	1360 (60.55%)	36 (57.14%)	28 (68.29%)	39 (55.71%)
Yes n (%)	886 (39.45%)	27 (42.86%)	13 (31.71%)	31 (44.29%)

TPOAb, anti-thyroperoxidase antibody. Continuous values are expressed as mean (SD). Statistical analyses were performed for each group in comparison with the < 16.0 IU/mL group. Age was analyzed using Student's t-test, and all other variables were analyzed using Fisher's exact test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

Table 4. Multivariate regression analysis for determinants of thyroid function

	Model 1				Model 2		
	logTSH	fT4	fT3		logTSH	fT4	fT3
R^2	0.039	0.062	0.209		0.096	0.107	0.212
Intercept	−0.023	1.362	3.036		0.199	1.257	3.016
Age (year)	0.004****	−0.002****	−0.007****		0.004****	−0.002****	−0.007****
Sex (male to female)	−0.023***	0.034****	0.128****		0.002	0.020****	0.122****
BMI (kg/m^2)	0.001	0.001	0.017****		0.005**	−0.001	0.016****
Increased blood flow					0.063**	−0.037**	−0.018
Diffuse FDG uptake					0.026	−0.031*	0.007
Thyroid volume (g)					−0.014****	0.009****	0.005*
TPOAb ≥ 128.0 IU/mL					0.062***	0.009	0.019

R^2: coefficient of determination. Results are represented as estimates. "log" denotes the base-10 logarithm.

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

Table 5. Characteristics and thyroid size on ultrasonography in the Thyroid-healthy cohort

	All (n = 1840)	Male (n = 1187)	Female (n = 653)	p
Age	51.7 (10.9)	52.1 (10.6)	51.1 (11.5)	0.0780
Height (cm)	167.1 (8.5)	171.8 (5.8)	158.4 (5.5)	< 0.0001
Weight (kg)	67.3 (14.1)	74.0 (11.6)	55.0 (9.3)	< 0.0001
BMI (kg/m^2)	23.9 (3.9)	25.1 (3.5)	21.9 (3.6)	< 0.0001
Thyroid function				
TSH (μIU/mL)	1.782 (0.924)	1.727 (0.896)	1.881 (0.967)	0.0007
fT4 (ng/dL)	1.291 (0.161)	1.314 (0.164)	1.248 (0.147)	< 0.0001
fT3 (pg/mL)	3.11 (0.38)	3.22 (0.35)	2.91 (0.36)	< 0.0001
Thyroid nodule				
No n (%)	1226 (66.63%)	844 (71.10%)	382 (58.50%)	< 0.0001
Yes n (%)	614 (33.37%)	343 (28.90%)	271 (41.50%)	
< 5.0 mm	318 (17.28%)	197 (16.60%)	121 (18.53%)	
5.0–9.9 mm	296 (16.09%)	146 (12.30%)	150 (22.97%)	
Isthmus (mm)				
Thickness	2.3 (0.8)	2.5 (0.8)	1.9 (0.7)	< 0.0001
Right lobe (mm)				
Transverse diameter	15.9 (2.9)	16.7 (2.8)	14.4 (2.5)	< 0.0001
Anteroposterior diameter	13.9 (2.7)	14.8 (2.5)	12.2 (2.1)	< 0.0001
Longitudinal diameter	49.7 (5.5)	50.2 (5.3)	48.6 (5.7)	< 0.0001
Left lobe (mm)				
Transverse diameter	15.5 (2.8)	16.0 (2.8)	14.5 (2.3)	< 0.0001
Anteroposterior diameter	12.1 (2.8)	13.0 (2.6)	10.5 (2.4)	< 0.0001
Longitudinal diameter	47.2 (6.0)	47.9 (5.7)	46.0 (6.3)	< 0.0001
Thyroid volume (g)	10.6 (3.9)	11.9 (3.8)	8.3 (2.8)	< 0.0001

Continuous variables are expressed as mean (SD). Comparisons between male and female groups were performed using the Student's *t*-test or Fisher's exact test, as appropriate. TSH values were log-transformed for statistical analyses due to their log-normal distribution.

Table 6. Reference ranges for individual thyroid dimensions measured by ultrasonography

	Male	Female
Isthmus (mm)		
Thickness	1.1–4.4	0.9–3.8
Right lobe (mm)		
Transverse diameter	11.8–22.8	9.7–18.7
Anteroposterior diameter	10.0–20.5	8.1–16.4
Longitudinal diameter	38.8–59.1	37.0–58.1
Left lobe (mm)		
Transverse diameter	10.9–21.7	10.3–18.9
Anteroposterior diameter	7.8–18.2	6.0–15.5
Longitudinal diameter	36.1–58.1	33.0–57.5
Thyroid volume (g)	6.0–20.8	4.1–13.7

The 2.5th–97.5th percentile was calculated in the Thyroid-healthy cohort, excluding subjects with nodules 5.0–9.9 mm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6