Cumulative Genetic Risk for Asthma Contributes to Disease Severity in Children with Asthma

Matthew Dapas[1], William Wentworth-Sheilds[2], Emma E. Thompson[2], Rajesh Kumar[3], Elizabeth Lippner[3], Robert A. Wood[4], George T. O'Connor[5], Gurjit K. Khurana Hershey[6], Rebecca S. Gruchalla[7], Andrew H. Liu[8], Edward M. Zoratti[9], Leonard B. Bacharier[10], Stephanie Lovinsky-Desir[11], Michele A. Gill[12], William J. Sheehan[13], Shilpa J. Patel[14], Matthew C. Altman[15], James E. Gern[16], Cynthia M. Visness[17], Peter J. Gergen[18], Patrice M. Becker[18], Daniel J. Jackson[16*], Carole Ober[2*]

1. Division of Rheumatology, Department of Medicine; Center for Human Immunobiology; Center for Genetic Medicine; Northwestern University Feinberg School of Medicine, Chicago, IL
2. Department of Human Genetics, University of Chicago, Chicago, IL
3. Ann and Robert H. Lurie Children's Hospital; Division of Allergy and Immunology, Department of Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, IL
4. Department of Pediatrics, Johns Hopkins University Medical Center, Baltimore, MD
5. Pulmonary Center, Boston University School of Medicine, Boston, MA
6. Division of Asthma Research, Cincinnati Children's Hospital, Cincinnati, OH
7. Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX
8. Children's Hospital Colorado; Department of Allergy and Immunology, University of Colorado School of Medicine, Aurora, CO
9. Departments of Medicine and Public Health Sciences, Henry Ford Health and Michigan State University College of Human Medicine, Detroit, MI.
10. Monroe Carell Jr. Children's Hospital at Vanderbilt University Medical Center, Nashville, TN
11. Department of Pediatrics and Environmental Health Sciences, Columbia University, New York, NY
12. Department of Pediatrics, Washington University School of Medicine, St. Louis, MO
13. Children's National Hospital; Division of Allergy and Immunology, George Washington University School of Medicine and Health Sciences, Washington, DC
14. Division of Emergency Medicine, Children's National Hospital, Washington, DC
15. Department of Allergy and Infectious Diseases, University of Washington, Seattle, WA
16. Departments of Pediatrics and Medicine, University of Wisconsin School of Medicine and Public Health, Madison, WI
17. Rho Inc., Durham, NC
18. National Institute of Allergy and Infectious Diseases, Bethesda, MD

* These authors jointly supervised this work.

ABSTRACT

Background: Childhood-onset asthma is highly heritable, with nearly 200 risk loci identified in genome-wide association studies. Aggregated polygenic risk scores can be used to quantify genetic predisposition to asthma, but their power to predict asthma severity in multi-ancestral groups has not been previously evaluated.

Objective: Our aim was to examine the predictive power of biobank-derived asthma polygenic risk scores in four multi-ancestry asthma study cohorts of children living in U.S. urban environments.

Methods: We generated polygenic risk scores for asthma, derived from a large-scale genome-wide association meta-analysis, and assessed genetic predictions across different subphenotypes of asthma and tested for associations between genetic asthma risk and measures of asthma severity.

Results: Polygenic risk score prediction was significantly stronger for more symptomatic asthma phenotypes ($P<0.001$), and scores were significantly higher in difficult-to-control vs. easy-to-control asthma ($P=0.02$). Genetic risk was also significantly associated with more frequent exacerbations ($P=0.03$), higher blood eosinophil levels ($P=0.01$), and lower lung function ($P<0.001$).

Conclusion: Cumulative genetic risk for asthma is associated with disease severity and exacerbation risk in children with asthma.

Conflict of Interest Disclosures

M. Altman, P. Becker, M. Dapas, P. Gergen, M. Gill, and E. Thompson have nothing to disclose outside the submitted work. J. Gern reports consulting fees from Arrowhead Pharmaceuticals and Meissa Vaccines Incorporated, stock options from Meissa Vaccines Incorporated, and two patents issued related to the methods to enhance the production of rhinoviruses, outside the submitted work. W. Sheehan reports consulting fees from Regeneron and Syneos Health, and research funding from Merck, outside the submitted work. L. Bacharier reports royalties from Elsevier, consulting and/or personal fees from GlaxoSmithKline, Genentech/Novartis, DBV Technologies, Kinaset, OM Pharma, Recludix, Teva, AstraZeneca, WebMD/Medscape, Vectura and American Board of Allergy and Immunology and consulting fees, honoraria and medical writing fees from Sanofi/Regeneron and non-financial support from American Academy of Allergy Asthma & Immunology, outside the submitted work. G.K.K. Hershey reports grants from Adare during the study. C. Ober reports personal fees from American Association of Asthma, Allergy, and Immunology, outside the submitted work. R. Wood reports grants from DBV, Aimmune, Regeneron, Genentech, Novaris, FARE and royalties from Up to Date, outside the submitted work. S. Lovinsky-Desir reports grants or contracts from the Health Effects Institute and participation on the Clean Air Scientific Advisory Committee, outside the submitted work. D. Jackson reports grants or contracts from NHLBI, NIH office of the director, and Regeneron, consulting fees from AbbVie, and participation on Upstream Bio, AstraZeneca, GlaxoSmithKline, Regeneron, Sanofi, Areteia, and Apogee, outside the submitted work. A. Liu reports grants from ResMed, AstraZeneca, Phadia/ThermoFisher Scientific, Revenio, Labcorp, ResMed/Propeller Health, and Revenio, grants from OM Pharma, outside the submitted work.

Key Messages

- Polygenic risk prediction is stronger in more symptomatic phenotypes
- Cumulative genetic risk for asthma is associated with greater asthma severity, higher exacerbation frequency, lower lung function, and increased eosinophil levels in children with asthma

Capsule Summary

We demonstrate that cumulative genetic risk for asthma is associated with disease severity in children with asthma living in urban environments.

Key Words (up to 10)

Asthma; genetics; polygenic risk score

Abbreviations

APIC, Asthma Phenotypes in Inner-City Children; AUC, area under the curve; CAAPA, Consortium on Asthma in African-ancestry Populations in the Americas; FEV_1, Forced expiratory volume in one second; FVC, Forced vital capacity; GBMI, Global Biobank Meta-analysis Initiative; GWAS, Genome-wide association study; ICD, International Classification of Diseases; IQR, interquartile range; MUPPITS, Mechanisms Underlying Asthma Exacerbations Prevented and Persistent with Immune Based Therapy: A Systems Approach; PRS, Polygenic risk score; ROC, receiver operating characteristic; URECA, Urban Environment and Childhood Asthma

INTRODUCTION

Asthma is a heterogenous disease with multiple endotypes and phenotypes, with severe forms of asthma accounting for a disproportionate share of the disease's morbidity, mortality, and healthcare costs[1]. Identifying factors that contribute to disease severity could help improve risk stratification and guide personalized management strategies. Genetics contributes significantly to the overall risk of developing asthma, as evidenced by nearly 200 risk loci uncovered in large genome-wide association studies (GWASs)[2-5]. However, the relationship between cumulative genetic asthma risk and disease severity has not been fully explored[6].

Genetic risk can be modeled using aggregated polygenic risk scores (PRSs). Many studies have developed predictive PRSs for asthma[7], and phenome-wide association studies with asthma PRSs have suggested that genetic risk may correlate with asthma severity[6,7]. To directly test this relationship, we generated PRSs for asthma in four multi-ancestry asthma study cohorts of children living in low-income, urban environments, who face an excess burden of asthma morbidity and mortality[8-10]. We then assessed genetic predictions for different subphenotypes of asthma and tested for associations between genetic asthma risk and multiple measures of asthma severity.

METHODS

Cohorts

We analyzed samples and phenotypes from several National Institutes of Allergy and Infectious Diseases (NIAID)-funded asthma studies conducted by the Inner-City Asthma Consortium (ICAC)[11]: the Asthma Phenotypes in the Inner City (APIC) study[12,13], the Urban Environment and Childhood Asthma (URECA) birth cohort study[14], and the Mechanisms Underlying Asthma Exacerbations Prevented and Persistent with Immune Based Therapy: A Systems Approach (MUPPITS 1 & 2) studies[8,15]. Non-asthma, population-based controls from the Consortium on Asthma among African-ancestry Populations in the Americas (CAAPA) study[16] was used as a comparator group to assess the distribution of the PRSs in an unselected healthy population with similar ancestry to the APIC, URECA, and MUPPITS participants. Ethical approval for this work was granted by a central institutional review board, WGC IRB[17].

APIC

APIC was a prospective, observational study in children with asthma age 6-17 years old from low-income areas (≥20% of residents below federal poverty level) of nine U.S. cities. Hierarchical clustering was performed on a large collection of baseline demographic, clinical, and environmental variables to identify

different phenotypes of childhood-onset asthma. Previously, five clusters were described (clusters A-E)

based on varying symptoms, biomarkers, and disease severity[13]. APIC participants were also previously

grouped based on required asthma controller treatment levels. Easy-to-control asthma was defined as

requiring ≤100mcg/day of fluticasone at a majority of study visits, whereas difficult-to-control asthma was

defined as requiring ≥500mcg/day[12]. APIC participants were genotyped using whole-genome sequencing

(WGS) as previously described[18].

URECA

URECA is a prospective, observational study of a birth cohort enriched for children with at least one parent

with asthma or allergic disease in four U.S. cities[14]. All families lived in neighborhoods where at least 20%

of the population had incomes below the poverty level. Clinical measurements at age 10 were used when

available; otherwise, the most recent measurement after age 5 was used. URECA participants were

genotyped using WGS as previously described[18].

MUPPITS 1 & 2

The MUPPITS studies investigated exacerbations in children with exacerbation-prone eosinophilic asthma

who were living in urban environments across nine U.S. cities. Participants in MUPPITS were 6-17 years

old, had blood eosinophils of ≥150 cells/μL, required at least twice-daily treatment with inhaled

corticosteroids, and had at least two exacerbations treated with systemic corticosteroids in the previous

year. All participants lived in census tracts where at least 10% of families had incomes below the poverty

level[19]. The MUPPITS-1 study was a 6-month prospective, longitudinal study that compared transcriptome

profiles in children who experienced cold symptoms between those that subsequently experienced an

exacerbation and those that did not[15]. The MUPPITS-2 study was a randomized controlled trial to assess

the safety and efficacy of mepolizumab for reducing exacerbations in this population[8]. Genotypes for

MUPPITS 1 & 2 were determined in 461 subjects with available DNA using the Illumina Genome Diversity

Array (GDA, 8v1.0, hg19) at the University of Chicago. Individuals were removed if they met any of the

following quality control (QC) criteria: genotyped vs. self-reported sex mismatches, unexpected sample

duplication, >2% missing genotypes across all variants, or autosomal heterozygosity >5 standard deviations

from the mean. Variants were excluded under the the following QC criteria: >4% missing genotypes across

all samples or exact Hardy-Weinberg p-value $<5\times10^{-8}$. A total of 1,661,770 directly genotyped single-

nucleotide variants remained following QC. Variants were lifted from human reference hg19 to hg38 using

CrossMap[20]. Eagle[21] was used for genotype phasing and minimac4[22] was used for genotype imputation.

Imputation was performed using the TOPMed reference panel (r2 version 1.0.0) of >100,000 whole genome

sequences (human genome build 38) and the Michigan Imputation Server[23]. Following QC, 428 samples

from MUPPITS 1 & 2 remained for analysis.

CAAPA

CAAPA[16] performed WGS on approximately 1,000 individuals with African ancestry from 19 populations across the Americas and Caribbean as previously described[24]. The WGS data is available through dbGAP (accession phs001123.v2.p1). In this study, we utilized WGS data from CAAPA non-asthma control samples from among the African American populations (n=205). Variant QC followed the same criteria as applied to APIC and URECA WGS data[18].

Modeling Genetic Asthma Risk

Polygenic risk for asthma was modeled on published association summary statistics from a large-scale, multi-ancestry GWAS conducted as part of the Global Biobank Meta-analysis Initiative (GBMI)[2]. Summary statistics were available via https://www.globalbiobankmeta.org. The GWAS was a meta-analysis across 18 biobanks, consisting of 153,763 cases and 1,647,022 controls. Primary ancestry among asthma cases was 79.3% European, 12.1% East Asian, 3.3% African, 2.6% Admixed American, 2.6% Central and Southern Asian, and 0.01% Middle Eastern[2]. Asthma case and control statuses were assigned primarily according to "PheCodes" mapped from International Classification of Disease (ICD) codes, but several biobanks used self-reported data.

To generate individual PRSs, we applied the polygenic risk model published by Wang and colleagues[25] from the GBMI multi-ancestry GWAS, which was generated using PRS-CS[26] and deposited in the Polygenic Score Catalog (PGS001782)[27]. PRS-CS applies a Bayesian regression framework with continuous shrinkage priors on variant effect sizes from all tested variants. The "PRS-CS-auto" algorithm was used with default parameters to estimate the global and local shrinkage parameters directly from the GBMI GWAS summary statistics for each autosomal chromosome. Linkage disequilibrium was modeled using the EUR reference panel constructed by the PRS-CS developers with 1000 Genomes (1KG) Project data[25]. We aggregated the posterior variant effects into per-individual scores across the intersection of genotyped single-nucleotide variants present in all genotyped cohorts using the Plink (v2.00a5.8LM AVX2 Intel, 21 Nov 2023) "score" function[28]. Missing genotypes in individuals were imputed as the product of the variant's posterior effect size and its effect-allele frequency[29]. Cumulative PRSs were converted to Z scores (across all genotyped cohorts) for analysis.

Ancestry estimation

Principal components (PCs) of genetic ancestry were used to control for population stratification. Ancestry PCs were calculated on the intersect of high-quality single-nucleotide variants genotyped in the asthma cohorts and population reference panels from the 1KG Project (n=156)[30] and the Human Genome Diversity (HGD) Project (n=52)[31], as in our previous study[18]. Ancestry PCs were calculated accounting for subject

relatedness using PC-Air[32] and PC-Relate[33], with initial kinship estimates derived using KING[34]. Kinship

was estimated in iterations until resultant PCs stabilized (n=3).

Association Testing

Differences in PRS distributions between cohorts were assessed using Wilcoxon rank-sum tests for pairwise

comparisons and Kruskal-Wallis tests for more than two groups. PRS asthma prediction was conducted

with the CAAPA non-asthma samples as controls. Statistical differences in receiver operating characteristic

(ROC) areas under the curve (AUC) were evaluated using DeLong's test. Differences in PRS distributions

between difficult-to-control and easy-to-control treatment groups[12] in APIC were assessed using logistic

regression, adjusting for age, sex, and the first three ancestry PCs. PRS effects on exacerbation rates in

MUPPITS-2 were determined using negative binomial generalized linear models, adjusting for age, sex,

and the first three PCs of global ancestry. PRS associations with quantitative traits across multiple studies

were assessed using a two-stage approach[18,35,36]. First, traits were regressed on age, sex, study, and the first

three ancestry PCs. Then, model residuals were rank-normalized and regressed again on the same covariates

with the addition of the PRS as a predictor. Adjusted trait values for plotting PRS associations were derived

by adding first-stage model intercepts to residuals.

RESULTS

We utilized genome-wide genotype data from four multi-ancestry asthma cohorts of children living in low-

income, urban environments (APIC[13], URECA[14], MUPPITS-1[15], and MUPPITS-2[8]), as summarized in

Table 1, including 1,108 children with childhood-onset asthma. Nearly all of the participants in these

studies were parent-identified as Black (66%) or Hispanic (26%), and the cohorts featured similar overall

distributions of reported race/ethnicity (**Table 1**) and genetic ancestry (**Figure 1**).

Table 1. Demographic characteristics of childhood-onset asthma cohorts

Characteristic	All	APIC	URECA	MUPPITS 1 & 2
Number	1462	507	527	428
Age, years, median (range)	10 (6-17)	11 (6-17)	10 (7-10)	11 (6-17)
Female sex	665 (45%)	216 (43%)	261 (50%)	188 (44%)
Race/Ethnicity				
Black (non-Hispanic)	966 (66%)	318 (63%)	377 (72%)	269 (63%)
White (non-Hispanic)	23 (2%)	7 (1%)	7 (1%)	9 (2%)
Hispanic	378 (26%)	154 (30%)	104 (20%)	120 (28%)
Other/mixed	93 (6%)	26 (5%)	38 (7%)	29 (7%)
Unknown	4 (<1%)	2 (<1%)	1 (<1%)	1 (<1%)
Site				
Baltimore	233 (16%)	84 (17%)	149 (28%)	-
Boston	228 (16%)	65 (13%)	124 (24%)	39 (9%)
Chicago	97 (7%)	62 (12%)	-	35 (8%)
Cincinnati	84 (6%)	45 (9%)	-	39 (9%)
Dallas	86 (6%)	38 (7%)	-	48 (11%)
Denver	103 (7%)	59 (12%)	-	44 (10%)
Detroit	106 (7%)	50 (10%)	-	56 (13%)
New York	227 (16%)	65 (13%)	99 (19%)	63 (15%)
St. Louis	207 (14%)	-	155 (29%)	52 (12%)
Washington, D.C	91 (6%)	39 (8%)	-	52 (12%)
Clinical traits				
Asthma Diagnosis	1108 (75%)	507 (100%)	173 (33%)	428 (100%)
Eosinophils/μL, median (IQR)	300 (190-500)	300 (160-485.5)	200 (100-350)	400 (300-600)
FEV_1/FVC, median (IQR)	0.80 (0.73-0.86)	0.79 (0.72-0.85)	0.84 (0.79-0.88)	0.76 (0.68-0.82)

APIC, Asthma Phenotypes in Inner-City Children; FEV1, Forced expiratory volume in one second; FVC, Forced vital capacity; IQR, interquartile range; MUPPITS, Mechanisms Underlying Asthma Exacerbations Prevented and Persistent with Immune Based Therapy: A Systems Approach; URECA, Urban Environment and Childhood Asthma

Figure 1. Principal components of ancestry by cohort. The top three principal components (PCs) of ancestry are plotted for sequenced APIC, URECA, MUPPITS, and CAAPA (non-asthma) participants, colored by study, along with the four ancestry reference populations used for determining ancestry. 1KG, 1000 Genomes Project; HGD, Human Genome Diversity Project.

PRS distributions differed significantly by cohort (P<0.001, Kruskal-Wallis), with higher median PRSs
observed in cohorts characterized by more severe forms of asthma (**Figures 2A, 2B**). Notably, the PRSs
were significantly higher in URECA children without asthma, who were enriched for family history of
asthma, than in population-based controls without asthma (CAAPA), likely owing to inherited disease risk
(P=0.03, Wilcoxon). The PRS predicted asthma diagnosis more accurately in cohorts characterized by
greater symptom burden, according to relative ROC AUCs (**Figure 2C**). Overall, the PRS yielded
significantly better prediction for moderate-severe asthma (APIC clusters D and E, MUPPITS) (AUC=0.65)
than for the other groups (AUC=0.53; P_Δ=<0.001, **Figure 2D**). These results demonstrate that genetic
asthma prediction is stronger for more symptomatic phenotypes and further suggest that genetic risk
correlates with increasing disease severity in children.

Figure 2. Genetic asthma prediction is stronger in more symptomatic phenotypes. A) Distributions and median values of asthma PRSs by genotyped cohort, including APIC's previously defined sub-phenotypes (clusters A-E)[13]. B) Schematic heatmap showing the relative degrees of asthma severity for each cohort. C) Receiver operating characteristic (ROC) curves with areas under the curve (AUC) for predicting asthma using the PRS for each cohort. D) Prediction of asthma in the combined moderate-to-severe asthma cohorts (APIC D, E & MUPPITS) was significantly more accurate than for the milder asthma subphenotypes (APIC A, B, C), combined.

To more directly test whether polygenic asthma risk correlates with disease severity, we tested PRS
associations with cohort-specific severity metrics. In APIC, PRSs were significantly higher in difficult-to-
control vs. easy-to-control asthma[12] (P=0.02; **Figure 3A**), indicating that children with more genetic risk
alleles tended to require higher treatment levels to control their asthma. In the MUPPITS-2 trial, which
investigated the effect of an anti-IL-5 inhibitor (mepolizumab) in children with exacerbation-prone
eosinophilic asthma[8], polygenic risk was significantly associated with higher exacerbation rates. However,
the association was only observed in the placebo-treated group (P=0.03; **Figure 3B**). The correlation was

not significant in the study arm treated with mepolizumab (P=0.53), suggesting that the biologic effectively
reduced the asthma genetic effect on the exacerbation rate.

Figure 3. Polygenic risk is associated with asthma severity and exacerbation risk. A) Distributions and median values of asthma PRSs in APIC by required treatment step[12] Sample sizes are shown below the x-axis labels. B) Exacerbation rate vs. PRSs in MUPPITS-2, split by treatment arm. Predicted exacerbation rates by PRS are shown for each treatment arm (solid lines) with 95% confidence intervals (shaded areas).

We next assessed associations between the PRS and quantitative biomarkers of asthma severity[18]. In children with asthma across all cohorts, the PRS was significantly correlated with higher blood eosinophil levels (P=0.01; **Figure 4A**) and lower lung function (P<0.001; **Figure 4C**), as measured by the ratio of forced expiratory volume in one second to forced vital capacity (FEV_1/FVC). Comparing the highest to lowest PRS deciles, these effects amounted to significant median differences of 101 eosinophils per μL (P=0.02; **Figure 4B**), and -3.0% FEV_1/FVC (P=0.005; **Figure 4D**). These findings further demonstrate that genetic asthma risk is correlated with quantitative measures relevant to asthma severity.

Figure 4. Polygenic risk is associated with quantitative biomarkers of asthma severity. A) Eosinophil count per μL, adjusted for age, sex, study, and ancestry, plotted against the PRS in a combined sample from APIC, URECA, and MUPPITS (n=1,062). The modeled correlation fit with 95% confidence interval is drawn in red. B) Distributions and median values of adjusted eosinophil counts in the lowest and highest PRS deciles. C) Adjusted FEV$_1$/FVC plotted against the PRS in a combined sample from APIC, URECA, and MUPPITS (n=1,098). D) Distributions and median values of adjusted FEV$_1$/FVC in the lowest and highest PRS deciles.

DISCUSSION

Genetic disease risk depends on phenotype definitions, ancestry, and environmental exposures. The GBMI asthma GWAS that we used for modeling our PRS drew predominantly from European and East Asian adult cohorts and defined asthma according to specific International Classification of Diseases (ICD)-9 and ICD-10 codes from participant electronic health records[37]. Therefore, the asthma phenotypes, ancestries, and environments captured in the GBMI GWAS differed overall from those in the four cohorts studied here, which focused on childhood-onset asthma in urban environments in the United States, consisting mostly of Black and Hispanic populations. It has been shown that PRS predictive performance deteriorates across ancestries[38], and we previously demonstrated how modeling genetic risk from different asthma GWASs yielded PRSs with different trait associations[7]. Furthermore, asthma cases in the GBMI GWAS consisted largely of adult-onset disease[2,7], and the genetic architectures of childhood- and adult-onset asthma are partly distinct, with reported genetic correlations ranging from 0.67-0.78[2,39]. Although the GBMI-derived PRS was found to be the best performing asthma PRS in individuals of African ancestry[40] and predicted asthma in this study with accuracy comparable to its performance in non-African populations[7,40], the GBMI PRS is not a perfect proxy for genetic asthma risk in this study's cohorts. Thus, the actual correlation between genetic asthma risk and disease severity is likely stronger than the effects we observed here.

Among patients with physician-diagnosed asthma according to ICD-10 codes, the vast majority have mild-to-moderate asthma, with reported prevalences of severe asthma ranging from 2-14%[41-44], depending on the country and definition used for severe asthma. Asthma in the GBMI GWAS was primarily defined according to ICD codes in the constituent biobanks. Therefore, it is unlikely that the PRS correlations we observed with asthma severity are due to asthma in GBMI being disproportionately severe. Instead, our results indicated that the cumulative burden of genetic risk variants for asthma in general contributes to disease severity. This conclusion is supported by GWASs of moderate-to-severe asthma, which found that 88% of association signals that reached genome-wide significance had been previously reported in studies that predominantly assessed samples from patients with mild asthma[45,46]. Our findings expand on those from a population-based longitudinal study in European ancestry individuals, in which asthma cases with a higher polygenic burden calculated from 17 risk variants were more likely to manifest atopy, airway hyperresponsiveness, and incompletely reversible airflow obstruction, and more often experienced hospitalizations for breathing problems[47]. Our results also align with another study that found polygenic risk for asthma was significantly associated with asthma exacerbations in European ancestry individuals[48]. Similar observations, where cumulative genetic risk from low-effect variants were associated with disease severity, have been reported in other complex diseases, as well, including cardiovascular disease[49] and

autoimmune disease[50]. Such genetic effects may combine with effects from rare variants[45,51] and

environmental exposures[52] to contribute to disease severity through shared or independent pathways[9].

In summary, the results of our study indicate that the cumulative effect of asthma-associated genetic variants

not only increase disease risk but also contribute to disease severity. By applying a GBMI-trained PRS to

four multi-ancestry childhood-onset asthma cohorts from low-income urban environments, we demonstrate

that the effects observed in this study transcend ancestry, asthma phenotypes, severity metrics, and

environments. Genetic risk models that are more closely aligned to demographic and clinical profiles of

study populations would likely more precisely capture genetic influences on severity. Future studies should

consider incorporating genetic risk together with other predictive factors for developing more

comprehensive and accurate prediction models of asthma severity to inform strategies for clinical

prevention, intervention, and management.

ACKNOWLEDGEMENTS

URECA V has been funded in whole or in part with Federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, and Department of Health and Human Services under Cooperative Agreement number 1UM1AI160040, HHSN272200900052C, HHSN272201000052I, 1UM1AI114271-01, and UM2AI117870. Additional support was provided by the National Center for Research Resources, National Center for Advancing Translational Sciences, and National Institutes of Health, under grants NCRR/NIHRR00052, NCRR/NIHM01RR00533, NCRR/NIH1UL1RR025771, NCRR/NIHM01RR00071, 1UL1RR024156, NCRR/NIH RR00052, UL1TR001079, NCRR/NIHM01RR00533, NCRR/NIH1UL1RR025771, NCRR/NIHM01RR00071, NCRR/NIH1UL1RR024156, NCATS/NIHUL1TR000040, 5UL1RR024992-02, NCATS/NIHUL1TR001079, NCATS/NIH1UL1TR001430, NCATS/NIHUL1TR001873, and NCATS/NIH UL1TR002345.

The APIC project has been funded in whole or in part with Federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, and Department of Health and Human Services under Contract numbers HHSN272200900052C, HHSN272201000052I, 1UM1AI114271-01, and UM2AI117870. Additional support was provided by the National Center for Research Resources, and National Center for Advancing Translational Sciences, National Institutes of Health, under grants NCATS/NIH UL1TR000150, NCRR/NIH UL1TR000451, UL1TR001105, NCRR/NIH 1UL1RR025780 (CCTSI is supported in part by Colorado CTSA Grant UL1TR000154 from NCATS/NIH UL1TR001082), NCATS/NIH UL1TR000040, NCATS/NIH UL1TR000075, NCRR/NCAT/NIH UL1TR000077-04. GlaxoSmithKline (GSK) provided Ventolin HFA, Flonase, Flovent 50 mcg, Flovent 100 mcg, Flovent 250 mcg , Advair 250/50 mcg, Advair 500/50 mcg under a clinical trial agreement with NIH NIAID. None of these companies had a role in the development or approval of the protocol, conduct of the trial, data analysis, manuscript preparation, or the decision to submit for publication.

The MUPPITS-1 and MUPPITS-2 projects have been funded in whole or in part with Federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, and Department of Health and Human Services under Contract numbers 1UM1AI114271-01 and UM2AI117870. Additional support was provided by the National Center for Research Resources, National Center for Advancing Translational Sciences, and National Institutes of Health, under grants UL1TRG01422 (05/19/08 – 04/29/12), CTSA Grant UL1RR025741 (04/30/12-08/11/15), CTSA Grant UL1TR000150 (08/12/15-03/31/19), CTSA Grant UL1TR001422, NIH/NCATS Colorado CTSA UL1 TR002535, NCATS/NIH UL1TR001876, and NIH/CTSA 5UL1TR001425- 03.

Dr. Becker's and Gergen's co-authorship of this report does not necessarily represent the views of the

National Institute of Allergy and Infectious Diseases, the National Institutes of Health or any other agency

of the United States Government.

DATA AVAILABILITY

Phenotype, genotype, GWAS summary statistics, and whole-genome sequencing files are available in

dbGaP under accession phs002921 for the APIC and URECA studies and under accession phs004273 for

the MUPPITS studies.

REFERENCES

1. Frent S-M. Severe Asthma: Phenotypes, Endotypes and Therapeutic Response. In: Özdemir Ö, editor. Asthma - Diagnosis, Management and Comorbidities. London: IntechOpen; 2025.

2. Tsuo K, Zhou W, Wang Y, Kanai M, Namba S, Gupta R, et al. Multi-ancestry meta-analysis of asthma identifies novel associations and highlights the value of increased power and diversity. Cell Genom. 2022;2(12):100212.

3. Han Y, Jia Q, Jahani PS, Hurrell BP, Pan C, Huang P, et al. Genome-wide analysis highlights contribution of immune system pathways to the genetic architecture of asthma. Nat Commun. 2020;11(1):1776.

4. Pividori M, Schoettler N, Nicolae DL, Ober C, Im HK. Shared and distinct genetic risk factors for childhood-onset and adult-onset asthma: genome-wide and transcriptome-wide studies. Lancet Respir Med. 2019;7(6):509-22.

5. Olafsdottir TA, Theodors F, Bjarnadottir K, Bjornsdottir US, Agustsdottir AB, Stefansson OA, et al. Eighty-eight variants highlight the role of T cell regulation and airway remodeling in asthma pathogenesis. Nat Commun. 2020;11(1):393.

6. Namjou B, Lape M, Malolepsza E, DeVore SB, Weirauch MT, Dikilitas O, et al. Multiancestral polygenic risk score for pediatric asthma. J Allergy Clin Immunol. 2022;150(5):1086-96.

7. Dapas M, Lee YL, Wentworth-Sheilds W, Im HK, Ober C, Schoettler N. Revealing polygenic pleiotropy using genetic risk scores for asthma. HGG Adv. 2023;4(4):100233.

8. Jackson DJ, Bacharier LB, Gergen PJ, Gagalis L, Calatroni A, Wellford S, et al. Mepolizumab for urban children with exacerbation-prone eosinophilic asthma in the USA (MUPPITS-2): a randomised, double-blind, placebo-controlled, parallel-group trial. Lancet. 2022;400(10351):502-11.

9. Liu AH, Babineau DC, Krouse RZ, Zoratti EM, Pongracic JA, O'Connor GT, et al. Pathways through which asthma risk factors contribute to asthma severity in inner-city children. J Allergy Clin Immunol. 2016;138(4):1042-50.

10. Poowuttikul P, Saini S, Seth D. Inner-City Asthma in Children. Clin Rev Allergy Immunol. 2019;56(2):248-68.

11. Togias A, Fenton MJ, Gergen PJ, Rotrosen D, Fauci AS. Asthma in the inner city: the perspective of the National Institute of Allergy and Infectious Diseases. J Allergy Clin Immunol. 2010;125(3):540-4.

12. Pongracic JA, Krouse RZ, Babineau DC, Zoratti EM, Cohen RT, Wood RA, et al. Distinguishing characteristics of difficult-to-control asthma in inner-city children and adolescents. J Allergy Clin Immunol. 2016;138(4):1030-41.

13. Zoratti EM, Krouse RZ, Babineau DC, Pongracic JA, O'Connor GT, Wood RA, et al. Asthma phenotypes in inner-city children. J Allergy Clin Immunol. 2016;138(4):1016-29.

14. Gern JE, Visness CM, Gergen PJ, Wood RA, Bloomberg GR, O'Connor GT, et al. The Urban Environment and Childhood Asthma (URECA) birth cohort study: design, methods, and study population. BMC Pulm Med. 2009;9:17.

15. Altman MC, Gill MA, Whalen E, Babineau DC, Shao B, Liu AH, et al. Transcriptome networks identify mechanisms of viral and nonviral asthma exacerbations in children. Nat Immunol. 2019;20(5):637-51.

16. Mathias RA, Taub MA, Gignoux CR, Fu W, Musharoff S, O'Connor TD, et al. A continuum of admixture in the Western Hemisphere revealed by the African Diaspora genome. Nat Commun. 2016;7:12522.

17. Stoffel B, Sorkness C, Pech C. Use of a Single, Independent IRB: Case Study of an NIH Funded Consortium. Contemp Clin Trials Commun. 2017;8:114-21.

271   18.   Dapas M, Thompson EE, Wentworth-Sheilds W, Clay S, Visness CM, Calatroni A, et al. Multi-
272   omic association study identifies DNA methylation-mediated genotype and smoking exposure
273   effects on lung function in children living in urban settings. PLoS Genet. 2023;19(1):e1010594.

274   19.   Altman MC, Flynn K, Rosasco MG, Dapas M, Kattan M, Lovinsky-Desir S, et al. Inducible
275   expression quantitative trait locus analysis of the MUC5AC gene in asthma in urban populations of
276   children. J Allergy Clin Immunol. 2021;148(6):1505-14.

277   20.   Zhao H, Sun Z, Wang J, Huang H, Kocher JP, Wang L. CrossMap: a versatile tool for coordinate
278   conversion between genome assemblies. Bioinformatics. 2014;30(7):1006-7.

279   21.   Loh PR, Palamara PF, Price AL. Fast and accurate long-range phasing in a UK Biobank cohort. Nat
280   Genet. 2016;48(7):811-6.

281   22.   Howie B, Fuchsberger C, Stephens M, Marchini J, Abecasis GR. Fast and accurate genotype
282   imputation in genome-wide association studies through pre-phasing. Nat Genet. 2012;44(8):955-9.

283   23.   Das S, Forer L, Schonherr S, Sidore C, Locke AE, Kwong A, et al. Next-generation genotype
284   imputation service and methods. Nat Genet. 2016;48(10):1284-7.

285   24.   Daya M, Rafaels N, Brunetti TM, Chavan S, Levin AM, Shetty A, et al. Association study in
286   African-admixed populations across the Americas recapitulates asthma risk loci in non-African
287   populations. Nat Commun. 2019;10(1):880.

288   25.   Wang Y, Namba S, Lopera E, Kerminen S, Tsuo K, Lall K, et al. Global Biobank analyses provide
289   lessons for developing polygenic risk scores across diverse cohorts. Cell Genom.
290   2023;3(1):100241.

291   26.   Ge T, Chen CY, Ni Y, Feng YA, Smoller JW. Polygenic prediction via Bayesian regression and
292   continuous shrinkage priors. Nat Commun. 2019;10(1):1776.

293   27.   Lambert SA, Gil L, Jupp S, Ritchie SC, Xu Y, Buniello A, et al. The Polygenic Score Catalog as an
294   open database for reproducibility and systematic evaluation. Nat Genet. 2021;53(4):420-5.

295   28.   Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK:
296   rising to the challenge of larger and richer datasets. Gigascience. 2015;4.

297   29.   Collister JA, Liu X, Clifton L. Calculating Polygenic Risk Scores (PRS) in UK Biobank: A
298   Practical Guide for Epidemiologists. Front Genet. 2022;13:818574.

299   30.   Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A global
300   reference for human genetic variation. Nature. 2015;526(7571):68-74.

301   31.   Bergstrom A, McCarthy SA, Hui R, Almarri MA, Ayub Q, Danecek P, et al. Insights into human
302   genetic variation and population history from 929 diverse genomes. Science. 2020;367(6484).

303   32.   Conomos MP, Miller MB, Thornton TA. Robust inference of population structure for ancestry
304   prediction and correction of stratification in the presence of relatedness. Genet Epidemiol.
305   2015;39(4):276-93.

306   33.   Conomos MP, Reiner AP, Weir BS, Thornton TA. Model-free Estimation of Recent Genetic
307   Relatedness. Am J Hum Genet. 2016;98(1):127-48.

308   34.   Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. Robust relationship
309   inference in genome-wide association studies. Bioinformatics. 2010;26(22):2867-73.

310   35.   McCaw ZR, Lane JM, Saxena R, Redline S, Lin X. Operating characteristics of the rank-based
311   inverse normal transformation for quantitative trait analysis in genome-wide association studies.
312   Biometrics. 2020;76(4):1262-72.

313   36.   Sofer T, Zheng X, Gogarten SM, Laurie CA, Grinde K, Shaffer JR, et al. A fully adjusted two-stage
314   procedure for rank-normalization in genetic association studies. Genet Epidemiol. 2019;43(3):263-
315   75.

316   37.   Zhou W, Kanai M, Wu KH, Rasheed H, Tsuo K, Hirbo JB, et al. Global Biobank Meta-analysis
317   Initiative: Powering genetic discovery across human disease. Cell Genom. 2022;2(10):100192.

38. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current polygenic
risk scores may exacerbate health disparities. Nat Genet. 2019;51(4):584-91.
39. Ferreira MAR, Mathur R, Vonk JM, Szwajda A, Brumpton B, Granell R, et al. Genetic
Architectures of Childhood- and Adult-Onset Asthma Are Partly Distinct. Am J Hum Genet.
2019;104(4):665-84.
40. Natarajan R, Szczesny B, Kanchan K, Esquinca E, Boorgula MP, Chavan S, et al. Mediation of
Polygenic Asthma Risk Through Gene Expression. Allergy. 2025. doi: 10.1111/all.70101.
41. Sato K, Ohno T, Ishii T, Ito C, Kaise T. The Prevalence, Characteristics, and Patient Burden of
Severe Asthma Determined by Using a Japan Health Care Claims Database. Clin Ther.
2019;41(11):2239-51.
42. Chupp G, Murphy KR, Gandhi HN, Gilbert I, Bleecker ER. Asthma control in the United States:
Relationships between short-acting beta(2)-agonist and systemic corticosteroid use. Ann Allergy
Asthma Immunol. 2024;133(3):302-9.
43. Gokhale M, Hattori T, Evitt L, Lenney W, Nordstrom B, Collins J, et al. Burden of asthma
exacerbations and health care utilization in pediatric patients with asthma in the US and England.
Immun Inflamm Dis. 2020;8(2):236-45.
44. Hansen S, von Bulow A, Sandin P, Ernstsson O, Janson C, Lehtimaki L, et al. Prevalence and
management of severe asthma in the Nordic countries: findings from the NORDSTAR cohort. ERJ
Open Res. 2023;9(2).
45. Chang D, Hunkapiller J, Bhangale T, Reeder J, Mukhyala K, Tom J, et al. A whole genome
sequencing study of moderate to severe asthma identifies a lung function locus associated with
asthma risk. Sci Rep. 2022;12(1):5574.
46. Shrine N, Portelli MA, John C, Soler Artigas M, Bennett N, Hall R, et al. Moderate-to-severe
asthma in individuals of European ancestry: a genome-wide association study. Lancet Respir Med.
2019;7(1):20-34.
47. Belsky DW, Sears MR, Hancox RJ, Harrington H, Houts R, Moffitt TE, et al. Polygenic risk and
the development and course of asthma: an analysis of data from a four-decade longitudinal study.
Lancet Respir Med. 2013;1(6):453-61.
48. Moll M, Sordillo JE, Ghosh AJ, Hayden LP, McDermott G, McGeachie MJ, et al. Polygenic risk
scores identify heterogeneity in asthma and chronic obstructive pulmonary disease. J Allergy Clin
Immunol. 2023;152(6):1423-32.
49. Sherafati A, Norland K, Naderian M, Schaid DJ, Kullo IJ. Polygenic Risk and Coronary Artery
Disease Severity. Circ Genom Precis Med. 2024;17(5):e004470.
50. Reid S, Alexsson A, Frodlund M, Morris D, Sandling JK, Bolin K, et al. High genetic risk score is
associated with early disease onset, damage accrual and decreased survival in systemic lupus
erythematosus. Ann Rheum Dis. 2020;79(3):363-9.
51. Clay S, Alladina J, Smith NP, Visness CM, Wood RA, O'Connor GT, et al. Gene-based association
study of rare variants in children of diverse ancestries implicates TNFRSF21 in the development of
allergic asthma. J Allergy Clin Immunol. 2024;153(3):809-20.
52. Lezmi G, Lejeune S, Pin I, Blanchon S, Bouazza N, Jolaine V, et al. Factors Associated with
Asthma Severity in Children: Data from the French COBRAPed Cohort. J Allergy Clin Immunol
Pract. 2021;9(5):1969-79.