A Human Glomerular Disease Atlas defines the APOL1-JAK-STAT feed forward loop in focal segmental glomerulosclerosis

Authors

Phillip J. McCown[1*], Charlotte Boys[2*], Somenath Datta[3*], Viji Nair[1,4], Edgar Otto[1], Damian Fermin[1], John Hartman[1], Francesca Annese[5], Fadhl Alakwaa[1], Khun Zaw Latt[1], Cathy Smith[1], Bradley Godfrey[1], Maria Larkina[1], Alison E. Ashley-Koch[3], Melanie E. Garrett[3], Savannah Moore[6], Karen Soldano[3], Mathew Alaba[1], Laura H. Mariani[1], Abhijit S. Naik[1], Robin Fallegger[2], Sean Eddy[1#], Julio Saez-Rodriguez[2,7#], Opeyemi A. Olabisi[3#], Matthias Kretzler[1,4#], for the Nephrotic Syndrome Study Network (NEPTUNE)

*These authors contributed equally to this work; #shared senior authors

[1]Division of Nephrology, Department of Internal Medicine, University of Michigan, Ann Arbor, Michigan, USA.

[2]Heidelberg University, Faculty of Medicine, and Heidelberg University Hospital, Institute for Computational Biomedicine, Heidelberg, Germany.

[3]Division of Nephrology, Duke Molecular Physiology Institute, Duke University School of Medicine, Durham, North Carolina, USA.

[4]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA

[5]Department of Precision and Regenerative Medicine and Ionian Area – Nephrology, Dialysis and Transplantation Unit, University of Bari "Aldo Moro", Bari, Italy.

[6]Division of Nephrology, Massachusetts General Hospital, Boston Massachusetts, USA

[7]European Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Genome Campus, Hinxton, Cambridge, United Kingdom.

[#]**Corresponding authors:**

Sean Eddy
Associate Research Scientist,
Department of Internal Medicine, Medical School, University of Michigan
1150 West Medical Center Drive,
Ann Arbor, MI 48109 USA
seaneddy@med.umich.edu

Matthias Kretzler
Division of Nephrology, Department of Internal Medicine,
University of Michigan,
1150 W. Medical Center Drive, Ann Arbor, MI 48109, USA
Kretzler@med.umich.edu

Opeyemi A. Olabisi
Division of Nephrology, Department of Medicine,
Duke Molecular Physiology Institute,
Duke University School of Medicine, 300 N. Duke St., Ofc#50-104, Durham, NC 27701, USA
opeyemi.olabisi@duke.edu

Julio Saez-Rodriguez
European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI)
Genome Campus, Hinxton, CB101SD, UK
saezlab@ebi.ac.uk

Abstract (limit 150 words – current 145)

To map transcriptional programs in rare glomerular diseases, single-nucleus RNA sequencing (snRNAseq) on kidney biopsies (N=120) from the Nephrotic Syndrome Study Network were integrated with snRNAseq and single-cell sequencing (scRNAseq) of reference kidney tissue (N=50) to create the Omnibus of CElls And Nuclei (OCEAN). Unsupervised analysis of multi-cellular programs identified that JAK-STAT pathway in podocytes and endothelial cells was associated with clinical measures of disease severity. JAK-STAT pathway activity was strongly correlated with apolipoprotein1 (APOL1) mRNA transcript expression and high risk APOL1 variant genotype, a major risk factor for FSGS. These findings were confirmed in an independent study of Black participants where loss of APOL1 function decreased JAK-STAT pathway activation in patient derived podocyte *ex vivo* models. The findings are consistent with a feed forward loop regulating the JAK-STAT-APOL1 driven tissue damage, providing mechanistic support for the JUSTICE Phase II trial targeting JAK activation in APOL1-mediated kidney disease.

Introduction

Kidney diseases are a global health threat[1]. Among these, primary glomerular diseases, which include focal segmental glomerulosclerosis (FSGS), minimal change disease (MCD), and IgA nephropathy (IgAN), though individually rare, are associated with debilitating symptoms and contribute substantially to the kidney failure pandemic[2]. Glomerular diseases are often characterized by nephrotic syndrome (NS), hallmarked by proteinuria, hypoalbuminemia, hyperlipidemia, and edema. Diagnosis is based on histological features related to patterns of injury in kidney biopsies that determine treatment course [3]. The only FDA-approved medical treatments are prednisone and adrenocorticotrophic hormone, which do not prevent progression to kidney failure in most patients and cause severe side effects. Guideline-based[3] clinical care includes off-label use of second line immunosuppressive medications. The variability in response to these immunosuppressives, including steroids, adds to the disease burden. While many therapies are currently under development[4,5] none have been approved specifically for FSGS and, until recently[6-8], no therapies were approved specifically for any glomerular disease.

The heterogeneity of etiology and disease progression, and paucity of treatment options, highlight the essential prerequisite of a comprehensive knowledge environment defining the clinical, structural and molecular determinants of glomerular diseases to develop and implement effective treatments. To address this challenge, the Nephrotic Syndrome Study Network (NEPTUNE) has prospectively enrolled and followed participants with glomerular diseases over the last 17 years[9]. The NEPTUNE study has expanded our understanding of molecular mechanisms in FSGS[10,11] and APOL1-mediated kidney disease (AMKD)[12,13].

The presence of select apolipoprotein L1 (APOL1) single nucleotide polymorphisms (SNPs) in people of African ancestry is known to contribute to increased risk of developing AMKD,

with worse outcomes for those with FSGS[14-17]. Increased expression of APOL1 variants (G1 and G2) and not the reference APOL1 allele (G0) in podocytes, specialized cells in the glomerular filter compartment of kidneys, caused dose-dependent injury[18,19] and FSGS[20]. Meanwhile, a specific APOL1 variant (N264K) has been identified to exhibit kidney protective effects[19,21,22]. However, only 20% of carriers of the high-risk APOL1 genotype are estimated to develop AMKD[14,23]. The basis of this incomplete penetrance is unknown.

The advent of single nucleus RNA sequencing (snRNAseq) and single cell RNA sequencing (scRNAseq) technologies offers an unprecedented opportunity to understand molecular mechanisms of disease and identify novel cell-states and niches. Multi-institutional research consortia, such as the Human Cell Atlas[24,25] and the Kidney Precision Medicine Project[26,27] have provided key insights into human health and disease at a cellular-level. NEPTUNE now deploys such mechanistic disease definitions to develop and test targeted therapies[28].

In this study, kidney biopsies contributed by NEPTUNE participants and kidney donor samples from the Human Kidney Transplant Transcriptomic Atlas (HKTTA), as reference tissue, were used to generate and integrate snRNAseq and scRNAseq profiles to create the Omnibus of CElls And Nuclei (OCEAN) dataset (https://neptune-ocean.cxg.miktmc.org/). An unsupervised analysis of the OCEAN data identified multi-cellular programs underlying glomerular disease heterogeneity and enabled cell-type specific explorations in FSGS leading to the characterization of JAK-STAT signaling and APOL1 interactions. Transcriptomic, genetic and proteomic analyses followed by validation using human podocytes derived from induced pluripotent stem cells (IPSCs) enabled detailed investigations into the molecular interactions in glomerular diseases, particularly in the context of the APOL1 risk allele (**Fig. 1**). Importantly, loss of APOL1 function was discovered to be associated with attenuation of JAK-STAT activation,

consistent with a feed-forward regulatory loop between APOL1 and JAK-STAT. These findings further support the JUSTICE trial[29], part of the NEPTUNE Match precision medicine trial platform[28], evaluating JAK inhibitors in people with APOL1 risk variants. This study is illustrative of the power of the OCEAN dataset to drive hypothesis generation for further mechanistic studies and clinical trials in rare glomerular diseases.

Results

Building the OCEAN - representing a comprehensive collection of glomerular disease kidney cell types

We derived the OCEAN dataset by sequencing kidney tissues of 120 participants obtained at the time of enrollment (baseline diagnostic biopsy) into NEPTUNE (**Table 1**) and stored in RNAlater for up to 10 years. The number of nuclei captured was not affected by time in storage. Reference data in OCEAN comprise kidney tissue profiles (scRNAseq and snRNAseq) of 50 participants in HKTTA (living donor, LD, **Table 1**). NEPTUNE recruited participants prior to biopsy based on clinical profile, capturing a diverse group of participants as subsequently defined by descriptive histopathology adjudication. Participants were predominantly diagnosed with FSGS (N = 33), MCD (N = 24) or proteinuric IgAN (N = 29)[9]. The dataset also includes biopsies from 34 participants with other proteinuric diseases or those pending histological adjudication. snRNAseq and/or scRNAseq profiles were integrated into an atlas of 1,047,781 nuclei and cells (**Fig. 2a**) that formed 17 major cell clusters (**Fig.2b**) and 28 discrete clusters (**SFig. S1a, i**), representing all the nephron cell types from both glomerular and tubulointerstitial compartments of the kidney. The number of nuclei contributed were similar for FSGS, MCD, and IgAN, and cell numbers were proportionate to the number of participants in each group (**Fig. 2c**). Clusters were preserved across all major diagnoses and LD sample types (**SFig. S1b-h**). Since biopsies are typically obtained from the kidney cortex, as expected, there were more cells in OCEAN from the tubular compartment, with the most from the proximal tubule (PT) and thick ascending limb (TAL) (**Fig 2d**). The distribution of tubular cells by cell types in the main groups (LD, FSGS, MCD and IgAN) were similar with all groups contributing cells to each cell type with a slightly increased contribution of PT cells from MCD (**Fig 2e**). However, there were

differences in the distribution of glomerular cells. The parietal epithelial (PEC), podocyte (POD) and a cluster that contained both PEC and POD (PEC/POD) cell type markers (claudin 1, CLDN1; nephrin, NPHS1; respectively), that account for 4% of the total number of cells, were differently distributed across the LD, FSGS, MCD, and IgA groups (**Fig. 2f**). Containing both PEC and POD markers, the PEC/POD cluster is suggestive of a transitioning state between the two cell types.

Discovery of disease pathways - Unsupervised analysis of OCEAN transcriptomic data provides molecular context to patient heterogeneity in glomerular diseases

Multicellular factor analysis (MFA)[30], a variance decomposition method which can be conceptualized as an extension of principal component analysis (PCA) across multiple data 'views', was used to quantify inter-patient variation in the molecular data. Data views included gene expression profiles aggregated at the major cell type level (**STable S1**) and inferred ligand-receptor interactions between cell types. MFA identified a set of latent factors, each describing a major axis of variation across participants. Patient samples received a score for each factor, which were characterized by gene weights within each cell type and ligand-receptor weights for sender-receiver cell pairs[31] (**Fig. 3a**). This analysis was carried out on the subset of snRNAseq samples within OCEAN profiled on the 10x platform comprising 93 NEPTUNE and 16 reference samples across 846,129 sequenced nuclei (**Fig. 3b**).

MFA with 20 latent factors explained 70–80% of inter-sample variance in gene expression per cell type (**Fig. 3c**). We categorized these latent factors into four groups: 1) batch factors, which primarily distinguished samples processed in different experimental batches,

likely due to technical variation in sequencing or sample handling; 2) outlier factors, which captured variability driven by a single sample deviating from the cohort; 3) technical factors, which separated samples based on expected clinical distinctions; and 4) biological factors, which accounted for 15–30% of variance and reflected disease-related gene expression changes. Among the six biological factors, two latent factors, Factor 3 and Factor 5, were significantly associated with kidney function at the time of biopsy as measured by eGFR ($r = -0.57$, p-adj $=$ 1.28e-08 and $r = 0.47$, p-adj $=$ 3.65e-06 respectively), as well as other clinical covariates (**Fig.3d**). Thus, the unsupervised exploration distinguished biological from technical variability and enabled further investigations into variations related to disease.

Discovery of disease pathways - Two axes of variation reveal a "cell state map" of kidney function decline in glomerular disease

To explore the biological and clinical relevance of these factors, we examined their relationship to participant demographics and kidney function. Factor 3 and Factor 5 scores were orthogonal, representing two distinct axes of variation. A two-dimensional map, with participant coordinates based on their Factor 3 and Factor 5 scores, helped visualize heterogeneity in disease severity (**Fig. 3e**). Together, they define a molecular spectrum of kidney function in glomerular failure, ranging from younger individuals with preserved kidney function to older individuals with progressive decline in eGFR (**Fig. 3e**). We stratified the two-dimensional participant map into four quadrants: HighF3LowF5, LowF3LowF5, LowF3HighF5, and HighF3HighF5 groups, where 'High' corresponds to a factor score > 0 and 'Low' to a factor score < 0. Young participants with higher eGFR (age < 18 years, eGFR ≥ 45 ml/min/1.73m²) mainly clustered in the LowF3HighF5 group (18 out of 28). In contrast, adults with low eGFR (age ≥ 18 years,

eGFR < 45 ml/min/1.73m²) were predominantly (18 out of 24) grouped in the HighF3LowF5

quadrant (**Fig. 3f**). Overall, eGFR decreased significantly and progressively between the

LowF3HighF5 group (best kidney function), to LowF3LowF5 (middle kidney function), and

HighF3LowF5 (worst kidney function) (**Fig. 3g**). Crucially, the LowF3HighF5 vs HighF3LowF5

stratification remained statistically significant ($p < 0.05$) not only for the demographic extremes

(age < 18 years or eGFR < 45 ml/min/1.73m²), but also among adults with preserved eGFR (age

≥ 18 years, eGFR ≥ 45 ml/min/1.73m²) (**SFig. S2a**). These findings suggest that Factor 3 and

Factor 5 provide a molecular readout of the cell states of the kidney at different levels of eGFR

and may be useful for stratifying patients along a trajectory of kidney function decline.

Discovery of disease pathways - Multicellular program of kidney function decline: decreased
metabolic activity and the establishment of a profibrotic state

Factor 5, which positively correlated with kidney function at the time of diagnostic biopsy,

explained the variance in gene expression **(Fig. 3h)** primarily in immune cells (13.73%),

endothelial cells (EC; 12.00%), fibroblasts (FIB; 11.50%), and vascular smooth

muscle/mesangial cells (vSMC/MC; 9.63%).

Pathway analysis[32] showed a significant increase in JAK-STAT as Factor 5 decreased,

particularly in endothelial cells (pval < 1e-9), together with general activation of mitogen-

activated protein kinase (MAPK), nuclear factor kappa B (NF-kB) and transforming growth

factor beta (TGF-β) pathways (**Fig. 3i**). Overrepresentation analysis of gene loadings revealed

that genes involved in oxidative phosphorylation and xenobiotic metabolism were more highly

expressed in high-Factor 5 participants and progressively diminished in low-Factor 5 participants

(**Fig. 3j**). In fibroblasts, genes related to fibroblast activation and extracellular matrix remodeling were upregulated as Factor 5 score decreased, suggesting a shift toward a profibrotic state.

In immune cells, genes associated with allograft rejection were downregulated in high-Factor 5 samples, possibly reflecting immunosuppressive effects of the more common steroid treatment in younger participants with preserved kidney function prior to kidney biopsy, who cluster within the LowF3HighF5 quadrant (**Fig. 3f**). Although Factor 5 score is associated with steroid treatment (**Fig. 3d**), this association does not persist after stratification for pediatric participants (**SFig. S2b**). Together, these results suggest that Factor 5 primarily captures transcriptional programs associated with kidney cellular metabolism which are more active in pediatric participants and diminish as kidney function declines. Further, this loss of metabolic activity occurs in coordination with the establishment of a profibrotic state.

Factor 3, negatively correlated with kidney function, described transcriptional changes in the tubular compartment and explained the variance, particularly within the DCT (8.64%) and PT (8.93%) (**Fig. 3h**). As Factor 3 score increased, the injury markers WAP four-disulfide core domain protein 2 (WFDC2), prominin 1 (PROM1), thymosin beta 4 x-linked (TMSB4X), thymosin beta 10 (TMSB10), vimentin (VIM), vascular cell adhesion molecule 1 (VCAM1), and hepatitis A virus cellular receptor 1 (HAVCR1) were upregulated in PT (**Fig. 3k**), consistent with a transition towards a stress and repair phenotype. In DCT, increased Factor 3 scores were associated with upregulation of many of these injury markers (WFDC2, PROM1, TMSB4X) and downregulation of differentiation markers uromodulin (UMOD) and transient receptor potential cation channel, subfamily M, member 6 (TRPM6), supporting a loss of epithelial identity.

Beyond intrinsic tubular injury signatures, Factor 3 was strongly associated with fibroblast-mediated collagen signaling in tubular cell types (**Fig. 3k**). Overrepresentation

analysis revealed widespread upregulation of epithelial-mesenchymal transition (EMT)-associated genes across all cell types with an increasing Factor 3 score (**Fig. 3m**). Pathway enrichment analysis further indicated that fibrotic and inflammatory pathways—JAK-STAT, NF-κB, TGF-β, and tumor necrosis factor alpha (TNF-α)—were significantly upregulated across all cell types in high-Factor 3 participants (**Fig. 3i**). The Factor 3 score was also positively correlated with immune cell infiltration, particularly monocyte/macrophage abundance (R = 0.73, p < 2.2e-16) and natural killer/NK T cell abundance (R = 0.67, p = 4.4e-14) (**Fig. 3l**). These findings suggested that Factor 3 reflected an injury-driven, multicellular transcriptional program characterized by tubular stress, immune activation, and progressive fibrosis, all associated with worsening kidney function.

Mapping cell-specific effects - APOL1 expression is associated with kidney function in FSGS

Unsupervised analysis of the OCEAN snRNAseq data established JAK-STAT activation as a key disease pathway associated with kidney function decline. Since JAK-STAT activation is known to induce APOL1 mRNA expression[33,34], which was not originally included as a feature in MFA model training, but is one of the strongest genetic risk factors for developing nephrotic syndrome, we examined how APOL1 expression progressed from LowF3High5 through to HighF3LowF5. APOL1 expression and distribution across cell types in OCEAN identified the highest level of APOL1 expression in the nuclei of podocytes, followed by endothelial cells, and with lower levels detected in nuclei of other cell types (**Fig. 4a**). Within podocytes, APOL1 expression was highest in nuclei from participants with FSGS (Log$_2$FC=1.05, padj<0.0001, **Fig. 4b**）compared to podocytes from reference samples and in pseudobulk podocyte profiles, where sufficient sequenced nuclei were available to make a reliable gene quantitation (≥20, **STable**

S2) (Log$_2$FC=1.1, p<0.05) (**Fig. 4b, SFig3a**). Among the pseudobulk podocyte profiles from participants with FSGS, APOL1 increased significantly between LowF3HighF5 (higher kidney function; lower JAK-STAT activation) and HighF3LowF5 (lower kidney function; higher JAK-STAT activation) groups (**Fig. 4c**).

Mapping cell-specific effects - APOL1 expression is correlated with JAK-STAT activation in participants with FSGS, and is elevated in those with two APOL1 high risk alleles

As we previously observed JAK-STAT activation in patients with FSGS and IgAN[11,35], and observed the association between Factor 3/Factor 5 scores with both JAK-STAT activity and APOL1 expression in FSGS podocytes, we anticipated an association between JAK-STAT activity and APOL1 expression in bulk RNAseq profiles of glomeruli from participants with FSGS (participant clinical and demographic information included in **STable S3**) that could be assessed using the previously published transcriptional fingerprint JAK-STAT pathway activity score (PAS)[11,35]. As APOL1 was originally a component of the JAK-STAT PAS, we recomputed the JAK-STAT PAS without APOL1 to enable correlation with APOL1 mRNA expression. The computed JAK-STAT PAS with APOL1 was virtually identical to JAK-STAT PAS computed without APOL1 (r=0.998, p<0.0001, **SFig3b**), thus JAK-STAT PAS (without APOL1) performed as a reasonable to our originally described JAK-STAT PAS and was used for all subsequent analyses.

As expected, JAK-STAT PAS was strongly correlated with APOL1 expression in glomerular RNAseq profiles from participants with FSGS (r = 0.70, p < 0.0001, **Fig. 4d**). Participants with 2 APOL1 risk alleles had higher APOL1 mRNA expression and JAK-STAT PAS and therefore

investigated potential association between risk allele status and concordant activation of JAK-STAT and APOL1. At the nuclei level and pseudobulk podocyte level, there was no allele specific difference detectable, possibly a consequence of the small sample size captured with a sufficient number of podocytes in OCEAN. However, in bulk RNAseq profiles available in NEPTUNE[10], participants with 2 risk alleles (n = 16) had the highest APOL1 mRNA expression (p < 0.05) compared to those with 1 (n = 13) or 0 risk alleles (n = 54) (**Fig. 4e**). Participants with 0 and 1 risk alleles had similar APOL1 mRNA expression levels, in pairwise comparisons, participants with 2 risk alleles (n = 16) had significantly higher expression than participants with 0 risk alleles (n = 54) (p < 0.05) (**SFig3c**). Similarly, the JAK-STAT PAS was highest in participants with 2 risk alleles (p < 0.05) (**Fig. 4f**), with significant difference between 0 and 2 risk alleles (p < 0.01) and an intermediate score for a single risk allele (**SFig2d**). Taken together, high APOL1 expression in podocytes in FSGS, particularly among those in the HighF3LowF5 group, is coupled with JAK-STAT activity suggestive of coordinated regulation of JAK-STAT activity and APOL1 regulation in FSGS.

Genetic loss of APOL1 function via N264K mutation reduced JAK-STAT target genes in podocytes, endothelial cells and glomerular pseudobulk profiles

To test the JAK-STAT-APOL1 interaction in the context of the N264K protective mutation[21] that blocks the pore-forming function of APOL1[36], we identified podocytes from two participants in OCEAN with FSGS and the APOL1 N264K variant. In these two cases, genes downstream of JAK-STAT activation including APOL1, interferon regulatory factor 1 (IRF1), one of the key transcriptional regulators of APOL1 expression[37] and intracellular adhesion molecule 1 (ICAM1) (all three were originally developed as key drivers of the JAK-STAT

PAS[11,35]) and signal transducer and activator of transcription 1 (STAT1) were assessed. At the combined nuclei level, all four were lower in podocytes with APOL1 HR and N264K compared to podocytes with APOL1 HR without N264K (p-adj<0.0001 for all 4 genes, **Fig. 4g**). Pseudobulk analysis was precluded by the small sample size and uneven nuclei capture across participants (**STable S2**).

Based on observing the gene expression and genetic analyses supportive of a JAK-STAT-APOL1 regulatory loop where both JAK-STAT activity was associated with APOL1 expression and APOL1 activity suggested positive regulation of JAK-STAT activity, we proceeded to test this hypothesis directly in a podocyte model system.

Functional validation in cell models: Clinical and demographic characteristics of APOL1 study participants

To test the mechanistic link between APOL1 and JAK-STAT signaling and investigate the role of this mechanism in the incomplete penetrance of APOL1-mediated kidney disease (AMKD), we established an independent case-control cohort for an iPSC-derived podocyte (iPOD) model of APOL1-mediated FSGS. We enrolled Black adults with biopsy-proven FSGS who also carried the APOL1 HR genotype, HR Cases (N = 27) and healthy Black volunteers (N=30) without diabetes, proteinuria (UPCR > 0.2 g/g)[38,39] or loss in kidney function (eGFR < 60 ml/min, (**Fig. 5a, Table 2, STable S4**) for the HR control group. To increase the likelihood that an HR control is a true control, we preferentially enrolled participants 50 years or older, since APOL1-mediated FSGS becomes evident by age 50 in most cases[20,40] (**Table 2; STable S4**). By contrast, HR cases were younger, with a mean age of 41 years, consistent with early age of onset

of APOL1-mediated FSGS. HR cases also had significant proteinuria and varying degrees of eGFR decline. The high frequency of G1 allele and low frequency of G2 allele among both HR cases and HR controls were aligned with prior publications[20,41] (**Table 2; STable S4**). There were slightly more men among the HR cases while more women had been enrolled as HR controls (**Table 2**).

Functional validation in cell models - iPSC-derived Podocytes (iPODs) Express Podocyte Markers and Respond to IFN-γ

To discover the dysregulated signaling pathways that distinguish podocytes of HR cases from those of HR controls, we differentiated participant derived induced pluripotent stem cells (iPSCs) from both groups into podocyte-like cells (iPODs)[42]. iPODs retained the complete genetic endowment of each participant, enabling capture of participant-specific molecular signatures with high fidelity. iPODs of both HR Cases and HR controls also expressed key podocyte markers, supporting their use as podocytes surrogates (**Fig. 5b**). Similar to human primary podocytes, stimulation of iPODs with interferon gamma (IFN-γ), a known inducer of APOL1 expression via JAK1/2 kinase activation, caused upregulation of APOL1 expression across all samples (**Fig. 5b**), with clustering of untreated and IFN-γ-treated iPODs into two distinct groups (**Fig. 5c**). Quality control assessment separated iPODs into Group A and Group B based on marker expression profiles, with particular attention to podocalyxin like (PODXL), phospholipase A2 receptor 1 (PLA2R1), integrin subunit alpha 3 (ITGA3). Group B iPODs had low expression of podocyte markers (PLA2R1, PODXL and ITGA3) and high expression of olfactomedin 3 (OLFM3), suggesting Group B iPODs were poorly differentiated (**Fig. 5c**). Therefore, the Group B iPODs were excluded from further analysis. Taken together, these results

demonstrated that the iPODs expressed characteristic podocyte markers, responded to IFN-γ stimulation, and exhibited variability in APOL1 expression influenced by genetic risk factors.

Functional validation in cell models - JAK-STAT Signaling is Potentiated in HR Cases and leads to higher APOL1 expression

To investigate the differential response to IFN-γ stimulation, gene expression profiles were analyzed in HR cases and HR controls under both baseline and stimulated conditions. Comparative analysis showed that under IFN-γ stimulation, there were 381 differentially expressed genes (DEGs, padj <0.05 and fold change ≥ 2) in HR cases of which 362 genes were upregulated and 19 genes downregulated. In HR controls, 281 DEGs with 277 genes upregulated and 4 genes downregulated were observed (**Fig. 5d**). Of these, 272 DEGs were common to both HR Cases and HR Controls. Importantly, 109 DEGs were unique to HR cases, of which 94 genes were upregulated and 15 downregulated. Pathway analysis of the 94 upregulated genes using KEGG demonstrated significant associations with activation of immune and inflammatory signaling pathways, including upregulation of pro-inflammatory cytokines, cytokine receptors and chemokines (**Fig. 5e**). These include CXCL2 (MIP-2a) and TNFRSF1B (TNFR2) reported to be associated with faster progression of kidney disease among individuals with high-risk APOL1 genotypes[43]. Further analysis of the 94 upregulated genes using ENCODE TF ChIP-Seq data revealed STAT1 (padj < 0.0006) and STAT2 (padj < 0.003), key components of JAK-STAT pathway (**Fig. 5f**), as the top enriched transcription factors. Consistent with this finding, individual analysis of key validated STAT1-regulated genes (dashed box, Fig. 5g)[11,44] and JAK-STAT pathway-associated genes—STAT1, STAT2, IRF1, IRF2, and IRF8—revealed higher expression levels in HR cases relative to HR controls (**Fig. 5g**). Notably, a trend towards higher

APOL1 expression was evident in HR cases compared to HR controls further supporting JAK-STAT driving activation in APOL1 HR cases.

Baseline analysis revealed that the levels of JAK-STAT components, including total STAT1, were similar between HR cases and HR controls, as shown in **Fig. 5h**. Comparative analysis of the whole genome of HR cases and HR controls did not identify variants in JAK-STAT genes or other genes that reached genome-wide significance (**SFig. S4**). However, upon stimulation with IFN-γ, HR cases exhibited higher STAT1 and IRF1 mRNA and protein (**Fig. 5i and j, SFig.S5**). Notably, the phosphorylation of STAT1, a critical activation step, did not appear to be markedly increased in HR cases versus controls, pointing to alternative mechanism(s) for the enhanced transcriptional activation of target genes in HR cases. Further investigation demonstrated elevated expression of IFN-γ-responsive genes, including CXCL11, reinforcing the heightened sensitivity of HR cases to IFN-γ signaling (**Fig. 5j**). Notably, treatment with the JAK inhibitor baricitinib effectively reversed the IFN-γ -induced expression of STAT1, IRF1, and APOL1 in HR cases (**Fig. 5i and j**), thereby confirming that the observed effects are triggered downstream of JAK1/2 kinase activation.

Functional validation in cell models - APOL1 Protein Localization and Monovalent Cation Transport in iPODs

We and others previously reported that APOL1 G1 or G2-mediated transport of K☐ and Na☐ across the plasma membrane is the proximal driver of APOL1-mediated cytotoxicity including podocytopathy[45-47]. Here, we determined that the APOL1 protein localized prominently on the plasma membrane of iPODs derived from HR cases, correlating with the elevated APOL1

expression levels (**Fig. 5k**). To study ion transport dynamics, rubidium (Rb▢) was employed as a tracer for potassium (K▢) due to its physicochemical similarity and suitability for transport assays[48,49]. We ascertained that iPODs of HR cases displayed significantly higher levels of Rb▢ efflux compared to HR control samples, indicating enhanced ion transport activity associated with increased APOL1 expression (**Fig. 5l**). Furthermore, treatment with baricitinib, partially reversed the elevated Rb▢ efflux in HR case iPODs (**Fig. 5l**). This suggested that the APOL1-mediated ion transport is modulated, at least in part, by JAK-dependent signaling pathways, presenting a potential therapeutic target for addressing APOL1-associated cellular dysfunction in podocytes.

Functional validation in cell models - APOL1-knockout in HR iPODs attenuated JAK-STAT signaling and abolished IFNγ-induced Rb$^+$ efflux.

To explore the hypothesis that APOL1 risk variants activate JAK-STAT signaling, we used CRISPR/Cas9 to knock out APOL1 in iPODs generated from a participant with G1G2 APOL1 genotype with diagnosis of FSGS **(Fig. 6a)**. We confirmed lack of IFN-γ-induced APOL1 protein expression by immunofluorescence and immunoblot (**Fig. 6b and c**). IFN-γ induced robust expression of APOL1 and other interferon regulated genes that are components of the JAK-STAT pathway as confirmed by qPCR (**Fig. 6d**) and RNAseq (**Fig. 6e**). Consistent with lack of APOL1 protein, the APOL1-knockout iPODs failed to efflux rubidium compared to APOL1-competent, isogenic control iPODs (**Fig. 6f**).

Validation of potentiated JAK-STAT pathway in HR cases by comparing with NEPTUNE cohort.

Lastly, we evaluated the 94 genes up-regulated in IFN-γ-induced cases, enriched for JAK-STAT activation pathway in the iPSC system (Fig. 5e) for concordant expression changes in the glomerular biopsy transcriptome of patients with FSGS; 87 genes were detected in the glomerular transcriptome of patients with FSGS and 26 were differentially regulated in FSGS compared to LD (p-adj<0.05). Of the 26 differentially regulated genes, 19 (73%) were concordantly regulated (ie. genes that were selectively up-regulated in IFN-γ iPODs from APOL1 HR cases were also up-regulated in FSGS compared to LD), while only 7 genes were discordantly regulated (ie. genes that were selectively up-regulated in IFN-γ iPODs from APOL1 HR cases were down-regulated in FSGS compared to LD). These data support translation of the IFN-γ stimulation in iPSC podocytes *ex vivo* to the APOL1-JAK-STAT interaction observed *in vivo*.

Taken together, these results strongly support the model where cytokine activation of JAK-STAT induces APOL1, which further acts in a feed forward manner to potentiate JAK-STAT signaling. Ablation of APOL1 activity, either genetically through N264K mutation, or through targeted knockdown of APOL1, suppresses JAK-STAT dependent signaling consistent with this model (**Fig. 7**).

Discussion

Here, we report the generation of OCEAN, a large-scale single cell and single-nucleus transcriptomic rare glomerular diseases atlas for the community to investigate cell-level effects to address a variety of different questions, including cellular changes with disease trajectory and across disease entities leading to potential cell selective markers for evaluating kidney health. This dataset and the subsequent use case with APOL1 demonstrate the power of molecular disease definitions in rare glomerular disease to bring novel therapies into the clinic. We illustrated the utility of coupling large scale single nucleus-level profiling with existing transcriptomic data and analysis suites to glean insight into underlying mechanisms in rare kidney diseases and association with clinical and genetic phenotypes. The study also details technical variation such as known batch effects so that OCEAN can be effectively used as a resource for the kidney research community to enhance mechanistic understanding of disease.

To define the key underlying determinants of glomerular disease on a cell resolved level we deployed a multicellular factor analysis (MFA) framework[30]. MFA allowed unbiased mapping of the core determinants of observed variance in OCEAN in terms of multi-cellular factors within different cell-types. These factors were interpreted based on the underlying molecular mechanisms such as pathways and cell-cell communication. By correlating these factors to patient-level phenotypic and clinical characteristics, we identified general and etiology-specific drivers of disease progression. Factors associated with kidney damage, showed enrichment of molecular programs known to be active in progressive loss of kidney function, more broadly, but now resolved at the cellular level in rare glomerular diseases. Demonstrating the activation of these disease progression programs to be in a disease stage but not disease-type specific manner is a critical insight for the glomerular disease community, as it provides the

rational to include primary glomerular diseases into umbrella trials cutting across conventional disease categories, rather targeting shared mechanism of disease (NCT05003986)[28,50].

Localization of the expression of APOL1 predominantly in podocytes and endothelial cells led to the identification of the JAK-STAT pathway, characterized by activation of its downstream molecular signature, and its strong association with APOL1 expression in the glomeruli, particularly in patients with FSGS, prompted further cell-type specific mechanistic evaluations. iPSC-derived podocytes from people with APOL1-associated FSGS showed increased JAK-STAT activation in response to IFN-γ stimulation, suggesting that potentiated JAK-STAT signaling may be a modifier of AMKD. Furthermore, CRISPR/Cas9-mediated knockout of APOL1 in iPODs, and the presence of N264K in podocytes from patients with FSGS, known to disrupt APOL1 pore function in vivo[36], both attenuated JAK-STAT activation, suggesting an APOL1 pore-dependent, feed forward mechanism.

Chronic JAK-STAT activation in the context of APOL1 could explain the elevated APOL1 expression observed in participants with HR APOL1 compared to those without such that once activated, it becomes a disease driving feed-forward loop. The limited self-renewing capacity of nephron progenitor cells into podocytes, compared to endothelial or tubular epithelial cells[51,52], may explain the unique vulnerability of podocytes to this pathological feed-forward loop.

What is clear from these findings and from epidemiological studies[41,53] is that APOL1 HR genotype is not sufficient for AMKD. Presumably other genetic, epigenetic, or environmental factors leading to activated JAK-STAT pathways might also be necessary. It remains to be seen if other cytokine or inflammatory pathways (e.g. IL-1β or TNF activation) could also serve as initiators of disease activity leading to a chronic activation state. It also

remains to be seen if elevated APOL1 expression itself may be a contributor to AMKD in the absence of APOL1 HR, but in the presence of JAK-STAT activation.

While single-cell and single-nucleus data from participants are observational and correlative in nature, the introduction of iPODs, and prior studies in kidney organoids,[10,33] demonstrate a clear causal relationship between IFN-γ stimulation to JAK-STAT activation to induction of APOL1 mRNA expression. Although iPODs may not fully recapitulate mature podocyte behavior *in vivo*, the recapitulation of human disease biology strongly suggests that, at least in relation to the JAK-STAT-APOL1 regulatory loop, the model functions as intended. Further, the marked differences between iPODs from APOL1 HR cases and controls suggests potential applications in patient-centric AMKD risk stratification or drug-response assessment in precision nephrology, not unlike patient-derived xenografts in oncology[54].

To further strengthen this feed-forward model, it will be important to test whether APOL1 risk variants alone are sufficient to drive JAK/STAT activation in the absence of cytokine stimulation. Experimental strategies such as CRISPRa-mediated induction or overexpression of G1/G2 versus G0 alleles could directly address this question. Importantly, published *in vivo* work[19,55] provides supportive evidence: doxycycline-induced expression of APOL1 risk variants in mouse podocytes, independent of exogenous interferon, was sufficient to enhance JAK/STAT signaling and promote glomerular injury. These findings align with our proposed model and highlight APOL1 risk variants as active amplifiers of JAK/STAT signaling rather than passive downstream effectors. The data presented herein supports the mechanistic use of JAK-STAT[56] and APOL1 inhibitors[57] in APOL1 high risk patients. The ongoing JUSTICE trial (NCT05237388) of baricitinib in AMKD is likely to reveal whether inhibition of JAK-STAT signaling is an effective therapeutic strategy.

In summary, we have created a comprehensive kidney tissue transcriptional atlas focused on glomerular diseases that is freely available to the glomerular disease research community to explore (Neptune-ocean.cxg.miktmc.org). This atlas provides a foundation for treatment target and disease biomarker discovery in patients with MCD, FSGS, and IgAN. Specifically, we demonstrated the utility of the atlas in identifying disease-associated transcriptional networks in FSGS by providing a human cell–type resolved mechanistic link between APOL1 risk variants and JAK/STAT-driven podocyte injury in FSGS. This mechanistic link is now being targeted in the ongoing JUSTICE trial, part of the precision medicine trial platform of NEPTUNE Match[28], which closes the loop from molecular profiling to targeted therapeutic intervention for glomerular diseases.

Methods

Participant cohorts - NEPTUNE and HKTTA

NEPTUNE (NCT01209000) is a prospective observational study that enrolls children and adults with proteinuria at the time of their first clinically indicated kidney biopsy from 21 participating sites[9]. Human Kidney Transplant Transcriptomic Atlas (HKTTA) is a prospective cohort of kidney transplant recipients, who were willing to provide an additional biopsy core for research purposes at the time of clinical biopsy pre- and post-transplantation. OCEAN included snRNAseq and scRNAseq data from kidney biopsies of 120 NEPTUNE participants and 50 HKTTA participants. An additional 92 NEPTUNE participants with FSGS with previously published bulk-RNAseq profiles[10] were included in the analysis. All clinical and research activities reported here are consistent with the principles of the Declaration of Istanbul.

Duke University and Partners HealthCare study participants for functional validation

In accordance with protocols approved by Partners HealthCare System and Duke University Health System IRB, self-identified Black or African American adults (age 18 years or older) with biopsy-proven focal segmental glomerulosclerosis (FSGS, N = 51) were consented and enrolled from nephrology clinics in Boston, Massachusetts and Durham, North Carolina between 2019 and 2023. Among these participants, 27 were found to have a APOL1 high-risk (HR) genotype. We labeled these participants high-risk (HR) cases. We consented healthy, self-identified Black or African American adults (and 50 years or older) with no known kidney disease (N=218). We enrolled the 30 healthy volunteers (N=30) found to have high-risk APOL1 genotype as high-risk (HR) controls. We excluded volunteers with an eGFR less than 60ml/min/1.73 m^2, a UPCR > 0.2, a low-risk APOL1 genotype, or found to have diabetes mellitus. The minimal enrollment age of 50 years for controls was informed by prior reports that

indicate that APOL1-associated, clinically significant kidney disease typically manifests by age 50[45,58,59]. All, except one HR control participant, met this age cutoff. Individuals in the HR control group were also excluded if they had proteinuria (urine protein-to-creatinine ratio [UPCR] > 0.2 g/g) or an estimated glomerular filtration rate (eGFR) < 60 mL/min/1.73 m², or diabetes mellitus.

Whole genome sequencing of NEPTUNE participants

Blood samples provided by consenting NEPTUNE participants were subjected to whole genome sequencing; reads were obtained using Illumina HiSeq. Sequencing was performed to an average read depth of at least 30X. FASTQ files were then aligned to the GRCh38 draft of the human genome using bwa-mem (version 0.7.17). Joint variant calling and filtering was performed using GATK (version 4.1.8.0) following recommended best practices (https://gatk.broadinstitute.org/hc/en-us/articles/360035535932-Germline-short-variant-discovery-SNPs-Indels). The genotype calls for the following APOL1 variant were then extracted for participants having snRNAseq data: G1: rs73885319, rs60910145, G2: rs71785313, and N264K: rs73885316. Participants without any calls for the G1 or G2 variants were noted as having 0 risk alleles.

Isolation of single nuclei for snRNAseq and single cells for scRNAseq

Nuclei from RNAlater-preserved biopsies of 120 NEPTUNE and 11 HKTTA participants, were processed as described previously[10]. Cells from Cryostor-preserved biopsies of 50 HKTTA participants, were isolated according to a published protocol[27]). All HKTTA biopsy samples, including the 11 HKTTA samples that were processed for snRNAseq (dx.doi.org/10.17504/protocols.io.86khzcw), were processed for scRNAseq

(). Some participants in HKTTA contributed kidney

biopsy samples at different time points within a year.

10x sample processing

10x snRNAseq and scRNAseq were performed following standard protocols, using the 10x

Chromium Single-Cell 3′ Reagent Kit v3 kit. Sample demultiplexing, barcode processing, and

gene expression quantifications were conducted using the 10x Cell Ranger pipeline (v.3-7.1)

using the GRCh38 (hg38) reference genome. For 10x snRNAseq, introns were included in Cell

Ranger analyses.

Quality control of 10XsnRNAseq and scRNAseq data

For 10x snRNAseq, nuclei and cognate barcodes that passed 10x Cell Ranger filters were used

for downstream analysis in a Seurat environment (Seurat v.4.3.0.1)[60-62]. SoupX (v.1.0.0-1.6.0)[63]

was conducted on all samples at default parameters to remove ambient RNA contamination in

samples. Nuclei containing greater than or equal to 10% of all transcripts originating from the

mitochondrial genome were removed. Nuclei that had an nFeatures count <500 or >5000 were

also removed to minimize severely damaged nuclei or multiplets. DoubletFinder (v.2.0.0-

2.0.3)[64], was used at default parameters. While the 10x scRNAseq processing used SoupX and

DoubletFinder as described above, cells that contained greater than or equal to 50% of all

transcripts originating from the mitochondrial genome were removed. 10x scRNAseq nFeatures

criteria were identical to snRNAseq processing during integration.

Parse sample processing

Nuclei that passed the Parse quality control pipeline were read into a Seurat environment. Nuclei

that contained an nFeatures count less than 200 or greater than 5000 were removed. Nuclei

containing only protein-coding mitochondrial transcripts greater than 10% of all transcripts were also removed.

Integration and clustering analysis

All snRNAseq and scRNAseq datasets were integrated into one Seurat object via RPCA-based integration. Integration anchors were selected from datasets that are conditional exemplars of the entire dataset. Here, snRNAseq from FSGS, MCD, and IgAN backgrounds were used in conjunction with sn and scRNAseq in LD backgrounds (**supplementary methods**). The IntegrateData function in Seurat required k.weight=75, since several scRNAseq samples contained low cell numbers. The resulting integrated object was scaled and subjected to principal components analysis to determine the number of clusters. Each cluster was subjected to the FindAllMarkers function in Seurat, with resulting cluster markers compared, manually, to known markers for kidney cell types to infer cluster labels[27,65]. Clinical metadata, including APOL1 risk allele calls, were appended to the Seurat object using the package tidyseurat (v.0.8.0). UMAPs and imaging of differentially expressed genes was conducted either with the Seurat package or with the VIP plugin (v.3.0.0) for cellxgene (v.1.1.1)[66] .

Pseudobulking OCEAN snRNAseq, Bulk-RNAseq data and correlations of APOL1 and JAK-STAT activity

Glomerular cell types, namely podocytes, transitioning parietal epithelium, parietal epithelium, vascular smooth muscle/mesangium, glomerular endothelium mixed with the peritubular endothelium, and afferent/efferent arterioles, were pooled from select FSGS patients and a pseudobulked dataset was constructed using the AggregateExpression function in Seurat. Correlations were conducted as previously published[11]. JAK-STAT transcriptional activity was computed by aggregating Z-transformed gene expression profiles of JAK-STAT target genes

into an average Z-score. Removal of APOL1 from the JAK-STAT gene list was conducted when a direct comparison to APOL1 was examined.

Post-processing of OCEAN data for unsupervised analysis of multicellular programs

OCEAN data were processed for multicellular factor analysis (MFA)[30] in Python using scanpy[67]. Cluster annotations were mapped to their major cell types (**STable S1**). The dataset was subsetted to single-nucleus RNAsequencing data. Cell and gene filtering were performed using scanpy.pp.filter_cells (min_genes=200, default parameters otherwise) and scanpy.pp.filter_genes (min_cells=3, default parameters otherwise).

Creation of multi-view pseudobulk expression object

The filtered AnnData object was converted to a MuData object containing aggregated pseudobulk expression profiles for each cell type using functionality of the decoupler package[68]. To ensure high-quality pseudobulk profiles, a minimum of 50 cells per cell type in one sample was required. Cell clusters cycDTL and tPC-IC were excluded from exploratory analysis as they reached this threshold in fewer than half of the samples. Single-nucleus gene expression was aggregated into sample-level pseudobulk profiles using decoupler.get_pseudobulk (mode='sum', min_cells=50, min_counts=100, min_smpls=3). The pseudobulk expression profiles were further filtered at the gene level using decoupler.filter_by_expr (min_count=10, min_total_count=15, large_n=5, min_prop=0.05). Sample filtering ensured that each sample was represented in at least two cell-type views. After gene and sample filtering, pseudobulk profiles were normalized using scanpy.pp.normalize_total (target_sum=10,000, default parameters otherwise). A variance-stabilizing transformation was applied using scanpy.pp.log1p with default parameters.

Cell-cell communication inference

Putative ligand-receptor interaction scores were calculated using LIANA+[31]. For consistency with the multi-view pseudobulk expression data, cycDTL and tPC-IC clusters were also excluded from cell-cell communication inference. Putative ligand-receptor interactions were identified based on co-expression using liana.mt.rank_aggregate.by_sample (expr_prop=0.1, use_raw=False, n_perms=100, return_all_lrs=False, default parameters otherwise). The resulting ligand-receptor scores were converted into a multi-view object using liana.multi.lrs_to_views (score_key='magnitude_rank', lr_prop=0.1, lrs_per_sample=20, lrs_per_view=20, samples_per_view=10, min_variance=0, lr_fill=0, default parameters otherwise).

Feature selection for MulticellularFactor Analysis (MFA)

Samples in OCEAN were collected over four sub-projects, which were found to be a source of variance within the snRNAseq data. To mitigate this, we applied batch-aware highly variable gene selection as a soft batch-correction method, focusing on variation within batches rather than inter-batch variance. Highly variable genes were identified for each cell type from the pseudobulk expression profiles using sc.pp.highly_variable_genes, with the sub-project ('Project1') as the batch_key, and default parameters otherwise. In ligand-receptor views, features were retained when at least one component (ligand or receptor) was identified as a batch-aware highly variable gene in the corresponding sender or receiver cell type.

Variance decomposition using Multicellular Factor Analysis (MFA)

Following feature selection, the cell type pseudobulk expression and ligand-receptor multi-view objects were concatenated into a single MuData object for input to Multicellular Factor Analysis, which was performed using fucntions of the package for MultiOmics Factor Analysis (MOFA)[69].

MOFA was run using muon.tl.mofa (use_obs='union', convergence_mode='medium', seed=1337), with 20 Factors. As feature selection had already been applied, use_var was set to None. Where not otherwise specified, default parameters were used. Factor scores and variable loadings (i.e., genes and ligand-receptor pairs) were extracted for downstream analysis using liana.utils.get_factor_scores and liana.utils.get_variable_loadings, respectively. Total R^2 values across all samples were computed using calculate_variance_explained from the mofax package (per_factor=True, default parameters otherwise). Per-sample R^2 values were calculated using the same function with sample as group_label (' OCEAN_MS_Exp_ID ').

CLR transformation

To prepare cell type compositional data for statistical testing, a centered log-ratio (CLR) transformation was applied to cell type proportions per sample using clr and multiplicative_replacement from the skbio.stats.composition module[70].

Enrichment analysis

Pathway activity inference on latent factor gene expression loadings were conducted using decoupler.run_ulm. Pathway activities were inferred using the PROGENy [32] prior knowledge on pathways' target genes, retrieved via decoupler.get_progeny (organism='human', top=500, default parameters otherwise). Overrepresentation analysis (ORA) on Hallmark gene sets[71] was performed using decoupler.get_ora_df, with gene sets obtained from decoupler.get_resource('MSigDB') and filtered for the Hallmark collection.

Data visualization for unsupervised patient variability analysis

All unsupervised patient variability visualizations for Figure 3 were generated in R using ggplot2[72], ggpubr[73], patchwork[74] and ComplexHeatmap[75].

Low pass whole genome sequencing (LP-WGS) of Duke University and Partners HealthCare study participants

Low-pass whole-genome sequencing (LP-WGS) was performed in two batches by Genewiz, Azenta Life Sciences, on DNA samples from 57 African American individuals carrying APOL1 high-risk (HR) genotypes. Sequencing reads were aligned to the GRCh38 human reference genome, and variant imputation was performed by Genewiz. Variant call files (VCFs) were merged, and variants failing quality thresholds based on Genome Analysis Toolkit (GATK) Best Practices were excluded. Subsequent data processing and quality control (QC) steps were conducted using PLINK[76]. Variants with missingness >15% and those with minor allele frequency (MAF) <0.05 were excluded. To further assess sample quality and relatedness, we generated a linkage disequilibrium-pruned dataset comprising biallelic variants with MAF ≥0.30. Identity-by-descent (IBD) analysis was conducted to detect duplicate or highly related samples. Given the admixed ancestry of the cohort (predominantly African with some European ancestry), we implemented an ancestry-aware genome-wide association study (GWAS) using Tractor[77]. Local ancestry inference (LAI) was performed with RFMix[78] and ancestry-specific karyograms were constructed to visualize African and European ancestry tracts across the genome. Tractor was then used to perform GWAS stratified by local ancestry, producing separate association results for African and European segments. For comparison, a conventional GWAS was also performed using PLINK. In both models, sequencing batch was included as a covariate. Manhattan plots were generated in R for visualization of GWAS results. To explore the functional relevance of top-associated variants, significant SNPs were queried against the NephQTL database[79], which contains expression quantitative trait loci (eQTLs) derived from glomerular and tubulointerstitial tissues in the NEPTUNE cohort.

Reprogramming of PBMCs to Induced pluripotent stem cells (iPSCs)

Peripheral blood was collected from 15 APOL1 high-risk (HR) cases and 12 HR controls. Peripheral blood mononuclear cells (PBMCs) were isolated using SepMate-50 tubes (Cat# 85460) and Lymphoprep (Cat# 07801) following manufacturer's protocol. PBMCs were reprogrammed into iPSCs using the CytoTune™ 2.0 Sendai Reprogramming Kit (Thermo Fisher Scientific, Cat# A16517), following the manufacturer's protocol and established guidelines. On day 0, $5 \times 10\square$ PBMCs were seeded in 1 mL of PBMC Expansion Medium and transduced with Sendai virus vectors encoding the reprogramming factors OCT3/4, SOX2, KLF4, and c-MYC (designated as KOS, hc-MYC, and hKLF4) at multiplicities of infection (MOI) of either 5:5:3 or 10:10:10. Following viral transduction, cells were spinoculated by centrifugation at 2,250 rpm for 90 minutes at room temperature. After 24 hours, the culture medium was replaced with fresh PBMC Expansion Medium. On day 1, $2 \times 10\square$ irradiated mouse embryonic fibroblasts (MEFs) were plated onto Geltrex-coated plates as a feeder layer. MEF medium was composed of DMEM/F12 + GlutaMAX, supplemented with 1% embryonic stem (ES) cell-qualified fetal bovine serum, 1% MEM non-essential amino acids (NEAA), 1% penicillin-streptomycin (all from Thermo Fisher Scientific), and 1 μL/mL of 55 mM β-mercaptoethanol (Sigma-Aldrich). On day 3, transduced PBMCs were transferred onto the MEF feeder layer, where they attached and subsequently formed iPSC colonies.

Characterization of iPSCs

Selected iPSC lines were characterized for pluripotency markers, trilineage differentiation potential, and genomic stability.

Immunocytochemistry for pluripotency markers

Pluripotency-associated markers—SSEA4, OCT4, SOX2, and TRA-1-60—were assessed by immunocytochemistry using the PSC 4-Marker Immunocytochemistry Kit (Thermo Fisher Scientific), following the manufacturer's protocol. Briefly, iPSCs were washed with Dulbecco's phosphate-buffered saline (DPBS), fixed in fixative solution for 15 minutes at room temperature (RT), and permeabilized for 15 minutes. Following a 30-minute incubation in blocking solution, cells were incubated with primary antibodies for 3 hours at RT. After three washes, cells were incubated with the appropriate secondary antibody for 1 hour at RT, followed by nuclear counterstaining with NucBlue Fixed Cell Stain (Thermo Fisher Scientific).

Trilineage differentiation via embryoid body (EB) formation and Scorecard assay

To evaluate trilineage differentiation capacity, embryoid bodies were generated by dissociating iPSC colonies into clumps using Versene (Thermo Fisher Scientific) and transferring them to ultra-low attachment flat-bottom plates (Costar, catalog number: [insert part number]). Cells were cultured in medium containing DMEM/F12 + GlutaMAX, 20% KnockOut Serum Replacement (KSR), 10% fetal bovine serum (FBS), 1% MEM non-essential amino acids, 1% penicillin-streptomycin, and 1 μL/mL 55 mM β-mercaptoethanol (Sigma-Aldrich). Revitacell™ Supplement (Thermo Fisher Scientific) was added for the first 24 hours. Media were changed every other day for 7–10 days, until EB clumps were visibly formed. Aggregates were filtered through a 30 μm filter (Miltenyi Biotec) to remove single cells. Total RNA was extracted using the RNeasy Mini Kit (Qiagen) and converted to cDNA using the SuperScript IV First-Strand Synthesis System (Thermo Fisher Scientific). Expression of lineage-specific genes was assessed using the TaqMan hPSC Scorecard Panel (384-well format; Thermo Fisher Scientific) following the manufacturer's instructions.

Karyotype analysis

Genomic stability of iPSC lines was assessed using the KaryoStat™ assay (Thermo Fisher Scientific), a microarray-based platform for detecting chromosomal abnormalities at high resolution.

CRISPR–Cas9–mediated knockout of APOL1 in human iPSCs

CRISPR–Cas9–mediated knockout of APOL1 in human iPSCs was performed using the Synthego Gene KO Kit v2 (Synthego), following the manufacturer's protocol. Three chemically modified synthetic sgRNAs targeting APOL1 were used: sgRNA #1: 5′-

TAAAGCTCTGGACAACCTTG-3′; sgRNA #2: 5′-AAGGCCAGCAGTACAGAAAC-3′;

sgRNA #3: 5′-AGCTTGAGGATAACATAAGA-3′. The sgRNAs were pooled in equimolar

ratios and complexed with recombinant Cas9 protein (2:1 molar ratio of sgRNA:Cas9) to form

ribonucleoprotein (RNP) complexes. The RNPs were incubated at room temperature for 10–15

minutes to allow complex formation. iPSCs were harvested at ~70–80% confluency and

resuspended in Neon Resuspension Buffer R. For electroporation, $2 \times 10\square$ cells were mixed with

the pre-formed RNPs and transfected using the Neon Transfection System (Invitrogen) with the

following parameters optimized for human iPSCs: 1,100 V, 30 ms pulse width, and 1 pulse.

Following electroporation, cells were immediately transferred into vitronectin-coated plates

containing pre-warmed Stem-Flex medium (Life Technologies; A3349401) supplemented with

10 µM Y-27632 (ROCK inhibitor) and incubated at 37°C with 5% $CO\square$. 72 hours post-

electroporation cells were re-plated in multiple 96-well plates in such a way that each well of 96-

wells has one single cell and allows them to grow for several days. Multiple clones were selected

and confirmed the APOL1 knock-out using Sanger sequencing and western blotting. CRISPR

editing efficiency and indel spectra were analyzed using the ICE (Inference of CRISPR Edits)

analysis tool (Synthego). Finally, possible CRISPR OFF-target effects were eliminated using

whole-genome sequencing of the selected clones and differentiated in kidney podocytes (iPODs)

using the methods described below.

Differentiation of iPSCs into iPODs

Differentiation of human iPSCs into mature kidney podocytes was performed using a directed,

multi-stage protocol adapted from Musah et al.[42,80]. iPSCs were first induced toward the

intermediate mesoderm lineage by treatment with CHIR99021 (a GSK-3β inhibitor) in basal

media, promoting WNT signaling activation. After approximately 4 days, cells were transitioned

to nephron progenitor specification media containing Activin A and BMP7 to support renal lineage commitment. Following nephron progenitor induction, cells were exposed to a podocyte maturation medium containing a combination of VEGF, retinoic acid, and a reduced concentration of CHIR99021 to promote differentiation into mature podocytes. Over the course of 14–21 days, differentiated cells acquired typical podocyte morphology, and expressed key podocyte markers such as synaptopodin (SYNPO), nephrin (NPHS1), and WT1, confirmed via immunofluorescence staining and qRT-PCR.

RNA isolation and quantitative real-time PCR (qPCR)

Total RNA was extracted from cultured cells using the RNeasy Mini Kit (Qiagen, Cat# 74104) according to the manufacturer's instructions. Briefly, cells were lysed in RLT buffer containing β-mercaptoethanol, followed by homogenization through pipetting. An equal volume of ethanol was added to the lysate, and the mixture was applied to a spin column for RNA binding. After sequential washes, RNA was eluted in RNase-free water and quantified using a NanoDrop spectrophotometer (Thermo Fisher Scientific). RNA quality was confirmed by A260/A280 ratios between 1.8 and 2.1. For cDNA synthesis, 1 µg of total RNA was reverse transcribed using the High-Capacity cDNA Reverse Transcription Kit with RNase Inhibitor (Applied Biosystems, Cat# 4374966), following the manufacturer's thermal cycling protocol: 25°C for 10 min, 37°C for 120 min, and 85°C for 5 min. Synthesized cDNA was stored at −20°C until further analysis. Quantitative PCR (qPCR) was conducted using TaqMan™ Fast Advanced Master Mix (Applied Biosystems, Cat# 4444557) and TaqMan™ Gene Expression Assays (FAM-labeled) specific to the genes of interest and housekeeping genes (e.g., GAPDH or ACTB). Each 20 µL reaction contained 10 µL master mix, 1 µL 20× gene expression assay, 2 µL cDNA (~100 ng RNA equivalent), and 7 µL nuclease-free water. Reactions were performed in optical 96-well plates on

a QuantStudio™ 6 Real-Time PCR System (Applied Biosystems) using the following thermal cycling conditions: 50°C for 2 min, 95°C for 2 min, followed by 40 cycles of 95°C for 1 sec and 60°C for 20 sec. All reactions were run in technical triplicates. Gene expression was analyzed using the ΔCt method, normalizing Ct values to the housekeeping gene. Fold changes in gene expression were calculated using the $2^{-\Delta Ct}$ or $2^{-\Delta\Delta Ct}$ method where appropriate.

Western blotting

Western blot analysis was performed as previously described[81]. Following experimental treatment, cells were washed with PBS and lysed in 1× Cell Lysis Buffer (Cell Signaling Technology, #9803). Lysates were sonicated and centrifuged at 10,000 rpm for 10 minutes at 4°C to collect clarified protein supernatants. Protein concentrations were quantified using a BCA Protein Assay Kit (Pierce, 23225). Equal amounts of protein (20–30 μg) were denatured in 1× Laemmli sample buffer containing β-mercaptoethanol at 100 °C for 5 minutes, separated via SDS-PAGE on precast gels (Bio-Rad), and transferred onto PVDF membranes using the Trans-Blot Turbo Transfer System (Bio-Rad). Membranes were blocked in 5% (w/v) nonfat dry milk (LabScientific, M0842) in 1× TBST (Boston BioProducts, IBB-181) and incubated overnight at 4°C with primary antibodies. The following primary antibodies were used: APOL1 (rabbit anti-human, Genentech, 3.1C1 & 3.7D6, 1:5000, 0.05 μg/mL; mouse anti-human, Genentech, 4.17A5, 4 μg/mL for IF), sodium–potassium ATPase (Abcam, ab76020; IF 1:500, WB 1:100,000), GAPDH (mouse anti-human, Santa Cruz, sc-47724, 1:200), vinculin (mouse anti-human, Sigma-Aldrich, V9131, 1:200), Stat1 (D1K9Y) (Rabbit anti-human, CST #14994T), Phospho Stat1 (Tyr701) (58D6) (Rabbit anti-human, CST #9167S), IRF-1 (D5E4) (Rabbit anti-human, CST# 8478T), and NEPH1 (Santa Cruz Biotechnology, sc-373787). All antibodies were used at manufacturer-recommended dilutions. After washing, membranes were incubated for 1

hour at room temperature with HRP-conjugated secondary antibodies (anti-mouse IgG, CST #7076; anti-rabbit IgG, CST #7074). Signal was developed using SuperSignal West Dura Extended Duration Substrate (Thermo Fisher Scientific, #34075) and imaged with the Bio-Rad ChemiDoc MP Imaging System. Band intensities were quantified by densitometry using AlphaView SA software (ProteinSimple).

Immunofluorescence staining

Cells were washed with phosphate-buffered saline (PBS) and fixed in 4% paraformaldehyde (PFA) in PBS for 15 minutes at room temperature. Following fixation, cells were washed three times with PBS and incubated in blocking buffer (1% bovine serum albumin in PBS) for 1 hour. Primary antibodies were applied overnight at 4°C. After washing, appropriate fluorescently labeled secondary antibodies were applied for 1 hour at room temperature. Nuclei were counterstained and mounted using ProLong™ Glass Antifade Mountant with NucBlue™ Stain (Thermo Fisher Scientific, Cat# P3698). Fluorescent images were acquired using either an ECHO Revolve microscope or a Zeiss LSM 780 upright confocal microscope.

Cell surface protein biotinylation assay

Cell surface protein labeling and isolation were performed using the Pierce™ Cell Surface Biotinylation and Isolation Kit (Thermo Fisher Scientific, #A44390) according to the manufacturer's instructions and described previously[45]. Briefly, cells were seeded on 6-well plates and grown to ~95% confluency. Following IFN-γ (10ng/ml) treatment, cells were washed with PBS and incubated with EZ-Link™ Sulfo-NHS-SS-Biotin for 10 minutes at room temperature to selectively label surface-exposed primary amines. Excess reagent was quenched with ice-cold 1× TBS (provided in the kit). Cells were subsequently lysed in ice-cold lysis buffer (provided in the kit) for 30 minutes. Total lysates were collected, and 5% of the input was

reserved for comparison in immunoblotting. Biotinylated proteins were isolated using NeutrAvidin™ resin, and elution of enriched surface proteins was carried out following the kit protocol. Western blotting was then performed using both total protein input (5%) and eluted biotin-labeled cell surface fractions. APOL1 expression was assessed in both fractions, with Na⊡/K⊡ ATPase used as a plasma membrane loading control and GAPDH used as a control for total input lysates.

Measurement of Rb⊡ flux in iPODs

Rb⊡ flux, a surrogate for potassium transport, was measured in human iPSC-derived podocytes (iPODs) using X-ray fluorescence (XRF)–based quantification. A total of 25,000 iPODs were seeded onto laminin-521–coated 96-well plates and treated with interferon-γ (IFN-γ, 10⊡ng/mL) in the presence or absence of baricitinib (10⊡μM) for 45 hours. Cells were subsequently incubated with 5⊡mM rubidium chloride (RbCl), a potassium analog, for 3 hours to allow intracellular uptake. After incubation, cells were washed three times with modified 1× Earle's Balanced Salt Solution (EBSS) lacking Ca²⊡ and K⊡ to remove extracellular Rb⊡, then air-dried. Intracellular Rb⊡ content was quantified using XRpro® X-ray fluorescence technology, as previously described[45].

Data Availability

All biopsy samples processed directly for inclusion in this study will be uploaded to the Gene Expression Omnibus (GEO) under accession number and publicly available upon publication. The constructed data object is currently hosted online at Neptune-ocean.cxg.miktmc.org, and will be deposited in the CellXGene server upon publication[82]. Biopsy samples previously uploaded to GEO that were included in this composite dataset and manuscript can be found at

GSE213030 (snRNAseq) and GSE254957 (Bulk RNAseq). Whole genome sequencing calls of

APOL1 will be made available.

References

Kidney disease: a global health priority. *Nat Rev Nephrol* **20**, 421-423 (2024). https://doi.org:10.1038/s41581-024-00829-x

Time to sound the alarm about the hidden epidemic of kidney disease. *Nature* **628**, 7-8 (2024). https://doi.org:10.1038/d41586-024-00961-5

Rovin, B. H. *et al.* Executive summary of the KDIGO 2021 Guideline for the Management of Glomerular Diseases. *Kidney Int* **100**, 753-779 (2021). https://doi.org:10.1016/j.kint.2021.05.015

Egbuna, O. *et al.* Inaxaplin for Proteinuric Kidney Disease in Persons with Two APOL1 Variants. *N Engl J Med* **388**, 969-979 (2023). https://doi.org:10.1056/NEJMoa2202396

Rheault, M. N. *et al.* Sparsentan versus Irbesartan in Focal Segmental Glomerulosclerosis. *N Engl J Med* **389**, 2436-2445 (2023). https://doi.org:10.1056/NEJMoa2308550

Heerspink Hiddo, J. L. *et al.* Atrasentan in Patients with IgA Nephropathy. *New England Journal of Medicine* **392**, 544-554 (2025). https://doi.org:10.1056/NEJMoa2409415

Heerspink, H. J. L. *et al.* Sparsentan in patients with IgA nephropathy: a prespecified interim analysis from a randomised, double-blind, active-controlled clinical trial. *The Lancet* **401**, 1584-1594 (2023). https://doi.org:10.1016/S0140-6736(23)00569-X

Cheung, C. K. & Barratt, J. The Rapidly Changing Treatment Landscape of IgA Nephropathy. *Seminars in Nephrology* **44**, 151573 (2024). https://doi.org:https://doi.org/10.1016/j.semnephrol.2025.151573

Gadegbeku, C. A. *et al.* Design of the Nephrotic Syndrome Study Network (NEPTUNE) to evaluate primary glomerular nephropathy by a multidisciplinary approach. *Kidney Int* **83**, 749-756 (2013). https://doi.org:10.1038/ki.2012.428

Mariani, L. H. *et al.* Precision nephrology identified tumor necrosis factor activation variability in minimal change disease and focal segmental glomerulosclerosis. *Kidney Int* **103**, 565-579 (2023). https://doi.org:10.1016/j.kint.2022.10.023

Tao, J. *et al.* JAK-STAT signaling is activated in the kidney and peripheral blood cells of patients with focal segmental glomerulosclerosis. *Kidney Int* **94**, 795-808 (2018). https://doi.org:10.1016/j.kint.2018.05.022

McNulty, M. T. *et al.* A glomerular transcriptomic landscape of apolipoprotein L1 in Black patients with focal segmental glomerulosclerosis. *Kidney Int* **102**, 136-148 (2022). https://doi.org:10.1016/j.kint.2021.10.041

Sampson, M. G. *et al.* Integrative Genomics Identifies Novel Associations with APOL1 Risk Genotypes in Black NEPTUNE Subjects. *J Am Soc Nephrol* **27**, 814-823 (2016). https://doi.org:10.1681/ASN.2014111131

Friedman, D. J. & Pollak, M. R. APOL1 Nephropathy: From Genetics to Clinical Applications. *Clin J Am Soc Nephrol* **16**, 294-303 (2021). https://doi.org:10.2215/CJN.15161219

Genovese, G., Friedman, D. J. & Pollak, M. R. APOL1 variants and kidney disease in people of recent African ancestry. *Nat Rev Nephrol* **9**, 240-244 (2013). https://doi.org:10.1038/nrneph.2013.34

Ng, D. K. *et al.* APOL1-associated glomerular disease among African-American children: a collaboration of the Chronic Kidney Disease in Children (CKiD) and

Nephrotic Syndrome Study Network (NEPTUNE) cohorts. *Nephrol Dial Transplant* **32**, 983-990 (2017). https://doi.org:10.1093/ndt/gfw061

Reidy, K. J., Hjorten, R. & Parekh, R. S. Genetic risk of APOL1 and kidney disease in children and young adults of African ancestry. *Curr Opin Pediatr* **30**, 252-259 (2018). https://doi.org:10.1097/MOP.0000000000000603

Wen, H. *et al.* APOL1 risk variants cause podocytes injury through enhancing endoplasmic reticulum stress. *Biosci Rep* **38** (2018). https://doi.org:10.1042/bsr20171713

Beckerman, P. *et al.* Transgenic expression of human APOL1 risk variants in podocytes induces kidney disease in mice. *Nat Med* **23**, 429-438 (2017). https://doi.org:10.1038/nm.4287

Kopp, J. B. *et al.* APOL1 genetic variants in focal segmental glomerulosclerosis and HIV-associated nephropathy. *J Am Soc Nephrol* **22**, 2129-2137 (2011). https://doi.org:10.1681/ASN.2011040388

Gupta, Y. *et al.* Strong protective effect of the APOL1 p.N264K variant against G2-associated focal segmental glomerulosclerosis and kidney disease. *Nat Commun* **14**, 7836 (2023). https://doi.org:10.1038/s41467-023-43020-9

McCarthy, G. M. *et al.* Recessive, gain-of-function toxicity in an APOL1 BAC transgenic mouse model mirrors human APOL1 kidney disease. *Dis Model Mech* **14** (2021). https://doi.org:10.1242/dmm.048952

Dummer, P. D. *et al.* APOL1 Kidney Disease Risk Variants: An Evolving Landscape. *Semin Nephrol* **35**, 222-236 (2015). https://doi.org:10.1016/j.semnephrol.2015.04.008

Regev, A. *et al.* The Human Cell Atlas. *Elife* **6** (2017). https://doi.org:10.7554/eLife.27041

Rood, J. E. *et al.* The Human Cell Atlas from a cell census to a unified foundation model. *Nature* **637**, 1065-1071 (2025). https://doi.org:10.1038/s41586-024-08338-4

de Boer, I. H. *et al.* Rationale and design of the Kidney Precision Medicine Project. *Kidney Int* **99**, 498-510 (2021). https://doi.org:10.1016/j.kint.2020.08.039

Lake, B. B. *et al.* An atlas of healthy and injured cell states and niches in the human kidney. *Nature* **619**, 585-594 (2023). https://doi.org:10.1038/s41586-023-05769-3

Trachtman, H. *et al.* Rationale and design of the Nephrotic Syndrome Study Network (NEPTUNE) Match in glomerular diseases: designing the right trial for the right patient, today. *Kidney Int* **105**, 218-230 (2024). https://doi.org:10.1016/j.kint.2023.11.018

Olabisi, O. A. *et al.* Design and Rationale of the Phase 2 Baricitinib Study in Apolipoprotein L1-Mediated Kidney Disease (JUSTICE). *Kidney Int Rep* **9**, 2677-2684 (2024). https://doi.org:10.1016/j.ekir.2024.06.033

Ramirez Flores, R. O., Lanzer, J. D., Dimitrov, D., Velten, B. & Saez-Rodriguez, J. Multicellular factor analysis of single-cell data for a tissue-centric understanding of disease. *Elife* **12** (2023). https://doi.org:10.7554/eLife.93161

Dimitrov, D. *et al.* LIANA+ provides an all-in-one framework for cell-cell communication inference. *Nat Cell Biol* **26**, 1613-1622 (2024). https://doi.org:10.1038/s41556-024-01469-w

Schubert, M. *et al.* Perturbation-response genes reveal signaling footprints in cancer gene expression. *Nat Commun* **9**, 20 (2018). https://doi.org:10.1038/s41467-017-02391-6

Juliar, B. A. *et al.* Interferon-gamma induces combined pyroptotic angiopathy and APOL1 expression in human kidney disease. *Cell Rep* **43**, 114310 (2024). https://doi.org:10.1016/j.celrep.2024.114310

Nystrom, S. E. *et al.* JAK inhibitor blocks COVID-19 cytokine-induced JAK/STAT/APOL1 signaling in glomerular cells and podocytopathy in human kidney organoids. *JCI Insight* **7** (2022). https://doi.org:10.1172/jci.insight.157432

Tao, J. *et al.* JAK-STAT Activity in Peripheral Blood Cells and Kidney Tissue in IgA Nephropathy. *Clin J Am Soc Nephrol* **15**, 973-982 (2020). https://doi.org:10.2215/CJN.11010919

Hung, A. M. *et al.* Genetic Inhibition of APOL1 Pore-Forming Function Prevents APOL1-Mediated Kidney Disease. *J Am Soc Nephrol* **34**, 1889-1899 (2023). https://doi.org:10.1681/ASN.0000000000000219

Nichols, B. *et al.* Innate immunity pathways regulate the nephropathy gene Apolipoprotein L1. *Kidney Int* **87**, 332-342 (2015). https://doi.org:10.1038/ki.2014.270

Appel, L. J. *et al.* Intensive blood-pressure control in hypertensive chronic kidney disease. *N Engl J Med* **363**, 918-929 (2010). https://doi.org:10.1056/NEJMoa0910975

Chen, T. K. *et al.* APOL1 Risk Variants, Incident Proteinuria, and Subsequent eGFR Decline in Blacks with Hypertension-Attributed CKD. *Clin J Am Soc Nephrol* **12**, 1771-1777 (2017). https://doi.org:10.2215/CJN.01180117

Kopp, J. B. *et al.* Clinical Features and Histology of Apolipoprotein L1-Associated Nephropathy in the FSGS Clinical Trial. *J Am Soc Nephrol* **26**, 1443-1448 (2015). https://doi.org:10.1681/ASN.2013111242

Friedman, D. J., Kozlitina, J., Genovese, G., Jog, P. & Pollak, M. R. Population-based risk assessment of APOL1 on renal disease. *J Am Soc Nephrol* **22**, 2098-2105 (2011). https://doi.org:10.1681/ASN.2011050519

Musah, S., Dimitrakakis, N., Camacho, D. M., Church, G. M. & Ingber, D. E. Directed differentiation of human induced pluripotent stem cells into mature kidney podocytes and establishment of a Glomerulus Chip. *Nat Protoc* **13**, 1662-1685 (2018). https://doi.org:10.1038/s41596-018-0007-8

Nadkarni, G. N. *et al.* Plasma biomarkers are associated with renal outcomes in individuals with APOL1 risk variants. *Kidney Int* **93**, 1409-1416 (2018). https://doi.org:10.1016/j.kint.2018.01.026

Brosius, F. C., Tuttle, K. R. & Kretzler, M. JAK inhibition in the treatment of diabetic kidney disease. *Diabetologia* **59**, 1624-1627 (2016). https://doi.org:10.1007/s00125-016-4021-5

Datta, S. *et al.* APOL1-mediated monovalent cation transport contributes to APOL1-mediated podocytopathy in kidney disease. *J Clin Invest* **134** (2024). https://doi.org:10.1172/JCI172262

Giovinazzo, J. A. *et al.* Apolipoprotein L-1 renal risk variants form active channels at the plasma membrane driving cytotoxicity. *Elife* **9** (2020). https://doi.org:10.7554/eLife.51185

Olabisi, O. A. *et al.* APOL1 kidney disease risk variants cause cytotoxicity by depleting cellular potassium and inducing stress-activated protein kinases. *Proc Natl Acad Sci U S A* **113**, 830-837 (2016). https://doi.org:10.1073/pnas.1522913113

Relman, A. S. The physiological behavior of rubidium and cesium in relation to that of potassium. *Yale J Biol Med* **29**, 248-262 (1956).

Terstappen, G. C. Functional analysis of native and recombinant ion channels using a high-capacity nonradioactive rubidium efflux assay. *Anal Biochem* **272**, 149-155 (1999). https://doi.org:10.1006/abio.1999.4179

Eddy, S. *et al.* Framework for precision medicine in focal segmental glomerulosclerosis: Translation of sparsentan-responsive genes in a rat model to kidney disease associated proteins in biofluids. *medRxiv* (2025). https://doi.org:10.1101/2025.02.26.25322958

Puelles, V. G. & Moeller, M. J. Postnatal podocyte gain: Is the jury still out? *Semin Cell Dev Biol* **91**, 147-152 (2019). https://doi.org:10.1016/j.semcdb.2018.07.007

Ronconi, E. *et al.* Regeneration of glomerular podocytes by human renal progenitors. *J Am Soc Nephrol* **20**, 322-332 (2009). https://doi.org:10.1681/ASN.2008070709

Grams, M. E. *et al.* Race, APOL1 Risk, and eGFR Decline in the General Population. *J Am Soc Nephrol* **27**, 2842-2850 (2016). https://doi.org:10.1681/ASN.2015070763

Blanchard, Z., Brown, E. A., Ghazaryan, A. & Welm, A. L. PDX models for functional precision oncology and discovery science. *Nat Rev Cancer* **25**, 153-166 (2025). https://doi.org:10.1038/s41568-024-00779-3

Wu, J. *et al.* The key role of NLRP3 and STING in APOL1-associated podocytopathy. *J Clin Invest* **131** (2021). https://doi.org:10.1172/JCI136329

(!!! INVALID CITATION !!! 29).

(!!! INVALID CITATION !!! 4).

Freedman, B. I. *et al.* Association of APOL1 variants with mild kidney disease in the first-degree relatives of African American patients with non-diabetic end-stage renal disease. *Kidney Int* **82**, 805-811 (2012). https://doi.org:10.1038/ki.2012.217

Kanji, Z. *et al.* Genetic variation in APOL1 associates with younger age at hemodialysis initiation. *J Am Soc Nephrol* **22**, 2091-2097 (2011). https://doi.org:10.1681/ASN.2010121234

Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018). https://doi.org:10.1038/nbt.4096

Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015). https://doi.org:10.1038/nbt.3192

Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019). https://doi.org:10.1016/j.cell.2019.05.031

Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet-based single-cell RNA sequencing data. *Gigascience* **9** (2020). https://doi.org:10.1093/gigascience/giaa151

McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337 e324 (2019). https://doi.org:10.1016/j.cels.2019.03.003

Menon, R. *et al.* Single cell transcriptomics identifies focal segmental glomerulosclerosis remission endothelial biomarker. *JCI Insight* **5** (2020). https://doi.org:10.1172/jci.insight.133267

Li, K. *et al.* Cellxgene VIP unleashes full power of interactive visualization and integrative analysis of scRNA-seq, spatial transcriptomics, and multiome data. *bioRxiv*, 2020.2008.2028.270652 (2022). https://doi.org:10.1101/2020.08.28.270652

Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**, 15 (2018). https://doi.org:10.1186/s13059-017-1382-0

Badia, I. M. P. *et al.* decoupleR: ensemble of computational methods to infer biological activities from omics data. *Bioinform Adv* **2**, vbac016 (2022). https://doi.org:10.1093/bioadv/vbac016

Argelaguet, R. *et al.* MOFA+: a statistical framework for comprehensive integration of multi-modal single-cell data. *Genome Biol* **21**, 111 (2020). https://doi.org:10.1186/s13059-020-02015-1

Rideout, J. R. *et al. scikit-bio/scikit-bio: scikit-bio 0.6.3 (0.6.3)*, < https://doi.org/10.5281/zenodo.14640761> (2025).

Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-425 (2015). https://doi.org:10.1016/j.cels.2015.12.004

Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

ggpubr: 'ggplot2' Based Publication Ready Plots (https://rpkgs.datanovia.com/ggpubr/, 2025).

Pedersen, T. L. {patchwork: The Composer of Plots. (2025).

Gu, Z. Complex heatmap visualization. *Imeta* **1**, e43 (2022). https://doi.org:10.1002/imt2.43

Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**, 559-575 (2007). https://doi.org:10.1086/519795

Atkinson, E. G. *et al.* Tractor uses local ancestry to enable the inclusion of admixed individuals in GWAS and to boost power. *Nat Genet* **53**, 195-204 (2021). https://doi.org:10.1038/s41588-020-00766-y

Maples, B. K., Gravel, S., Kenny, E. E. & Bustamante, C. D. RFMix: a discriminative modeling approach for rapid and robust local-ancestry inference. *Am J Hum Genet* **93**, 278-288 (2013). https://doi.org:10.1016/j.ajhg.2013.06.020

Gillies, C. E. *et al.* An eQTL Landscape of Kidney Tissue in Human Nephrotic Syndrome. *Am J Hum Genet* **103**, 232-244 (2018). https://doi.org:10.1016/j.ajhg.2018.07.004

Burt, M., Bhattachaya, R., Okafor, A. E. & Musah, S. Guided Differentiation of Mature Kidney Podocytes from Human Induced Pluripotent Stem Cells Under Chemically Defined Conditions. *J Vis Exp* (2020). https://doi.org:10.3791/61299

Datta, S. *et al.* Kidney Disease-Associated APOL1 Variants Have Dose-Dependent, Dominant Toxic Gain-of-Function. *J Am Soc Nephrol* **31**, 2083-2096 (2020). https://doi.org:10.1681/ASN.2020010079

Megill, C. *et al.* cellxgene: a performant, scalable exploration platform for high dimensional sparse matrices. *bioRxiv*, 2021.2004.2005.438318 (2021). https://doi.org:10.1101/2021.04.05.438318

Table 1 - Clinical and demographic characteristics of OCEAN participants

Characteristic	FSGS	MCD	IgAN	Other protein-uric diseases	Transplanted kidneys		Total
					Living donor	Deceased donor	
Number	33	24	29	34	18	31	169
Mean age in years ± SD (range)	33.6±21.7 (3-76)	24.4±21.7 (2-68)	35.1±14.2 (6-63)	33.2±23 (2-73)	47±10 (30-66)	40.4±11.9 (17-56)	34.6±19.5 (2-76)
% Female	42.4	37.5	34.4	38.2	61.1	29.0	34.9
% Pediatric[a]	39.4	58.3	10.3	41.2	0.0	3.2	26.6
Race[b]							
Asian/Asian American	0	0	8	1	N/A	N/A	9
Black/African American	26	11	1	16	1	2	57
Multi-racial	0	5	1	1	N/A	N/A	8
White/ Caucasian	5	5	17	13	16	15	71
Not reported or Other	2	3	2	3	1	14	25
Clinical							
% APOL1 high risk	45.5	8.3	6.9	32.4	0.0	0.0	17.8
Mean GFR[c] at baseline ± SD(mL/min/ 1.73m^2)	64.3±35.5	102±43.3	71.8±27.1	76.7±34.7	85.2±5.1	100±14.2	78.9±35.9
Median UPCR[IQR] at baseline (mg/mg)	2.5 [4.7]	3.4 [11]	2.7 [2.9]	4.9 [4.8]	0.1 [0.0]	0.1 [0.02]	2.8 [4.8]

[a]Defined as 17years and below. [b]Race options provided were: Asian/Asian American; Black/African American; Multi-Racial; Native American/Alaskan Native/First Nation; Native Hawaiian/Other Pacific Islander; White/Caucasian; Unknown. Missing and unknown were combined under "not reported"; other was a category used for the living donors, transplant, or deceased transplant donor groups.[c]eGFR except for living donor and transplant/deceased donor where values are measured GFR.

Abbreviations: FSGS- focal segmental glomerulosclerosis ;MCD- minimal change disease ; IgAN -IgA nephropathy ; LD – living donor ; NA-not applicable; SD – standard deviation; IQR – interquartile range; APOL1- apolipoproteinL1; GFR – glomerular filtration rate; UPCR – urine protein to creatinine ratio

Table 2: Summarized Characteristics of High-Risk Cases and High-Risk Controls

Characteristics	High-Risk Cases (N=27)	High-Risk Controls (N=30)
Mean age in years (SD, range)	41.1 (13.4, 21-62)	61.2 (9.8, 35-74)
Percent Female	44.4%	70%
Self-identified Black race	100%	100%
APOL1 Genotype (%)		
G1G1	(N=16; 59.3%)	(N=13; 43.3%)
G1G2	(N= 9; 33.3%)	N=14; 46.7%)
G2G2	(N= 2; 7.4%)	(N=3; 10%)
Mean UPCR g/g (SD, range)	3.1 (4.8, 0.2-18)	0.09 (0.04, 0.05-0.2)
Mean eGFR (SD, range)	35.8 (32.0, 3-103)	90.6 (16.6, 64-124)

Abbreviations: SD – standard deviation; APOL1- apolipoproteinL1; eGFR – estimated glomerular filtration rate; UPCR – urine protein to creatinine ratio

Figures with Figure Legends

Figure 1. Overview of study's analytical approach. This figure presents the high-level analysis framework for "A Human Glomerular Disease Atlas defines the APOL1-JAK-STAT feed forward loop in focal segmental glomerulosclerosis".

Figure 2. OCEAN composition and distribution. a. UMAP of 1,047,781 nuclei and cells within OCEAN reveals all nephron and several non-nephron cell types. We inferred the cell type that represented the majority membership of each Seurat cluster, based on cell type markers from the KPMP and PREMIERE. **b**. Cell types were identified based on common transcriptomic signatures associated with cells seen in KPMP and PREMIERE. **c.** Proportional distribution of the nuclei of OCEAN based on participants with FSGS, MCD, IgAN, LD, or other proteinuric diseases. **d.** Overall proportional composition of OCEAN containing both snRNAseq and scRNAseq datasets. **e.** Proportional composition of nuclei within OCEAN that were found within LD and participants with FSGS, MCD, or IgAN. **f.** Proportional composition of PEC, PEC/POD, and POD nuclei within LD and participants with FSGS, MCD, or IgAN.

Abbreviations: KPMP- Kidney Precision Medicine Project; PREMIERE - PREcision Medicine through IntErrogation of Rna in the kidnEy. FIB = fibroblast, PEC = parietal epithelium, PEC/POD = transitioning parietal epithelium, POD = podocyte, vSMC/MC = vascular smooth muscle and mesangium, PT = proximal tubule, DTL = descending thin limb, aPT = adaptive proximal tubule, cycDTL = cycling

descending thin limb, ATL = ascending thin limb, TAL = thick ascending limb, DCT = distal convoluted tubule, CNT = connecting tubule, IC = intercalated cell, tPC-IC = transitioning principal/intercalated cell, PC = principal cell, EC = endothelial cell, Immune = immune cells. FSGS- focal segmental glomerulosclerosis ;MCD- minimal change disease ; IgAN -IgA nephropathy ; LD – living donor.

Figure 3. Unsupervised exploratory analysis of OCEAN reveals molecular detail of kidney function decline

a. Schematic of the unsupervised analysis using Multicellular Factor Analysis (MFA)[30] to define the overall Factor space and characterize it in terms of pathways[32] and cell-cell interactions[31]. **b.** Overview of samples included in the unsupervised analysis. **c.** Total variance explained (R2) across all factors for each of the cell-type views, categorized by source of variance. **d.** Post-hoc association with clinical metadata of interest. *- FDR-adjusted p-value < 0.01. **e.** A two-dimensional map of MFA Factors 3 and 5 puts OCEAN biopsies on a spectrum of eGFR (ml/min/1.73m2) and age (years) at biopsy. **f-g.** Age and eGFR distribution of biopsies by Factor 3-Factor 5 stratification group. **h.** Variance explained per cell-type view for Factor 3 and Factor 5. **i.** PROGENy pathway activity enrichment on gene loadings of Factor 3 and Factor 5. *- FDR-adjusted p-value < 0.01. **j.** Hallmark gene set enrichment on Factor 5 loadings. Score indicates -log10(FDR.p.value) on terms enriched among positive loadings, and log10(FDR.p.value) on terms enriched among negative loadings. **k.** Selected loadings of Factor 3 in transcriptomic views (left) and ligand-receptor views (right). **l.** Post-hoc association between Factor 3 score and center log-ratio transformed cell type proportions of cells annotated MAC_MON and NKC_NKT. **m.** Hallmark gene set enrichment on Factor 3 loadings. Score indicates -log10(FDR.p.value) on terms enriched among positive loadings, and log10(FDR.p.value) on terms enriched among negative loadings.

Figure 4. APOL1 expression is highest in podocytes from participants with FSGS and with 2 APOL1 high risk alleles. **a.** APOL1 expression, per nephron cell type, from the 10XsnRNAseq data of all participants included in OCEAN. **b.** APOL1 expression from the snRNAseq data of podocytes from living donors and participants with FSGS, IgAN, and MCD. **c.** APOL1 expression from the pseudobulked snRNAseq data of podocytes from participants with FSGS, across the LowF3HighF5, LowF3LowF5 and HighF3LowF5 groups. **d.** Pearson correlation of bulk glomerular RNAseq data from NEPTUNE participants whereby a JAK-STAT pathway activity score (PAS)[9,22] was positively correlated with APOL1 mRNA expression. **e.** Glomerular bulk RNAseq data of APOL1 mRNA expression and **f.** JAK-STAT PAS_V2 as stratified from NEPTUNE participants with 0-1 APOL1 high risk alleles vs 2 APOL1 high risk alleles, as defined in Table 2. * = p < 0.05. **g.** From the snRNAseq data of OCEAN, the expression of genes associated with JAK-STAT target genes in participants with FSGS, in the presence of N264K (Yes) or absence of N264K (No).

Figure 5. Multimodal analysis reveals increased JAK-STAT-APOL1 signaling in iPSC-derived podocytes (iPODS) of HR cases compared to HR controls.
a. Overview of study design showing PBMC collection from participants with high-risk APOL1 genotype and biopsy-proven FSGS (HR cases, N=27) and healthy controls with high-risk APOL1 genotype but without kidney disease (HR controls, N=30). iPODs were generated from the first 15 HR cases and 12 HR controls, followed by whole genome RNA sequencing. Whole genome sequencing was performed on all 57 samples. **b.** Heat map of key podocyte-marker genes shows expression across representative iPODS of HR cases and HR controls; expression of APOL1 is induced by interferon gamma (IFNγ) treatment. **c.** UMAP partitions iPODS into two groups—successfully differentiated iPODS, group A, have good expression of podocyte markers, while poorly differentiated group B express low levels of podocyte markers. **d.** IFNγ induces 381 and 281 DEGs (padj < 0.05; fold change ≥ 2) in HR cases and HR controls, respectively, with 272 overlapping DEGs. **e.** Bar chart of top enriched terms in KEGG_2021_Human gene set library. The top 10 enriched terms for the input gene set are displayed based on the -log10(p-value). The top five enriched terms have the most significant overlap with the input query gene set, with p-values of 0.00005, 0.0002, 0.0007, 0.0001, 0.002, and 0.003, respectively. **f.** Bar chart of top enriched terms from the ENCODE_TF_ChIP-seq_2015 gene set library. The top 10 enriched terms for the input gene set are displayed based on the -log10(p-value). The top three enriched terms have the most significant overlap with the input query gene set, with p-values of 8.5^{-7}, 9.6^{-6}, 1.3^{-5}, respectively. **g,** IFNγ induces higher mRNA fold-expression of APOL1, components of JAK-STAT signaling, and well-established STAT1-regulated genes (in dotted box) which reflects STAT1 activity in iPODs of HR cases compared to HR controls. **h.** In the first two HR cases and HR controls, the basal protein levels of JAK-STAT components are similar. **i.** However, IFNγ induces higher protein levels of APOL1, STAT1 and IRF1 in iPODs of HR cases, which are reversed by JAK inhibitor, baricitinib. **j.** Consistent with protein expression, IFNγ induces higher mRNA levels of APOL1, STAT1, IRF1 and CXCL11 in HR cases relative to HR controls (n= 2 biological replicates per group).The effects were reversed by baricitinib. **k.** membrane protein biotinylation assay shows that IFNγ-treated iPODs of HR case1 exhibit increased APOL1 localization to the plasma membrane compared to HR control1. **l.** Consistent with this finding, IFNγ-induced APOL1-mediated rubidium (Rb+) efflux in iPODs of HR cases, which were partially or completely reversed by baricitinib.

Figure 6. CRISPR/Cas9-mediated knockout of APOL1 attenuates JAK-STAT signaling and rubidium efflux in iPOD of a HR case.

a. Overview of CRISPR/Cas9-mediated knockout of APOL1 in iPSC derived from HR Case1.
b. Compared with isogenic control, iPODs derived from APOL1 knockout (KO) iPSC lack detectable APOL1 by immunofluorescence staining following treatment with IFNγ. **c.** IFNγ-induced STAT phosphorylation and IRF1 expression is reduced in APOL1 KO iPOD.
d. Consistently, IFNγ-induced expression of STAT1 and IRF1 mRNA is also attenuated in APOL1 KO iPOD. **e.** Heat map shows reduction in IFNγ-induced, JAK-STAT regulated genes (mRNA). **f.** Unlike isogenic control iPODs in which IFNγ-induced APOL1 causes Rb+ efflux, there are no Rb+ efflux APOL1-KO IPODs (n= 10 biological replicates per group).

Figure 7. Summary schematic. Proposed model of JAK-STAT induced APOL1 expression feed forward loop on progression towards APOL1 mediated kidney disease. "JAK-STAT-APOL1 Feed Forward Model" created in BioRender. by Subramanian, L. and is licensed under CC BY 4.0.

Acknowledgement

We would like to thank Lalita Subramanian for help with writing, editing and graphics for this paper. Library prep and next-generation sequencing was carried out in the Advanced Genomics Core at the University of Michigan.

Funding

The Nephrotic Syndrome Study Network (NEPTUNE) is part of the Rare Diseases Clinical Research Network (RDCRN), which is funded by the National Institutes of Health (NIH) and led by the National Center for Advancing Translational Sciences (NCATS) through its Division of Rare Diseases Research Innovation (DRDRI). NEPTUNE is funded under grant number U54DK083912 as a collaboration between NCATS and the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Additional funding and/or programmatic support is provided by the University of Michigan, NephCure Kidney International, Alport Syndrome Foundation, and the Halpin Foundation. RDCRN consortia are supported by the RDCRN Data Management and Coordinating Center (DMCC), funded by NCATS and the National Institute of Neurological Disorders and Stroke (NINDS) under U2CTR002818. This study was also supported by research grants from Maze therapeutics, the NEPTUNE Public private partnership involving Travere, Boehringer Ingelheim, Sanofi and Dimerix. The human stem cell-derived podocyte model of APOL1 nephropathy project was funded by NIH Director New Innovators Award to OAO, grant number 1DP2DK124891 and by 5R01MD016401. Additional funding was provided by Massachusetts General Hospital, Duke University, and the German Science Foundation (DFG) funding through the Clinical Research Unit 5011 InteraKD (Project ID: 445703531 to JSR). For computational analysis, the authors acknowledge support by the state of

Baden-Württemberg through bwHPC and the German Research Foundation (DFG) through grant INST 35/1597-1 FUGG, as well as the data storage service SDS@hd supported by the Ministry of Science, Research and the Arts Baden-Württemberg (MWK) and the German Research Foundation (DFG) through grant INST 35/1503-1 FUGG.

Author contributions

SE, PJM, CB, SD, OAO, JS-R, MK conceived and planned the study; SE, PJM, EAO, SD, SM, KS, DF, JH, FA, FAA, BG, LHM, ASN performed the experiments or acquired data; SE, PJM, EAO, FAA, VN, KZL, MA, CB, RF, AEA-K, MEG, SD, OAO, ML analyzed the data; SE, PJM, CB, SD, OAO, VN, JS-R interpreted the findings; SE, PJM, CB, SD, OAO, VN, JH, JS-R, MK edited and wrote the manuscript. All authors reviewed and approved the manuscript.

Competing interest declaration

MK reports grants and contracts through the University of Michigan outside of this work from Chan Zuckerberg Initiative, AstraZeneca, NovoNordisk, Eli Lilly, Boehringer-Ingelheim, European Union Innovative Medicine Initiative, Certa Therapeutics, RenalytixAI, Regeneron, Novo Nordisk, Sanofi, Dimerix, Travere and Vera Therapeutics. He has received consulting fees through the University of Michigan from NovoNordisk, Alexion, Novartis, Roche Diagnostics and Vera Therapeutics. MK and VN have a patent PCT/EP2014/073413 "Biomarkers and methods for progression prediction for chronic kidney disease" licensed. MK has served on the NIH-NCATS council, is the committee chair for the American Society of Nephrology Program and is on the board of Nephcure Kidney International. SE receives research funding through the University of Michigan from AstraZeneca PLC, Eli Lilly and Company, NovoNordisk, and Travere Therapeutics, Gilead Sciences Inc, Moderna Inc, and IONIS Pharmaceuticals Inc. OAO reports grants and contracts through Duke University with the NIH. Also received research funding through Duke University from Icagen, and an investigational study drug from Eli Lilly. He received consulting fees from Maze and Podium Bio. JS-R reports in the last 3 years funding from GSK and Pfizer and fees/honoraria from Travere Therapeutics, Stadapharm, Astex, Owkin, Pfizer, Grunenthal, Tempus and Moderna.

List of Supplementary Materials

- List of contributors for the NEPTUNE study

- Supplementary Tables

  - Supplementary Table S1. Cluster annotation for the major cell types in the MOFA view

  - Supplementary Table S2 – Number of podocytes sequenced per patient with FSGS

  - Supplementary Table S3 - Characteristics of NEPTUNE participants with FSGS and bulk glomerular transcriptomic data

  - Supplementary Table S4 – Demographic and clinical characteristics of case-control cohort for iPSC-derived podocyte (iPOD) model of APOL1 mediated FSGS

- Supplementary Figures

  - Supplementary Figure S1 - UMAPs by disease category/group and Seurat specific cluster output

  - Supplementary Figure S2 – Clinical information from NEPTUNE participants stratified by Factor 3-Factor 5 status.

  - Supplementary Figure S3 - JAK-STAT-APOL1 expression in podocytes of participants with FSGS.

  - Supplementary Figure S4 - Lack of genetic modifiers in HR cases and HR controls

  - Supplementary Figure S5 - Interferon regulated downstream effects in iPODS from HR cases and HR controls