Standardized Multicenter Critical Care Database Integrating Minute-Level Vital Signs,

Laboratory Tests, Interventions, and Outcomes: Profile of the OneICU Database

Authors

Takahiro Kinoshita, MD, MPH[1], Yutaka Umemura, MD, PhD[2], Makoto Watanabe, MD, PhD[3],

Kenichiro Uchida, MD, PhD[4], Yuji Nishimoto, MD[5], Shinshu Katayama, MD, PhD[6], Yoshiaki

Inoue, MD, PhD[7], Hiroshi Kurosawa, MD, PhD[8], Yutaka Igarashi, MD, PhD[9], Akira Kodate,

MD[10], Keita Iyama, MD, PhD[11], Shigeaki Inoue, MD, PhD, MSCI[12], Hirotada Kobayashi, MD,

MSSc[13], Yasushi Nakamori, MD, PhD[14]

1. MeDiCU, Inc., Osaka, Japan

2. Department of Emergency and Critical Care, Osaka General Medical Center, Osaka, Japan

3. Department of Emergency Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan

4. Department of Traumatology and Critical Care Medicine, Osaka Metropolitan University

Graduate School of Medicine, Osaka, Japan

5. Department of Cardiology, Toyonaka Municipal Hospital, Toyonaka, Japan

6. Department of Anesthesiology and Critical Care Medicine, Jichi Medical University Saitama

Medical Center, Saitama, Japan

7. Department of Emergency and Critical Care Medicine, Faculty of Medicine, University of

Tsukuba, Tsukuba, Japan

8. Center for Pediatric Critical Care Medicine, Hyogo Prefectural Kobe Children's Hospital,

Kobe, Japan

9. Department of Emergency and Critical Care Medicine, Nippon Medical School, Tokyo, Japan

10. Department of Emergency and Critical Care Medicine, Sapporo City General Hospital,

Sapporo, Japan

11. Acute and Critical Care Center, Nagasaki University Hospital, Nagasaki, Japan

12. Department of Emergency and Critical Care Medicine, Wakayama Medical University

Hospital, Wakayama, Japan

13. Department of Intensive Care Medicine, Kameda Medical Center, Kamogawa, Japan

14. Department of Emergency and Critical Care Medicine, Kansai Medical University Medical

Center, Osaka, Japan

Corresponding author contact information:

Takahiro Kinoshita, MD, MPH

MeDiCU, Inc., 1-15-23, Higashinakamoto, Higashinari-ku, Osaka 537-0021, Japan

Tel: +81-6-6981-0148

E-mail: takahiro.kinoshita@medicu.co.jp

Reprints request:

Reprints of this manuscript are not required.

Conflicts of Interests and Source of Funding:

Takahiro Kinoshita is the founder of MeDiCU, Inc., which develops and manages the database

described in this study. The other authors declare no conflicts of interest. No financial support

was received for this study.

Keywords: artificial intelligence, clinical decision support system, database, data collection, data

science, electronic☐health☐records

Abstract

Introduction: Standrdized intensive care unit (ICU) databases with high☐frequency data

collection from multicenter electronic medical records remain scarce. We aimed to describe the

profile of newly developed OneICU database, which includes minute-level recordings of vital

signs, laboratory values, interventions, and diagnosis codes, and to evaluate the importance of

vital☐sign measurement frequency and laboratory data completeness for developing

machine☐learning–based clinical decision support.

Methods: This retrospective, multicenter observational study collected critically ill patient data

from 12 tertiary care hospitals from 2013 to 2025. Patient demographics, measurement frequency

of vital signs and laboratory tests, as well as the availability of the Sequential Organ Failure

Assessment (SOFA) score components were compared across three large ICU databases:

OneICU, Medical Information Mart for Intensive Care (MIMIC)-IV and eICU. We then

evaluated the prediction accuracy of five machine☐learning models to forecast hypotensive

events 60–120 minutes in advance, defined as a median invasive mean arterial pressure (MAP) <

65 mmHg or vasopressor initiation, using different MAP sampling frequencies: OneICU

1☐minute, OneICU 5☐minute, OneICU hourly, MIMIC☐IV hourly, and eICU 5☐minute.

Results: OneICU currently includes 152,269 ICU stays from 127,757 unique patients. Compared

with MIMIC-IV and eICU, OneICU captured more frequent vital signs (minute-level vs. hourly

in MIMIC-IV and every five minutes in eICU) and provided broader availability of SOFA score

components. In particular, the respiratory component was available for 73.6☐% of stays in

OneICU versus 37.8☐% in MIMIC☐IV and 30.9☐% in eICU, and the liver component for

93.3☐% versus 45.2☐% and 41.3☐%, respectively. The test-set area under the receiver operating

characteristic curve was highest for the OneICU 1-minute model (0.942), followed by OneICU

5-minute model (0.939), OneICU hourly model (0.901), eICU 5-minute model (0.899), and

MIMIC☐IV hourly model (0.799).

Conclusions: A high☐resolution, multicenter ICU database integrating minute☐level vital sign

recordings with comprehensive SOFA score coverage is feasible and was associated with

superior hypotension☐prediction performance. OneICU enables detailed analyses of ICU

trajectories and addresses the current scarcity of large☐scale ICU data from Asian populations.

Background

The use of comprehensive, high-resolution time-series data from electronic medical records (EMRs) has become integral to advancing the management of critically ill patients in the intensive care unit (ICU). Continuous streams of vital signs, laboratory results, and interventions recorded at minute-level intervals provide opportunities for precise assessment of disease severity, early detection of clinical deterioration, and personalized therapeutic strategies. Recent studies suggest that large-scale time-series databases can facilitate data-driven approaches and artificial intelligence (AI) models, leading to improvements in ICU care and patient outcomes (1–5).

Despite these advantages, the development of large scale time-series ICU databases remains challenging due to the heterogeneity of EMR systems, the technical complexity of data standardization, and privacy considerations. Prominent databases such as the Medical Information Mart for Intensive Care (MIMIC-IV) (6), the eICU Collaborative Research Database (eICU-CRD) (7), Amsterdam University Medical Center database (AmsterdamUMCdb) (8), and High time-resolution intensive care unit dataset (HiRID) (3) have fueled numerous studies; however, vital-sign sampling frequencies differ markedly, ranging from every 2 min in HiRID to approximately hourly in MIMIC☐IV, complicating direct comparisons of datasets and the models developed from them. Moreover, most available datasets originate from single institutions in Western countries, limiting their generalizability. Asian ICU databases remain scarce, and initiatives that integrate high resolution data from geographically diverse hospitals are virtually nonexistent. This lack of representative data from Asian populations can lead to gaps in evidence-based practice and preclude the development of fair and context-appropriate AI.

To address these issues, we developed OneICU, a large, multicenter ICU database that

aggregates data from various Japanese tertiary care hospitals. In this first report, we outline the

database design, data processing workflow, and patient characteristics, and present

machine☐learning models for hypotension prediction, evaluating prediction accuracy against

MIMIC☐IV and eICU.

Methods

Database development

OneICU is a large-scale, continuously evolving ICU database established within a separate

multicenter retrospective study whose primary objective is to develop a machine-learning model

to predict sudden deterioration in acute care settings. The present manuscript profiles the

database—its design, contents, and data quality—and report early modeling works; the results of

the final model using the full dataset will be published separately upon the study completion. As

of December 2025, 37 tertiary care hospitals in Japan were participating in the study, with 12

having completed data collection and cleaning. The protocol for the original study was approved

by the Kansai Medical University Ethics Committee (reference number: 2024015). Given the

inclusion of historical admissions, the Institutional Review Board (IRB) approved reliance on

institutional public notification, rather than direct re☐contact, contingent upon the posting of

prominent opt☐out notices at each site. Each participating institution displayed an opt☐out

notice to allow patients or their families to decline participation. All sites had submitted their

opt☐out notices to their institutional IRBs before the first data transfer, and any records flagged

by opt☐out were excluded from the analytic database before anonymization.

Patients were eligible if they were admitted to medical, surgical, emergency, pediatric, neonatal

ICUs, or coronary care, stroke care, or high care units and had at least one vital sign

measurement. Data collection was defined to begin on April 1, 2010, or on the date each ICU-

specific EMR system was implemented, whichever was later. Accordingly, for the currently

participating hospitals, 2013 is the earliest year of data contribution. The IRB also approved the

secondary use of de-identified data.

Data Standardization Workflow

A key challenge for developing the OneICU database was standardizing the substantial volume

of data generated by heterogeneous EMR systems into a single schema of tables. We therefore

adopted an extract–load–transform (ELT) paradigm rather than the traditional extract–transform–

load pipeline (9). In the ELT design, de☐identified yet still unstructured data are first loaded in

bulk into a secure cloud environment, and only then are they transformed to the target schema.

The data are categorized into three types: static, point time-series, and interval time-series

variables. Static variables are those without repeated measurements, including age, sex, and

diagnosis active upon ICU discharge. Point time-series variables capture repeated observations at

distinct time points, such as vital signs and laboratory test results. Interval time-series variables

require both a start and end time to reflect real-world clinical practice, such as infusion and

ventilator settings. This strategy facilitated high-volume data processing and minimized local

computational bottlenecks. Detailed procedures for each step (extraction, loading, and

transformation) are provided in the Supplementary Methods.

Relational Structure

The OneICU database comprises four tiers—patients, hospitals, hospital_admissions, and

icu_stays (Figure 1). Patient level attributes are stored in patients (key: *subject_id*) and

hospital characteristics are stored in hospitals (key: *hospital_id*). Each hospitalization is

represented in hospital_admissions and linked to patients and hospitals via *subject_id* and

hospital_id. Admission level diagnoses are stored in hospital_admission_diagnoses and

reference hospital_admissions via *hospital_admission_id*. Each ICU encounter is represented in

icu_stays (key: *icu_stay_id*), which serves as the parent table for ICU diagnoses (icu_diagnoses),

physiological measurements (e.g., vital_measurements, body_weight_measurements),

therapeutic interventions (e.g., infusions, mechanical_ventilations), and laboratory and clinical

assessments (e.g., laboratory_tests_blood, gcs); these tables join to icu_stays by *icu_stay_id*.

Descriptions of all variables across the tables are provided in the Supplementary Methods. We

also generated 24 derived tables, including those providing the Charlson Comorbidity Index and

hourly Sequential Organ Failure Assessment (SOFA) scores. The SQL code is publicly available

in the project GitHub repository (https://github.com/medicu-inc/one-icu).

Illustrative Example of Time-Series Data

To illustrate the granularity of OneICU data, we plotted vital signs, medication/infusion intervals,

and severity scores (SOFA, Japanese Association for Acute Medicine Disseminated

Intravascular Coagulation [JAAM DIC]), which were calculated for all patients on cloud

environment, for an arbitrary patient admitted with sepsis. Minute-level vital signs (heart rate

[HR], blood pressure [BP], respiratory rate [RR], oxygen saturation [SpO], body temperature

[BT]) were extracted from the vital measurements table, while medications, fluids, and ventilator

settings were accessed from interval time-series tables. Severity scores were plotted at 6-hour

intervals to reflect clinical changes.

Profile analysis

We compared the demographics of the OneICU database with two globally recognized and

extensively studied benchmark ICU databases, MIMIC-IV and eICU. Comparisons included the

number of centers, number of unique ICU stays, number of patients, data availability periods,

patient age, sex, and ICU length of stays. We also assessed the availability and summary

statistics of the SOFA score and the Acute Physiology and Chronic Health Evaluation (APACHE)

II, III, and IVa scores. We quantified use of vasopressors (norepinephrine, epinephrine,

vasopressin, dopamine, phenylephrine), invasive mechanical ventilation, noninvasive positive

pressure ventilation (NPPV), high☐flow oxygen therapy, and continuous renal replacement

therapy (CRRT). Among norepinephrine users, we compared the initial dose standardized to

µg/kg/min. Mortality rates in the emergency room (ER), ICU, and in-hospital were investigated

respectively. In some participating hospitals, patient records at ER of the severest cases were

managed in the ICU-EMR system; therefore, we excluded patients who died in the ER from ICU

and in-hospital mortality calculations.

The comprehensiveness of OneICU was evaluated by comparing the availability and frequency

of key variables against MIMIC-IV and eICU. Specifically, we investigated the granularity of

vital sign measurements by calculating the average measurements per hour for HR, RR, invasive

and non-invasive BP, SpO☐, and BT among patients who stayed more than 24 hours. For

laboratory tests, we calculated the average number of daily measurements of blood gas (pH,

lactate), complete blood count (white blood cell count [WBC], hemoglobin [Hb]), chemistry

(sodium, albumin), and coagulation (prothrombin time international normalized ratio, D-dimer).

Availability of SOFA score components, respiratory, liver, cardiovascular, coagulation, renal,

and central nervous system parameters, was also investigated.

Machine learning

To assess the impact of vital-sign granularity on prediction model performance, we conducted

comparative modeling across OneICU, MIMIC☐IV, and eICU to predict hypotensive events

occurring 1–2 hours after the current observation window. Eligible ICU stays required recorded

age and sex and at least one measurement of HR, RR, invasive MAP, and SpO☐. Each ICU stay

was partitioned into consecutive, non☐overlapping 60☐minute windows. For each window, we

computed median HR, RR, and SpO☐ and extracted the most recent available blood gas results

(pH, partial pressure of oxygen [PaO☐], partial pressure of carbon dioxide [PaCO☐], base

excess, lactate, sodium, potassium).

From OneICU, we extracted invasive MAP at three sampling frequencies: 60 values at 1☐minute

intervals, 12 values at 5☐minute intervals, and 1 value at hourly intervals. In addition, we

appended database☐specific MAP sequences for the other two databases: 1 value at hourly

intervals (MIMIC☐IV) and 12 values at 5☐minute intervals (eICU). The outcome was a

hypotensive event occurring in the window 60–120 minutes after the index window, defined as

either (i) median invasive MAP < 65 mmHg or (ii) initiation of a vasopressor. We excluded

windows with missing invasive MAP, windows for which the outcome occurring 60–120

minutes after the index window could not be defined or ascertained, and all windows at or after

the first vasopressor initiation to avoid treatment☐related label leakage; this was particularly

required because eICU lacks infusion end times.

Within each database, data were randomly split into a training set (80%) and a test set (20%) at

the ICU-stay level. In the training partition, we used H2O AutoML (H2O.ai, USA) to fit multiple

algorithms with automated hyperparameter tuning (generalized linear models, gradient boosting

machines, distributed random forests, extremely randomized trees, deep neural networks, and

stacked ensembles), eliminating investigator-driven hyperparameter selection. To maintain

computational efficiency, cross-validation was disabled and the training partition was further

divided into mutually exclusive subsets for base training (80%), blending (10%), and the

leaderboard (10%). Stacked ensembles were trained using the blending subset, and all candidate

models were ranked by the area under the receiver operating characteristic curve (AUROC)

computed on the reserved leaderboard subset. The final model was defined as the top‐ranked

algorithm on this leaderboard, for the combination of database and sample frequency. For

reporting, we retained a logistic‐regression baseline and the best‐performing machine-learning

model. Performance on the held‐out test set was summarized by ROC curves with AUROC. We

also evaluated specificity, positive predictive value (PPV), and negative predictive value (NPV)

at the threshold corresponding to 90% sensitivity, and sensitivity, PPV, and NPV at the threshold

corresponding to 90% specificity. All analytic steps described above were rigorously

standardized across databases; only the sampling frequency of invasive MAP features differed.

Continuous variables are presented as medians (25–75 percentiles), and categorical variables as

numbers (%). All analyses were performed using Google BigQuery (Google LLC, USA), R 4.3.1

(Institute for Statistics and Mathematics, Austria), and Python 3.12.3 (Python Software

Foundation, USA). All code can be accessed through the following GitHub repository:

https://github.com/takapion/OneICU_profile_paper.

Results

Figure 2 displays the progression of vital signs, interventions, and severity scores during the ICU

stay of a patient with septic shock caused by acute obstructive suppurative cholangitis. HR,

systolic and diastolic BP, RR, and SpO_2 fluctuated during the first 90 minutes, reflecting clinical

instability and subsequent response to therapy. Norepinephrine was started to stabilize

hemodynamics, followed by vasopressin for additional support, while mechanical ventilation

was implemented based on the patient's respiratory status. SOFA score and JAAM DIC score

were elevated at ICU admission and both scores declined over time after antibiotic and

anticoagulation medications.

Table 1 compares key characteristics of the OneICU, MIMIC-IV, and eICU. Both OneICU and

eICU collected data from multiple institutions, whereas MIMIC-IV drew from a single center.

Among the three databases, eICU contained the largest number of ICU stays and unique patients,

followed by OneICU, with MIMIC-IV having the smallest sample size. Patient severity, as

measured by the Acute Physiology Score III, could be compared between OneICU and MIMIC-

IV and appeared higher in OneICU. The proportions receiving vasopressors, invasive mechanical

ventilation, and CRRT were higher in OneICU and MIMIC-IV than in eICU; NPPV and high-

flow oxygen therapy were not evaluable in eICU. ICU and in-hospital mortality rates were

highest in MIMIC-IV compared to the other two databases. The median ICU length of stay was

slightly longer in MIMIC-IV than in OneICU and eICU.

Vital signs were measured most frequently in OneICU, followed by eICU and then MIMIC-IV

(Figure 3A). Blood gas measurements (pH, lactate) were also obtained more frequently in

OneICU than in the other databases, whereas complete blood count measurements (WBC, Hb)

were relatively similar (Figure 3B). Albumin was measured most frequently in OneICU, while

coagulation tests were performed less frequently in eICU than the other two databases.

Cardiovascular, coagulation, renal, and central nervous system components of the SOFA score

were available in over 90% of patients in all three databases (Figure 3C). Respiratory (OneICU:

73.6%; MIMIC-IV: 37.8%; eICU: 30.9%) and liver (OneICU: 93.3%; MIMIC-IV: 45.2%; eICU:

41.3%) components were captured more comprehensively in OneICU than in MIMIC-IV or

eICU.

For the machine‐learning analysis, eligible hourly windows were approximately 2.8 million in

OneICU, 4.0 million in MIMIC‐IV, and 1.1 million in eICU, representing 75,337, 75,668, and

32,471 unique patients, respectively. Patients were randomly assigned at the ICU-stay level to

training and test sets: 60,269 (training) and 15,068 (test) in OneICU; 60,534 (training) and

15,134 (test) in MIMIC‐IV; and 25,976 (training) and 6,495 (test) in eICU.

Across all three datasets, AutoML selected stacked ensembles as the best‐performing models

based on the highest internal validation AUROC. Figure 4 displays ROC curves for logistic

regression and the best‐performing machine‐learning model in each dataset's test set. The

1‐minute interval model developed in OneICU achieved the highest AUROC for both

approaches; the best‐performing machine‐learning model in OneICU achieved an AUROC of

0.942, the highest among all models evaluated.

The prevalence of hypotensive events was 12.8% in OneICU, 12.0% in MIMIC‐IV, and 11.3%

in eICU. When sensitivity was fixed at 90%, specificity was highest for the OneICU 1‐minute

interval model, followed by the OneICU 5‐minute interval model, the eICU 5‐minute interval

model, the OneICU hourly interval model, and the MIMIC‐IV hourly interval model (Table 2).

When specificity was fixed at 90%, sensitivity was highest for the OneICU 1‐minute interval

model, followed by the OneICU 5□minute interval model, the OneICU hourly interval model,

the eICU 5□minute interval model, and the MIMIC□IV hourly interval model (Table 3). Among

the three OneICU models, PPV and NPV were higher for models trained with more granular

MAP measurements; comparisons of PPV and NPV across databases were not appropriate

because these metrics depend on outcome prevalence.

Discussion

The current study describes the design and early findings of OneICU, a multicenter database that

leverages minute-level ICU data captured from heterogeneous EMR systems. The high-

resolution data enables the precise reconstruction of ICU clinical practice, offering insights into

how patients respond to treatments and how interventions evolve over time. Figure 3 indicates

that OneICU has the record of the vital signs (e.g., HR, RR, and SpO_2) every minute in most

patients, whereas the typical measurement frequency is every five minutes in eICU and every

hour in MIMIC-IV. The high-frequency acquisition of core parameters allows detailed analyses

of clinical trajectories, which is particularly relevant for AI-based predictive modeling.

A well-known barrier to effectively using health information technologies is the lack of EMR

interoperability across multiple institutions (10,11). Differences in system architectures, data

formats, and terminologies can impede the efficient data exchange among hospitals. EMR-based

research databases face similar challenges due to variations in documentation principles and

medical practices across institutions (12,13). For example, a study of the eICU database

identified 8,565 intubated patients in the treatment table; however, records of endotracheal tube

insertion and removal were available for only 1,004 of them (14). This issue likely contributes to

the more frequent use of single-institution datasets such as MIMIC-III and IV, even though eICU

encompasses a larger number of ICU stays overall (15).

One of the key strengths of the OneICU database is its ability to integrate diverse datasets from

multiple institutions and standardize them through a streamlined ELT pipeline. By loading

unstructured data onto a secure cloud-based SQL server before transformation, OneICU

effectively manages variations in EMR formats, table structures, and documentation practices.

This approach preserves the granular, time-stamped nature of each record while iterative

transformations and standardization procedures ensure consistency across static, point time-

series, and interval time-series variables. The result is a flexible, scalable schema that can

accommodate continuous data growth from additional institutions, regardless of EMR vendor or

hospital network, without compromising data integrity or interoperability.

We also observed that OneICU maintains a low rate of missing data for vital signs and laboratory

tests despite collecting data from multiple institutions. Notably, the respiratory and liver

components of the SOFA score are more commonly available in OneICU; PaO_2 and total

bilirubin are more comprehensively recorded in OneICU (16). Similarly, diagnosing DIC is far

more feasible using OneICU, since fibrinogen degradation products and D-dimer, either of which

is necessary for the International Society on Thrombosis and Haemostasis (ISTH)-overt DIC and

JAAM DIC scoring, are rarely assessed in MIMIC-IV and eICU (17). Observed differences in

availability could arise from both clinical practice and extraction process. For example, the

higher computability of the SOFA liver subscore suggests that total bilirubin is measured more

routinely in Japanese ICUs. By contrast, the absence of D-dimer in eICU is most consistent with

non☐extraction rather than universal non☐measurement.

We also reported that vital□sign sampling frequency and measurement completeness were key

determinants of model accuracy in hypotension prediction. We used a strictly standardized

pipeline across OneICU, MIMIC□IV, and eICU, with identical inclusion and exclusion criteria,

a common outcome definition, and a fixed predictor set; the only planned difference was the

sampling frequency of invasive MAP. Among three models developed using OneICU, higher

MAP sampling frequency was associated with higher AUROC and better threshold□based

metrics, including (i) specificity, PPV, and NPV at the threshold chosen to achieve 90%

sensitivity and (ii) sensitivity, PPV, and NPV at the threshold chosen to achieve 90% specificity.

Across datasets, AUROC was highest for the OneICU 1□minute interval model, intermediate for

the eICU 5□minute interval model, and lowest for the MIMIC□IV hourly interval model,

consistent with each dataset's native MAP resolution. Although direct cross□database

comparisons are challenging, AUROC and threshold□based metrics were higher for the OneICU

5□minute interval model than for the eICU 5□minute interval model, which may reflect

differences in measurement completeness of other predictors, including blood gas results.

Overall, our findings suggest that higher□frequency MAP measurements, together with greater

measurement completeness, substantially improve the performance of machine□learning models,

highlighting the value of high□granularity data for model development and evaluation.

Developing a large-scale Asian ICU database is crucial for promoting fairness and reducing

algorithmic biases in critical care research. Two recently published AI guidelines, Transparent

Reporting of a multivariable model for Individual Prognosis Or Diagnosis (TRIPOD)+AI and

Fairness, Universality, Traceability, Usability, Robustness, and Explainability (FUTURE)-AI,

emphasize the importance of representativeness and diversity in AI model development to

prevent the creation or exacerbation of health inequalities (18,19). Most established ICU

databases originate from Western institutions, limiting their applicability to Asian healthcare

settings characterized by different patient demographics, disease prevalence, and resource

allocation. Ensuring that AI models and clinical decision-support tools are trained and validated

on data representative of Asian populations not only enhances their performance in these settings

but also helps prevent the perpetuation of health disparities through biased models. By

systematically aggregating data from multiple institutions in Japan, OneICU takes a significant

step toward equitable AI development, addressing the growing demand for robust, generalizable

clinical decision-support tools that benefit critically ill patients across diverse geographical and

ethnic contexts.

The present study has several limitations. First, OneICU currently includes only patients from

Japanese tertiary care hospitals, the majority of whom are ethnically Japanese. Consequently, the

generalizability of findings derived from this dataset to other populations may be limited. Second,

the absence of imaging data and continuous waveform signals restricts research on advanced

diagnostic approaches and physiological parameter monitoring. Third, while OneICU's

sequential opt-out consent process and thorough data de-identification protect patient privacy,

these measures also eliminate certain demographic details and potentially relevant clinical

nuances. For example, patients over 100 years old and those with rare diseases are excluded to

maintain confidentiality, certain subpopulations remain underrepresented. Fourth, the models

developed to compare machine☐learning performance across datasets are not necessarily optimal

for any individual dataset, as dataset-specific features and context may have been removed by

the standardized pipeline. Moreover, the generalizability of the developed models remains

unclear because cross☐dataset testing was not feasible; models trained on higher☐granularity

inputs cannot be applied to datasets with lower sampling resolution. Collectively, these

considerations underscore the need to interpret findings derived from OneICU with caution and

to pursue ongoing efforts aimed at expanding and refining this multicenter database.

Conclusions

The OneICU database offers a high-resolution data platform for examining critical care practices

and outcomes, integrating minute-level data from heterogeneous EMR systems across multiple

institutions via a streamlined ELT pipeline. This capability enables detailed reconstruction of

ICU workflows and advances fairness by better representing Asian populations underrepresented

in existing databases.

Declarations

Ethics approval and consent to participate

The protocol for the original study was approved by the Kansai Medical University Ethics

Committee. The current study involves the secondary use of anonymized data.

Consent for publication

The Institutional Review Board waived the requirement for informed consent due to the

anonymized nature of the data.

Availability of data and material

The authors' agreement with the OneICU project does not permit public sharing of the data used

in this manuscript. However, reasonable requests for data access will be considered by the

OneICU administrator upon submission of a request through the corresponding author.

Competing interests

TK is the founder of MeDiCU, Inc., which develops and manages the database described in this study. The other authors declare no competing interests.

Funding

No financial support was received for this study.

Authors' contributions

TK conceived, designed, and coordinated the study. TK wrote the initial draft of the manuscript. YU, MW, KU, YN, SK, YI, HK, YI, AK, OT, SI, HK, and YN contributed to the interpretation of data and assisted in the preparation of the manuscript and critically reviewed the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

None.

Reference

1. Adams R, Henry KE, Sridharan A, Soleimani H, Zhan A, Rawat N, et al. Prospective, multi-

site study of patient outcomes after implementation of the TREWS machine learning-based

early warning system for sepsis. Nat Med. 2022 Jul;28(7):1455–60.

2. Rahman A, Chang Y, Dong J, Conroy B, Natarajan A, Kinoshita T, et al. Early prediction of

hemodynamic interventions in the intensive care unit using machine learning. Crit Care.

2021 Dec;25(1):388.

3. Hyland SL, Faltys M, Hüser M, Lyu X, Gumbsch T, Esteban C, et al. Early prediction of

circulatory failure in the intensive care unit using machine learning. Nat Med. 2020

Mar;26(3):364–73.

4. Komorowski M, Celi LA, Badawi O, Gordon AC, Faisal AA. The Artificial Intelligence

Clinician learns optimal treatment strategies for sepsis in intensive care. Nat Med. 2018

Nov;24(11):1716–20.

5. Fleuren LM, Klausch TLT, Zwager CL, Schoonmade LJ, Guo T, Roggeveen LF, et al.

Machine learning for the prediction of sepsis: a systematic review and meta-analysis of

diagnostic test accuracy. Intensive Care Med. 2020 Mar;46(3):383–400.

6. Johnson AEW, Bulgarelli L, Shen L, Gayles A, Shammout A, Horng S, et al. MIMIC-IV, a

freely accessible electronic health record dataset. Sci Data. 2023 Jan 3;10(1):1.

7. Pollard TJ, Johnson AEW, Raffa JD, Celi LA, Mark RG, Badawi O. The eICU Collaborative

Research Database, a freely available multi-center database for critical care research. Sci

Data. 2018 Sep 11;5(1):180178.

8. Thoral PJ, Peppink JM, Driessen RH, Sijbrands EJG, Kompanje EJO, Kaplan L, et al.

Sharing ICU Patient Data Responsibly Under the Society of Critical Care

Medicine/European Society of Intensive Care Medicine Joint Data Science Collaboration:

The Amsterdam University Medical Centers Database (AmsterdamUMCdb) Example*.

Critical Care Medicine. 2021 Jun;49(6):e563–77.

9. Fleuren LM, Dam TA, Tonutti M, De Bruin DP, Lalisang RCA, Gommers D, et al. The

Dutch Data Warehouse, a multicenter and full-admission electronic health records database

for critically ill COVID-19 patients. Crit Care. 2021 Dec;25(1):304.

10. Reisman M. EHRs: The Challenge of Making Electronic Data Usable and Interoperable. P T.

2017 Sep;42(9):572–5.

11. Shull JG. Digital Health and the State of Interoperable Electronic Health Records. JMIR Med

Inform. 2019 Nov 1;7(4):e12712.

12. Weiskopf NG, Weng C. Methods and dimensions of electronic health record data quality

assessment: enabling reuse for clinical research. J Am Med Inform Assoc. 2013;20(1):144–

51.

13. Lewis AE, Weiskopf N, Abrams ZB, Foraker R, Lai AM, Payne PRO, et al. Electronic

health record data quality assessment and tools: a systematic review. J Am Med Inform

Assoc. 2023 Jun 30;30(10):1730–40.

14. Essay P, Shahin TB, Balkan B, Mosier J, Subbian V. The Connected Intensive Care Unit

Patient: Exploratory Analyses and Cohort Discovery From a Critical Care Telemedicine

Database. JMIR Med Inform. 2019 Jan 24;7(1):e13006.

15. Kallout J, Lamer A, Grosjean J, Kerdelhué G, Bouzillé G, Clavier T, et al. Contribution of

Open Access Databases to Intensive Care Medicine Research: Scoping Review. J Med

Internet Res. 2025 Jan 9;27:e57263.

16. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA

(Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure: On

behalf of the Working Group on Sepsis-Related Problems of the European Society of

Intensive Care Medicine. Intensive Care Med. 1996 Jul;22(7):707–10.

17. Gando S, Iba T, Eguchi Y, Ohtomo Y, Okamoto K, Koseki K, et al. A multicenter,

prospective validation of disseminated intravascular coagulation diagnostic criteria for

critically ill patients: Comparing current criteria*: Critical Care Medicine. 2006

Mar;34(3):625–31.

18. Collins GS, Moons KGM, Dhiman P, Riley RD, Beam AL, Van Calster B, et al.

TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use

regression or machine learning methods. BMJ. 2024 Apr 16;e078378.

19. Lekadir K, Frangi AF, Porras AR, Glocker B, Cintas C, Langlotz CP, et al. FUTURE-AI:

international consensus guideline for trustworthy and deployable artificial intelligence in

healthcare. BMJ. 2025 Feb 5;e081554.

Table 1. Baseline characteristics and clinical features of ICU stays in OneICU, MIMIC☐IV, and
eICU

	OneICU	MIMIC-IV	eICU
Number of centers	12	1	208
Unique ICU stays	152,269	94,458	200,859
Unique patients	127,757	65,366	139,367
Time period	2013–2025	2005–2023	2014–2015
Age	69 (52–78)	66 (55–77)	65 (53–76)
missing	0 (0.0%)	0 (0.0%)	95 (0.0%)
Sex			
female	61,800 (40.6%)	41,583 (44.0%)	92,303 (46.0%)
male	90,447 (59.4%)	52,875 (56.0%)	108,379 (54.0%)
unknown	22 (0.0%)	0 (0.0%)	177 (0.1%)
SOFA first 24hr	4 (2–7)	3 (2–6)	N/A
Acute Physiology Score II	15 (10–21)	N/A	N/A
Acute Physiology Score III	60 (40–96)	39 (29–53)	N/A
APACHE IVa score	N/A	N/A	51 (37–68)
Vasopressor	38,479 (25.3%)	26,886 (28.5%)	34,449 (17.2%)
Initial norepinephrine dose (µg/kg/min)	0.066 (0.040–0.112)	0.090 (0.050–0.180)	0.074 (0.039–0.150)
Invasive mechanical ventilation	50,495 (33.2%)	39,353 (41.7%)	41,161 (20.5%)
NPPV	3,130 (2.1%)	5,153 (5.5%)	N/A
High-flow oxygen therapy	9,027 (5.9%)	3,829 (4.1%)	N/A
CRRT	6,080 (4.0%)	3,597 (3.8%)	1,510 (0.8%)
ER death	2,644 (1.7%)	N/A	N/A
missing	138 (0.1%)		
ICU mortality			
Alive	139,235 (93.1%)	86,698 (91.8%)	189,918 (94.6%)
Expired	9,036 (6.0%)	7,746 (8.2%)	10,907 (5.4%)
Unknown	1354 (0.9%)	14 (0.0%)	34 (0.0%)

In-hospital mortality			
Alive	131,320 (87.8%)	83,108 (88.0%)	181,104 (90.2%)
Expired	13,297 (8.9%)	11,350 (12.0%)	18,004 (9.0%)
Unknown	5,008 (3.3%)	0 (0.0%)	1,751 (0.9%)
ICU length of stay	1.5 (0.7–4.2)	2.0 (1.1–3.9)	1.6 (0.8–3.0)
missing	0 (0.0%)	14 (0.0%)	0 (0.0%)

Summary of the three ICU databases, OneICU, MIMIC-IV, and eICU. Categorial variables are

expressed as numbers (percentage), and continuous variables are presented as medians (25−75

percentiles).

Abbreviations: MIMIC-IV, Medical Information Mart for Intensive Care-IV; ICU, Intensive

Care Unit; SOFA, Sequential Organ Failure Assessment; APACHE, Acute Physiology and

Chronic Health Evaluation; NPPV, Noninvasive Positive Pressure Ventilation; CRRT,

Continuous Renal Replacement Therapy; ER, Emergency Room

Table 2. Threshold-based performance at 90% sensitivity for hypotension prediction models

model	threshold	sensitivity	specificity	PPV	NPV	N: test person-hours
OneICU 1-minute model	0.084	0.900	0.833	0.441	0.983	539659
OneICU 5-minute model	0.081	0.900	0.824	0.430	0.983	539659
OneICU hourly model	0.055	0.900	0.703	0.308	0.980	539659
MIMIC-IV (hourly model)	0.073	0.900	0.381	0.165	0.966	799779
eICU (5-minute model)	0.060	0.900	0.708	0.283	0.982	229628

Performance is reported on the held□out test set at the operating point where sensitivity was set

to 0.900 for predicting hypotensive events occurring 60–120 minutes after the index 60□minute

window. For each model, the corresponding probability threshold and resulting specificity, PPV,

and NPV are shown. OneICU models were evaluated using invasive MAP sampled at 1□minute,

5□minute, and hourly intervals; MIMIC□IV used approximately hourly MAP; and eICU used

5□minute MAP. N indicates the number of test person□hours (60□minute windows).

Abbreviations: MAP, mean arterial pressure; PPV, positive predictive value; NPV, negative

predictive value

Table 3. Threshold-based performance at 90% specificity for hypotension prediction models

model	threshold	sensitivity	specificity	PPV	NPV	N: test person-hours
OneICU 1-minute model	0.166	0.836	0.900	0.551	0.974	539659
OneICU 5-minute model	0.171	0.828	0.900	0.549	0.973	539659
OneICU hourly model	0.209	0.734	0.900	0.519	0.958	539659
MIMIC-IV (hourly model)	0.295	0.523	0.900	0.415	0.933	799779
eICU (5-minute model)	0.204	0.708	0.900	0.476	0.960	229628

Performance is reported on the held☐out test set at the operating point where specificity was set

to 0.900 for predicting hypotensive events occurring 60–120 minutes after the index 60☐minute

window. For each model, the corresponding probability threshold and resulting sensitivity, PPV,

and NPV are shown. OneICU models were evaluated using invasive MAP sampled at 1☐minute,

5☐minute, and hourly intervals; MIMIC☐IV used approximately hourly MAP; and eICU used

5☐minute MAP. N indicates the number of test person☐hours (60☐minute windows).

Abbreviations: MAP, mean arterial pressure; PPV, positive predictive value; NPV, negative

predictive value

Figure 1. Entity–relationship diagram of the OneICU database

Rectangles represent tables; listed fields are example column names and data types. One patient
(*subject_id*) can have multiple hospital admissions (*hospital_admission_id*), and each admission
can include multiple ICU stays (*icu_stay_id*). Each admission is associated with a single hospital
(*hospital_id*). The icu_stays table anchors ICU diagnoses table and time□stamped child tables
for physiological measurements, therapeutic interventions, and laboratory and clinical
assessments, all linked by *icu_stay_id*.

Abbreviations: ICU, Intensive Care Unit

Figure 2. Example Patient's ICU Stay: Vital Signs, Interventions, and Severity Scores

The figure illustrates the clinical trajectory of a patient admitted to the ICU with acute

obstructive suppurative cholangitis and septic shock. A minute-by-minute plot of vital signs

captures dynamic changes in the patient's clinical condition. The time-series data show

fluctuations in vital signs, reflecting clinical instability and response to interventions.

Vasopressor administration (noradrenaline and vasopressin), as well as the initiation and

termination of mechanical ventilation, are indicated. Severity scores (SOFA and JAAM DIC) are

plotted at 6-hour intervals, demonstrating a gradual improvement following treatment initiation.

Abbreviations: ICU, Intensive Care Unit; HR, Heart Rate; SBP, Systolic Blood Pressure; DBP,

Diastolic Blood Pressure; RR, Respiratory Rate; SpO□, Oxygen Saturation; BT, Body

Temperature; SOFA, Sequential Organ Failure Assessment; JAAM DIC, Japanese Association

for Acute Medicine Disseminated Intravascular Coagulation

Figure 3. Comparison of Measurement Frequencies Across OneICU, MIMIC-IV, and eICU

(A) Distribution of measurement frequencies for vital signs, including heart rate, respiratory rate,

invasive and non-invasive blood pressure, oxygen saturation, and body temperature. OneICU

records these parameters with minute-level granularity, whereas MIMIC-IV and eICU have

lower measurement frequencies.

(B) Comparison of laboratory test frequencies, including blood gas (pH, lactate), complete blood

count (WBC, Hb), chemistry (sodium, albumin), and coagulation markers (PT-INR, D-dimer).

OneICU captures these variables more frequently than MIMIC-IV and eICU.

(C) Availability of SOFA score components across databases. The respiratory and liver SOFA

components are measured more frequently in OneICU compared to the other databases.

Abbreviations: ICU, Intensive Care Unit; MIMIC-IV, Medical Information Mart for Intensive

Care-IV; HR, Heart Rate; RR, Respiratory Rate; BP, Blood Pressure; SpO□, Oxygen Saturation;

BT, Body Temperature; WBC, White Blood Cell count; Hb, Hemoglobin; Na, Sodium; Alb,

Albumin; PT-INR, Prothrombin Time International Normalized Ratio; SOFA, Sequential Organ

Failure Assessment; CNS, Central Nervous System

Figure 4. Comparison of hypotension prediction model performance across OneICU MAP

sampling frequencies, MIMIC☐IV, and eICU

(A) Logistic regression models using the common predictor set and database□specific invasive
MAP sequences (OneICU: 1□minute, 5□minute, and hourly; MIMIC□IV: approximately
hourly; eICU: 5□minute)

(B) Best AutoML models (top□ranked by AUROC on an internal leaderboard; stacked
ensembles in all models) using the same predictor sets and MAP sequences as in (A)

The outcome was a hypotensive event in the 60–120□minute prediction window after the index
60□minute window (median invasive MAP < 65 mmHg or vasopressor initiation). Data were
split at the ICU-stay level into 80% training and 20% test sets; analytic steps were identical
across databases and OneICU frequency settings except for MAP sampling frequency.

Abbreviations: ICU, Intensive Care Unit; MIMIC-IV, Medical Information Mart for Intensive
Care-IV; ROC, receiver operating characteristic; AUROC, area under the ROC curve; MAP,
mean arterial pressure