Changing contact patterns in Newfoundland and

Labrador, Canada in response to public health

measures during the COVID-19 pandemic

Renny Doig[1*], Amy Hurford[2,3*], Suzette Spurrell[4], Andrea Morrissey[4], Liangliang

Wang[1], Caroline Colijn[5]

[1]Department of Statistics and Actuarial Science, Simon Fraser University, 8888

University Drive, Burnaby, BC, Canada

[2]Department of Biology, Memorial University of Newfoundland, 45 Artic Ave,

St. John's, NL, Canada

[3]Department of Mathematics and Statistics, Memorial University of

Newfoundland, 45 Artic Ave, St. John's, NL, Canada

[4]Newfoundland and Labrador Health Services, Government of Newfoundland and

Labrador, 363 Conception Bay Highway, Holyrood, NL, Canada

[5]Department of Mathematics, Simon Fraser University, 8888 University Drive,

Burnaby, BC, Canada

[*]Corresponding author(s') e-mail(s): rennyd@sfu.ca; ahurford@mun.ca

Abstract

The provincial government of Newfoundland and Labrador, Canada implemented

a contact tracing program as part of a containment strategy during the COVID-19

pandemic. A high proportion of cases were detected and contact traced, and our

analysis provides insights into secondary case distributions and contact patterns in

Newfoundland and Labrador. We used a heuristic approximation of secondary

cases to account for ambiguities in who infected whom. These approximate values

provide an empirical distribution of secondary cases. These distributions are

compared against the stringency of public health measures. Additionally, we

visualised age- and contact-based patterns and compared these patterns with

respect to stringency. The maximum number of contacts traced per week was

4,645 and the mean number of contacts traced per case was 12.5. Approximate

95% CIs of the effective reproduction number under Alert levels 2-4 were

(1.02,1.21), (0.99,1.39), (0.84,1.06), and (1.20,1.47). We find that this level of

contact tracing was sufficient, in combination with other public health

interventions, to contain pandemic SARS-CoV-2 spread in Newfoundland and

Labrador prior to the establishment of the Omicron variant. Understanding age-

based contact patterns is necessary to describe disease spread and the risk of

severe outcomes. A successful containment strategy requires that contact tracing

capacity is not exceeded, making it necessary to understand the behaviour of high-

contact individuals.

Keywords: contact tracing; effective reproduction number; stringency; contact

heterogeneity

Introduction

The province of Newfoundland and Labrador (NL), Canada, announced its first

COVID-19 case on March 14, 2020. In response to this the provincial government

implemented a containment strategy to address the risk posed by the pandemic [1,

p11-12]. The objective of a containment strategy is to prevent the spread of a

disease by minimizing the risk of transmission. Containment strategies consist of

several policies that test, trace, and isolate infectious individuals, that restrict both

the importation of new cases and infection spread, and that are intended to interact

synergistically. These strategies are distinguished from elimination and

eradication strategies, which seek to reduce transmission to zero, and mitigation

strategies which aim to reduce overall impact while accepting some community

transmission [2-4]. Containment strategies are typically only viable during stages

of a pandemic with low case counts and limited community spread. This was the

case in NL until the Omicron variant of the virus was detected on December 15,

2021 [1, p59], at which point the existing containment strategies were deemed to

no longer be effective. By early January 2022 the provincial government had

switched to a mitigation strategy [1, p61].

Contact tracing involves identifying individuals that were potentially exposed by

an infectious individual and is an important part of a containment strategy.

Contact tracing allows for the detection of asymptomatic and pre-symptomatic

individuals, and the potential isolation of infectious individuals before they can

generate secondary cases. The effectiveness of contact tracing, as a containment

measure, often depends on both the number of contacts being traced and the

proportion of cases that are successfully identified, and as capacity is reached,

containment no longer becomes feasible [5-7]. Other public health measures that

limit the number of people contacted by an infectious individual, or that prevent

infection spread by other means, have a role in determining the efficacy of a

contact tracing program, and in protecting contact tracing capacity [7-8].

In NL, contact tracing began with the first reported case on March 14, 2020 and

consistent contact tracing was performed until January 4, 2022 [9], although data

continued to be collected until June 2, 2022 in the pursuance of a mitigation

strategy. After the public health emergency was declared on March 18, 2020 [1,

p10], the NL government issued changes to public health measures via Special

Measures Orders (SMOs), where these measures may have contributed to contact

tracing efficacy and protecting contact tracing capacity as part of a containment

strategy. The initial response to the COVID-19 pandemic involved moving classes

online for grades kindergarten to twelve, travel measures, closing recreational

facilities, imposing limitations on indoor gatherings, and physical distancing

recommendations [1, p14]. Subsequent SMOs relaxed or further restricted these

measures in response to case counts and, later, vaccination rates.

As of April 30, 2020, the NL government introduced the Alert system, which

bundled together many policies affecting local transmission. Alert level 5 was the

strictest set of policies and consisted of small gathering limits and closures of

many activities and non-essential businesses. Alert level 2 was the most relaxed

set of policies that was implemented; it permitted larger gatherings and businesses

to operate conditional on physical distancing and masking guidelines [1, p17-19].

Alert level 1 was a return to a "new normal" and was never implemented, as

instead the Public Health Emergency was repealed on March 14, 2022 [1, p71].

The Alert level system did not encompass all of the component policies of NLs

containment strategy. Travel measures, suspension of in-person kindergarten to

grade twelve school, and mandates on indoor masking were issued in separate

SMOs and were not directly linked to the Alert levels. Additionally, while

gathering limits were not fixed for each Alert level, the stringency of the gathering

limits tended to correspond to the stringency of the Alert level.

The implementation of stricter public health measures may change age-group

contact patterns, and other aspects of contact assortativity. Characterizing such

changes helps to accurately assess the risk of severe infection, which is greatest

for older individuals. Assortativity patterns, such as individuals with a high

number of contacts also contacting other individuals with a large number of

contacts, affect the risk of an outbreak that grows quickly due to consecutive

super-spreading events. Understanding the risk of quickly-growing large

outbreaks is necessary to understand whether contact tracing capacity is likely to

be exceeded.

We use the data collected by NL through their contact tracing program to

understand the efficacy of the containment strategy implemented by the NL

government during the COVID-19 pandemic. We report contacts traced per week

and contacts per case for different stringencies of public health measures (Alert

levels) to understand the capacity of NL's contact tracing program. We report the

number of travel-related and community cases, and estimate distributions of

secondary infections and cluster sizes to evaluate the efficacy of NL's

containment strategy. We visualize contact patterns for age-groups and

assortativity based on the number of contacts to better understand the outbreaks

that occurred in the NL population.

Methods

114   **Methods**

Data collection

Data collection

The contact tracing program in NL was carried out by a core team of 15 public

health officials. When an eligible individual was tested, a public health official

would call the individual to inform them of their test result. If the test result was

positive, a list of recent contacts was obtained, contacts being defined as any

individuals who the case had close contact with within 72 hours of the case's

episode date (episode date being date of symptom onset if the case was

symptomatic, or date of specimen collection if the case was asymptomatic) [10].

Later in the pandemic, individuals could learn of their test result by checking an

online system. When results were communicated online, public health officials

monitored the test results database for the purpose of initiating contact tracing as

quickly as possible when positive cases were reported. Contacts of a case were

instructed by either a letter and/or a telephone call to test and isolate for 14 days.

If the individual tested positive, then the duration of the isolation period would be

reduced to 10 days from symptom onset, or date of positive test for those

asymptomatic. Contacts of a case were usually contacted by the contact tracers

within 24 hours. During periods of many cases, additional staff and third parties

were contracted to support the core team of contact tracers. During the study

period, all contacts were notified as described, such that contact tracing was

conducted consistently. On December 15, 2021, it was stated by the Chief

Medical Officer of Health that due to the short generation time of the Omicron

variant, contact tracing could no longer be completed with the same efficiency as

had occurred prior, and this is how we defined the end of our study period.

For the duration of the study period, symptomatic individuals in NL could seek

testing if they met testing criteria at that moment in time. Testing criteria shifted

throughout the study period; at times one symptom could screen clients in for a

COVID-19 test while at other times two symptoms were required, one of which

had to be fever or cough. Furthermore, some asymptomatic individuals were also

required to complete tests: rotational workers, with some modifications to the

corresponding self-isolation rules [1, p24-25], close contacts of a confirmed case,

and individuals that were asked to complete testing due to publicly issued

exposure notifications corresponding to a venue, flight, or ferry. There were also

testing requirements for entering some long-term care facilities, personal care

homes, assisted living facilities [1, p17], and health care facilities.

For positive cases, the source is recorded as being related to travel, a close contact

of a traveler, or a local source. Additionally, geographic information is recorded

in the form of the Forward Sortation Area (FSA) of the individual. Due to the

relatively small amount of local spread during the focal period of this study, the

contact tracing networks observed are assumed to be mostly complete.

Data preparation and cleaning

We organized the contact tracing data into distinct contact networks, which are

defined in this paper as a collection of individuals who are connected by at least

one contact event. A contact event occurs between individuals A and B if A lists

B as a close contact or vice versa. The contact event alone does not indicate who

infected whom . Compiling connected contact events into clusters results in two

categories of cases: those that are only travel-related and those that eventually

result in local spread. If a contact network contains only individuals who are

recorded as having a history of travel or as being a close contact of a traveler, then

all individuals in the cluster are labeled "travel-related." Otherwise, a network has

at least one instance of a contact event which constitutes local spread. Since the

policies of interest in this work are the Alert level measures, which target local

spread, travel-related networks are removed for the analysis. Within each contact

network is a case cluster, which contains all of the individuals in the network that

tested positive.

The Alert level that was in place at the time and location of each contact event is

not recorded during the contact tracing process. To compare the structure of the

contact networks between Alert levels, each contact event was linked to an Alert

level based on the FSA in which the contact happened. The Newfoundland and

Labrador, Department of Health and Community Services Public Service

Advisories placed certain geographic regions under particular Alert levels at

different times. Each geographic region was assigned a FSA that most closely

matched it. The Alert level for each contact event was the Alert level assigned to

the FSA at the time the contact was reported to have occurred. It is noted that

since SMOs affecting the Alert level were not issued based on FSA there may be

minor discrepancies between the assigned Alert level and the true Alert level for

some contact events.

During the data cleaning process, any cases that occurred outside of the focal

period were removed. Cases that were travel-related were removed as well as any

entries that had missing or invalid FSA information.

Modeling the number of secondary infections

The effective reproduction number, R_t, is the average number of secondary

infections resulting from one index case in a partially susceptible population at

time t. When complete contact tracing data is available, R_t can be estimated

directly as secondary case counts are available for each case. In cases where

available data is limited to time series of reported cases or symptom onset, direct

estimation of R_t is not possible and some model structure needs to be applied to

leverage some information about secondary case distributions from the data.

Mechanistic models of transmission dynamics are common [11], as are models

that have a more limited structure, assuming only a probability distribution over

the interval of time between cases [12,13]. In both cases, the lack of direct

observation of contact events between individuals necessitates some additional

assumptions/structure. The contact tracing data used in this study is partially

complete in that we observe contact events, but not directly who infected whom.

As such, we adopt a heuristic approach, described below, which makes use of the

partial information available without having to rely on a structured model

requiring additional parameters to be estimated or assumptions to be made.

The number of secondary cases generated by a particular index case, i, is denoted

I_i. To account for heterogeneity in infectiousness, I_i is often modeled as a

negative binomial random variable with mean R_t and precision parameter k [14].

Under this parameterization of the negative binomial distribution the number of

secondary cases has variance $R_t + R_t^2/k$. As the estimated value of the precision

parameter for our data was large relative to R_t^2, we also consider a Poisson

distribution with mean R_t, as it is the limiting distribution of a negative binomial

as the precision parameter gets very large.

Observing the number of secondary cases requires knowledge of the infector and

the infectee in each contact event. As an example of a contact network for which

the infector and infectee labels can be assigned, consider a contact network

containing two individuals, A and B, such that A travels and tests positive, then

lists B as a contact who then tests positive without coming into contact with

anyone else. Clusters including multiple travellers or more interconnected

individuals do not admit a clear infector/infectee label for each individual,

therefore direct observations of I_i are not available. To compensate for this, an

approximate number of secondary cases, $\widetilde{I}_i$, is computed. Each individual in the

dataset will either be an index case or they will have at least one other individual

listing them as a contact. The number of cases listing an individual as a contact is

the number of potential infectors that the individual has. Take three individuals,

A, B, and C, who have n_A, n_B, and n_C potential infectors, respectively. Now

consider a fourth individual, D, who is a potential infector of all of A, B, and C.

Individual D constitutes $1/n_A$-th of A's potential infectors, and similarly for B

and C. So, to approximate the number of secondary cases attributable to D, we

define $\widetilde{I}_D = 1/n_A + 1/n_B + 1/n_C$. These $\widetilde{I}_i$'s are then modeled using a continuous

version of the above negative binomial and Poisson distributions, swapping any

factorials in the probability mass function for the appropriate gamma function.

This weighting procedure is intended to be a heuristic approach, used so that a

parametric estimate of the effective reproduction number can be obtained. A point

estimate and an approximate 95% confidence interval about R_t is obtained via

maximum likelihood estimation.

Visualizing contact patterns

Two types of plots are used to visualize the contact structure within the clusters.

The first is an age-stratified contact matrix. Each individual is placed into an age

group: [0,5), [5,18), [18,30), [30,40), [40,50), [50,60), [60,70), and 70+. The age

groups were chosen to reflect pre-school-age ([0,5)), school-age ([5,18)), and then

10-year bins after the age of 18. The grouping together of school-age individuals

is supported by a study conducted in Tianjin, China [15], as well as contact

matrices estimated in Canada [16] and the United Kingdom [17], which show

very similar contact patterns for these ages. The number of contacts between each

age group are counted and plotted as a heat map that shows the total number of

interactions between age groups within the observed contact networks. Counts

below five are shown as being exactly five to preserve anonymity. The second

type of plot shows the relationship between the number of contacts reported by

each pair of cases connected by a contact event. For each pair of connected

individuals, the number of contacts reported by each was computed and plotted as

a point, with the initial case on the horizontal axis and the contacted case on the

vertical axis. Rather than plotting the individual points, this pattern is shown by

applying 2-dimensional kernel smoothing to these points and plotting the contours

of the smoothed density. The darker contours represent regions of case/contact

pairs where there were a relatively higher number of contact events with that

pairing of reported cases.

Both visualizations are computed for all data in the focal period disaggregated

into the Alert level during which the contact event took place. In addition to these

aggregate plots, the same plots are computed for two individual clusters. The two

clusters that are selected contain the most contact events. All analysis is done

using the R programming language [18]; the ggplot2 package [19] is used to

generate all figures.

259   Results

An overview of the history of the disease between the period of March 14, 2020

and December 15, 2021 is given in Figure 1, which shows weekly aggregate cases

in the province. Reported cases are separated into those that are travel-related and

those constituting local spread; local spread is then further separated into the Alert

level in effect when the contact event occurred. Changes to provincial masking

mandates are shown separately from the Alert levels, as these policies were kept

separate. The period between March 14, 2020 and April 30, 2020 is shaded grey

and corresponds to the period prior to the Alert level system; bars in this period

are coloured black because there is no Alert level to which they correspond. After

an SMO that placed the province at Alert level 5, subsequent relaxations brought

the Alert level province-wide down to level 2 by July 2020 when there were no

instances of community spread. In late 2020 and early 2021, when the entire

province was at Alert level 2 and there was an indoor mask mandate in effect,

there were several weeks with reported cases. However, most of these cases

remained travel-related and did not result in community spread. In February 2021,

when a large amount of community spread was detected on the Avalon peninsula

of Newfoundland, the entire province moved to Alert level 5. The implementation

of measures at the community-level, rather than province-wide, started shortly

after this on February 27, 2021; the province continued to manage community-

level policy changes for the remainder of our focal period. The case peak in early

summer 2021 was largely restricted to travel-related cases, while the peak in late

summer/early fall, which began shortly after the mask mandate had been lifted,

was dominated by community spread and a mask mandate was reissued a few

weeks later. Protecting contact tracing capacity was one stated rationale for re-

issuing the mandatory mask requirement, and at this time more than 1,000

contacts were being traced per week (Figure 1, top panel). At two points within

the focal period, February 2021 and September 17, 2021 [1, p49; 17], the NL

government reported a strain on contact tracing capacity. The core contact-tracing

team in eastern Newfoundland consisted of fifteen public health officials [20].

Figure 1 shows the weekly total of contacts traced during the period (top panel).

Peaks in the number of contacts traced tend to correspond to peaks in reported

cases, but on a larger scale.

Fig. 1 Weekly cases (bottom) and contacts traced (top) in Newfoundland and Labrador until

December 15, 2021 with colours indicating if the case is travel-related or if not, and which Alert

level it occurred under. Vertical dashed lines indicate province-wide mandates on masking

indoors.

Between March 14, 2020 and December 15, 2021, 22,626 contacts were traced.

Of these contacts, 2,111 were cases, 1,522 of which were related to local spread.

Analyses of the Alert levels focused on data occurring after April 30, 2020,

removing 203 cases. A further 7 cases were removed for having incorrect or

missing FSA data. A numerical summary of the contact tracing efforts is supplied

in Table 1. The table shows the total number of contacts traced, the total number

of cases reported, the maximum number of contacts traced in one week, and the

average number of contacts traced per person; these results are shown for the

entire focal period as well as a breakdown by Alert level. Of the 1,522 cases,

1,299 (85.3%) reported symptoms, 71 (4.66%) were hospitalized, and 17 (1.12%)

died.

Table 1 Numerical summaries of contact tracing during focal period

	Overall	Alert 2	Alert 3	Alert 4	Alert 5
Contacts traced	19269	5740	1484	3199	6004
Cases reported	1522	484	137	349	342
Max. contacts traced in a week	4645	1217	406	996	4645
Mean contacts traced per person	12.5	11.9	10.8	9.17	17.6

Observed transmission clusters

There were 166 distinct clusters containing more than one person during the focal period. The largest cluster contained 276 cases. The median cluster size was 3, with 90% of the clusters containing at most 11 cases. The distribution of cluster sizes across the whole focal period is illustrated in Figure 2. Both plots show a distribution of cluster sizes concentrated on small numbers of cases with a long tail.

Fig. 2 Histogram of the cluster sizes across the focal period; inset contains the corresponding empirical cumulative distribution function, truncated to 15.

The largest contact network during the focal period is shown in Figure 3. In addition to the case cluster itself, shown with the larger opaque points, Figure 3 also shows individuals who were traced but did not test positive in the smaller, translucent points. This network is linked to an outbreak in eastern Newfoundland between February and April, 2021. Of the contact events that comprise the network, 38.7% occurred under Alert level 5, 60.8% occurred under Alert level 4,

and 0.5% occurred under Alert level 2. At the centre of the cluster most of the

contacts fall within the 5-17 age group. There are also several contacts that are in

the 30-59 age groups. The smaller points indicate contact events which did not

produce secondary cases either because transmission did not occur or because

contact tracing contained the infection.

Fig. 3 Graph of the largest contact network in the dataset. Large, opaque points are the cases

which comprise the cluster itself and the smaller, translucent points are contacts which were traced

but did not result in further local spread. The opacity of the edges corresponds to the number of

subsequent contacts resulting from that infection.

Effective reproduction number

The smoothed empirical distributions of secondary cases are visualized in Figure

4 for the period from October 2020 until the end of the focal period. The values of

approximated secondary infections, $\widetilde{I}_l$, are smoothed over weekly periods and are

disaggregated by Alert level. The corresponding plot of weekly cases is shown

above. There are three predominant peaks visible during this period; the first

occurring under Alert levels 4 and 5, the second under Alert levels 2 and 4, and

the final mostly under Alert levels 2 and 3. For most weeks the distribution of

secondary cases appears to be centred around 1, skewing more towards 0 as the

outbreaks taper off.

Fig. 4 Weekly cases between October 2020 and December 15, 2021 (top) and smoothed empirical

distribution of secondary cases, pooled over a weekly period, (bottom).

The secondary cases were then aggregated across the entire focal period for each

Alert level; the corresponding histograms are shown in Figure 5. Both the

negative binomial and Poisson distributions were fit to the data in each Alert level

via maximum likelihood and compared using the Akaike Information Criterion

(AIC). The fitted Poisson distribution resulted in a lower AIC for all Alert levels,

so the estimates below correspond to the Poisson distribution. It is noted that the

point estimates for the reproduction number were very similar under the negative

binomial distribution, however the likelihood function was very insensitive to the

value of the precision parameter. Maximum likelihood estimates (MLEs) for the

reproduction numbers and precision parameters, as well as the AIC for each Alert

level are shown in Table 2. The MLEs of R_t for each Alert level are also shown in

Figure 5 along with an approximate 95% confidence interval (CI). The probability

mass functions, computed using the MLEs, are drawn over the histograms.

Fig. 5 Histogram of the estimated number of secondary cases caused by each index case under

each Alert level. Probability mass functions evaluated at the MLEs are shown with a solid line.

Approximate 95% CIs for R_t are shown in the top right corner.

Table 2 Point estimates and 95% CIs for R_t under each Alert level

	Poisson		Negative Binomial		
Alert Level	$\widehat{R_t}$	AIC	$\widehat{R_t}$	$\hat{k}$	AIC
2	1.11	1148	1.11	194.0	1151
3	1.19	298	1.19	58.7	301
4	0.95	680	0.95	107.0	684

| 5 | 1.33 | 825 | 1.33 | 46.8 | 826 |

Contact patterns

The visualizations of contact patterns within the clusters are shown for all clusters

within each Alert level as well as for the two largest clusters observed after April

30, 2020, with respect to the number of contact events. The first cluster is the

largest observed cluster with 605 contact events and 276 cases, occurring in the

former eastern health region (Newfoundland) between February and April, 2021

and the second largest cluster contained 261 contact events and 91 cases and

occurred in the former central health region (Newfoundland) between May and

June, 2021. Of the contact events that comprise the central cluster, 65.9% of the

contacts occurred under Alert level 4 and 34.1% occurred under Alert level 2.

Subsequent references to these clusters will refer to the region in which they

occurred. The four regional health authorities of Newfoundland and Labrador

were amalgamated in April 2023.

Age contact matrices are shown in Figure 6. In Alert levels 2 and 3 the contacts

are fairly dispersed across age groups and show minimal clustering around the

diagonal. Alert level 4 shows a relatively large proportion of contacts occurring

within and between the 5-59 age groups. Alert level 5 shows strong interaction

between the 5-17 and 40-49 age groups. All four plots show the highest number of

contacts occurring in the 5-17 age group.

The eastern health cluster shows predominantly cases in the 5-17 and 40-49 age

groups with interaction between these age groups. Other ages are present, in

particular a collection of contacts in the 30-59 age groups. There are some

similarities between the patterns present in the eastern health cluster and those for

Alert level 5. Given the relative scales of the two plots, an explanation for this

could be that the eastern cluster is driving the pattern for the entire Alert level.

The plot for the central health cluster shows an arrow-shaped pattern with a high

concentration in the [18,30) age group as well as interactions between this group

and the [30,60) age groups. Notably, there is much less interaction between the

[30,60) age groups than there is between them and [18,30). There is also a more

uniform pattern occurring between the [5,18) and [18,30) age groups. The pattern

present for the central cluster does not bear a strong resemblance to any of the

Alert level patterns. This is an example of a case where population-level trends in

age assortativity for an Alert level are not reflected in individual clusters.

Fig. 6 Contact matrices indicating the number of contacts that occur within and between age

groups. The panel on the left shows matrices computed for each of the Alert levels separately,

aggregated over all contacts occurring under that particular Alert level. On the right are the same

matrices computed from the eastern and central clusters.

The density plots of the high-contact/low-contact interaction patterns are shown in

Figure 7. Alert levels 2 and 3 both show patterns with a single point of

concentration and some dispersion away from the diagonal as the number of

contacts increases. In Alert level 2 the contact frequencies are centred about the

point (6,6), meaning that for many of the cases with around 6 contacts, each

412 contact themselves produced around 6 contacts. Alert level 3 shows a similar

413 pattern, but more dispersed and slightly more skewed towards high-contacts/high-

414 contact pairs. Alert levels 4 and 5 differ in that they have multiple points of

415 concentration and less consistent dispersion. Alert level 4 has one point of

416 concentration around (5,5), but with another localized concentration at a high-

417 contact/high-contact point on the diagonal. The number of contacts per-person in

418 Alert level 5 is quite dispersed. In all cases, most individuals have fewer than 20

419 contacts. The eastern cluster shows two concentrations: one in the low-contact

420 corner and another at a higher number of contacts. This plot shows a clear

421 interaction between high-contact and low-contact individuals (paler areas off the

422 diagonal). The central cluster has three areas of concentration. The highest

423 concentration is around 40 contacts per person, with a lower concentration at the

424 low-contact/low-contact corner of the plot, and low-contact/high-contact mode

425 located symmetrically about the diagonal. While the central cluster had few cases,

426 it tends to show a much higher number of contacts per person.

428 **Fig. 7** Joint distribution over the number of contacts in each case/contact pair; each region (i.e.

429 each colour) contains all pairs with similar estimated density values, as indicated in the quantile

430 legend. The left panels show the contours constructed from data aggregated for each Alert level.

431 The panels on the right show the contours for the eastern cluster and the central cluster.

Discussion

Between March 20, 2020 and December 15, 2021, the NL government

implemented a contact tracing program as part of a COVID-19 containment

strategy. The data provides information about secondary cases and contact

patterns in the population. Secondary case distributions can indicate how effective

policy measures are at suppressing local spread. Analyzing contact patterns under

varying levels of policy stringency can either validate existing assumptions of

contact homogeneity or provide a basis for more accurate models of disease

transmission. NL's public health measures until the establishment of the Omicron

variant were effective at containing SARS-CoV-2 spread (Figures 1 and 4), and

the contact tracing effort to achieve these results is summarized in Figure 1 and

Table 1.

The distributions of secondary cases (Figures 4 and 5) and the estimates of R_t

(Figure 5) offer insight into the efficacy of the Alert level system. The shapes of

the secondary case distributions in Figure 4 are skewed above 1 during periods of

increasing incidence and skewed below 1 as peaks taper off. The estimates of R_t

for each Alert level are mostly close to 1 with 95% lower confidence bounds of

1.02, 0.99, 0.84, and 1.20. This is consistent with local spread being contained

until late December 2021. The higher value for Alert level 5 could be a

consequence of low compliance with public health measures prior to the February

2021 outbreak in the eastern health region [1, p28], and delays between

identifying the outbreak and the implementation of Alert level policies. The

results here should be evaluated while keeping in mind that the heuristic nature of

the estimation procedure precludes rigorous statistical interpretation. Furthermore,

Figure 1 shows that the Alert levels are not distributed uniformly across the focal

period; a consequence of this is that factors such as variant or vaccination rates

could be affecting our estimates of R_t.

Statistical evidence supporting the use of a Poisson distribution to model

secondary cases over the negative binomial distribution typically coincides with

low heterogeneity in transmission, which has not been characteristic of COVID-

19 [21,22], though high uncertainty in the precision parameter has been observed

in settings with very stringent control measures [23]. Additionally, it has been

observed that imposing control measures can reduce overdispersion [24].

Therefore, one plausible explanation for this deviation from the existing COVID-

19 literature, with respect to the secondary case distribution, is that control

measures in place, even those with lower stringency, have reduced the occurrence

of superspreader events and resulted in a more homogenous pattern of infectivity

for our dataset.

The age contact matrices shown in Figure 6 show that the 5-17 age group had the

most contacts and many of those contacts occur between individuals in that age

group. Additionally, the matrix for the eastern cluster showed interaction between

this age group and the 40-49 age group. These observations could be supported by

transmission between high school students which either is preceded by or

proceeded from transmission between students and their parents/guardians

[10,17]. These patterns can also be seen in Figure 3, with many individuals in the

[5,18) age group at the centre of the cluster and smaller clumps of individuals in

the [40,50) age group. The pattern present in the central cluster suggests a form of

transmission facilitated by the [18,30) age group with minimal interaction

between the older affected age groups. This represents a very different form of

interaction than the former cluster. Overall, trends in interaction between age

groups are most pronounced under the most stringent set of public health

measures with trends becoming less apparent as stringency decreases. Comparing

the contact matrices of the eastern and central clusters to the Alert-level clusters

illustrates how contact patterns aggregated across multiple clusters may not be

representative of patterns within particular clusters. This suggests that during the

early stages of an outbreak, models which reflect contact patterns specific to the

outbreak setting may be a more reliable basis for local policy decisions than

aggregate models.

Age contact matrices have been estimated for all of Canada during the period May

2020 to December 2020 [16]. These showed a relatively high number of contacts

occurring within the same age group with the most contacts occurring within the

5-17 age group. These contact matrices also show older age groups have more

between-age interactions with a higher number of interactions among adults aged

18-60 and a lower number for those of retirement age. These patterns are similar

to, but less pronounced than, those found in the POLYMOD study [17]. These

Canada-wide patterns are most similar to those observed in NL during Alert level

2 (Figure 6). The block-like behaviour of the 18-60 age groups is also found in

NL during Alert level 4. However, both the patterns observed during Alert level 5

and those observed in the two selected clusters have no clear resemblance to either

the Canada-wide study or the POLYMOD study. This underlines the importance

of understanding local contact patterns when modelling the spread of an infectious

disease in small populations, or during periods of low incidence.

The interaction between low-contact and high-contact individuals was shown in

Figure 7. These plots provide some information about how the distribution of the

number of contacts per case varies across the population. A containment strategy

requires that contact tracing capacity not be exceeded, as such it is valuable to

investigate potential pressure on contact tracing demand arising from high-contact

individuals. Alert levels 2 and 3 both show a pattern of high concentration in one

area of the plot with greater dispersion as the number of contacts per case

increases. This suggests that given an individual has a high number of contacts,

there will be higher variance in the number of secondary contacts associated with

them, compared to an individual with a small number of contacts. This pattern

could be explained by each individual's number of contacts being drawn from a

common distribution with one mode and a skew to the right. On the other hand,

Alert levels 4 and 5 exhibit less consistent behaviour with multiple points of

concentration. This could be explained by the more stringent measures both

limiting gathering sizes (low-contact modes) while still permitting some essential,

high-contact jobs to be performed (higher-contact modes). That we observe a

concentration of low-contact cases in each Alert level is another indication that

the Alert level system was effective in limiting community spread. The results for

the two example clusters are less neatly concentrated patterns. In the eastern

cluster there appear to be two main concentrations of contacts per case, with an

additional, smaller concentration reflected off the diagonal. Comparing this plot to

the corresponding age contact matrix suggests a potential relationship between the

age of a case and the distribution of the number of contacts per case. The patterns

of the central cluster are quite similar to the eastern cluster, although at a slightly

greater magnitude. Contrasting this with the age-based contact patterns, which

were different between the two example clusters, suggests that whatever

demographic or environmental variables drive one form of contact patterns do not

affect the other in the same way. The difference in magnitude between the eastern

and central clusters also provides additional support for the efficacy of the Alert

system since many of the individuals in the central network had a high number of

contacts, even though the resulting cluster had fewer cases than the eastern

cluster.

Many infectious disease transmission models rely on strong assumptions about the

contact patterns, e.g. homogeneous mixing, or on contact patterns which are

estimated *a priori*. The observed contact patterns do not support a model of

homogeneously mixing populations and supports previously made observations

[15] that contact patterns change considerably in response to public health

measures. This degree of change, which goes well beyond simply having overall

contact volumes rise and fall with control measures, should raise notes of caution

for infectious disease transmission modelling that relies on network and contact

structure. Contact tracing provides a double benefit to infectious disease

modellers and public health officials. In addition to its role in effective

containment strategies, clear contact tracing data can provide modellers with

information about person-to-person contact patterns which enable more effective

modelling of low-incidence disease outbreaks, such as those in small populations

and during the early stages of an outbreak.

Conclusion

Estimated effective reproduction numbers are consistent with periods of contained

local spread. Further analysis needs to be done to determine the individual effects

of the Alert system, vaccination rates, and variant, as well as the interactions

between them. The data revealed heterogeneity in contact patterns within the

population. The patterns varied between contact networks as well as between

individual networks and patterns aggregated over multiple networks. These

suggest that population-specific age-based contact patterns are important to

modelling disease spread in small populations. Additionally, given the importance

of contact tracing to a containment strategy, understanding the behaviour of high-

contact individuals may be crucial to keeping cases within contact tracing

capacity.

Abbreviations

AIC: Akaike Information Criterion

CI: Confidence interval

FSA: Forward Sortation Area

MLE: Maximum likelihood estimate

NL: Newfoundland and Labrador

SMO: Special Measures Order

References

1. Government of Newfoundland and Labrador. Report to the House of Assembly on
the COVID-19 Public Health Emergency. 2022. Available from:
https://www.assembly.nl.ca/business/electronicdocuments/ReporttoHOACOVID-
19PublicHealthEmergency2022.pdf

2. Martignoni MM, Arino J, Hurford A. Is SARS-CoV-2 elimination or mitigation best?
Regional and disease characteristics determine the recommended strategy. Royal
Society Open Science. 2024

3. Hurford A, Martignoni MM, Loredo-Osti JC, Anokye F, Arino J, Husain BS, et al.
Pandemic modelling for regions implementing an elimination strategy. Journal of
Theoretical Biology. 2023; 561.

4. Baker M G, Wilson N, Blakely T. Elimination could be the optimal response strategy
for covid-19 and other emerging pandemic disease. BMJ. 2020.

5. Tupper P, Otto SP, Colijn C. Fundamental limitations of contact tracing for COVID-
19. Facets. 2021; 6:1993-2001.

6. Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, Munday JD,

Kucharski AJ, Edmunds WJ, Centre for the Mathematical Modelling of Infectious

Diseases COVID-19 Working Group, Funk S, Eggo RM. Feasibility of controlling

COVID-19 outbreaks by isolation of cases and contacts. Lancet Global Health. 2020;

8:488-496. https://doi.org/10.1016/S2214-109X(20)30074-7

7. Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in

emerging epidemics. PLoS ONE. 2006; 1(1). doi:10.1371/journal.pone.0000012

8. Martignoni MM, Renault J, Baafi J, Hurford A. Downsizing of COVID-19 contact

tracing in highly immune populations. PLOS ONE., M. M. 2022; 17(6).

https://doi.org/10.1371/journal.pone.0268586

9. Government of Newfoundland and Labrador. Public Advisory: Update on COVID-

19 in Newfoundland and Labrador [press release]. 2022 Jan 4.

10. Nunn A, Morrissey A, Crocker A, Patterson K, Stares J, Smith K, et al. Community-

based COVID-19 outbreak of the B.1.1.7 (Alpha) variant of concern in

Newfoundland, February to March 2021. Can Commun Dis Rep. 2023; 49(2/3):59-

66. https://doi.org/10.14745/ccdr.v49i23a05

11. Cintrón-Arias A, Castillo-Chávez C, Bettencourt LM, Lloyd AL, Banks HT. The

estimation of the effective reproduction number from disease outbreak data.

Mathematical Biosciences and Engineering. 2009; 6(2):261-282.

https://doi.org/10.3934/mbe.2009.6.261

12. Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory

syndrome reveal similar impacts of control measures. American Journal of

Epidemiology. 2004; 160(6):509-516.

13. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to

estimate time-varying reproduction numbers during epidemics. American Journal of

Epidemiology. 2013; 178(9):1505-1512. https://doi.org/10.1093/aje/kwt133 Lloyd-

Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of

variation on disease emergence. Nature. 2005; 438(17):355-359.

https://doi.org/10.1038/nature04153

14. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of

variation on disease emergence. Nature. 2005; 438(17):355-359.

https://doi.org/10.1038/nature04153

15. Zhang J, Litvinova M, Liang Y, Wang Y, Wang W, Zhao S, et al. Changes in contact
619   patterns shape the dynamics of the COVID-19 outbreak in China. Science. 2020;
620   368:1481-6.

621   16. Brankston G, Merkley E, Fisman DN, Tuite AR, Poljak Z, Loewen PJ, Greer AL.
622   Quantifying contact patterns in response to COVID-19 public health measures in
623   Canada. 2021; 21(2040). https://doi.org/10.1186/s12889-021-12080-1

624   17. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social
625   contacts and mixing patterns relevant to the spread of infectious diseases. PLoS Med.
626   2008; 5(3). doi:10.1371/journal.pmed.0050074

627   18. R Core Team. R: A language and environment for statistical computing. Vienna,
628   Austria: R Foundation for Statistical Computing; 2022.

629   19. Wickam H. ggplot2: Elegant graphics for data analysis. New York: Springer-Verlag;
630   2016.

631   20. 'Dealing with a totally new virus': inside the outbreak with eastern health's top
632   contact tracer. Canadian Broadcasting Corporation, Newfoundland and Labrador
633   [Internet]. 2021 April 6. Available from:
634   https://www.cbc.ca/news/canada/newfoundland-labrador/suzette-spurrell-contact-
635   tracing-q-a-1.5976417

636   21. Zhang Y, Li Y, Wang L, Li M, Zhou X. Evaluating transmission heterogeneity and
637   super-spreading event of COVID-19 in a metropolis of China. International Journal
638   of Environmental Research and Public Health. 2020; 17.

639   22. Wegehaupt O, Endo A, Vassall A. Superspreading, overdispersion and their
640   implications in the SARS-CoV-2 (COVID-19) pandemic: a systematic review and
641   meta-analysis of the literature. BMC Public Health. 2023; 23(1).
642   https://doi.org/10.1186/s12889-023-15915-1

643   23. Kwok KO, Chan HHH, Huang Y, Hui DSC, Tambyah PA, Wei WI, Chau PYK,
644   Wong SYS, Tang JWT. Inferring super-spreading from transmission clusters of
645   COVID-19 in Hong Kong, Japan, and Singapore. Journal of Hospital Infection.
646   2020; 105:682-5.

647   24. He D, Zhao S, Xu X, Lin Q, Zhuang Z, Cao P, Wang MH, Lou Y, Xiao L, Wu Y,

648   Yang L. Low dispersion in the infectiousness of COVID-19 cases implies difficulty

649   in control. BMC Public Health. 2020; 20(1558)

Declarations

Ethics approval and consent to participate

Ethics approval for this project was granted by the Newfoundland and Labrador

Health Research Ethics Board (reference # 2021.013).

Consent for publication

The Newfoundland and Labrador Health Research Ethics Board approved the

research involving secondary use data without requiring further consent to

participate from the participants (i.e., Article 5.5A of the Tri-Council Policy

Statement 2).

Availability of data and materials

Data requires ethics approval and can be requested from Newfoundland and

Labrador Health Services, Digital Health, the data custodian.

Competing interests

AH was a member of the Newfoundland and Labrador Predictive Analytics

modelling team and previously received funding from the Newfoundland and

Labrador Department of Health and Community Services.

Funding

This research was funded by the Canadian Network for Modelling Infectious

Diseases (560516-2020) and the Discovery grant (RGPIN-2019-06131) from the

Natural Sciences and Engineering Research Council of Canada.

Authors' contributions

RD performed the analysis, generated visuals, and wrote the first draft of the

manuscript. SS and AM lead the contact tracing process. RD, AH, LW, and CC

conceived of the project and designed the analytical approach. All authors edited

the manuscript.

Acknowledgements

The authors would like to acknowledge the substantial efforts of Newfoundland

and Labrador Public Health, the Department of Health and Community Services,

Regional Health Authorities and all individuals involved in contact tracing and the

recording of these data, as well as the Newfoundland and Labrador Health

Services, Digital Health as the custodians of these data.